


Anthony J. Viola  
Andre K. Cizmarik  
Jennifer L. Dereka  
Zachary W. Silverman  
EDWARDS WILDMAN PALMER LLP  
Attorneys for Plaintiffs  
Kowa Company, Ltd.,  
Kowa Pharmaceuticals America, Inc., and  
Nissan Chemical Industries, Ltd.  
750 Lexington Ave.  
New York, NY 10022  
(212) 308-4411

14 CV 2758



UNITED STATES DISTRICT COURT  
SOUTHERN DISTRICT OF NEW YORK

Kowa Company, Ltd.,  
Kowa Pharmaceuticals America, Inc., and  
Nissan Chemical Industries, Ltd.,

Plaintiffs,

v.

Amneal Pharmaceuticals, LLC,

Defendant.

Civil Action No. \_\_\_\_\_

**COMPLAINT**

Plaintiffs, Kowa Company, Ltd. ("KCL"), Kowa Pharmaceuticals America, Inc. ("KPA") (collectively, "Kowa"), and Nissan Chemical Industries, Ltd. ("NCI") by their undersigned counsel, for their Complaint against defendant Amneal Pharmaceuticals, LLC ("Amneal"), allege as follows:

**Jurisdiction and Venue**

1. This is an action for patent infringement arising under the patent laws of the United States, Title 35, United States Code and arising under 35 U.S.C. §§ 271(e)(2), 271(b), 271(c), and 281-283. Subject matter jurisdiction is proper under 28 U.S.C. §§ 1331 and 1338(a). Venue is proper under 28 U.S.C. §§ 1391(b)-(c) and 1400(b). Personal jurisdiction over the defendants in New York is proper under N.Y. C.P.L.R. §§ 301 and 302(a) and because defendants are doing business in this jurisdiction.

**Parties**

2. KCL is a Japanese corporation having its corporate headquarters and principal place of business in Aichi, Japan. KPA is a wholly owned U.S. subsidiary of KCL. KPA has its corporate headquarters and principal place of business in Montgomery, Alabama and is organized under the laws of Delaware.

3. NCI is a Japanese corporation having its corporate headquarters and principal place of business in Tokyo, Japan.

4. KCL and NCI are engaged in the business of research, developing, manufacturing, and marketing of a broad spectrum of innovative pharmaceutical products, including Livalo<sup>®</sup>.

5. Upon information and belief, Amneal is incorporated in Delaware having a place of business in Bridgewater, New Jersey. Upon information and belief, Amneal filed Abbreviated New Drug Application (“ANDA”) No. 20-5961.

6. Upon information and belief, Amneal is currently transacting business in the Southern District of New York, at least by making and shipping into this Judicial District, or by using, offering to sell or selling or by causing others to use, offer to sell or sell, pharmaceutical products into this Judicial District.

7. Upon information and belief, Amneal derives substantial revenue from interstate and/or international commerce, including substantial revenue from goods used or consumed or services rendered in the State of New York and the Southern District of New York. Amneal is registered with the N.Y. State Department of State, Division of Corporations, to do business as a foreign corporation in New York. By filing its ANDA, Amneal has committed, and unless enjoined, will continue to commit a tortious act without the state of New York, that Amneal expects or should reasonably expect to have consequences in the State of New York including in this Judicial District.

#### **The New Drug Application**

8. KPA sells drug products containing pitavastatin calcium (the “pitavastatin drug product”) under the trade name Livalo<sup>®</sup> in the United States pursuant to the United States Food and Drug Administration’s approval of a New Drug Application (“NDA”) held by KCL (NDA No. 22-363).

9. Livalo<sup>®</sup> is approved for use as an adjunctive therapy to diet to reduce elevated total cholesterol, low-density lipoprotein cholesterol, apolipoprotein B, triglycerides, and to increase HDL-C in adult patients with primary hyperlipidemia or mixed dyslipidemia.

10. The approval letter for Livalo<sup>®</sup>, with approved labeling, was issued by the FDA on August 3, 2009.

11. Certain amendments to the approved labeling for Livalo<sup>®</sup> have subsequently been approved.

#### **The Patents in Suit**

12. United States Patent No. 5,856,336 (“the ‘336 patent”), entitled “Quinoline Type Mevalonolactones,” a true and correct copy of which is appended hereto as **Exhibit A**, was duly

issued on January 5, 1999 to inventors Yoshihiro Fujikawa, Mikio Suzuki, Hiroshi Iwasaki, Mitsuaki Sakashita, and Masaki Kitahara, and assigned to plaintiff NCI. The '336 patent claims, inter alia, the pitavastatin drug product, and a method for reducing hyperlipidemia, hyperlipoproteinemia or atherosclerosis, which comprises administering an effective amount of the pitavastatin drug product.

13. Plaintiff NCI has been and still is the owner through assignment of the '336 patent, which expires on December 25, 2020 pursuant to a patent-term extension. KCL is NCI's licensee for the '336 patent and KPA holds a license from KCL for the '336 patent.

14. United States Patent No. 8,557,993 ("the '993 patent"), entitled "Crystalline Forms of Pitavastatin Calcium," a true and correct copy of which is appended hereto as **Exhibit B**, was duly issued on October 15, 2013 to inventors Paul Adriaan Van Der Schaaf, Fritz Blatter, Martin Szelagiewicz, and Kai-Uwe Schoening, and ultimately was assigned to plaintiff NCI. The '993 patent claims, inter alia, crystalline polymorphs or the amorphous form of pitavastatin or processes for preparing the same.

15. Plaintiff NCI has been and still is the owner through assignment of the '993 patent, which expires on February 2, 2024. KCL is NCI's licensee for the '993 patent and KPA holds a license from KCL for the '993 patent.

16. In accordance with its license, KPA sells the pitavastatin drug product under the trade name Livalo<sup>®</sup> in the United States. Sales of Livalo<sup>®</sup> are made pursuant to approval by the FDA of NDA No. 22-363.

17. Plaintiff KCL manufactures the Livalo<sup>®</sup> drug products as sold by KPA.

18. Plaintiffs Kowa and NCI will be substantially and irreparably harmed by infringement of either of the '336 or '993 patents (the "Livalo<sup>®</sup> patents"). There is no adequate remedy at law.

**COUNT I**

**INFRINGEMENT OF THE '336 PATENT UNDER 35 U.S.C. § 271(e)(2)(A)**

19. Plaintiffs repeat and incorporate herein by reference the allegations contained in each of the foregoing paragraphs.

20. Upon information and belief, defendant Amneal filed an Abbreviated New Drug Application ("ANDA") with the Food and Drug Administration ("FDA") under 21 U.S.C. § 355(j) (ANDA No. 20-5961) seeking approval to market 1 mg, 2 mg, and 4 mg tablets comprising pitavastatin calcium.

21. By this ANDA filing, Amneal has indicated that it intends to engage, and that there is substantial likelihood that it will engage, in the commercial manufacture, importation, use, offer for sale, and/or sale, or inducement thereof, of Plaintiffs' patented pitavastatin drug product immediately or imminently upon receiving FDA approval to do so. Also by its ANDA filing, Amneal has indicated that its drug product is bioequivalent to Plaintiffs' pitavastatin drug product.

22. By its ANDA filing, Amneal seeks to obtain approval to commercially manufacture, use, import, offer for sale, and/or sell, alleged generic equivalents of Plaintiffs' Livalo<sup>®</sup> pitavastatin drug product prior to the expiration date of the '336 patent.

23. By a letter dated March 5, 2014 (the "Notice Letter"), Amneal informed KCL and NCI that Amneal had filed a certification to the FDA, pursuant to 21 U.S.C. § 355(j)(2)(B)(iv)(I).

Upon information and belief, on or about March 7, 2014, KCL received the Notice letter. On or about March 7, 2014, NCI received the Notice Letter.

24. The Notice Letter, purporting to be Amneal's Notification Pursuant to 21 U.S.C. § 355(j)(2)(B)(ii)(I), asserts that in Amneal's opinion, the '336 patent purportedly is "unenforceable, invalid, and/or not infringed, either literally or under the doctrine of equivalents, by the manufacture, use, sale, offer for sale, and/or importation of the drug product for which ANDA No. 205961 has been submitted by Amneal."

25. Amneal's filing of ANDA No. 20-5961 for the purpose of obtaining FDA approval to engage in the commercial manufacture, use, importation, offer for sale and/or sale, or inducement thereof, of its proposed pitavastatin drug product before the expiration of the '336 patent is an act of infringement under 35 U.S.C. § 271(e)(2)(A).

26. Amneal's manufacture, use, importation, offer for sale, and/or sale, or inducement thereof, of its proposed pitavastatin drug product will directly infringe or induce infringement of at least one claim of the '336 patent under 35 U.S.C. § 271(e)(2)(A).

27. Upon information and belief, Amneal's proposed label for its pitavastatin drug product will include the treatment of at least one of hyperlipidemia, hyperlipoproteinemia, and atherosclerosis.

28. Unless Amneal is enjoined from infringing and inducing the infringement of the '336 patent, Plaintiffs will suffer substantial and irreparable injury. Plaintiffs have no adequate remedy at law.

**COUNT II**

**INFRINGEMENT OF THE METHOD CLAIM OF THE '336 PATENT  
UNDER 35 U.S.C. § 271(b)**

29. Plaintiffs repeat and incorporate herein by reference the allegations contained in each of the foregoing paragraphs.

30. Upon information and belief, approval of ANDA 20-5961 is substantially likely to result in the commercial manufacture, use, importation, offer for sale, and/or sale, or inducement thereof, of a pitavastatin drug product which is marketed and sold for use in a method claimed in one or more claims of the '336 patent, immediately or imminently upon approval of the ANDA, and prior to the expiration of the '336 patent.

31. Upon information and belief, Amneal's proposed label for its pitavastatin drug product will include the treatment of at least one of hyperlipidemia, hyperlipoproteinemia or atherosclerosis.

32. Upon information and belief, Amneal is aware or reasonably should be aware, of the widespread use of pitavastatin as an adjunctive therapy to diet to reduce elevated total cholesterol, low-density lipoprotein cholesterol, apolipoprotein B, triglycerides, and to increase HDL-C in adult patients with primary hyperlipidemia or mixed dyslipidemia. The beneficial effects of pitavastatin as an adjunctive therapy to diet to reduce elevated total cholesterol, low-density lipoprotein cholesterol, apolipoprotein B, triglycerides, and to increase HDL-C in adult patients with primary hyperlipidemia or mixed dyslipidemia would be readily apparent to customers of Amneal (e.g., including, without limitation, physicians, pharmacists, pharmacy benefits management companies, health care providers who establish drug formularies for their insurers and/or patients). Amneal will be marketing its pitavastatin drug product with specific

intent to actively induce, aid and abet infringement of the '336 patent. Amneal knows or reasonably should know that its proposed conduct will induce infringement of the '336 patent.

33. Unless Amneal is enjoined from infringing and inducing the infringement of the '336 patent, Plaintiffs will suffer substantial and irreparable injury. Plaintiffs have no adequate remedy at law.

### **COUNT III**

#### **INFRINGEMENT OF THE METHOD CLAIM OF THE '336 PATENT UNDER 35 U.S.C. § 271(c)**

34. Plaintiffs repeat and incorporate herein by reference the allegations contained in each of the foregoing paragraphs.

35. Upon information and belief, Amneal's proposed pitavastatin drug product comprises pitavastatin calcium as referenced in the claims of the '336 patent.

36. Upon information and belief, Amneal's proposed pitavastatin drug product will be especially made for use in a manner that is an infringement of the '336 patent.

37. Upon information and belief, Amneal knows that Amneal's proposed pitavastatin drug product will be especially made for use in a manner that is an infringement of the '336 patent.

38. Upon information and belief, sale of Amneal's proposed pitavastatin drug product will result in direct infringement of the '336 patent.

39. Upon information and belief, Amneal's proposed pitavastatin drug product is not a staple article or commodity of commerce which is suitable for a substantial noninfringing use.

40. Upon information and belief, Amneal knows that Amneal's proposed pitavastatin drug product is not a staple article or commodity of commerce which is suitable for substantial noninfringing use.



41. Upon information and belief, approval of ANDA 20-5961 is substantially likely to result in the commercial use, manufacture, offer for sale and/or sale (or the inducement thereof or contribution thereto) of a drug product which is especially made, adapted, marketed, sold, and approved exclusively for use in a method claimed in the '336 patent, immediately or imminently upon approval of the ANDA.

42. Plaintiffs will be substantially and irreparably harmed if defendants are not enjoined from contributing to the infringement of the '336 patent. Plaintiffs have no adequate remedy at law.

#### COUNT IV

##### INFRINGEMENT OF THE '993 PATENT UNDER 35 U.S.C. § 271(e)(2)(A)

43. Plaintiffs repeat and incorporate herein by reference the allegations contained in each of the foregoing paragraphs.

44. Amneal's Notice Letter, purporting to be Amneal's Notice of Certification under 21 U.S.C. § 355(j)(2)(B)(iv), indicates that Amneal intends to manufacture, use, sell, or offer for sale, its proposed pitavastatin drug product prior to the expiration of the '993 patent.

45. The Notice Letter asserts that in Amneal's opinion, the '993 patent purportedly is "unenforceable, invalid, and/or not infringed, either literally or under the doctrine of equivalents, by the manufacture, use, sale, offer for sale, and/or importation of the drug product for which ANDA No. 205961 has been submitted by Amneal."

46. Amneal's filing of ANDA No. 20-5961 for the purpose of obtaining FDA approval to engage in the commercial manufacture, use, importation, offer for sale and/or sale, or the inducement thereof, of its proposed pitavastatin drug product before the expiration of the '993 patent is an act of infringement under 35 U.S.C. § 271(e)(2)(A).

47. Amneal's manufacture, use, importation, offer for sale, sale, and/or importation of its proposed pitavastatin drug product will directly infringe or induce infringement of at least one claim of the '993 patent under 35 U.S.C. § 271(e)(2)(A).

48. Unless Amneal is enjoined from infringing the '993 patent, plaintiffs will suffer substantial and irreparable injury. Plaintiffs have no adequate remedy at law.

WHEREFORE, Plaintiffs request the following relief:

- (a) a declaratory judgment pursuant to 28 U.S.C. § 2201 et seq. that making, using, selling, offering to sell and/or importing Amneal's pitavastatin drug product for which it seeks FDA approval or any drug product containing pitavastatin will infringe at least one claim of one or more of the Livalo<sup>®</sup> patents;
- (b) a declaratory judgment pursuant to 28 U.S.C. § 2201 et seq. that the making, using, offering for sale, selling and/or importing of Amneal's pitavastatin drug product or any drug product containing pitavastatin, will induce the infringement at least one claim of one or more of the Livalo<sup>®</sup> patents;
- (c) a declaratory judgment pursuant to 28 U.S.C. § 2201 et seq. that the making, using, offering for sale, selling and/or importing of Amneal's pitavastatin drug product or any drug product containing pitavastatin, will contribute to the infringement of at least one claim of one or more of the Livalo<sup>®</sup> patents;
- (d) a declaratory judgment pursuant to 28 U.S.C. § 2201 et seq. and an order pursuant to 35 U.S.C. § 271(e)(4)(A) providing that the effective date of any FDA approval for Amneal to commercially make, use, sell, offer to sell or import its pitavastatin drug product or any drug product containing pitavastatin

- be no earlier than the date following the expiration date of the last to expire of the Livalo<sup>®</sup> patents (as extended, if applicable);
- (e) a permanent injunction restraining and enjoining against any infringement by defendants, their officers, agents, attorneys, employees, successors or assigns, or those acting in privity or concert with them, of the Livalo<sup>®</sup> patents, through the commercial manufacture, use, sale, offer for sale or importation into the United States of Amneal's pitavastatin drug product or any drug product containing pitavastatin, and/or any inducement of or contribution to the same;
- (f) Attorneys' fees in this action under 35 U.S.C. § 285; and
- (g) Such further and other relief in favor of Plaintiffs and against defendants as this Court may deem just and proper.

Dated: New York, New York  
April 17, 2014

Kowa Company, Ltd.,  
Kowa Pharmaceuticals America, Inc., and  
Nissan Chemical Industries, Ltd.

By their attorneys,



---

Anthony J. Viola  
Andre K. Cizmarik  
Jennifer L. Dereka  
Zachary W. Silverman  
EDWARDS WILDMAN PALMER LLP  
750 Lexington Avenue  
New York, NY 10022  
(212) 308-4411

David G. Conlin (to be admitted *pro hac vice*)  
Kathleen B. Carr (to be admitted *pro hac vice*)  
Adam P. Samansky  
EDWARDS WILDMAN PALMER LLP  
111 Huntington Avenue  
Boston, MA 02199  
(617) 239-0100

# EXHIBIT A



US005856336A

**United States Patent** [19]  
**Fujikawa et al.**

[11] **Patent Number:** **5,856,336**  
 [45] **Date of Patent:** **Jan. 5, 1999**

[54] **QUINOLINE TYPE MEVALONOLACTONES**

[75] **Inventors:** **Yoshihiro Fujikawa; Mikio Suzuki;**  
**Hiroshi Iwasaki**, all of Funabashi;  
**Mitsuaki Sakashita; Masaki Kitahara**,  
 both of Shiraoka-machi, all of Japan

[73] **Assignee:** **Nissan Chemical Industries Ltd.**,  
 Tokyo, Japan

[21] **Appl. No.:** **883,398**

[22] **Filed:** **May 15, 1992**

**Related U.S. Application Data**

[62] Division of Ser. No. 631,092, Dec. 19, 1990, which is a continuation of Ser. No. 233,752, Aug. 19, 1988.

[30] **Foreign Application Priority Data**

Aug. 20, 1987	[JP]	Japan	.....	62-207224
Jan. 26, 1988	[JP]	Japan	.....	63-15585
Aug. 3, 1988	[JP]	Japan	.....	63-193606

[51] **Int. Cl.<sup>6</sup>** ..... **A61K 31/47; C07D 215/12**

[52] **U.S. Cl.** ..... **514/311; 546/173**

[58] **Field of Search** ..... **546/173; 514/311**

[56] **References Cited****U.S. PATENT DOCUMENTS**


5,753,675 5/1998 Wattanasin ..... 514/311

*Primary Examiner*—Laura L. Stockton

*Attorney, Agent, or Firm*—Oblon, Spivak, McClelland, Maier & Neustadt, P.C.

[57] **ABSTRACT**

A compound of the formula



[A]

Z=—CH(OH)—CH<sub>2</sub>—CH(OH)—CH<sub>2</sub>—COO.½Ca  
 have HMG—CoA inhibiting effects, making them useful as inhibitors of cholesterol biosynthesis. The compound may be prepared as a pharmaceutical for reducing hyperlipidemia, hyperlipoproteinemia or atherosclerosis.

**2 Claims, No Drawings**

5,856,336

1

## QUINOLINE TYPE MEVALONOLACTONES

This is a division, of application Ser. No. 07/631,092, filed on Dec. 19, 1990, which is a continuation of 07/233,752, filed Aug. 19, 1988.

The present invention relates to novel mevalonolactones having a quinoline ring, processes for their production, pharmaceutical compositions containing them and their pharmaceutical uses particularly as anti-hyperlipidemic, hypolipoproteinemic and anti-atherosclerotic agents, and intermediates useful for their production and processes for the production of such intermediates.

Some fermentation metabolic products such as compactine, CS-514, Mevinolin or semi-synthetic derivatives or fully synthetic derivatives thereof are known to be inhibitors against HMG-CoA reductase which is a rate limiting enzyme for cholesterol biosynthesis. (A. Endo J. Med Chem., 28(4) 401 (1985))


CS-514 and Mevinolin have been clinically proved to be potentially useful anti-hyperlipoproteinemic agents, and they are considered to be effective for curing or preventing diseases of coronary artery sclerosis or atherosclerosis. (IXth Int. Symp. Drugs Affect. Lipid Metab., 1986, p30, p31, p66)

However, with respect to fully synthetic derivatives, particularly hetero aromatic derivatives of inhibitors against HMG-CoA reductase, limited information is disclosed in the following literatures:

WPI ACC NO. 84-158675, 86-028274, 86-098816, 86-332070, 87-124519, 87-220987, 88-07781, 88-008460, 88-091798 and 88-112505.


The present inventors have found that mevalonolactone derivatives having a quinoline ring, the corresponding dihydroxy carboxylic acids and salts and esters thereof have high inhibitory activities against cholesterol biosynthesis wherein HMG-CoA reductase acts as a rate limiting enzyme. The present invention has been accomplished on the basis of this discovery.

The novel mevalonolactone derivatives of the present invention are represented by the following formula I:



wherein  $R_1, R_2, R_3, R_4$  and  $R^9$  are independently hydrogen,  $C_{1-6}$  alkyl,  $C_{2-6}$  cycloalkyl,  $C_{1-3}$  alkoxy, n-butoxy, i-butoxy, sec-butoxy,  $R^7R^8N-$  (wherein  $R^7$  and  $R^8$  are independently hydrogen or  $C_{1-3}$  alkyl), trifluoromethyl, trifluoromethoxy, difluoromethoxy, fluoro, chloro, bromo, phenyl, phenoxy, benzyloxy, hydroxy, trimethylsilyloxy, diphenyl-t-butylsilyloxy, hydroxymethyl or  $-O(CH_2)_lOR^{19}$  (wherein  $R^{19}$  is hydrogen or  $C_{1-3}$  alkyl, and  $l$  is 1, 2 or 3); or when located at the ortho position to each other,  $R^1$  and  $R^2$ , or  $R^3$  and  $R^4$  together form  $-CH=CH-CH=CH-$ ; or when located at the ortho position to each other,  $R^1$  and  $R^2$  together form  $-OC(R^{15})(R^{16})O-$  (wherein  $R^{15}$  and  $R^{16}$  are independently hydrogen or  $C_{1-3}$  alkyl);  $Y$  is  $-CH_2-$ ,  $-CH_2CH_2-$ ,  $-CH=CH-$ ,  $-CH_2-CH=CH-$  or  $-CH=CH-CH_2-$ ; and  $Z$  is  $-Q-CH_2WCH_2-CO_2R^{12}$ ,

2



(wherein  $Q$  is  $-C(O)-$ ,  $-C(OR^{13})_2-$  or  $-CH(OH)-$ ;  $W$  is  $-C(O)-$ ,  $-C(OR^{13})_2-$  or  $-C(R^{14})(OH)-$ ;  $R^{11}$  is hydrogen or  $C_{1-3}$  alkyl;  $R^{12}$  is hydrogen or  $R^{14}$  (wherein  $R^{14}$  is physiologically hydrolyzable alkyl or  $M$  (wherein  $M$  is  $NH_4$ , sodium, potassium,  $\frac{1}{2}$  calcium or a hydrate of lower alkylamine, di-lower alkylamine or tri-lower alkylamine)); two  $R^{13}$  are independently primary or secondary  $C_{1-6}$  alkyl; or two  $R^{13}$  together form  $-(CH_2)_2-$  or  $-(CH_2)_3-$ ;  $R^{17}$  and  $R^{18}$  are independently hydrogen or  $C_{1-3}$  alkyl; and  $R^5$  is hydrogen,  $C_{1-6}$  alkyl,  $C_{2-3}$  alkenyl,  $C_{3-6}$  cycloalkyl,



(wherein  $R^9$  is hydrogen,  $C_{1-4}$  alkyl,  $C_{1-3}$  alkoxy, fluoro, chloro, bromo or trifluoromethyl), phenyl- $(CH_2)_m-$  (wherein  $m$  is 1, 2 or 3),  $-(CH_2)_nCH(CH_3)-$ phenyl or phenyl- $(CH_2)_nCH(CH_3)-$  (wherein  $n$  is 0, 1 or 2).

Various substituents in the formula I will be described in detail with reference to specific examples. However, it should be understood that the present invention is by no means restricted by such specific examples.

$C_{1-6}$  alkyl for  $R^1, R^2, R^3, R^4, R^6$  and  $R^9$  includes, for example, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, sec-butyl and t-butyl.  $C_{1-3}$  alkoxy for  $R^1, R^2, R^3, R^4$  and  $R^6$  includes, for example, methoxy, ethoxy, n-propoxy and i-propoxy.

$C_{1-3}$  alkyl for  $R^{11}$  includes, for example, methyl, ethyl, n-propyl and i-propyl.

$C_{1-3}$  alkyl for  $R^{13}$  includes, for example, methyl, ethyl, n-propyl and i-propyl.

Alkyl for  $R^{14}$  includes, for example, methyl, ethyl, n-propyl, i-propyl, n-butyl and i-butyl.

$M$  is a metal capable of forming a pharmaceutically acceptable salt, and it includes, for example, sodium and potassium.

$CO_2M$  includes, for example,  $-CO_2NH_4$  and  $-CO_2H$ . (primary to tertiary lower alkylamine such as trimethylamine).

$C_{1-6}$  alkyl for  $R^5$  includes, for example, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, sec-butyl, t-butyl, n-pentyl and n-hexyl.

$C_{3-6}$  cycloalkyl for  $R^5$  includes, for example, cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

$C_{2-3}$  alkenyl for  $R^5$  includes, for example, vinyl and i-propenyl.

Phenyl- $(CH_2)_m-$  for  $R^5$  includes, for example, benzyl,  $\beta$ -phenylethyl and  $\gamma$ -phenylpropyl.

Phenyl- $(CH_2)_nCH(CH_3)-$  for  $R^5$  includes, for example,  $\alpha$ -phenylethyl and  $\alpha$ -benzylethyl.

$C_{1-3}$  alkyl for  $R^7$  and  $R^8$  includes, for example, methyl, ethyl, n-propyl and i-propyl.

Further, these compounds may have at least one or two asymmetric carbon atoms and may have at least two to four

5,856,336

3

optical isomers. The compounds of the formula I include all of these optical isomers and all of the mixtures thereof.

Among compounds having carboxylic acid moieties falling outside the definition of  $-\text{CO}_2\text{R}^{12}$  of the carboxylic acid moiety of substituent Z of the compounds of the present invention, those which undergo physiological hydrolysis, after intake, to produce the corresponding carboxylic acids (compounds wherein the  $-\text{CO}_2\text{R}^{12}$  moiety is  $-\text{CO}_2\text{H}$ ) are equivalent to the compounds of the present invention.

Now, preferred substituents of the compounds of the present invention will be described.

In the following preferred, more preferred still further preferred and most preferred examples, the numerals for the positions of the substituents indicate the positions on the quinoline ring. For example, N' shown by e.g. 1' or 2' indicates the position of the substituent on the phenyl substituted at the 4-position of the quinoline ring (the carbon connected to the quinoline ring is designated as 1'). The meanings of the respective substituents are the same as the above-mentioned meanings.

Preferred substituents for  $\text{R}^1$ ,  $\text{R}^2$  and  $\text{R}^6$  are hydrogen, fluoro, chloro, bromo,  $\text{C}_{1-3}$  alkyl,  $\text{C}_{1-3}$  alkoxy,  $\text{C}_{3-6}$  cycloalkyl, dimethylamino, hydroxy, hydroxymethyl, hydroxyethyl, trifluoromethyl, trifluoromethoxy, difluoromethoxy, phenoxy and benzyloxy.

Further, when  $\text{R}^6$  is hydrogen, it is preferred that  $\text{R}^1$  and  $\text{R}^2$  together form methylenedioxy.


As preferred examples for  $\text{R}^3$  and  $\text{R}^4$ , when  $\text{R}^4$  is hydrogen,  $\text{R}^3$  is hydrogen, 3'-fluoro, 3'-chloro, 3'-methyl, 4'-methyl, 4'-chloro and 4'-fluoro.

Other preferred combinations of  $\text{R}^3$  and  $\text{R}^4$  include 3'-methyl-4'-chloro, 3',5'-dichloro, 3',5'-difluoro, 3',5'-dimethyl and 3'-methyl-4'-fluoro.

Preferred examples for  $\text{R}^5$  include primary and secondary  $\text{C}_{1-6}$  alkyl and  $\text{C}_{3-6}$  cycloalkyl.

Preferred examples for Y include  $-\text{CH}_2-\text{CH}_2-$  and  $-\text{CH}=\text{CH}-$ .

Preferred examples for Z include



$-\text{CH}(\text{OH})\text{CH}_2\text{CH}_2(\text{OH})\text{CH}_2\text{CO}_2\text{R}^{12}$ ,  $-\text{CH}(\text{OH})\text{CH}_2\text{C}(\text{O})\text{CH}_2\text{CO}_2\text{R}^{12}$  and  $-\text{CH}(\text{OH})\text{CH}_2\text{C}(\text{OR}^{13})_2\text{CH}_2\text{CO}_2\text{R}^{12}$ .

Now, more preferred substituents of the compounds of the present invention will be described.

As more preferred examples for  $\text{R}^1$ ,  $\text{R}^2$  and  $\text{R}^6$ , when both  $\text{R}^2$  and  $\text{R}^6$  are hydrogen,  $\text{R}^1$  is hydrogen, 5-fluoro, 6-fluoro, 7-fluoro, 8-fluoro, 5-chloro, 6-chloro, 7-chloro, 8-chloro, 5-bromo, 6-bromo, 7-bromo, 8-bromo, 5-methyl, 6-methyl, 7-methyl, 8-methyl, 5-methoxy, 6-methoxy, 7-methoxy, 8-methoxy, 5-trifluoromethyl, 6-trifluoromethyl, 7-trifluoromethyl, 8-trifluoromethyl, 6-trifluoromethoxy, 6-difluoromethoxy, 8-hydroxyethyl, 5-hydroxy, 6-hydroxy, 7-hydroxy, 8-hydroxy, 6-ethyl, 6-n-butyl and 7-dimethylamino.

When  $\text{R}^6$  is hydrogen,  $\text{R}^1$  and  $\text{R}^2$  together represent 6-chloro-8-methyl, 6-bromo-7-methoxy, 6-methyl-7-chloro, 6-chloro-8-hydroxy, 5-methyl-2-hydroxy, 6-methoxy-7-chloro, 6-chloro-7-methoxy, 6-hydroxy-7-chloro, 6-chloro-7-hydroxy, 6-chloro-8-bromo, 5-chloro-6-hydroxy, 6-bromo-8-chloro, 6-bromo-8-hydroxy, 5-methyl-8-chloro, 7-hydroxy-8-chloro, 6-bromo-8-hydroxy, 6-methoxy-7-methyl, 6-chloro-8-bromo, 6-methyl-8-bromo, 6,7-difluoro, 6,8-difluoro, 6,7-methylenedioxy, 6,8-dichloro, 5,8-

4

dimethyl, 6,8-dimethyl, 6,7-dimethoxy, 6,7-diethoxy, 6,7-dibromo or 6,8-dibromo.

When  $\text{R}^1$ ,  $\text{R}^2$  and  $\text{R}^6$  are not hydrogen, they together represent 5,7-dimethoxy-8-hydroxy, 5,8-dichloro-6-hydroxy, 6,7,8-trimethoxy, 6,7,8-trimethyl, 6,7,8-trichloro, 5-fluoro-6,8-dibromo or 5-chloro-6,8-dibromo.

As more preferred examples for  $\text{R}^3$  and  $\text{R}^4$ , when  $\text{R}^3$  is hydrogen,  $\text{R}^4$  is hydrogen, 4'-methyl, 4'-chloro or 4'-fluoro. When both  $\text{R}^3$  and  $\text{R}^4$  are not hydrogen, they together represent 3',5'-dimethyl or 3'-methyl-4'-fluoro.

As more preferred examples for  $\text{R}^5$ , the above-mentioned preferred examples of  $\text{R}^5$  may be mentioned.

As preferred examples for Y,  $-\text{CH}_2-\text{CH}_2-$  and (E)- $-\text{CH}=\text{CH}-$  may be mentioned. As more preferred examples for Z, the above preferred examples for Z may be mentioned.

Now, still further preferred substituents of the compounds of the present invention will be described. As examples for  $\text{R}^1$ ,  $\text{R}^2$  and  $\text{R}^6$ , when both  $\text{R}^2$  and  $\text{R}^6$  are hydrogen,  $\text{R}^1$  is hydrogen, 6-methyl, 6-ethyl, 6-trifluoromethyl, 6-hydroxy, 6-methoxy, 6-chloro, 6-bromo, 6-n-butyl and 7-dimethylamino.

When only  $\text{R}^6$  is hydrogen,  $\text{R}^1$  and  $\text{R}^2$  represent 6,8-dichloro, 5,8-dimethyl, 6,8-dimethyl, 6,7-dimethoxy, 6,7-diethoxy, 6,7-dibromo, 6,8-dibromo, 6,7-difluoro and 6,8-difluoro.

As still further preferred examples for  $\text{R}^3$  and  $\text{R}^4$ , when  $\text{R}^3$  is hydrogen,  $\text{R}^4$  is hydrogen, 4'-chloro or 4'-fluoro, or  $\text{R}^3$  and  $\text{R}^4$  together represent 3'-methyl-4'-fluoro.

Still further preferred examples for  $\text{R}^5$  include ethyl, n-propyl, i-propyl and cyclopropyl.

Still further preferred examples for Y include (E)- $-\text{CH}=\text{CH}-$ .

As still further preferred examples for Z, the above-mentioned preferred example for Z may be mentioned.

Now, the most preferred substituents for the compounds of the present invention will be described.

As the most preferred examples for  $\text{R}^1$ ,  $\text{R}^2$  and  $\text{R}^6$ , when both  $\text{R}^2$  and  $\text{R}^6$  are hydrogen,  $\text{R}^1$  is hydrogen, 6-methyl or 6-chloro.

When only  $\text{R}^6$  is hydrogen,  $\text{R}^1$  and  $\text{R}^2$  together represent, for example, 6,7-dimethoxy.

As the most preferred examples for  $\text{R}^3$  and  $\text{R}^4$ ,  $\text{R}^3$  is hydrogen and  $\text{R}^4$  is hydrogen, 4'-chloro or 4'-fluoro.

The most preferred examples for  $\text{R}^5$  include i-propyl and cyclopropyl. The most preferred example for Y may be (E)- $-\text{CH}=\text{CH}-$ .

As the most preferred examples for Z, the above-mentioned preferred examples for Z may be mentioned.

Now, particularly preferred specific compounds of the present invention will be presented. The following compounds (a) to (z) are shown in the form of carboxylic acids. However, the present invention include not only the compounds in the form of carboxylic acids but also the corresponding lactones formed by the condensation of the carboxylic acids with hydroxy at the 5-position, and sodium salts and lower alkyl esters (such as methyl, ethyl, i-propyl and n-propyl esters) of the carboxylic acids, which can be physiologically hydrolyzed to the carboxylic acids.

(a) (E)-3,5-dihydroxy-7-[4'-(4"-fluorophenyl)-2'-(1"-methylethyl)-quinolin-3'-yl]-hept-6-enoic acid

(b) (E)-3,5-dihydroxy-7-[4'-(4"-fluorophenyl)-2'-(1"-methylethyl)-6'-chloro-quinolin-3'-yl]-hept-6-enoic acid

(c) (E)-3,5-dihydroxy-7-[4'-(4"-fluorophenyl)-2'-(1"-methylethyl)-6'-methyl-quinolin-3'-yl]-hept-6-enoic acid

(d) (E)-3,5-dihydroxy-7-[4'-(4"-fluorophenyl)-2'-(1"-methylethyl)-6',7'-dimethoxy-quinolin-3'-yl]-hept-6-enoic acid



5,856,336

5

(e) (E)-3,5-dihydroxy-7-[4'-(4"-fluorophenyl)-2'-cyclopropyl-quinolin-3'-yl]-hept-6-enoic acid

(f) (E)-3,5-dihydroxy-7-[4'-(4"-fluorophenyl)-2'-cyclopropyl-6'-chloro-quinolin-3'-yl]-hept-6-enoic acid

(g) (E)-3,5-dihydroxy-7-[4'-(4"-fluorophenyl)-2'-cyclopropyl-6'-methyl-quinolin-3'-yl]-hept-6-enoic acid

(h) (E)-3,5-dihydroxy-7-[4'-(4"-fluorophenyl)-2'-cyclopropyl-6',7'-dimethoxy-quinolin-3'-yl]-hept-6-enoic acid

(i) (E)-3,5-dihydroxy-7-[4'-(4"-chlorophenyl)-2'-(1"-methylethyl)-quinolin-3'-yl]-hept-6-enoic acid

(j) (E)-3,5-dihydroxy-7-[4'-(4"-chlorophenyl)-2'-(1"-methylethyl)-6'-chloro-quinolin-3'-yl]-hept-6-enoic acid

(k) (E)-3,5-dihydroxy-7-[4'-(4"-chlorophenyl)-2'-(1"-methylethyl)-6'-methyl-quinolin-3'-yl]-hept-6-enoic acid

(l) (E)-3,5-dihydroxy-7-[4'-(4"-chlorophenyl)-2'-(1"-methylethyl)-6',7'-dimethoxy-quinolin-3'-yl]-hept-6-enoic acid

(m) (E)-3,5-dihydroxy-7-[4'-(4"-chlorophenyl)-2'-cyclopropyl-quinolin-3'-yl]-hept-6-enoic acid

(n) (E)-3,5-dihydroxy-7-[4'-(4"-chlorophenyl)-2'-cyclopropyl-6'-chloro-quinolin-3'-yl]-hept-6-enoic acid

(o) (E)-3,5-dihydroxy-7-[4'-(4"-chlorophenyl)-2'-cyclopropyl-6'-methyl-quinolin-3'-yl]-hept-6-enoic acid

(p) (E)-3,5-dihydroxy-7-[4'-(4"-chlorophenyl)-2'-cyclopropyl-6',7'-dimethoxy-quinolin-3'-yl]-hept-6-enoic acid

(q) (E)-3,5-dihydroxy-7-[4'-phenyl-2'-(1"-methylethyl)-quinolin-3'-yl]-hept-6-enoic acid

(r) (E)-3,5-dihydroxy-7-[4'-phenyl-2'-(1"-methylethyl)-6'-chloro-quinolin-3'-yl]-hept-6-enoic acid

(s) (E)-3,5-dihydroxy-7-[4'-phenyl-2'-(1"-methylethyl)-6'-methyl-quinolin-3'-yl]-hept-6-enoic acid

(t) (E)-3,5-dihydroxy-7-[4'-phenyl-2'-(1"-methylethyl)-6',7'-dimethoxy-quinolin-3'-yl]-hept-6-enoic acid

(u) (E)-3,5-dihydroxy-7-[4'-phenyl-2'-cyclopropyl-quinolin-3'-yl]-hept-6-enoic acid

(v) (E)-3,5-dihydroxy-7-[4'-phenyl-2'-cyclopropyl-6'-chloro-quinolin-3'-yl]-hept-6-enoic acid


(w) (E)-3,5-dihydroxy-7-[4'-phenyl-2'-cyclopropyl-6'-methyl-quinolin-3'-yl]-hept-6-enoic acid

(x) (E)-3,5-dihydroxy-7-[4'-phenyl-2'-cyclopropyl-6',7'-dimethoxy-quinolin-3'-yl]-hept-6-enoic acid

(y) (E)-3,5-dihydroxy-7-[4'-(4"-fluorophenyl)-2'-(1"-methylethyl)-6'-methoxy-quinolin-3'-yl]-hept-6-enoic acid






(z) (E)-3,5-dihydroxy-7-[4'-(4"-fluorophenyl)-2'-cyclopropyl-6'-methoxy-quinolin-3'-yl]-hept-6-enoic acid

The mevalonolactones of the formula I can be prepared by the following reaction scheme. The enal III can also be prepared by processes K, L and M.




60

6  
-continued



55

5,856,336



In the above reaction scheme, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup>, R<sup>6</sup> and R<sup>12</sup> are as defined above with respect to the formula I, and

5,856,336

9

R<sup>21</sup> and R<sup>22</sup> independently represent C<sub>1-4</sub> lower alkyl such as methyl, ethyl, n-propyl, i-propyl or n-butyl.

Step A represents a reduction reaction of the ester to a primary alcohol. Such reduction reaction can be conducted by using various metal hydrides, preferably diisobutylaluminum hydride, in a solvent such as tetrahydrofuran or toluene at a temperature of from -20° to 20° C., preferably from -10° to 10° C.

Step B represents an oxidation reaction of the primary alcohol to an aldehyde, which can be conducted by using various oxidizing agents. Preferably, the reaction can be conducted by using pyridinium chlorochromate in methylene chloride at a temperature of from 0° to 25° C., or by using oxalyl chloride, dimethyl sulfoxide and a tertiary amine such as triethylamine (Swern oxidation), or by using a sulfur trioxide pyridine complex.

Step C represents a synthesis of a 3-ethoxy-1-hydroxy-2-propene derivative, which can be prepared by reacting a compound V to lithium compound which has been preliminarily formed by treating cis-1-ethoxy-2-(tri-n-butylstannyl) ethylene with butyl lithium in tetrahydrofuran.

As the reaction temperature, it is preferred to employ a low temperature at a level of from -60° to -78° C.

Step D represents a synthesis of an enal by acidic hydrolysis. As the acid catalyst, it is preferred to employ p-toluene sulfonic acid, hydrochloric acid or sulfuric acid, and the reaction may be conducted in a solvent mixture of water and tetrahydrofuran or ethanol at a temperature of from 10° to 25° C. The 3-ethoxy-1-hydroxy-2-propene derivative obtained in Step C can be used in Step D without purification i.e. by simply removing tetra-n-butyl tin formed simultaneously.

Step E represents a double anion condensation reaction between the enal III and an acetoacetate. Such condensation reaction is preferably conducted by using sodium hydride and n-butyl lithium as the base in tetrahydrofuran at a temperature of from -80° to 0° C., preferably from -30° to -10° C.

Step F represents a reduction reaction of the carbonyl group, which can be conducted by using a metal hydride, preferably sodium borohydride in ethanol at a temperature of from -10° to 25° C., preferably from -10° to 5° C.

Further, the reduction reaction may be conducted by using zinc borohydride in dry ethyl ether or dry tetrahydrofuran at a temperature of -100° to 25° C., preferably from -80° to -50° C.

Step G is a step for hydrolyzing the ester. The hydrolysis can be conducted by using an equimolar amount of a base, preferably potassium hydroxide or sodium hydroxide, in a solvent mixture of water and methanol or ethanol at a temperature of from 10° to 25° C. The free acid hereby obtained may be converted to a salt with a suitable base.

Step H is a step for forming a mevalonolactone by the dehydration reaction of the free hydroxy acid I-2. The dehydration reaction can be conducted in benzene or toluene under reflux while removing the resulting water or by adding a suitable dehydrating agent such as molecular sieve.

Further, the dehydration reaction may be conducted in dry methylene chloride by using a lactone-forming agent such as carbodiimide, preferably a water soluble carbodiimide such as N-cyclohexyl-N'-[2'-(methylmorpholinium)ethyl] carbodiimide p-toluene sulfonate at a temperature of from 10° to 35° C., preferably from 20° to 25° C.

Step J represents a reaction for hydrogenating the double bond connecting the mevalonolactone moiety and the quinoline ring. This hydrogenation reaction can be conducted by using a catalytic amount of palladium-carbon or rhodium-

10

carbon in a solvent such as methanol, ethanol, tetrahydrofuran or acetonitrile at a temperature of from 0° to 50° C., preferably from 10° to 25° C.

Step K represents a reaction for the synthesis of an  $\alpha,\beta$ -unsaturated carboxylic acid ester, whereby a trans-form  $\alpha,\beta$ -unsaturated carboxylic acid ester can be obtained by a so-called Horner-Wittig reaction by using an alkoxy carbonylmethyl phosphonate. The reaction is conducted by using sodium hydride or potassium t-butoxide as the base in dry tetrahydrofuran at a temperature of from -30° to 0° C., preferably from -20° to -15° C.


Step L represents a reduction reaction of the  $\alpha,\beta$ -unsaturated carboxylic acid ester to an allyl alcohol. This reduction reaction can be conducted by using various metal hydrides, preferably diisobutylaluminumhydride, in a solvent such as dry tetrahydrofuran or toluene at a temperature of from -10° to 10° C., preferably from -10° to 0° C.

Step M represents an oxidation reaction of the allyl alcohol to an enal. This oxidation reaction can be conducted by using various oxidizing agents, particularly active manganese dioxide, in a solvent such as tetrahydrofuran, acetone, ethyl ether or ethyl acetate at a temperature of from 0° to 100° C., preferably from 15° to 50° C.

Step N represents a reaction for the synthesis of an  $\alpha,\beta$ -unsaturated ketone by the selective oxidation of the dihydroxy carboxylic acid ester. This reaction can be conducted by using activated manganese dioxide in a solvent such as ethyl ether, tetrahydrofuran, benzene or toluene at a temperature of from 20° to 80° C., preferably from 40° to 80° C.

In addition to the compounds disclosed in Examples given hereinafter, compounds of the formulas I-2 and I-5 given in Table 1 can be prepared by the process of the present invention. In Table 1, i- means iso, sec- means secondary and c- means cyclo. Likewise, Me means methyl, Et means ethyl, Pr means propyl, Bu means butyl, Pent means pentyl, Hex means hexyl and Ph means phenyl.

TABLE 1


						
I-2 (R <sup>12</sup> = H)						
I-5 (R <sup>12</sup> = Na)						
R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>6</sup>	
6-OMe	H	H	H	i-Pr	H	
6-OMe	H	4-F	H	i-Pr	H	
6-Br	H	4-F	H	i-Pr	H	
6-Me	S-Me	4-F	H	i-Pr	H	
7-OMe	8-OMe	4-F	H	i-Pr	H	
6-Br	H	2-F	H	i-Pr	H	
6,7		4-F	H	i-Pr	H	



5,856,336

11

TABLE 1-continued

R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>6</sup>
H	H	4-F	H		H
H	H	4-Ph	H	i-Pr	H
H	H	4-PhCH <sub>2</sub>	H	i-Pr	H
6-Cl	H	4-F	H	c-Pr	H
6-Cl	H	4-F	H	sec-Bu	H
6-OCH <sub>2</sub> Ph	H	4-F	H	i-Pr	H
H	H	4-F	H	i-Bu	H
H	H	4-F	H	c-Pent	H
6-Cl	H	4-F	H	c-Pent	H
6-Me <sub>2</sub> N	H	4-F	H	i-Pr	H
6-Me	H	4-F	H	c-Pr	H
6-i-Pr	H	4-F	H	i-Pr	H
7-Me	H	4-F	H	c-Pr	H
6-OMe	H	4-F	H	c-Pr	H
6-Br	H	4-F	H	c-Pr	H
6-i-Pr	H	4-F	H	c-Pr	H
6-Cl	8-Cl	4-F	H	c-Pr	H
5-F	6-Br	4-F	H	i-Pr	8-Br
6-OMe	7-OMe	4-F	H	i-Pr	8-OMe
6-Me	7-Me	4-F	H	i-Pr	8-Me
6-Cl	7-Cl	4-F	H	i-Pr	8-Cl
H	H	4-F	H	c-Bu	H
H	H	4-F	H	c-Hex	H
6-OMe	7-OMe	H	H	i-Pr	H
6-OMe	7-OMe	4-Cl	H	i-Pr	H
6-OMe	7-OMe	H	H	c-Pr	H
6-OMe	7-OMe	4-Cl	H	c-Pr	H
6-OMe	7-OMe	4-F	H	c-Pr	H
6-Me	H	H	H	i-Pr	H
6-Me	H	4-Cl	H	i-Pr	H
6-Me	H	H	H	c-Pr	H
6-Me	H	4-Cl	H	c-Pr	H
6-Me	H	4-F	H	c-Pr	H
6-Cl	H	H	H	i-Pr	H
6-Cl	H	4-Cl	H	i-Pr	H
6-Cl	H	H	H	c-Pr	H
6-Cl	H	4-Cl	H	c-Pr	H
6-Cl	H	4-F	H	c-Pr	H
H	H	H	H	i-Pr	H
H	H	4-Cl	H	i-Pr	H
H	H	H	H	c-Pr	H
H	H	4-Cl	H	c-Pr	H
H	H	4-F	H	c-Pr	H

Further, pharmaceutically acceptable salts such as potassium salts or esters such as ethyl esters or methyl esters of these compounds can be prepared in the same manner.

The compounds of the present invention exhibit high inhibitory activities against the cholesterol biosynthesis wherein HMG-CoA reductase acts as a rate limiting enzyme, as shown by the test results given hereinafter, and thus are capable of suppressing or reducing the amount of cholesterol in blood as lipoprotein. Thus, the compounds of the present invention are useful as curing agents against hyperlipidemia, hyperlipoproteinemia and atherosclerosis.

12

They may be formulated into various suitable formulations depending upon the manner of the administration. The compounds of the present invention may be administered in the form of free acids or in the form of physiologically hydrolyzable and acceptable esters or lactones, or pharmaceutically acceptable salts.

The pharmaceutical composition of the present invention is preferably administered orally in the form of the compound of the present invention per se or in the form of powders, granules, tablets or capsules formulated by mixing the compound of the present invention with a suitable pharmaceutically acceptable carrier including a binder such as hydroxypropyl cellulose, syrup, gum arabic, gelatin, sorbitol, tragacanth gum, polyvinyl pyrrolidone or CMC-Ca, an excipient such as lactose, sugar, corn starch, calcium phosphate, sorbitol, glycine or crystal cellulose powder, a lubricant such as magnesium stearate, talk, polyethylene glycol or silica, and a disintegrator such as potato starch.

However, the pharmaceutical composition of the present invention is not limited to such oral administration and it is applicable for parenteral administration. For example, it may be administered in the form of e.g. a suppository formulated by using oily base material such as cacao butter, polyethylene glycol, lanolin or fatty acid triglyceride, a transdermal therapeutic base formulated by using liquid paraffin, white vaseline, a higher alcohol, Macrogol ointment, hydrophilic ointment or hydro-gel base material, an injection formulation formulated by using one or more materials selected from the group consisting of polyethylene glycol, hydro-gel base material, distilled water, distilled water for injection and excipient such as lactose or corn starch, or a formulation for administration through mucous membranes such as an ocular mucous membrane, a nasal mucous membrane and an oral mucous membrane.

Further, the compounds of the present invention may be combined with basic ion-exchange resins which are capable of binding bile acids and yet not being absorbed in gastrointestinal tract.

The daily dose of the compound of the formula I is from 0.05 to 500 mg, preferably from 0.5 to 50 mg for an adult. It is administered from once to three times per day. The dose may of course be varied depending upon the age, the weight or the condition of illness of the patient.

The compounds of the formulas II to VII are novel, and they are important intermediates for the preparation of the compounds of the formula I. Accordingly, the present invention relates also to the compounds of the formulas II to VII and the processes for their production.

Now, the present invention will be described in further detail with reference to Test Examples for the pharmacological activities of the compounds of the present invention, their Preparation Examples and Formulation Examples. However, it should be understood that the present invention is by no means restricted by such specific Examples.

#### PHARMACOLOGICAL TEST EXAMPLES

Test A: Inhibition of cholesterol biosynthesis from acetate in vitro

Enzyme solution was prepared from liver of male Wistar rat bilially cannulated and discharged bile for over 24 hours. Liver was cut out at mid-dark and microsome and supernatant fraction which was precipitable with 40-80% of saturation of ammonium sulfate (sup fraction) were prepared from liver homogenate according to the modified method of Knauss et. al.; Kuroda, M., et. al., Biochim. Biophys. Acta, 489, 119 (1977). For assay of cholesterol biosynthesis, microsome (0.1 mg protein) and sup fraction (1.0 mg protein) were incubated for 2 hours at 37° C. in 200  $\mu$ l of the

5,856,336

13

reaction mixture containing ATP; 1 mM, Glutathione; 6 mM, Glucose-1-phosphate; 10 mM, NAD; 0.25 mM, NADP; 0.25 mM, CoA; 0.04 mM and 0.2 mM [ $^{14}\text{C}$ ]sodium acetate (0.2  $\mu\text{Ci}$ ) with 4  $\mu\text{l}$  of test compound solution dissolved in water or dimethyl sulfoxide. To stop reaction and saponify, 1 ml of 15% EtOH-KOH was added to the reactions and heated at 75° C. for 1 hour. Nonsaponifiable lipids were extracted with petroleum ether and incorporated  $^{14}\text{C}$  radioactivity was counted. Inhibitory activity of compounds was indicated with IC50.

Test B: Inhibition of cholesterol biosynthesis in culture cells

Hep G2 cells at over 5th passage were seeded to 12 well plates and incubated with Dulbecco's modified Eagle (DME) medium containing 10% of fetal bovine serum (FBS) at 37° C., 5% CO<sub>2</sub> until cells were confluent for about 7 days. Cells were exposed to the DME medium containing 5% of lipoprotein deficient serum (LpDS) prepared by ultracentrifugation method for over 24 hours. Medium was changed to 0.5 ml of fresh 5% LpDS containing DME before assay and 10  $\mu\text{l}$  of test compound solution dissolved in water or DMSO were added. 0.2  $\mu\text{Ci}$  of [ $^{14}\text{C}$ ]sodium acetate (20  $\mu\text{l}$ ) was added at 0 hr(B-1) or 4 hrs(B-2) after addition of compounds. After 4 hrs further incubation with [ $^{14}\text{C}$ ]sodium acetate, medium was removed and cells were washed with phosphate buffered saline(PBS) chilled at 4° C. Cells were scraped with rubber policeman and collected to tubes with PBS and digested with 0.2 ml of 0.5N KOH at 37° C. Aliquot of digestion was used for protein analysis and remaining was saponified with 1 ml of 15% EtOH-KOH at 75° C. for 1 hour. Nonsaponifiable lipids were extracted with petroleum ether and  $^{14}\text{C}$  radioactivity was counted. Counts were revised by cell protein and indicated with DPM/mg protein. Inhibitory activity of compounds was indicated with IC50.

Test C: Inhibition of cholesterol biosynthesis in vivo

Male Sprague-Dawley rats weighing about 150 g were fed normal Purina chow diet and water ad libitum, and exposed to 12 hours light/12 hours dark lighting pattern (2:00 PM–2:00 AM dark) prior to use for in vivo inhibition test of cholesterol biosynthesis. Animals were separated groups consisting of five rats as to be average mean body weight in each groups. Test compounds at dosage of 0.02–0.2 mg/kg body weight (0.4 ml/100 g body weight), were dissolved in water or suspended or in 0.5% methyl cellulose and orally administered at 2–3 hours before mid-dark (8:00 PM), while cholesterol biosynthesis reaches to maximum in rats. As control, rats were orally administered only water or vehicle. At 90 minutes after sample administration, rats were injected intraperitoneally with 10  $\mu\text{Ci}$  of [ $^{14}\text{C}$ ]sodium acetate at volume of 0.2 ml per one. 2 Hours later, blood samples were obtained and serum were separated immediately. Total lipids were extracted according to the method of Folch et al. and saponified with EtOH-KOH. Nonsaponifiable lipids were extracted with petroleum ether and radio activity incorporated into nonsaponifiable lipids was counted.

Inhibitory activity was indicated as percent decrease of counts in testing groups (DPM/2 ml serum/2 hours) from that in control group.

With respect to the compounds of the present invention, the inhibitory activities against the cholesterol biosynthesis in which HMG-CoA reductase serves as a rate limiting enzyme, were measured by the above Test A and B. The results are shown in Tables, 2, 2-2, 3 and 3-2. Further, the results of the measurements by Test C are also presented.

14

TABLE 2

Inhibitory activities by Test A	
Compound	IC <sub>50</sub> (molar concentration)
(Compounds of the present invention)	
I-13	$1.25 \times 10^{-7}$
I-51	$1.0 \times 10^{-8}$
I-52	$7.1 \times 10^{-8}$
I-53	$1.9 \times 10^{-7}$
(Reference compounds)	
Mevinolin	$1.4 \times 10^{-8}$
CS-514	$9.0 \times 10^{-9}$


In Table 2-2, the relative activities are shown based on the activities of CS-514 being evaluated to be 1.

TABLE 2-2


Relative activities by Test A	
Compound	Relative activities
(Compounds of the present invention)	
I-16	1.75
I-116	2.25
I-117	0.37
I-120	3.21
I-522	0.76

Structures of reference compounds:

(1) Mevinolin



(2) CS-514



5,856,336

15

TABLE 3

Inhibitory activities by Test B-1	
Compound	I <sub>50</sub> (molar concentration)
(Compound of the present invention)	*
I-51 (Reference compound)	1 × 10 <sup>-7</sup>
CS-514	3.5 × 10 <sup>-7</sup>

In Table 3-2, the relative activities are shown based on the activities of CS-514 being evaluated to be 1.

TABLE 3-2

Relative activities by Test B-1	
Compound	Relative activities
I-116	19.4
I-520	20.0
II-20	20.8

Results of the measurement of the inhibitory activities by Test C

The percent decrease of counts after the oral administration of 0.05 mg/kg of compound I-520 was 55% relative to the measured value of the control group. The percent decrease of counts after the oral administration of 10 mg/kg of CS-514 was 55% under the same condition. The compounds of the present invention exhibited activities superior to the reference compound such as CS-514 or Mevinolin in Test A, and exhibited activities superior to CS-514 in Tests B and C.

Test D: Acute toxicity

A 0.5% CMC suspension of a test compound was orally administered to ICR male mice (group of three mice). The acute toxicity was determined based on the mortality after seven days. With compound I-57, I-58, I-59, I-511, I-512, I-513, I-514, I-515, I-517 and I-523 of the present invention, the mortality was 0% even when they were orally administered in an amount of 1000 mg/kg.

#### Example 1

Ethyl (E)-3,5-dihydroxy-7-[4'-(4'-fluorophenyl)-2'-(1'-methylethyl)-quinolin-3'-yl]-hept-6-enoate (compound I-11) (prepared by steps of Example 1-a through Example 1-q)

#### Example 1-a

Ethyl 4-(4'-fluorophenyl)-2-(1'-methylethyl)-quinolin-3-yl-carboxylate (compound VII-1)

The synthesis was conducted in accordance with the method disclosed in J. Org. Chem., 2899 (1966).

6.45 g (0.03 mol) of 2-amino-4'-fluorobenzophenone, 55 5.53 g (0.035 mol) of ethyl isobutyrylacetate and 0.1 ml of conc. sulfuric acid were dissolved in 30 ml of glacial acetic acid, and the mixture was heated at 100° C. for about 10 hours. After confirming the substantial disappearance of 2-amino-4'-fluorobenzophenone by thin layer chromatography, the reaction solution was cooled to room temperature, and a mixture of 45 ml of conc. aqueous ammonia and 120 ml of water cooled with ice, was gradually added thereto. A separated oily substance was solidified when left to stand overnight in a refrigerator. This solid was 65 recrystallized from a small amount of ethanol to obtain 6.47 g (55%) of white powder. Melting point: 68°-70.5° C.

16

#### Example 1-b

4-(4'-fluorophenyl)-3-hydroxymethyl-2-(1'-methylethyl)-quinoline (compound VI-1)

5.4 g (0.016 mol) of compound VII-1 was dissolved in dry 5 toluene under a nitrogen atmosphere and cooled in ice bath to 0° C. To this solution, 40 ml of a 16 wt % diisobutylaluminum hydride-toluene solution was dropwise added, and the mixture was stirred at 0° C. for two hours. After confirming the complete disappearance of compound VII-1 10 by thin layer chromatography, a saturated ammonium chloride solution was added thereto at 0° C. to terminate the reaction. Ethyl ether was added to the reaction mixture, and the organic layer was separated. A gelled product was dissolved by an addition of an aqueous sodium hydroxide 15 solution and extracted anew with ethyl ether. The ethyl ether extracts were put together, dried over anhydrous magnesium sulfate and filtered. The solvent was distilled off. The residual oil underwent crystallization when left to stand. It was recrystallized from ethyl acetate-n-hexane to obtain 3.3 20 g of white crystals. Yield: 70%. Melting point: 136°-137° C.

#### Example 1-c

4-(4'-fluorophenyl)-2-(1'-methylethyl)-quinolin-3-yl-carboxyaldehyde (compound V-1)

2.0 g (9.3 mmol) of pyridinium chlorochromate and 0.4 g 25 of anhydrous sodium acetate was suspended in 10 ml of dry dichloromethane. To this suspension, a solution obtained by dissolving 1 g (3.4 mmol) of compound VI-1 in 10 ml of dry dichloromethane, was immediately added at room temperature. The mixture was stirred for one hour. Then, 100 ml of 30 ethyl ether was added thereto, and the mixture was thoroughly mixed. The reaction mixture was filtered under suction through a silica gel layer. The filtrate was dried under reduced pressure. The residue was dissolved in the isopropyl 35 ether, and insoluble substances were filtered off. The filtrate was again dried under reduced pressure, and the residue was recrystallized from diisopropyl ether to obtain 0.7 g (Yield: 70%) of slightly yellow prism crystals. Melting point: 124°-126° C.

#### Example 1-d

3-(3'-ethoxy-1'-hydroxy-2'-propenyl)-4-(4'-fluorophenyl)-2-(1'-methylethyl)-quinoline (compound IV-1)

1.13 g (3.13 mmol) of cis-1-ethoxy-2-(tri-n-butylstannyl) 45 ethylene was dissolved in 8 ml of dry tetrahydrofuran, and the solution was cooled to -78° C. in a nitrogen stream. To this solution, 2 ml (3.2 mmol) of a 15 wt % n-butyllithium-n-hexane solution was dropwise added. The mixture was stirred for 45 minutes. Then, a solution prepared by dissolving 50 0.76 g (2.6 mmol) of compound V-1 in 10 ml of dry tetrahydrofuran was dropwise added thereto. The reaction mixture was stirred at -78° C. for two hours. Then, 2 ml of a saturated ammonium chloride solution was added thereto to terminate the reaction. The organic layer was extracted with diethyl ether, and the diethyl ether extract was washed with a saturated sodium chloride aqueous solution and dried over anhydrous magnesium sulfate. The solvent was distilled off under reduced pressure. The residue was separated with n-hexane and acetonitrile. The solvent was distilled off under reduced pressure from the acetonitrile layer, and an oily substance thereby obtained was purified by silica gel column chromatography (eluent: 2.5% methanol-chloroform) to obtain 0.91 g of the desired compound in a purified oily form.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ ppm: 1.1(t,3H,7Hz) 1.37(d,6H,J=7Hz) 3.7(m,1H); 3.7(q,2H,J=7Hz) 4.75(t,1H,7Hz) 5.7(m, 1H) 5.95(m,1H) 7.05-8.2(m,8H)

5,856,336

17

## Example 1-e

(E)-3-[4'-(4"-fluorophenyl)-2'-(1"-methylethyl)-quinolin-3'-yl]propanaldehyde (compound III-1)

0.91 g of compound IV-1 was dissolved in 20 ml of tetrahydrofuran, and 5 ml of water and 100 mg of p-toluenesulfonic acid were added thereto. The mixture was stirred at room temperature for 24 hours. The reaction solution was extracted with diethyl ether a few times. The extracts were washed with a saturated sodium chloride aqueous solution and dried over anhydrous magnesium sulfate. Then, the solvent was distilled off. The residue was purified by silica gel column chromatography (eluent: chloroform) to obtain the desired product as white prism crystals. 0.4 g (50%). Melting point: 127°–128° C.

## Example 1-f

Ethyl (E)-7-[4'-(4"-fluorophenyl)-2'-(1"-methylethyl)-quinolin-3'-yl]-5-hydroxy-3-oxohept-6-enoate (compound II-1)

50 mg of 60% sodium hydride was washed with dry petroleum ether and dried under a nitrogen stream, and then suspended in 5 ml of dry tetrahydrofuran. The suspension was cooled to -15° C. in a nitrogen atmosphere. Then, 120 mg (0.92 mmol) of ethyl acetoacetate was dropwise added thereto, and the mixture was stirred for 15 minutes. Then, 0.6 ml (0.92 mmol) of a 15 wt % n-butyllithium-n-hexane solution was dropwise added thereto, and the mixture was stirred for 30 minutes. Then, a solution prepared by dissolving 160 mg (0.5 mmol) of compound III-1 in dry tetrahydrofuran, was dropwise added thereto, and the mixture was stirred for one hour. To the reaction mixture, 1 ml of a saturated ammonium chloride aqueous solution was added at -15° C. Then, the mixture was extracted three times with diethyl ether. The diethyl ether solution was washed with a saturated sodium chloride aqueous solution and dried over anhydrous magnesium sulfate. The solution was evaporated to dryness under reduced pressure. The residue was recrystallized from diisopropyl ether to obtain 130 mg (yield: 59%) of white crystals. Melting point: 99°–101° C.

## Example 1-g

Ethyl (E)-3,5-dihydroxy-7-[4'-(4"-fluorophenyl)-2'-(1"-methylethyl)-quinolin-3'-yl]-hept-6-enoate (compound I-11)

110 mg (0.245 mmol) of compound II-1 was dissolved in 5 ml of ethanol in a nitrogen atmosphere, and the solution was cooled 0° C. Then, 10 mg (0.263 mmol) of sodium borohydride was added, and the mixture was stirred for one hour. Then, 1 ml of a 10% hydrochloric acid aqueous solution was added thereto, and the mixture was extracted three times with ethyl ether. The ethyl ether solution was washed with a saturated sodium chloride aqueous solution and dried over anhydrous magnesium sulfate. Then, the solution was evaporated to dryness under reduced pressure. The residual oil was purified by silica gel column chromatography (eluent: 5% methanol-chloroform) to obtain the desired product as a pure colorless oily substance. 70 mg (Yield: 64%)

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ ppm: 1.30(t,3H,J=8Hz) 1.39(d,6H,J=8Hz) 1.4–1.8(m,2H); 2.42(d,2H,J=7Hz) 3.0–3.8 (m,2H) 3.50(m,1H) 3.9–4.6(m,2H) 4.20(q,2H,J=8Hz) 5.35(m,1H) 6.59(m,1H) 7.10–8.18(m,8H)

## Example 2

Sodium salt of (E)-3,5-dihydroxy-7-[4'-(4"-fluorophenyl)-2'-(1"-methylethyl)-quinolin-3'-yl]-hept-6-enoic acid (compound I-51)

60 mg (0.133 mmol) of compound I-11 was dissolved in 3 ml of ethanol. Then, 0.26 ml of a 0.5N sodium hydroxide

18

aqueous solution was dropwise added thereto. The mixture was stirred at room temperature for further one hour, and ethanol was distilled off under reduced pressure. Then, 5 ml of water was added thereto, and the mixture was extracted with ethyl ether. The aqueous layer was freeze-dried to obtain 40 mg (67%) of hygroscopic white powder. Melting point: 207°–209° C. (decomposed).

## Example 3

(E)-3,5-dihydroxy-7-[4'-(4"-fluorophenyl)-2'-(1"-methylethyl)-quinolin-3'-yl]-hept-6-enoic acid (compound I-21)

110 mg (0.244 mmol) of compound I-11 was dissolved in 10 ml of ethanol. Then, 0.79 ml of a 0.5N sodium hydroxide aqueous solution was dropwise added thereto. The mixture was stirred at room temperature for further one hour, and ethanol was distilled off under reduced pressure. Then, 10 ml of water was added thereto, and the mixture was extracted with ethyl ether. The aqueous layer was weakly acidified (pH 4) with a dilute hydrochloric aqueous solution and extracted three times with ethyl ether. The ethyl ether layers were put together and dried over anhydrous magnesium sulfate. Then, the solvent was distilled off under reduced pressure to obtain 90 mg of slightly yellow oily substance.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ ppm: 1.36(d,6H,J=7Hz) 2.4(m,2H) 3.5(m,1H) 3.45(m,1H); 3.8–4.6(m,2H) 5.40(dd,1H,J<sub>1</sub>=19Hz,J<sub>2</sub>=8Hz) 6.55 (d,1H,J=19Hz) 7.0–8.3(m,8H)

## Example 4

(E)-6-[4'-(4"-fluorophenyl)-2'-(1"-methylethyl)-quinolin-3'-ylethynyl]-4-hydroxy-3,4,5,6-tetrahydro-2H-pyran-2-one (compound I-31)

90 mg of compound I-21 was dissolved in 10 ml of dry toluene, and the solution was refluxed under heating for 3 hours by means of a Dean Stark apparatus.

Toluene was distilled off under reduced pressure, and the residual solid was recrystallized from diisopropyl ether to obtain 40 mg of colorless prism crystals. Melting point: 182°–184° C.

By silica gel thin chromatography, the product gave two absorption spots close to each other attributable to the diastereomers. (Developing solvent: 3% methanol-chloroform)

These diastereomers were separated and isolated by silica gel thin layer chromatography. [Developing solvent: t-BuOMe/hexane/acetone=7/2/1 (v/v), R<sub>f</sub>=0.6 and 0.7 (obtained weight ratio: 1/2)]

R<sub>f</sub>=0.7: trans lactone

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ ppm: 1.40(d,6H,J=7Hz) 1.6(m,2H) 2.65(m,2H) 3.48(m,1H); 4.20(m,1H) 5.15(m,1H) 5.37(dd,1H,J<sub>1</sub>=18Hz,J<sub>2</sub>=7Hz) 6.68(d,1H, J=19Hz) 7.1–8.2(m,8H)

R<sub>f</sub>=0.6: cis lactone

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ ppm: 1.40(d,6H,J=7Hz) 1.6(m,2H) 2.65(m,2H) 3.48(m,1H); 4.20(m,1H) 4.65(m,1H) 5.40(dd,1H,J<sub>1</sub>=18Hz,J<sub>2</sub>=7Hz) 6.66(m,1H) 7.0–8.2(m,8H)

## Example 5

6-[4'-(4"-fluorophenyl)-2'-(1"-methylethyl)-quinolin-3'-ylethynyl]-4-hydroxy-3,4,5,6-tetrahydro-2H-pyran-2-one (compound I-41)

20 mg of a mixture of diastereomers of compound I-31 was dissolved in 5 ml of ethanol, and 10 mg of 5% palladium-carbon was added thereto. The mixture was stirred under a hydrogen atmosphere. After confirming the disappearance of the starting substance and the appearance of a new spot by thin layer chromatography, the palladium-carbon was filtered off, and ethanol was distilled off to obtain colorless oil.

5,856,336

19

This oil was purified by preparative thin layer chromatography to obtain 16 mg of the desired product as pure colorless oil.

MS(m/e): 408(M<sup>+</sup>+H), 407(M<sup>+</sup>), 366, 292, 278

In the same manner as in Example 1-a, compounds VII-2 to VII-27 were prepared. The physical properties of these compounds are shown in Table 4. (In the Table, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5</sup> and R<sup>21</sup> correspond to the substituents of compound VII.)

TABLE 4

(Compounds in this Table are compounds of the formula VII wherein R<sup>6</sup> is hydrogen.)

Compound	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>21</sup>	m.p. (°C.)
VII-2	H	H	4-F	H	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	121-122
VII-3	H	H	H	H	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	102-102.5
VII-4	H	H	H	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	85-85.5
VII-5	6-Cl	H	H	H	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	100.5-101.5
VII-6	6-Cl	H	H	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	105.5-106.5
VII-7	H	H	2-F	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	101.0-102.0
VII-8	7-Me	H	H	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	oil
VII-9	H	H	4-Cl	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	134.0-136.5
VII-10	H	H	4-OMe	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	88.0-89.0
VII-11	H	H	4-Me	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	108.5-109.5
VII-12	6-Cl	H	2-Cl	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	101.0-103.0
VII-13	H	H	4-CF <sub>3</sub>	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	117.5-119.0
VII-14	H	H	3-Me	4-F	i-Pr	C <sub>2</sub> H <sub>5</sub>	oil
VII-15	H	H	3-Me	5-Me	i-Pr	C <sub>2</sub> H <sub>5</sub>	oil
VII-16	6-OMe	7-OMe	4-F	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	96.0-98.0
VII-17	H	H	4-F	H	C <sub>2</sub> H <sub>5</sub>	CH <sub>3</sub>	139.0-139.5
VII-18	H	H	4-F	H	n-Pr	C <sub>2</sub> H <sub>5</sub>	oil
VII-19	6-Cl	H	4-F	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	94.5-95.5
VII-20	H	H	4-F	H	c-Pr	CH <sub>3</sub>	113.5-116.5
VII-21	H	H	4-OPh	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	oil
VII-22	6-Cl	8-Cl	4-F	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	96.0-98.0
VII-23	6-Cl	H	H	H	Ph	C <sub>2</sub> H <sub>5</sub>	118.8-119.5
VII-24	6-Cl	H	H	H	c-Pr	CH <sub>3</sub>	97.0-98.5
VII-25	H	H	4-F	H	sec-Bu	CH <sub>3</sub>	oil
VII-26	6-Me	H	4-F	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	109.0-111.0
VII-27	6-OMe	7-OMe	4-F	H	c-Pr	CH <sub>3</sub>	153.0-153.5

VII-8

H-NMR (in CDCl<sub>3</sub>) δ ppm: 0.92 (t,3H,J=7Hz), 1.41 (d,6H,J=6Hz); 2.47 (s,3H), 3.27 (Heptaplet,1H,J=6Hz) 3.96 (q,2H,J=7Hz), 7.0-7.8(m, 8H)

VII-14

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.01 (t,3H,J=7Hz), 1.42 (d,6H,J=6Hz); 2.38 (s,3H,J=3Hz), 3.25(Heptaplet, 1H,J=6Hz) 4.04 (q,2H,J=7Hz), 6.9-8.1(m,7Hz)

VII-15

H-NMR(in CDCl<sub>3</sub>) δ ppm: 0.97(t,3H,J=7Hz), 1.43 (d,6H,J=6Hz); 2.29 (s,6H) 3.25 (Heptaplet, 1H,J=6Hz) 4.00 (q,2H,J=7Hz), 6.8-8.0(m,7H)

VII-18

H-NMR (in CDCl<sub>3</sub>) δ ppm: 0.98 (t,3H,J=7Hz), 1.02 (t,3H,J=7Hz); 1.6-2.3(m,2H), 2.8-3.1(m,2H) 4.03 (q,2H,J=7Hz), 6.9-8.1(m,8H)

VII 21

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.03 (t,3H,J=7Hz), 1.41 (d,6H,J=6Hz); 3.25(Heptapet,1H,J=6Hz), 4.05(q,2H,J=7Hz), 6.8-8.1(m, 13H)

VII-25

H-NMR (in CDCl<sub>3</sub>) δ ppm: 0.97 (d,6H,J=6Hz), 2.0-2.6 (m,1H); 2.85 (d,2H,J=7Hz), 3.51(s,3H), 6.8-8.1 (m,8H)

In the same manner as in Example 1-b, compounds VI-2 to VI-27 were prepared. (In Table 5, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> and R<sup>5</sup> correspond to the substituents in compound VI.)

20

TABLE 5

(Compounds in this Table are compounds of the formula VI wherein R<sup>6</sup> is hydrogen.)

Compound	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	m.p. (°C.)
VI-2	H	H	p-F	H	CH <sub>3</sub>	—
VI-3	H	H	H	H	CH <sub>3</sub>	149-151
VI-4	H	H	H	H	i-Pr	130-130.5
VI-5	6-Cl	H	H	H	CH <sub>3</sub>	139-141
VI-6	6-Cl	H	H	H	i-Pr	168-169
VI-7	H	H	2-F	H	i-Pr	140.5-142.0
VI-8	7-Me	H	H	H	i-Pr	155.0-157.0
VI-9	H	H	4-Cl	H	i-Pr	192.0-195.0
VI-10	H	H	4-OMe	H	i-Pr	186.0-188.5
VI-11	H	H	4-Me	H	i-Pr	161.0-164.0
VI-12	6-Cl	H	2-Cl	H	i-Pr	122.0-124.0
VI-13	H	H	4-CF <sub>3</sub>	H	i-Pr	183.0-186.0
VI-14	H	H	3-Me	4-F	i-Pr	161.0-162.5
VI-15	H	H	3-Me	5-Me	i-Pr	137.0-138.0
VI-16	6-Me	7-OMe	4-F	H	i-Pr	164.0-165.0
VI-17	H	H	4-F	H	C <sub>2</sub> H <sub>5</sub>	141.5-143.5
VI-18	H	H	4-F	H	n-Pr	146.5-148.5
VI-19	6-Cl	H	4-F	H	i-Pr	171.0-172.0
VI-20	H	H	4-F	H	c-Pr	120-126
VI-21	H	H	4-OPh	H	i-Pr	153.0-154.0
VI-22	6-Cl	8-Cl	4-F	H	i-Pr	98.5-103
VI-23	6-Cl	H	H	H	Ph	171.5-172.5
VI-24	6-Cl	H	H	H	c-Pr	84.0-86.0
VI-25	H	H	4-F	H	sec-Bu	119.0-121.0
VI-26	6-Me	H	4-F	H	i-Pr	160.0-161.5
VI-27	6-OMe	7-OMe	4-F	H	c-Pr	162.0-163.0

In the same manner as in Example 1-c, compounds V-2 to V-27 were prepared. (In Table 6, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> and R<sup>5</sup> correspond to the substituents of compound of V.)

TABLE 6

(Compounds in this Table are compounds of the formula V wherein R<sup>6</sup> is hydrogen.)

Compound	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	m.p. (°C.)
V-2	H	H	p-F	H	CH <sub>3</sub>	125-128
V-3	H	H	H	H	CH <sub>3</sub>	143-146
V-4	H	H	H	H	i-Pr	92-93
V-5	6-Cl	H	H	H	CH <sub>3</sub>	220-222
V-6	6-Cl	H	H	H	i-Pr	140-140.5
V-7	H	H	2-F	H	i-Pr	121.5-124.0
V-8	7-Me	H	H	H	i-Pr	105.1-109.2
V-9	H	H	4-Cl	H	i-Pr	147.0-147.8
V-10	H	H	4-OMe	H	i-Pr	135.6-136.8
V-11	H	H	4-Me	H	i-Pr	119.4-120.4
V-12	6-Cl	H	2-Cl	H	i-Pr	105.8-106.9
V-13	H	H	4-CF <sub>3</sub>	H	i-Pr	163.7-164.2
V-14	H	H	3-Me	4-F	i-Pr	161.1-108.1
V-15	H	H	3-Me	5-Me	i-Pr	120.8-122.3
V-16	6-OMe	7-OMe	4-F	H	i-Pr	164.4-165.2
V-17	H	H	4-F	H	C <sub>2</sub> H <sub>5</sub>	143.1-144.2
V-18	H	H	4-F	H	n-Pr	150.2-155.3
V-19	6-Cl	H	4-F	H	i-Pr	164.5-165.3
V-20	H	H	4-F	H	c-Pr	150.1-151.6
V-21	H	H	4-OPh	H	i-Pr	106.9-107.7
V-22	6-Cl	8-Cl	4-F	H	i-Pr	135.0-135.7
V-23	6-Cl	H	H	H	Ph	174.8-175.3
V-24	6-Cl	H	H	H	c-Pr	157.5-158.0
V-25	H	H	4-F	H	sec-Bu	125.0-126.5
V-26	6-Me	H	4-F	H	i-Pr	155.0-157.0
V-27	6-OMe	7-OMe	4-F	H	c-Pr	200.0-200.5

In the same manner as in Example 1-d, compounds IV-2 to IV-6 were prepared. (In Table 7, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> and R<sup>5</sup> correspond to the substituents of compound IV.)



5,856,336

21

TABLE 7

(Compounds in this Table are compounds of the formula IV wherein R <sup>6</sup> is hydrogen.)						
Compound	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	m.p. (°C.)
IV-2	H	H	4-F	H	CH <sub>3</sub>	177-179
IV-3	H	H	H	H	CH <sub>3</sub>	—
IV-4	H	H	H	H	i-Pr	—
IV-5	6-Cl	H	H	H	CH <sub>3</sub>	—
IV-6	6-Cl	H	H	H	i-Pr	—

In the same manner as in Example 1-e, compounds III-2 to III-27 were prepared. (In Table 8, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> and R<sup>5</sup> correspond to the substituents of compound III.)

TABLE 8

(Compounds in this Table are compounds of the formula III wherein R <sup>6</sup> is hydrogen.)						
Compound	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	m.p. (°C.)
III-2	H	H	4-F	H	CH <sub>3</sub>	194-196
III-3	H	H	H	H	CH <sub>3</sub>	170-171.5
III-4	H	H	H	H	i-Pr	107-108.5
III-5	6-Cl	H	H	H	CH <sub>3</sub>	192-194
III-6	6-Cl	H	H	H	i-Pr	125.5-127
III-7	H	H	2-F	H	i-Pr	80.1-80.2
III-8	7-Me	H	H	H	i-Pr	121.1-122.3
III-9	H	H	4-Cl	H	i-Pr	148.0-149.1
III-10	H	H	4-OMe	H	i-Pr	137.4-140.1
III-11	H	H	4-Me	H	i-Pr	111.6-113.1
III-12	6-Cl	H	2-Cl	H	i-Pr	83.8-84.5
III-13	H	H	4-CF <sub>3</sub>	H	i-Pr	126.2-128.8
III-14	H	H	3-Me	4-F	i-Pr	124.8-126.4
III-15	H	H	3-Me	5-Me	i-Pr	117.6-120.3
III-16	6-OMe	7-OMe	4-F	H	i-Pr	147.8-150.9
III-17	H	H	4-F	H	C <sub>2</sub> H <sub>5</sub>	124.3-128.5
III-18	H	H	4-F	H	n-Pr	117.8-121.5
III-19	6-Cl	H	4-F	H	i-Pr	135.2-135.9
III-20	H	H	4-F	H	c-Pr	141.3-144.1
III-21	H	H	4-OPh	H	i-Pr	oil
III-22	6-Cl	8-Cl	4-F	H	i-Pr	117-122
III-23	6-Cl	H	H	H	Ph	142.8-144.3
III-24	6-Cl	H	H	H	c-Pr	161.0-161.5
III-25	H	H	4-F	H	sec-Bu	78.0-81.0
III-26	6-Me	H	4-F	H	i-Pr	137.0-137.5
III-27	6-OMe	7-OMe	4-F	H	c-Pr	189.5-191.0

III-22

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.40(d6H, J=7Hz), 3.44 (Heptaplet, 1H, J=7Hz); 5.93(dd, 1H, J=8Hz, J=16Hz), 6.8-8.1 (m, 14H) 9.34(d, 1H, J=8Hz)

In the same manner as in Example 1-f, compounds II-2 to II-27 were prepared. (In Table 9, R<sup>1</sup>, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup> and R<sup>5</sup> correspond to the substituents of compound II.)

TABLE 9

(Compounds in this Table are compounds of the formula of II wherein R <sup>6</sup> is hydrogen.)							
Compound	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>12</sup>	m.p. (°C.)
II-2	H	H	p-F	H	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	oil
II-3	H	H	H	H	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	105-106
II-4	H	H	H	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	88.5-90.5
II-5	6-Cl	H	H	H	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	77-82
II-6	6-Cl	H	H	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	96-98
II-7	H	H	2-F	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	oil
II-8	7-Me	H	H	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	68.5-74.0
II-9	H	H	4-Cl	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	91.0-94.0

22

TABLE 9-continued

(Compounds in this Table are compounds of the formula of II wherein R <sup>6</sup> is hydrogen.)							
Compound	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>12</sup>	m.p. (°C.)
II-10	H	H	4-OMe	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	78.0-78.5
II-11	H	H	4-OMe	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	75.0-78.0
II-12	6-Cl	H	2-Cl	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	oil
II-13	H	H	4-CF <sub>3</sub>	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	78.0-83.0
II-14	H	H	3-Me	4-F	i-Pr	C <sub>2</sub> H <sub>5</sub>	66.0-71.0
II-15	H	H	3-Me	5-Me	i-Pr	C <sub>2</sub> H <sub>5</sub>	oil
II-16	6-OMe	7-OMe	4-F	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	83.0-90.0
II-17	H	H	4-F	H	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>	94.0-97.0
II-18	H	H	4-F	H	n-Pr	C <sub>2</sub> H <sub>5</sub>	oil
II-19	6-Cl	H	4-F	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	111.0-113.5
II-20	H	H	4-F	H	c-Pr	C <sub>2</sub> H <sub>5</sub>	91.0-93.0
II-21	H	H	4-OPh	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	121.0-125.0
II-22	6-Cl	8-Cl	4-F	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	oil
II-23	6-Cl	H	H	H	Ph	C <sub>2</sub> H <sub>5</sub>	oil
II-24	6-Cl	H	H	H	c-Pr	C <sub>2</sub> H <sub>5</sub>	69.0-71.0
II-25	H	H	4-F	H	sec-Bu	C <sub>2</sub> H <sub>5</sub>	oil
II-26	6-Me	H	4-F	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	oil
II-27	6-OMe	7-OMe	4-F	H	c-Pr	C <sub>2</sub> H <sub>5</sub>	oil

25 II-7

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.21(t, 3H, J=7Hz), 1.32(d, 6H, J=6Hz); 2.2-2.4(m, 2H), 2.5-2.7(m, 1H) 3.28(s, 1H), 3.34 (Heptaplet, 1H, J=6Hz) 4.08(q, 2H, J=7Hz), 4.3-4.6(m, 1H) 5.28(dd, 1H, J=6Hz, J=15Hz), 6.53(dd, 1H, J=1.5Hz, J=15Hz), 6.9-8.0(m, 8H)

30 6.9-8.0(m, 8H)

II-12

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.25(t, 3H, J=7Hz), 1.33(d, 6H, J=6Hz); 2.2-2.4(m, 2H), 2.5-2.8(m, 1H); 3.32(s, 2H), 3.38 (Heptaplet, 1H, J=6Hz); 4.13(q, 2H, J=7Hz), 4.2-4.6(m, 1H); 5.34(dd, 1H, J=6Hz, J=15Hz), 6.53(dd, 1H, J=1.5Hz, J=15Hz), 7.0-8.0(m, 7H)

35 5.34(dd, 1H, J=6Hz, J=15Hz), 6.53(dd, 1H, J=1.5Hz, J=15Hz), 7.0-8.0(m, 7H)

II-15

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.23(t, 3H, J=7Hz), 1.35(d, 6H, J=6Hz); 2.2-2.4(m, 2H), 2.31(s, 6H); 2.6-2.8(m, 1H), 3.32(s, 2H); 3.35(Heptaplet, 1H, J=6Hz), 4.12(q, 2H, J=7Hz); 4.3-4.7 (m, 1H), 5.30(dd, 1H, J=6Hz, J=16Hz); 6.51(dd, 1H, J=1Hz, J=16Hz), 6.7-8.0(m, 7H)

40 II-18

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.00 (t, 3H, J=7Hz), 1.26(t, 3H, J=7Hz); 1.6-2.3(m, 2H), 2.42 (d, 2H, J=6Hz); 2.6-3.2(m, 3H), 3.35(s, 2H) 4.11(q, 2H, J=7Hz), 4.3-4.7(m, 1H) 5.27(dd, 1H, J=6Hz, J=16Hz) 6.46(dd, 1H, J=1.5Hz, J=16Hz), 6.9-8.0 (m, 8H)

45 II-22

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.26(t, 3H, J=7Hz), 1.33(d, 6H, J=6Hz); 2.43(d, 2H, J=6Hz), 2.6-2.9(m, 1H) 3.36(s, 2H), 3.44 (Heptaplet, 1H, J=6Hz) 4.13(q, 2H, J=7Hz), 4.3-4.7(m, 1H) 5.30(dd, 1H, J=6Hz, J=16Hz), 6.53(dd, 1H, J=1.5Hz, J=16Hz), 7.0-7.6(m, 6H)

50 II-23

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.23(t, 3H, J=7Hz), 2.21(d, 2H, J=6Hz); 2.4-2.6(m, 1H), 3.25(s, 2H) 4.09(q, 2H, J=7Hz), 4.1-4.4(m, 1H) 5.08(dd, 1H, J=6Hz, J=16Hz), 6.26(dd, 1H, J=1.5Hz, J=16Hz), 7.0-8.0 (m, 13H)

60 II-25

H-NMR (in CDCl<sub>3</sub>) δ ppm: 0.96(d, 6H, J=6Hz), 1.26(t, 3H, J=7Hz), 1.8-2.4(m, 1H), 2.43 (d, 2H, J=6Hz), 2.6-2.9(m, 1H), 2.88(d, 2H, J=7Hz), 3.36(s, 2H), 4.14(q, 2H, J=7Hz), 4.3-4.7 (m, 1H), 5.0-5.5(m, 1H), 6.3-6.7(m, 1H), 6.9-8.1(m, 3H)

65 II-26

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.25(t, 3H, J=7Hz), 1.32(d, 6H, J=6Hz), 2.32(s, 3H), 2.39(d, 2H, J=7Hz), 2.6-3.1(m, 1H),

5,856,336

23


3.36(s,2H), 3.41(Heptaplet,1H,J=6Hz), 4.11(q,2H,J=7Hz), 4.3-4.7(m,1H), 5.0-5.5(m,1H), 6.3-6.7(m,1H), 6.8-7.9(m,7H)

I-27

H-NMR (in CDCl<sub>3</sub>) δ ppm: 0.8-1.5(m,4H), 1.26(t,3H,J=7Hz), 2.0-2.9(m,4H), 3.42(s,2H), 3.71(s,3H), 4.00(s,3H), 4.20(q,2H,J=7Hz), 4.4-4.8(m,1H), 5.3-5.8(m,1H), 6.4-6.9(m,1H), 6.58(s,1H), 7.0-7.5(m,5H)

In the same manner as in Example 1-g, compounds I-12 to I-127 were prepared.

TABLE 10



Compound	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>12</sup>	m.p. (°C.) Mass spectrum
I-12	H	H	4-F	H	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	oil 423, 292 M/e 264, 249
I-13	H	H	H	H	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	92-105
I-14	H	H	H	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	97-100
I-15	6-Cl	H	H	H	CH <sub>3</sub>	C <sub>2</sub> H <sub>5</sub>	oil
I-16	6-Cl	H	H	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	oil
I-17	H	H	2-F	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	oil
I-18	7-Me	H	H	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	oil
I-19	H	H	4-Cl	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	98-104
I-110	H	H	4-OMe	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	94-98
I-111	H	H	4-Me	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	79-85
I-112	6-Cl	H	2-Cl	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	oil
I-113	H	H	4-CF <sub>3</sub>	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	117-128
I-114	H	H	3-Me	4-F	i-Pr	C <sub>2</sub> H <sub>5</sub>	85-92
I-115	H	H	3-Me	5-Me	i-Pr	C <sub>2</sub> H <sub>5</sub>	oil
I-116	6-OMe	7-OMe	4-F	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	gum
I-117	H	H	4-F	H	C <sub>2</sub> H <sub>5</sub>	C <sub>2</sub> H <sub>5</sub>	oil
I-118	H	H	4-F	H	n-Pr	C <sub>2</sub> H <sub>5</sub>	oil
I-119	6-Cl	H	4-F	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	79-82
I-120	H	H	4-F	H	c-Pr	C <sub>2</sub> H <sub>5</sub>	100-104
I-121	H	H	4-OPh	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	oil
I-222	6-Cl	8-Cl	4-F	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	133-143
I-123	6-Cl	H	H	H	Ph	C <sub>2</sub> H <sub>5</sub>	gum
I-124	6-Cl	H	H	H	c-Pr	C <sub>2</sub> H <sub>5</sub>	oil
I-125	H	H	4-F	H	sec-Bu	C <sub>2</sub> H <sub>5</sub>	oil
I-126	6-Me	H	4-F	H	i-Pr	C <sub>2</sub> H <sub>5</sub>	oil
I-127	6-OMe	7-OMe	4-F	H	c-Pr	C <sub>2</sub> H <sub>5</sub>	gum

I-17

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.29(t,3H,J=7Hz), 1.40(d,6H,J=6Hz); 1.4-1.7(m,2H), 2.3-2.5(m,2H) 2.9-3.2(m,1H), 3.49(Heptaplet,1H,J=6Hz) 3.5-3.8(m,1H), 3.9-4.5(m,2H) 4.20(q,2H,J=7Hz), 5.2-5.7(m,1H) 6.5-6.9(m,1H), 7.0-8.2(m,8H)

I-18

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.0-1.4(m,2H), 1.31(t,3H,J=7Hz); 1.39(d,6H,J=6Hz), 2.3-2.5(m,2H) 2.52(s,3H), 3.1-3.4 (m,1H) 3.48(Heptaplet,1H,J=6Hz),3.5-3.8(m,1H) 3.8-4.1(m,1H), 4.20(q,2H,J=7Hz) 4.2-4.5(m,1H), 5.2-5.6(m,1H) 6.4-6.8(m,1H), 7.0-8.0(m,8H)

I-19

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.29(t,3H,J=7Hz), 1.38(d,6H,J=6Hz); 1.4-1.8(m,2H), 2.3-2.5(m,2H) 3.2-3.4(m,1H),

24

3.49(Heptaplet,1H,J=6Hz) 3.6-3.8(m,1H), 3.9-4.2(m,1H) 4.20(q,2H,J=7Hz), 4.3-4.5(m,1H) 5.2-5.5(m,1H), 6.5-6.8(m,1H) 7.0-8.2(m,8H)

I-110

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.29(t,3H,J=7Hz), 1.40(d,6H,J=6Hz); 1.5-1.6(m,2H), 2.3-2.5(m,2H) 2.8-3.0(m,1H), 3.4-3.6(m,1H) 3.52(Heptaplet,1H,J=6Hz), 3.88(s,3H) 3.9-4.1(m,1H), 4.20(q,2H,J=7Hz) 4.3-4.5(m,1H), 5.3-5.5(m,1H) 6.5-6.7(m,1H), 6.9-8.1(m,8H)

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.30(t,3H,J=7Hz), 1.3-1.5(m,2H); 1.39(d,6H,J=6Hz), 2.3-2.5(m,2H) 2.43(s,3H), 2.8-3.0(m,1H) 3.50(Heptaplet,1H,J=6Hz), 3.5-3.7(m,1H) 3.9-4.2(m,1H), 4.19(q,2H,J=7Hz) 4.2-4.5(m,1H), 5.2-5.6(m,1H) 6.4-6.8(m,1H), 6.9-8.2(m,8H)

I-112

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.30(t,3H,J=7Hz), 1.3-1.6(m,2H); 1.37(d,6H,J=6Hz), 2.3-2.5(m,2H) 2.9-3.2(m,1H), 3.47(Heptaplet,1H,J=6Hz) 3.5-3.8(m,1H), 3.9-4.1(m,1H) 4.19(q,2H,J=7Hz), 4.2-4.5(m,1H) 5.3-5.7(m,1H), 6.5-6.8(m,1H) 7.1-8.1(m,7H)

I-113

H-NMR(in CDCl<sub>3</sub>) δ ppm: 1.0-1.3(m,2H), 1.30(t,3H,J=7Hz); 1.40(d,6H,J=6Hz), 2.3-2.4(m,2H) 3.3-3.5(m,1H), 3.49 (Heptaplet,1H,J=6Hz) 3.6-3.7(m,1H), 3.9-4.1(m,1H) 4.18(q,2H,J=7Hz), 4.2-4.5(m,1H) 5.1-5.5(m,1H), 6.5-6.8(m,1H) 7.2-8.2(m,8H)

I-114

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.2-1.4(m,2H), 1.30(t,3H,J=7Hz); 1.39(d,6H,J=6Hz), 2.32(bs,3H) 2.3-2.5(m,2H), 3.0-3.3(m,1H) 3.50(Heptaplet,1H,J=6Hz), 3.6-3.8(m,1H) 3.8-4.1(m,1H), 4.20(q,2H,J=7Hz) 4.3-4.6(m,1H), 5.2-5.6(m,1H) 6.5-6.8(m,1H), 7.0-8.2(m,7H)

I-115

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.1-1.4(m,2H), 1.30(t,3H,J=7Hz); 1.40(d,6H,J=6Hz), 2.2-2.5(m,2H) 2.35(s,6H), 2.7-3.1(m,1H) 3.51(Heptaplet, 1H,J=6Hz), 3.6-3.7(m,1H) 3.8-4.1(m,1H), 4.20 (q,2H,J=7Hz) 4.2-4.6(m,1H), 5.2-5.6(m,1H) 6.4-6.8(m,1H), 6.8-8.2(m,7H)

I-116

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.30(t,3H,J=7Hz), 1.37(d,6H,J=6Hz); 1.5-1.8(m,2H), 2.3-2.5(m,2H) 2.9-3.2(m,1H), 3.46 (Heptaplet,1H,J=6Hz) 3.6-3.8(m,1H), 3.75(s,3H) 3.9-4.1(m,1H), 4.07(s,3H) 4.20(q,2H,J=7Hz), 4.2-4.5(m,1H) 5.1-5.5(m,1H), 6.4-6.8(m,2H) 7.1-7.5(m,5H)

I-117

H-NMR(in CDCl<sub>3</sub>) δ ppm: 1.30(t,3H,J=7Hz), 1.37(t,3H,J=7Hz); 1.4-1.7 (m,2H), 2.2-2.6(m,2H) 2.8-3.2(m,3H), 3.6-3.9(m,1H) 3.9-4.7(m,4H), 5.2-5.7(m,1H) 6.3-6.7(m,1H) 7.0-8.2(m,8H)

I-118

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.01(t,3H,J=7Hz), 1.27(t,3H,J=7Hz); 1.4-2.1(m,4H), 2.3-2.6(m,2H); 2.8-3.3(m,3H), 3.6-3.3(m,1H); 3.9-4.1(m,1H), 4.18(q,2H,J=7Hz); 4.2-4.5(m,1H), 5.2-5.6(m,1H); 6.4-6.7(m,1H), 7.0-8.1(m,8H);

I-119

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.2-1.5(m,2H), 1.31(t,3H,J=7Hz); 1.37(d,6H,J=7Hz), 2.3-2.6(m,2H); 3.0-3.4(m,1H), 3.49(Heptaplet,1H,J=6Hz); 3.6-3.8(m,1H), 3.8-4.2(m,1H); 4.20(q,2H,J=7Hz), 4.3-4.5(m,1H); 5.2-5.6(m,1H), 6.4-6.8(m,1H); 7.0-8.1(m,7H);

I-120

H-NMR (in CDCl<sub>3</sub>) δ ppm: 0.8-1.8(m,6H), 1.30(t,3H,J=7Hz); 2.1-2.6(m,3H), 2.9-3.3(m,1H); 3.4-3.7(m,1H), 3.8-4.6(m,2H); 4.20(q,2H,J=7Hz), 5.4-5.8(m,1H); 6.4-6.3(m,1H), 6.8-8.0(m,8H);

I-121

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.29(t,3H,J=7Hz), 1.39(d,6H,J=6Hz); 1.4-1.9(m,2H), 2.3-2.5(m,2H); 2.7-3.2(m,1H),

5,856,336

25

3.51(Heptaplet,1H,J=6Hz); 3.6–3.8(m,1H), 3.9–4.2(m,1H); 4.19(q,2H,J=7Hz), 4.3–4.6(m,1H); 5.2–5.6(m,1H), 6.4–6.8(m,1H); 6.9–8.2(m,13H);

I-122

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.1–1.8(m,2H), 1.31(t,3H,J=7Hz); 1.41(d,6H,J=6Hz), 2.3–2.5(m,2H); 2.9–3.4(m,1H), 3.50(Heptaplet,1H,J=6Hz); 3.6–3.8(m,1H), 3.9–4.5(m,2H); 4.20(q,2H,J=7Hz), 5.2–5.6(m,1H); 6.4–6.8(m,1H), 7.1–7.3(m,5H); 7.72(d,1H,J=6Hz);

I-123

H-NMR (in CDCl<sub>3</sub>) δ ppm: 0.8–1.5(m,2H), 1.29(t,3H,J=7Hz); 2.2–2.4(m,2H), 2.6–2.9(m,1H); 3.2–3.6(m,1H), 3.7–4.3(m,2H); 4.17(q,2H,J=7Hz), 5.0–5.4(m,1H); 6.1–6.5(m,1H), 7.0–8.2(m,13H);

I-124

H-NMR (in CDCl<sub>3</sub>) δ ppm: 0.8–1.8(m,6H), 1.29(t,3H,J=7Hz), 2.2–2.6(m,3H), 2.8–3.2(m,1H), 3.3–3.7(m,1H), 3.9–4.5(m,2H), 4.19(q,2H,J=7Hz), 5.4–5.8(m,1H), 6.5–6.8(m,1H), 7.1–8.0(m,8H),

26

I-125

NMR (in CDCl<sub>3</sub>) δ ppm: 0.94(d,6H,J=6Hz), 1.0–1.7(m,3H), 1.27(t,3H,J=7Hz), 1.9–2.5(m,3H), 2.90(d,2H,J=7Hz), 3.3–4.4(m,3H), 4.12(q,2H,J=7Hz), 5.0–5.5(m,1H), 6.2–6.7(m,1H), 6.9–8.0(m,8H),

I-126

H-NMR (in CDCl<sub>3</sub>) δ ppm: 1.0–1.6(m,3H), 1.21(t,3H,J=7Hz), 1.34(d,6H,J=6Hz), 2.34(s,3H), 2.37(d,2H,J=7Hz), 2.9–3.7(m,2H), 3.8–4.5(m,2H), 4.15(q,2H,J=7Hz), 5.0–5.5(m,1H), 6.3–6.7(m,1H), 6.9–8.0(m,7H),


I-127

H-NMR (in CDCl<sub>3</sub>) δ ppm: 0.8–1.9(m,8H), 1.29(t,3H,J=7Hz),

2.1–2.6(m,3H), 2.8–3.2(m,1H), 3.72(s,3H), 4.02(s,3H), 4.19(q,2H,J=7Hz), 4.3–4.6(m,1H), 5.4–5.8(m,1H), 6.4–6.8(m,1H), 6.56(s,1H), 7.0–7.4(m,5H)

In the same manner as in Exmple 2, compounds I-52 to I-527 were prepared.

TABLE 11



Compound	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>12</sup>	m.p. (°C.)
I-52	H	H	4-F	H	CH <sub>3</sub>	Na	138–142 (decomposed)
I-53	H	H	H	H	CH <sub>3</sub>	Na	130–132 (decomposed)
I-54	H	H	H	H	i-Pr	Na	196–197 (decomposed)
I-55	6-Cl	H	H	H	CH <sub>3</sub>	Na	211–215 (decomposed)
I-56	6-Cl	H	H	H	i-Pr	Na	195–198 (decomposed)
I-57	H	H	2-F	H	i-Pr	Na	193–201 (decomposed)
I-58	7-Me	H	H	H	i-Pr	Na	170–175 (decomposed)
I-59	H	H	4-Cl	H	i-Pr	Na	193–202 (decomposed)
I-510	H	H	4-OMe	H	i-Pr	Na	178–193 (decomposed)
I-511	H	H	4-Me	H	i-Pr	Na	187–200 (decomposed)
I-512	6-Cl	H	2-Cl	H	i-Pr	Na	203–209 (decomposed)
I-513	H	H	4-CF <sub>3</sub>	H	i-Pr	Na	200–212 (decomposed)
I-514	H	H	3-Me	4-F	i-Pr	Na	195–200 (decomposed)
I-515	H	H	3-Me	5-Me	i-Pr	Na	192–197 (decomposed)
I-516	6-OMe	7-OMe	4-F	H	i-Pr	Na	239–245 (decomposed)
I-517	H	H	4-F	H	C <sub>2</sub> H <sub>5</sub>	Na	230–237 (decomposed)
I-518	H	H	4-F	H	n-Pr	Na	193–200 (decomposed)

5,856,336

27

28

TABLE 11-continued

I-5 (R<sup>12</sup> = Na)

Compound	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	R <sup>12</sup>	m.p. (°C.)
I-519	6-Cl	H	4-F	H	i-Pr	Na	193–198 (decomposed)
I-520	H	H	4-F	H	c-Pr	Na	197–199 (decomposed)
I-521	H	H	4-OPh	H	i-Pr	Na	180–189 (decomposed)
I-522	6-Cl	8-Cl	4-F	H	i-Pr	Na	183–187 (decomposed)
I-523	6-Cl	H	H	H	Ph	Na	190–196 (decomposed)
I-524	6-Cl	H	H	H	c-Pr	Na	204–210 (decomposed)
I-525	H	H	4-F	H	sec-Bu	Na	—
I-526	6-Me	H	4-F	H	i-Pr	Na	204–208 (decomposed)
I-527	6-OMe	7-OMe	4-F	H	c-Pr	Na	234–238 (decomposed)

I-57

H-NMR (in DMSO-d<sup>6</sup>) δ ppm: 0.9–1.2(m,2H), 1.37(d, 6H,J=7Hz); 1.6–2.1(m,2H), 3.48(Heptaplet,1H,J=6Hz); 3.7–4.3(m,4H), 5.3–5.6(m,1H); 6.4–6.7(m,1H), 7.1–8.1(m, 8H);

I-58

H-NMR (in DMSO-d<sup>6</sup>) δ ppm: 0.9–1.2(m,2H), 1.31(d, 6H,J=7Hz); 1.7–2.2(m,2H), 2.50(s,3H); 3.3–4.5(m,5H), 5.2–5.6(m,1H); 6.3–6.6(m,1H), 7.1–7.9(m,8H);

I-59

H-NMR (in DMSO-d<sup>6</sup>) δ ppm: 0.9–1.3(m,2H), 1.33(d, 6H,J=7Hz); 1.6–2.2(m,2H), 3.48(Heptaplet,1H,J=7Hz); 3.5–4.6(m,4H), 5.2–5.6(m,2H); 6.3–6.6(m,1H), 7.1–8.1(m, 8H);

I-510

H-NMR (in DMSO-d<sup>6</sup>) δ ppm: 1.0–1.3(m,2H), 1.32(d, 6H,J=7Hz); 1.6–2.2(m,2H), 3.0–3.8(m,4H); 3.86(s,3H), 4.0–4.3(m,1H); 5.3–5.6(m,1H), 6.3–6.6(m,1H); 6.9–8.1(m, 8H);

I-511

H-NMR (in DMSO-d<sup>6</sup>) δ ppm: 0.9–1.3(m,2H), 1.33(d, 6H,J=7Hz); 1.7–2.1(m,2H), 2.41(s,3H); 3.2–4.3(m,5H), 5.3–5.6(m,1H); 6.3–6.6(m,1H), 7.0–8.3(m,8H);

I-512

H-NMR (in DMSO-d<sup>6</sup>) δ ppm: 0.9–1.3(m,2H), 1.33(d, 6H,J=7Hz); 1.6–2.2(m,2H), 3.1–3.8(m,3H); 3.48(Heptaplet, 1H,J=7Hz), 3.9–4.2(m,1H); 5.3–5.7(m,1H), 6.3–6.7(m,1H); 7.0–8.1(m,7H);

I-513

H-NMR (in DMSO-d<sup>6</sup>) δ ppm: 0.8–1.3(m,2H), 1.34(d,6H, J=7Hz); 1.6–2.2(m,2H), 2.7–3.9(m,3H); 3.49(Heptaplet,1H, J=7Hz), 3.9–4.3(m,1H); 5.2–5.6(m,1H), 6.3–6.7(m,1H); 7.1–8.1(m,8H);

I-514

35 H-NMR (in DMSO-d<sup>6</sup>) δ ppm: 0.9–1.3(m,2H), 1.35(d, 6H,J=7Hz); 1.7–2.1(m,2H), 2.30(d,3H,J=2Hz); 3.0–3.8(m, 3H), 3.51(Heptaplet,1H,J=7Hz); 3.9–4.3(m,1H), 5.3–5.6(m, 1H); 6.3–6.6(m,1H), 6.9–8.1(m,7H);

II-515

40 H-NMR (in DMSO-d<sup>6</sup>) δ ppm: 1.0–1.2(m,2H), 1.35(d, 6H,J=7Hz); 1.6–2.2(m,2H), 2.35(s,6H); 3.0–3.8(m,3H), 3.51(Heptaplet,1H,J=7Hz); 4.0–4.3(m,1H), 5.3–5.6(m,1H); 6.3–6.6(m,1H), 6.8–8.0(m,7H);

I-516

45 H-NMR (in DMSO-d<sup>6</sup>) δ ppm: 0.9–1.3(m,2H), 1.31(d, 6H,J=7Hz); 1.7–2.0(m,2H), 3.2–3.7(m,4H); 3.62(s,3H), 3.9–4.2(m,1H); 3.94(s,3H), 5.1–5.5(m,1H); 6.2–6.6(m,1H), 7.0–7.5(m,6H);

I-517

50 H-NMR (in DMSO-d<sup>6</sup>) δ ppm: 0.9–1.5(m,2H), 1.34(t,3H, J=7Hz); 1.6–2.2(m,2H), 2.7–3.4(m,4H); 3.6–4.3(m,2H), 5.2–5.7(m,1H); 6.1–6.6(m,1H), 6.9–8.1(m,8H);

I-518

55 H-NMR (in DMSO-d<sup>6</sup>) δ ppm: 0.8–1.3(m,2H), 1.01(t,3H, J=7Hz); 1.6–2.1(m,4H), 2.7–3.8(m,5H); 3.9–4.3(m,1H), 5.2–5.7(m,1H); 6.3–6.6(m,1H), 7.1–8.1(m,8H);

I-519

60 H-NMR (in DMSO-d<sup>6</sup>) δ ppm: 0.9–1.3(m,2H), 1.33(d, 6H,J=7Hz); 1.6–2.2(m,2H), 2.9–3.9(m,3H); 3.49(Heptaplet, 1H,J=7Hz), 4.0–4.3(m,1H); 5.3–5.6(m,1H), 6.3–6.6(m,1H); 7.2–8.1(m,7H);

I-520

H-NMR (in DMSO-d<sup>6</sup>) δ ppm: 0.8–1.5(m,6H), 1.7–2.2 (m,2H); 2.3–2.7(m,1H), 3.0–3.9(m,3H); 4.0–4.3(m,1H), 5.5–5.8(m,1H); 6.4–6.7(m,1H), 7.2–8.0(m,8H);

I-521

65 H-NMR (in DMSO-d<sup>6</sup>) δ ppm: 0.9–1.5(m,2H), 1.36(d, 6H,J=7Hz); 1.7–2.3(m,2H), 3.0–3.9(m,3H); 3.50(Heptaplet,

5,856,336

29

1H, J=6Hz), 4.0–4.3(m, 1H); 5.2–5.6(m, 1H) 6.4–6.7(m, 1H); 7.0–8.1 (m, 13H);

I-522

H-NMR (in DMSO-d<sub>6</sub>) δ ppm: 0.8–1.3(m, 2H), 1.37(d, 6H, J=7Hz); 1.6–2.2(m, 2H), 3.1–3.9(m, 3H); 3.51(Heptaplet, 1H, J=7Hz), 4.0–4.3(m, 1H); 5.3–5.7(m, 1H), 6.3–6.7(m, 1H); 7.1–8.0(m, 6H);

I-523

H-NMR (in DMSO-d<sub>6</sub>) δ ppm: 0.8–1.4(m, 2H), 1.6–2.1 (m, 2H); 2.9–3.7(m, 3H), 3.7–4.1(m, 1H); 5.1–5.4(m, 1H), 6.1–6.4(m, 1H); 7.1–8.2(m, 13H);

I-524

H-NMR (in DMSO-d<sub>6</sub>) δ ppm: 0.8–1.5(m, 5H), 1.6–2.2 (m, 2H); 2.3–2.7(m, 2H), 3.0–3.8(m, 3H); 3.9–4.3(m, 1H), 5.4–5.8(m, 1H); 6.3–6.6(m, 1H), 7.0–8.0(m, 8H);

I-525

H-NMR (in DMSO-d<sub>6</sub>) δ ppm: 0.9–1.6(m, 2H) 0.96(d, 6H, J=6Hz); 1.7–2.6(m, 3H), 2.89(d, 2H, J=7Hz); 3.0–3.8(m, 3H), 3.9–4.2(m, 1H); 5.2–5.6(m, 1H), 6.2–6.6(m, 1H); 7.1–8.1(m, 8H);

I-526


H-NMR (in DMSO-d<sub>6</sub>) δ ppm: 1.30(d, 6H, J=7Hz), 1.7–2.0(m, 2H), 2.34(s, 3H), 2.4–2.6(m, 1H), 3.0–3.3(m, 2H), 3.3–3.8(m, 3H); 3.9–4.2(m, 1H), 5.2–5.6(m, 1H); 6.3–6.6(m, 1H), 7.0–8.0(m, 7H);

I-527

H-NMR (in DMSO-d<sub>6</sub>) δ ppm: 0.7–1.5(m, 5H), 1.8–2.2 (m, 2H), 2.2–2.6(m, 2H), 3.1–3.3(m, 2H), 3.59(s, 3H), 3.9–4.2 (m, 2H), 3.91(s, 3H), 5.4–5.7(m, 1H), 6.3–6.6(m, 1H), 6.52(s, 1H), 7.0–7.4(m, 5H);

In the same manner as in Example 3, compounds I-22 to I-26 can be prepared.

TABLE 12




Compound	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>
I-22	H	H	4-F	H	CH <sub>3</sub>
I-23	H	H	H	H	CH <sub>3</sub>
I-24	H	H	H	H	i-Pr
I-25	6-Cl	H	H	H	CH <sub>3</sub>
I-26	6-Cl	H	H	H	i-Pr

In the same manner as in Example 4, compounds I-32 to I-36 can be prepared.

30

TABLE 13



Compound	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>
I-32	H	H	4-F	H	CH <sub>3</sub>
I-33	H	H	H	H	CH <sub>3</sub>
I-34	H	H	H	H	i-Pr
I-35	6-Cl	H	H	H	CH <sub>3</sub>
I-36	6-Cl	H	H	H	i-Pr

## FORMULATION EXAMPLE 1

Tablets	
Compound I-51	1.0 g
Lactose	5.0 g
Crystal cellulose powder	8.0 g
Corn starch	3.0 g
Hydroxypropyl cellulose	1.0 g
CMC-Ca	1.5 g
Magnesium stearate	0.5 g
<b>Total</b>	<b>20.0 g</b>

The above components were mixed by a usual method and then tableted to produce 100 tablets each containing 10 mg of the active ingredient.

## FORMULATION EXAMPLE 2

Capsules	
Compound I-51	1.0 g
Lactose	3.5 g
Crystal cellulose powder	10.0 g
Magnesium stearate	0.5
<b>Total</b>	<b>15.0 g</b>

The above components were mixed by a usual method and then packed in No. 4 gelatin capsules to obtain 100 capsules each containing 10 mg of the active ingredient.

## FORMULATION EXAMPLE 3

Soft capsules	
Compound I-51	1.00 g
PEG (polyethylene glycol) 400	3.89 g
Saturated fatty acid triglyceride	15.00 g
Peppermint oil	0.01 g
Polysorbate 80	0.10 g
<b>Total</b>	<b>20.00 g</b>

The above components were mixed and packed in No. 3 soft gelatin capsules by a usual method to obtain 100 soft capsules each containing 10 mg of the active ingredient.

5,856,336

31

FORMULATION EXAMPLE 4

Ointment	
Compound I-51	1.0 g (10.0 g)
Liquid paraffin	10.0 g (10.0 g)
Cetanol	20.0 g (20.0 g)
White vaseline	68.4 g (59.4 g)
Ethylparaben	0.1 g (0.1 g)
L-menthol	0.5 g (0.5 g)
Total	100.0 g

The above components were mixed by a usual method to obtain a 1% (10%) ointment.

FORMULATION EXAMPLE 5

Suppository	
Compound I-51	1.0 g
Witepsol H15*	46.9 g
Witepsol W35*	52.0 g
Polysorbate 80	0.1 g
Total	100.0 g

\*Trademark for triglyceride compound

The above components were melt-mixed by a usual method and poured into suppository containers, followed by cooling for solidification to obtain 100 suppositories of 1 g each containing 10 mg of the active component.

FORMULATION EXAMPLE 6

Injection formulation	
Compound I-51	1 mg
Distilled water for injection formulation	5 ml

The formulation is prepared by dissolving the compound in the distilled water whenever it is required.

32


FORMULATION EXAMPLE 7

Granules	
Compound I-51	1.0 g
Lactose	6.0 g
Crystal cellulose powder	6.5 g
Corn starch	5.0 g
Hydroxypropyl cellulose	1.0 g
Magnesium stearate	0.5 g
Total	20.0 g

The above components were granulated by a usual method and packaged to obtain 100 packages each containing 200 mg of the granules so that each package contains 10 mg of the active ingredient.

We claim:

1. A compound of the formula,



[A]

Z = —CH(OH)—CH<sub>2</sub>—CH(OH)—CH<sub>2</sub>—COO.½Ca.

2. A method for reducing hyperlipidemia, hyperlipoproteinemia or atherosclerosis, which comprises administering an effective amount of the compound of formula A as defined in claim 1.

\* \* \* \* \*

# **EXHIBIT B**

(12) **United States Patent**  
**Van der Schaaf et al.**

(10) **Patent No.:** **US 8,557,993 B2**  
 (45) **Date of Patent:** **Oct. 15, 2013**

(54) **CRYSTALLINE FORMS OF PITAVASTATIN CALCIUM**

FOREIGN PATENT DOCUMENTS

(71) Applicant: **Nissan Chemical Industries, Ltd.**,  
 Tokyo (JP)

(72) Inventors: **Paul Adriaan Van der Schaaf**,  
 Hagenthal-le-Haut (FR); **Fritz Blatter**,  
 Reinach (CH); **Martin Szelagiewicz**,  
 Muenchenstein (CH); **Kai-Uwe**  
**Schoening**, Oberwil (CH)

(73) Assignee: **Nissan Chemical Industries Ltd.**,  
 Tokyo (JP)

(\* ) Notice: Subject to any disclaimer, the term of this  
 patent is extended or adjusted under 35  
 U.S.C. 154(b) by 0 days.

EP	0 304 063	2/1989
EP	0 520 406	12/1992
EP	1 099 694	5/2001
EP	1 472 227	11/2004
EP	1 472 228	11/2004
EP	1 697 326	9/2006
JP	61-171460	8/1986
JP	05-148237	5/1993
JP	6-92970	4/1994
JP	8-12674	1/1996
JP	2005-500382	1/2005
JP	2005-516064	6/2005
JP	2005-520814	7/2005
JP	2007-516952	6/2007
WO	03/016317	2/2003
WO	03/064382	8/2003
WO	WO 03/064392	8/2003
WO	WO 03/070717	8/2003
WO	03/087091	10/2003
WO	WO 2004/072040	8/2004

(21) Appl. No.: **13/664,498**

(22) Filed: **Oct. 31, 2012**

(65) **Prior Publication Data**

US 2013/0053413 A1 Feb. 28, 2013

**Related U.S. Application Data**

(63) Continuation of application No. 13/280,431, filed on  
 Oct. 25, 2011, now abandoned, which is a continuation  
 of application No. 12/331,086, filed on Dec. 9, 2008,  
 now abandoned, which is a continuation of application  
 No. 10/544,752, filed as application No.  
 PCT/EP2004/050066 on Feb. 2, 2004, now  
 abandoned.

(30) **Foreign Application Priority Data**

Feb. 12, 2003 (EP) ..... 03405080

(51) **Int. Cl.**  
**C07D 215/38** (2006.01)

(52) **U.S. Cl.**  
 USPC ..... **546/101**

(58) **Field of Classification Search**  
 USPC ..... 546/101  
 See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,175,944	A	3/1965	Hoeksema	
5,011,930	A *	4/1991	Fujikawa et al.	546/101
5,284,953	A	2/1994	Ohara et al.	
5,407,929	A	4/1995	Takahashi et al.	
5,473,075	A	12/1995	Ohara et al.	
5,514,804	A	5/1996	Ohara et al.	
5,856,336	A *	1/1999	Fujikawa et al.	514/311
5,872,130	A *	2/1999	Fujikawa et al.	514/311
5,939,552	A	8/1999	Ikeda et al.	
6,335,449	B1	1/2002	Ohara et al.	
6,528,661	B2	3/2003	Niddam et al.	
6,835,838	B2	12/2004	Chen et al.	
7,371,865	B2	5/2008	Acemoglu et al.	
2002/0099224	A1	7/2002	Niddam et al.	
2003/0105359	A1	6/2003	Van Der Schaaf et al.	
2003/0233001	A1	12/2003	Storz	
2004/0063961	A1	4/2004	Van Der Schaaf et al.	
2005/0130978	A1	6/2005	Yuda et al.	
2012/0245200	A1	9/2012	Ohara et al.	

OTHER PUBLICATIONS

Jan. 21, 2010 Submission of References in JP 2006-501997 (JP counterpart to present application) with English translation.  
 Mar. 3, 2010 Submission of References in JP 2006-501997 (JP counterpart to present application) with English translation.  
 Mar. 26, 2010 Submission of References in JP 2006-501997 (JP counterpart to present application) with English translation.  
 Jun. 29, 2010 Office Action in JP 2006-501997 (JP counterpart to present application) with English translation.  
 Aug. 23, 2010 Submission of References in JP 2006-501997 (JP counterpart to present application) with English translation.  
 Dec. 27, 2010 Submission of References in JP 2006-501997 (JP counterpart to present application) with English translation.  
 Jan. 21, 2010 Submission of References in JP 2006-520594 (JP counterpart to related U.S. Appl. No. 13/487,289) with English translation.  
 Mar. 3, 2010 Submission of References in JP 2006-520594 (JP counterpart to related U.S. Appl. No. 13/487,289) with English translation.  
 Mar. 26, 2010 Submission of References in JP 2006-520594 (JP counterpart to related U.S. Appl. No. 13/487,289) with English translation.  
 Apr. 12, 2011 Office Action in JP 2006-520594 (JP counterpart to related U.S. Appl. No. 13/487,289) with English translation.  
 Akiba et al., "Six-Month Repeated Oral Toxicity Study of NK-104 in Rats," The Journal of Toxicological Sciences, vol. 23, Supplement V, 713-720, 1998.  
 Aug. 26, 2004 International Search Report in EP 2004-707232 (EP counterpart to present application).  
 Mar. 14, 2006 Office Action in EP 2004-707232 (EP counterpart to present application).  
 Dec. 14, 2006 Third Party Submission in EP 2004-707232 (EP counterpart to present application).

(Continued)

Primary Examiner — D M Seaman

(74) *Attorney, Agent, or Firm* — Oblon, Spivak, McClelland, Maier & Neustadt, L.L.P.

(57) **ABSTRACT**

The present invention is directed to new crystalline forms of Pitavastatin hemicalcium salt, referred to hereinafter as polymorphic Forms A, B, C, D, E and F, as well as the amorphous form. Furthermore, the present invention is directed to processes for the preparation of these crystalline forms and the amorphous form and pharmaceutical compositions comprising these crystalline forms or the amorphous forms.

39 Claims, 9 Drawing Sheets



## US 8,557,993 B2

Page 2


(56)

## References Cited

## OTHER PUBLICATIONS

- Aug. 11, 2008 Third Party Submission in EP 2004-707232 (EP counterpart to present application).
- Feb. 17, 2010 Office Action in EP 2004-707232 (EP counterpart to present application).
- Sep. 29, 2010 Third Party Submission in EP 2004-707232 (EP counterpart to present application).
- Jan. 25, 2011 Office Action in EP 2004-707232 (EP counterpart to present application).
- Jul. 15, 2005 International Search Report in EP 2004-807807 (EP counterpart to related U.S. Appl. No. 13/487,289).
- Sorbera et al., "NK-104: Hypolipidemic HMG-CoA Reductase Inhibitor," *Drugs of the Future* 1998, 23(8), pp. 847-859.
- Nov. 14, 2005 International Preliminary Report on Patentability in EP 2004-807807 (EP counterpart to related U.S. Appl. No. 13/487,289).
- Jan. 19, 2007 Office Action in EP 2004-807807 (EP counterpart to related U.S. Appl. No. 13/487,289).
- Apr. 4, 2008 Office Action in EP 2004-807807 (EP counterpart to related U.S. Appl. No. 13/487,289).
- Aug. 2, 2010 Office Action in EP 2004-807807 (EP counterpart to related U.S. Appl. No. 13/487,289).
- Sep. 29, 2010 Third Party Submission in EP 2004-807807 (EP counterpart to related U.S. Appl. No. 13/487,289).
- Feb. 8, 2011 Office Action in EP 2004-807807 (EP counterpart to related U.S. Appl. No. 13/487,289).
- Apr. 11, 2011 Third Party Submission in EP 2004-807807 (EP counterpart to related U.S. Appl. No. 13/487,289).
- Byrn et al., *Solid State Chemistry of Drugs*, 2d ed., SSCI, Inc., 1998, pp. 59-64.
- The United States Pharmacopeia 2011, *The National Formulary*, USP 34 NF 29, vol. 1.
- Brittain. *Polymorphism in Pharmaceutical Solids*, 2e1 ed., Informa Healthcare USA, 1999, 2009.
- Berge et al., "Pharmaceutical Salts," *Journal of Pharmaceutical Sciences*, vol. 66, No. 1, Jan. 1977, pp. 1 to 18.
- Takahashi et al., "Synthesis of an Artificial HMG-CoA Reductase Inhibitor NK-104 via a Hydroboration-Cross-Coupling Reaction," *Bull. Chem. Soc. Jpn.*, 68, 2649-2656 (1995).
- Miyachi et al., "A Novel Synthetic Method of HMG-CoA Reductase Inhibitor NK-104 Via a Hydroboration-Cross Coupling Sequence," *Tetrahedron Letters*, vol. 34, No. 51, pp. 8267-8270, 1993.
- Bhattacharya, et al., "Thermoanalytical and Crystallographic Methods," *Polymorphism in Pharmaceutical Solids*, Brittain H. ed., 2d ed., Informa Healthcare USA, Inc., 2009, pp. 318-335.
- Ivanisevic, et al., "Uses of X-Ray Powder Diffraction in the Pharmaceutical Industry," *Pharm. Form. Qual.*, 2011, pp. 30-33.
- Evaluation Reports for Approval for Prescription Drug: Pitavastatin Calcium, LIVALO tablet 1 mg, LIVALO tablet 2 mg, <http://www.info.pmda.go.jp/>, made public Sep. 10, 2003, with partial English translation.
- Feb. 5, 2013 Office Action in JP 2011-260984 (JP counterpart to related U.S. Appl. No. 13/487,289), with partial English translation.
- Ogata, *How to Operate Chemical Experiment*, vol. 1, 1963, pp. 154-155, 185-199 (Feb. 5, 2013 Office Action in JP 2011-260984).
- Dec. 10, 2012 Submission of References in JP 2006-501997 (JP counterpart to present application) with English translation.
- Nov. 6, 2012 Submission of References in JP 2006-520594 (JP counterpart to related U.S. Appl. No. 13/487,289), with English translation.
- "Medical Supplies Interview Form: HMG-CoA reductase inhibitor designated drugs LIVALO Tablet 1 mg and LIVALO Tablet 2 mg," Japanese Society of Hospital Pharmacists, Sep. 2003, with English translation.
- Aug. 19, 2010 Submission of References in JP 2006-501997 (JP counterpart to present application) with partial English translation.
- Mar. 8, 2011 Office Action in JP 2006-501997 (JP counterpart to present application) with English translation.
- Apr. 26, 2011 Submission of References in JP 2006-501997 (JP counterpart to present application).
- Sep. 30, 2010 Submission of References in JP 2006-520594 (JP counterpart to related U.S. Appl. No. 13/487,289), with partial English translation.
- Introductory Chemistry Course 2, *Physical Chemistry*, pp. 321-341, Aug. 28, 1997 (Sep. 30, 2011 Submission of References in JP 2006-520594).
- Apr. 26, 2011 Submission of References in JP 2006-520594 (JP counterpart to related U.S. Appl. No. 13/487,289), with partial English translation.
- Japanese Pharmacopoeia, Fourteenth Edition, pp. 49-51 Mar. 30, 2011 (Apr. 26, 2011 Submission of References in JP 2006-520594).
- Mar. 27, 2012 Office Action in JP 2006-520594 (JP counterpart to related U.S. Appl. No. 13/487,289), with English translation.
- Information Offer Form dispatched Apr. 6, 2010 in Japanese Patent Application No. 2006-5001997 (with English-language translation).
- Information Offer Form dispatched Feb. 23, 2010 in Japanese Patent Application No. 2006-501997 (with English-language translation).
- Information Offer Form dispatched Apr. 27, 2010 in Japanese Patent Application No. 2006-501997 (with English-language translation).
- Official Action dispatched Jun. 29, 2010 in Japanese Patent Application No. 2006-501997 (with English-language translation).
- M. Suzuki et al, First Systematic Chiral Syntheses of Two Pairs of Enantiomers with 3,5-Dihydroxyheptenoic Acid Chain, Associated with a Potent Synthetic Statin NK-104, *Biorganic & Medicinal Chemistry Letters*, 9, (1999), 2977-2982.
- Third Party Observation Submitted on Aug. 21, 2010 in JP 2006-501977 (including excerpt from JP-A-2005-520814).
- English Language Translation of Aug. 21, 2010 Third Party Observation Submitted in JP 2006-501977 (including excerpt from WO 03/064932, which is a counterpart to JP-A-2005-520814).
- Suzuki, Mikio, Development Work for HMG-CoA Reductase Inhibitor NK-104, (2001), (partial English translation attached).
- Certificate for Library Material stored in the National Diet Library, Development Work for HMG-CoA Reductase Inhibitor NK-104, Published (2001) Volume Heisei-13, Chief Librarian of Kansai-kan of National Diet Library, Kazuyuki Yamaguchi (partial English translation attached).

\* cited by examiner



*Fig. 1*







Fig. 2



**Fig. 3A**



*Fig. 3B*



*Fig. 4*





Fig. 5



*Fig. 6*







Fig. 7A



**Fig. 7B**

US 8,557,993 B2

1


## CRYSTALLINE FORMS OF PITAVASTATIN CALCIUM

### CROSS REFERENCES TO RELATED APPLICATIONS

This application is a continuation of U.S. patent application Ser. No. 13/280,431, filed Oct. 25, 2011, which is a continuation of U.S. patent application Ser. No. 12/331,086, filed on Dec. 9, 2008, now abandoned; which is a continuation of U.S. patent application Ser. No. 10/544,752, filed on Aug. 8, 2005, now abandoned; which was a 371 of International Patent Application No. PCT/EP2004/050066, filed on Feb. 2, 2004, and claims priority to European Patent Application No. 03405080.7, filed on Feb. 12, 2003, all of which are incorporated herein by reference in their entireties.

The present invention is directed to new crystalline forms and the amorphous form of Pitavastatin calcium, processes for the preparation thereof and pharmaceutical compositions comprising these forms.

The present invention relates to new crystalline forms and the amorphous form of Pitavastatin calcium. Pitavastatin is also known by the names NK-104, Itavastatin and Nisvastatin. Pitavastatin calcium is known by the chemical name: (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt. Pitavastatin calcium has the following formula:



Pitavastatin calcium has recently been developed as a new chemically synthesized and powerful statin by Kowa Company Ltd, Japan. On the basis of reported data, the potency of Pitavastatin is dose-dependent and appears to be equivalent to that of Atorvastatin. This new statin is safe and well tolerated in the treatment of patients with hypercholesterolaemia. Significant interactions with a number of other commonly used drugs can be considered to be extremely low.

Processes for the preparation of Pitavastatin are described in EP-A-0304063 and EP-A-1099694 and in the publications by N. Miyachi et al. in *Tetrahedron Letters* (1993) vol. 34, pages 8267-8270 and by K. Takahashi et al. in *Bull. Chem. Soc. Jpn.* (1995) vol. 68,2649-2656. These publications describe the synthesis of Pitavastatin in great detail but do not describe the hemicalcium salt of Pitavastatin. The publications by L. A. Sorbera et al. in *Drugs of the Future* (1998) vol. 23, pages 847-859 and by M. Suzuki et al. in *Bioorganic & Medicinal Chemistry Letters* (1999) vol. 9, pages 2977-2982 describe Pitavastatin calcium, however, a precise procedure for its preparation is not given. A full synthetic procedure for the preparation of Pitavastatin calcium is described in EP-A-0520406. In the process described in this patent Pitavastatin calcium is obtained by precipitation from an aqueous solution as a white crystalline material with a melting point of 190-192 C. It is known that pharmaceutical substances can exhibit

2

polymorphism. Polymorphism is commonly defined as the ability of any substance to have two or more different crystal structures. Drug substances may also encapsulate solvent molecules when crystallized. These solvates or hydrates are referred to as pseudopolymorphs. It is also possible that the amorphous form is encountered. Different polymorphs, pseudopolymorphs or the amorphous form differ in their physical properties such as melting point, solubility etc. These can appreciably influence pharmaceutical properties such as dissolution rate and bioavailability. It is also economically desirable that the product is stable for extended periods of time without the need for specialized storage conditions. It is therefore important to evaluate polymorphism of drug substances. Furthermore, the discovery of new crystalline polymorphic forms of a drug enlarge the repertoire of materials that a formulation scientist has with which to design a pharmaceutical dosage form of a drug with a targeted release profile or other desired characteristics. We now have surprisingly found novel crystalline forms of Pitavastatin calcium, herein designated as form A, B, C, D, E and F, and the amorphous form of Pitavastatin calcium.

Accordingly, the present invention is directed to the polymorphic Forms A, B, C, D, E and F, and the amorphous form of Pitavastatin calcium salt (2:1).

One object of the invention is a crystalline polymorph of (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt, herein designated as Form A, which exhibits a characteristic X-ray powder diffraction pattern with characteristic peaks expressed in d-values (Å) and in 2θ as given in Table 1 (vs=very strong intensity, s=strong intensity, m=medium intensity, w=weak intensity, vw=very weak intensity).

TABLE 1

d-spacings and 2θ angles for Form A.

d-spacing [Å]	Angle [2θ]	Rel. Intensity
17.6	5.0	s
13.0	6.8	s
9.7	9.1	s
8.8	10.0	w
8.4	10.5	m
8.1	11.0	m
6.7	13.3	vw
6.5	13.7	s
6.3	14.0	w
6.0	14.7	w
5.57	15.9	vw
5.25	16.9	w
5.17	17.1	vw
4.82	18.4	m
4.64	19.1	w
4.27	20.8	vs
4.20	21.1	m
4.10	21.6	m
3.87	22.9	m
3.74	23.7	m
3.67	24.2	s
3.53	25.2	w
3.29	27.1	m
3.02	29.6	vw
2.95	30.2	w
2.63	34.0	w

Another object of the invention is a crystalline polymorph of (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt, herein designated as Form B, which exhibits a characteristic X-ray powder diffraction pattern with characteristic peaks expressed in d-values (Å) and in 2θ as given in Table 2.

## US 8,557,993 B2

3

TABLE 2

d-spacings and 2θ angles for Form B.		
d-spacing [Å]	Angle [2θ]	Rel. Intensity
19.0	4.6	w
16.6	5.3	vs
14.2	6.2	s
11.5	7.7	s
9.6	9.2	m
9.2	9.6	m
8.5	10.3	w
7.8	11.3	m
7.6	11.7	w
7.0	12.6	vw
6.8	13.0	w
6.4	13.9	m
6.0	14.7	vw
5.94	14.9	w
5.66	15.6	w
5.43	16.3	m
5.22	17.0	vw
5.10	17.4	vw
4.92	18.0	w
4.74	18.7	m
4.59	19.3	m
4.43	20.0	s
4.33	20.5	w
4.26	20.8	m
4.19	21.2	w, shoulder
4.13	21.5	m
3.97	22.4	m
3.83	23.2	s
3.73	23.6	m
3.64	24.4	vw
3.53	25.2	w, broad
3.42	26.0	w
3.37	26.4	vw
3.30	27.0	w
3.19	27.9	vw
3.09	28.9	w

Another object of the invention is a crystalline polymorph of (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt, herein designated as Form C, which exhibits a characteristic X-ray powder diffraction pattern with characteristic peaks expressed in d-values (Å) and in 2θ as given in Table 3.

TABLE 3

d-spacings and 2θ angles for Form C.		
d-spacing [Å]	Angle [2θ]	Rel. Intensity
21.6	4.1	m
15.9	5.6	s
11.4	7.8	m
10.6	8.3	m
8.6	10.3	m
7.7	11.6	w
5.06	17.5	w
4.95	17.9	w
4.74	18.7	m
4.55	19.5	s
4.31	20.6	m
4.13	21.5	vw
4.06	21.9	m
3.84	23.1	m
3.71	24.0	w
3.58	24.8	w

Another object of the invention is a crystalline polymorph of (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt, herein designated as Form D, which exhibits a characteristic X-ray powder diffraction pattern with characteristic peaks expressed in d-values (Å) and in 2θ as given in Table 4.

4

TABLE 4

d-spacings and 2θ angles for Form D.		
d-spacing [Å]	Angle [2θ]	Rel. Intensity
17.5	5.0	m
13.5	6.5	m
13.0	6.8	s
10.1	8.7	m
8.8	10.0	m
8.6	10.2	m
8.2	10.8	m
6.8	13.1	w
6.55	13.5	m
6.20	14.3	s
5.78	15.3	vw
5.52	16.1	m
5.28	16.8	w
4.87	18.2	w
4.80	18.5	m
4.66	19.0	w
4.46	19.9	m
4.34	20.5	m
4.23	21.0	vs
4.09	21.7	s
3.99	22.3	w
3.80	23.4	m
3.70	24.0	m
3.47	25.6	w
3.40	26.2	m

Another object of the invention is a crystalline polymorph of (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt, herein designated as Form E, which exhibits a characteristic X-ray powder diffraction pattern with characteristic peaks expressed in d-values (Å) and in 2θ as given in Table 5.

TABLE 5

d-spacings and 2θ angles for Form E.		
d-spacing [Å]	Angle [2θ]	Rel. Intensity
20.0	4.4	vw
17.7	5.0	s
13.4	6.6	s
13.1	6.8	s
10.0	8.9	s
8.8	10.0	m
8.6	10.3	s
8.2	10.8	m
6.6	13.3	s
6.5	13.6	m
6.3	14.0	s
5.84	15.2	vw
5.56	15.9	w
5.39	16.4	w
5.24	16.9	vw
4.99	17.8	vw
4.84	18.3	m
4.69	18.9	w
4.39	20.2	vs
4.34	20.4	m
4.30	20.7	m
4.24	20.9	m
4.21	21.1	vs
4.12	21.6	m
4.08	21.7	m
3.99	22.3	m
3.77	23.5	m
3.73	23.8	m
3.69	24.1	w
3.60	24.7	vw
3.50	25.4	vw
3.35	26.6	m

US 8,557,993 B2

5

TABLE 5-continued

d-spacings and 2θ angles for Form E.		
d-spacing [Å]	Angle [2θ]	Rel. Intensity
2.96	30.2	w
2.64	34.0	vw

Another object of the invention is a crystalline polymorph of (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt, herein designated as Form F, which exhibits a characteristic X-ray powder diffraction pattern with characteristic peaks expressed in d-values (Å) and in 2θ as given in Table 6.

TABLE 6

d-spacing and 2θ angles for Form F.		
d-spacing [Å]	Angle [2θ]	Rel. Intensity
17.2	5.1	m
15.8	5.6	w
12.6	7.0	s
10.0	8.8	m
9.2	9.6	s
8.7	10.2	w
8.1	10.9	m
7.8	11.3	w
7.4	11.9	m
7.1	12.5	m
6.8	13.0	s
6.5	13.7	m
6.2	14.4	s
8.04	14.7	m
5.79	15.3	vw
5.70	15.5	w
5.28	16.8	m
5.03	17.6	w
4.85	18.3	m
4.61	19.3	m
4.51	19.7	m
4.30	20.6	m
4.18	21.2	vs
4.08	21.8	s
3.90	22.8	s
3.84	23.1	w
3.74	23.8	w, shoulder
3.69	24.1	s
3.59	24.8	s
3.46	25.7	m
3.40	26.2	vw
3.35	26.6	m
3.31	26.9	w
3.14	28.4	w
3.02	29.5	w
3.00	29.8	vw
2.89	30.9	m

Small changes in the experimental details can cause small deviation in the d-values and 2θ of characteristic peaks in the X-ray powder diffraction patterns.

Another object of the invention is the amorphous form of (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt which exhibits characteristic X-ray powder diffraction patterns as depicted in FIG. 7.

Powder X-ray diffraction is performed on a Philips 1710 powder X-ray diffractometer using Cu k (α1) radiation (1.54060 Å); 2θ angles are recorded with an experimental error of ±0.1-0.2°. A discussion of the theory of X-ray powder diffraction patterns can be found in "X-ray diffraction procedures" by H. P. Klug and L. E. Alexander, J. Wiley, New York (1974).

6

Furthermore, the present invention is directed to processes for the preparation of Form A, B, C, D, E and F, and the amorphous form of Pitavastatin calcium.

Form A can be generally prepared from Pitavastatin sodium upon reaction with CaCl<sub>2</sub> in an aqueous reaction medium. Alternatively, Form A of the invention may also be obtained in situ from the free acid ((3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid) or the corresponding lactone with Ca(OH)<sub>2</sub>, advantageously also in an aqueous reaction medium. The aqueous reaction medium usually contains at least 80% b.w. of water; preferably it is water or water containing minor amounts of solvents and/or reactants from previous steps. Form A may contain up to 15% water, preferably about 3 to 12%, more preferably 9 to 11% of water.

Form B can be generally prepared by suspending form A in ethanol containing water as a co solvent. The amount of water is preferably about 1 to 50%.

Form C can be generally prepared by suspending form A in isopropanol containing water as a co solvent. The amount of water is preferably about 1 to 50%, especially 1 to 20% and more preferably about 5%. Form C can also be prepared from a mixture of isopropanol and a ketone solvent, containing water as a co solvent. Preferably, the ketone solvent is acetone, and the amount of ketone solvent are about 1 to 30%, more preferably about 10%. The amount of water is preferably about 1 to 20%, more preferably about 5%.

Form D can be generally prepared by suspending form A in absolute ethanol.

Form E can be generally prepared by suspending form A in 1,4-dioxane containing water as a co solvent. The amount of water is preferably about 1 to 50%.

Form F can be generally prepared by suspending form A in methanol containing water as a co solvent. The amount of water is preferably about 1 to 50%.

In the above mentioned processes small amounts of seeding crystals of the desired crystalline form may be added to the reaction mixture. Preferably small amounts are about 1 to 20 weight %, more preferably about 5 weight %. Seeding crystals may be added before or, where appropriate, after the step initiating the crystallization (e. g. cooling, addition of non-solvent etc. as described above). Addition before initiating the crystallization is of specific technical interest.

The amorphous form can be generally prepared by addition of a non-solvent to a concentrated solution of Pitavastatin calcium in an organic solvent. As non-solvent may be taken for example heptane or methyl tert-butyl ether, whereas examples for the organic solvent are 1,4-dioxane, tetrahydrofuran and ethyl methyl ketone. It is preferable that the non-solvent and solvent are miscible. The amorphous form can also be prepared by lyophilization of an aqueous solution of Pitavastatin calcium.

Preparations of polymorphic forms A, B, C, D, E, F as well as the amorphous form are usually done in substantially pure reaction systems, essentially consisting of the educt specified, preferably in substantially crystalline form, and solvents and/or non-solvents as given above.

Another object of the present invention are processes for the preparation of crystalline forms of Pitavastatin calcium essentially free of residual organic solvent.

Particularly, the present invention is related to processes for the preparation of crystalline forms of Pitavastatin calcium essentially free of residual organic solvent by exposing the crystalline form of Pitavastatin calcium to an atmosphere with a defined relative air humidity. More particularly, the present invention is directed to a process for the preparation of any crystalline form or amorphous form of Pitavastatin calcium which is essentially free of residual organic solvent.

US 8,557,993 B2

7

These can, for example, be prepared by exposing the crystalline form or amorphous form to an atmosphere with a relative air humidity of 5 to 100%. Preferably, these are prepared by exposure to an inert gas stream with a defined relative air humidity to exchange residual organic solvent with water. In general, a relative air humidity of 5 to 100%, especially 40 to 80%, is used.

Another object of the present invention are pharmaceutical compositions comprising an effective amount of crystalline polymorphic Form A, B, C, D, E or F or the amorphous form of Pitavastatin calcium, and a pharmaceutical acceptable carrier.

These polymorphic forms may be used as single component or as mixtures with other crystalline forms or the amorphous form.

As to the novel polymorphic forms and amorphous form of Pitavastatin calcium it is preferred that these contain 25-100% by weight, especially 50-100% by weight, of at least one of the novel forms, based on the total amount of Pitavastatin calcium. Preferably, such an amount of the novel polymorphic forms or amorphous form of Pitavastatin calcium is 75-100% by weight, especially 90-100% by weight. Highly preferred is an amount of 95-100% by weight.

The compositions of the invention include powders, granulates, aggregates and other solid compositions comprising at least one of the novel forms. In addition, the compositions that are contemplated by the present invention may further include diluents, such as cellulose-derived materials like powdered cellulose, microcrystalline cellulose, microfine cellulose, methyl cellulose, ethyl cellulose, hydroxyethyl cellulose, hydroxypropyl cellulose, hydroxypropylmethyl cellulose, carboxymethyl, cellulose salts and other substituted and unsubstituted celluloses; starch; pregelatinized starch; inorganic diluents like calcium carbonate and calcium diphosphate and other diluents known to the pharmaceutical industry. Yet other suitable diluents include waxes, sugars and sugar alcohols like mannitol and sorbitol, acrylate polymers and copolymers, as well as pectin, dextrin and gelatin.

Further excipients that are within the contemplation of the present invention include binders, such as acacia gum, pregelatinized starch, sodium alginate, glucose and other binders used in wet and dry granulation and direct compression tableting processes. Excipients that also may be present in the solid compositions further include disintegrants like sodium starch glycolat, crospovidone, low-substituted hydroxypropyl cellulose and others. In addition, excipients may include tableting lubricants like magnesium and calcium stearate and sodium stearyl fumarate; flavorings; sweeteners; preservatives; pharmaceutically acceptable dyes and glidants such as silicon dioxide.

The dosages include dosages suitable for oral, buccal, rectal, parenteral (including subcutaneous, intramuscular, and intravenous), inhalant and ophthalmic administration. Although the most suitable route in any given case will depend on the nature and severity of the condition being treated, the most preferred route of the present invention is oral. The dosages may be conveniently presented in unit dosage form and prepared by any of the methods well-known in the art of pharmacy.

Dosage forms include solid dosage forms, like tablets, powders, capsules, suppositories, sachets, troches and lozenges as well as liquid suspensions and elixirs. While the description is not intended to be limiting, the invention is also not intended to pertain to true solutions of Pitavastatin calcium whereupon the properties that distinguish the solid forms of Pitavastatin calcium are lost. However, the use of the

8

novel forms to prepare such solutions is considered to be within the contemplation of the invention.

Capsule dosages, of course, will contain the solid composition within a capsule which may be made of gelatin or other conventional encapsulating material. Tablets and powders may be coated. Tablets and powders may be coated with an enteric coating. The enteric coated powder forms may have coatings comprising phthalic acid cellulose acetate, hydroxypropylmethyl-cellulose phthalate, polyvinyl alcohol phthalate, carboxymethylethylcellulose, a copolymer of styrene and maleic acid, a copolymer of methacrylic acid and methyl methacrylate, and like materials, and if desired, they may be employed with suitable plasticizers and/or extending agents. A coated tablet may have a coating on the surface of the tablet or may be a tablet comprising a powder or granules with an enteric-coating.

Preferred unit dosages of the pharmaceutical compositions of this invention typically contain from 0.5 to 100 mg of the novel Pitavastatin calcium forms or mixtures thereof with each other or other forms of Pitavastatin calcium. More usually, the combined weight of the

Pitavastatin calcium forms of a unit dosage are from 2.5 mg to 80 mg, for example 5, 10, 20 or 40 mg.

The following Examples illustrate the invention in more detail. Temperatures are given in degrees Celsius.

## EXAMPLE 1

## Preparation of Form A

4.15 gr of (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid tert-butyl ester (Pitavastatin tert-butyl ester) was suspended in 52 ml of a mixture of methyl tert-butyl ether and methanol (10:3). To this mixture were added 2.17 ml of a 4M aqueous solution of NaOH, and the resulting yellowish solution was stirred for 2.5 hours at 50° C. The reaction mixture was cooled to room temperature followed by the addition of 50 ml water and stirring for an additional hour. The aqueous phase was separated and once extracted with 20 ml of methyl tert-butyl ether. To this aqueous solution were added a solution of 0.58 gr CaCl<sub>2</sub> in 80 ml of water over a period of 1 hour. The resulting suspension was stirred for about 16 hours at room temperature. The suspension was filtered and the obtained solid was dried at 40° C. and 50 mbar for about 16 hours. The obtained product is crystal Form A which is characterized by an X-ray powder diffraction pattern as shown in FIG. 1. Further characterization of the obtained Form A by thermogravimetry coupled with FT-IR spectroscopy revealed a water content of about 10%. Differential scanning calorimetry revealed a melting point of 95° C.

## EXAMPLE 2

## Preparation of Form B

100 mg Pitavastatin calcium Form A was suspended in 2 ml water and stirred at room temperature for 30 min, followed by the addition of 2 ml of ethanol and additional stirring for 18 hours. The suspension was filtered and dried in air, yielding 36 mg of Form B. The obtained crystal Form B is characterized by an X-ray powder diffraction pattern as shown in FIG. 2. Further characterization of the obtained Form B by thermogravimetry coupled with FT-IR spectroscopy revealed a water content of about 10%.

US 8,557,993 B2

9

## EXAMPLE 3

## Preparation of Form C

62 mg Pitavastatin calcium Form A was suspended in 2 ml isopropanol containing 5% water. This suspension was heated to 60° C., which led to almost complete dissolution of Form A, and again cooled to room temperature. At this temperature the suspension was stirred for 66 hours. The resulting suspension was filtered, once washed with some isopropanol containing 5% water, and dried in air. The obtained crystal Form C is characterized by an X-ray powder diffraction pattern as shown in FIG. 3. Further characterization of the obtained Form C by thermogravimetry coupled with FT-IR spectroscopy revealed that the sample contains about 6.3% isopropanol and a small amount of water.

## EXAMPLE 4

## Preparation of Form C

65 mg Pitavastatin calcium Form A was suspended in a mixture of 0.9 ml isopropanol, 0.1 ml acetone and 40 µl water. Stirring this suspension for about 1 hour led to nearly complete dissolution. Seeding with 4 mg of Form C (from example 3) and stirring for 2 hours led to the formation of a concentrated suspension. This suspension was diluted with the same amount of solvent mixture as above and stirred for an additional 40 hours. The suspension was filtered and the obtained solid was dried at 40° C. for about 10 min. Analysis by X-ray powder diffraction indicates the product to be crystal Form C as shown in FIG. 3.

## EXAMPLE 5

## Preparation of Form D

60 mg of Pitavastatin calcium Form A was suspended in 1 ml absolute ethanol and stirred at room temperature for 20 hours. The resulting suspension was filtered and dried in air. The obtained crystal Form D is characterized by an X-ray powder diffraction pattern as shown in FIG. 4.

## EXAMPLE 6

## Preparation of Form E

60 mg of Pitavastatin calcium Form A was suspended in a mixture of 1,4-dioxane and water (1:1), and stirred for 18 hours at room temperature. The resulting suspension was filtered and dried in air. The obtained crystal Form E is characterized by an X-ray powder diffraction pattern as shown in FIG. 5.

## EXAMPLE 7

## Preparation of Form F

60 mg of Pitavastatin calcium Form A was suspended in 3 ml methanol containing 20% water, and stirred at 40° C. for 1 hour. The resulting suspension was slowly cooled to room temperature and stirring was continued for 4 hours. The suspension was heated again to 40° C., stirred for 30 min, slowly cooled to room temperature and stirred for an additional 15 hours. The suspension was filtered and the obtained white

10

solid dried in air. The obtained crystal Form F is characterized by an X-ray powder diffraction pattern as shown in FIG. 6.

## EXAMPLE 8

## Preparation of the Amorphous Form

62 mg of Pitavastatin calcium Form A was dissolved in 0.3 ml 1,4-dioxane. To this stirred solution was slowly added 2.3 ml n-heptane at room temperature, and stirred for an additional 16 hours. The resulting suspension was filtered and dried in air. The obtained solid was amorphous as is shown by the X-ray diffraction pattern given in FIG. 7 (top).

## EXAMPLE 9

## Preparation of the Amorphous Form

60 mg of Pitavastatin calcium Form A was dissolved in 1.5 ml ethyl methyl ketone. To this solution was added in steps of 1 ml each 30 sec a total of 21 ml methyl tert-butyl ether. The resulting suspension was stirred at room temperature for about 16 hours. The suspension was filtered and the obtained solid was dried in air. An X-ray diffraction study on the product showed it to be amorphous, see FIG. 7 (bottom). Further characterization of the obtained product by thermogravimetry coupled with FT-IR spectroscopy revealed that the sample contained about 5.5% methyl tert-butyl ether. Differential scanning calorimetry showed the sample to have a glass transition temperature of about 68° C.

## BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a characteristic X-ray powder diffraction pattern for Form A.  
 FIG. 2 is a characteristic X-ray powder diffraction pattern for Form B.  
 FIGS. 3A and 3B are two characteristic X-ray powder diffraction patterns for Form C.  
 FIG. 4 is a characteristic X-ray powder diffraction pattern for Form D.  
 FIG. 5 is a characteristic X-ray powder diffraction pattern for Form E.  
 FIG. 6 is a characteristic X-ray powder diffraction pattern for Form F.  
 FIGS. 7A and 7B are two characteristic X-ray powder diffraction patterns for the amorphous form.

The invention claimed is:

1. A crystalline polymorph A, B, C, D, E, F, or the amorphous form, of (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt wherein

A) polymorph A exhibits a characteristic X-ray powder diffraction pattern with characteristic peaks expressed in 2θ at 5.0 (s), 6.8 (s), 9.1 (s), 10.0 (w), 10.5 (m), 11.0 (m), 13.3 (vw), 13.7 (s), 14.0 (w), 14.7 (w), 15.9 (vw), 16.9 (w), 17.1 (vw), 18.4 (m), 19.1 (w), 20.8 (vs), 21.1 (m), 21.6 (m), 22.9 (m), 23.7 (m), 24.2 (s), 25.2 (w), 27.1 (m), 29.6 (vw), 30.2 (w), 34.0 (w);  
 B) polymorph B exhibits a characteristic X-ray powder diffraction pattern with characteristic peaks expressed in 2θ at 4.6 (w), 5.3 (vs), 6.2 (s), 7.7 (s), 9.2 (m), 9.6 (m), 10.3 (w), 11.3 (m), 11.7 (w), 12.6 (vw), 13.0 (w), 13.9 (m), 14.7 (vw), 14.9 (w), 15.6 (w), 16.3 (m), 17.0 (vw), 17.4 (vw), 18.0 (w), 18.7 (m), 19.3 (m), 20.0 (s), 20.5 (w), 20.8 (m), 21.2 (w, shoulder), 21.5 (m), 22.4 (m),

US 8,557,993 B2

11

- 23.2 (s), 23.8 (m), 24.4 (vw), 25.2 (w, broad), 26.0 (w), 26.4 (vw), 27.0 (w), 27.9 (vw), 28.9 (w);
- C) polymorph C exhibits a characteristic X-ray powder diffraction pattern with characteristic peaks expressed in  $2\theta$  at 4.1 (m), 5.6 (s), 7.8 (m), 8.3 (m), 10.3 (m), 11.6 (w), 17.5 (w), 17.9 (w), 18.7 (m), 19.5 (s), 20.6 (m), 21.5 (vw), 21.9 (m), 23.1 (m), 24.0 (w), 24.8 (w);
- D) polymorph D exhibits a characteristic X-ray powder diffraction pattern with characteristic peaks expressed in  $2\theta$  at 5.0 (m), 6.5 (m), 6.8 (s), 8.7 (m), 10.0 (m), 10.2 (m), 10.8 (m), 13.1 (w), 13.5 (m), 14.3 (s), 15.3 (vw), 16.1 (m), 16.8 (w), 18.2 (w), 18.5 (m), 19.0 (w), 19.9 (m), 20.5 (m), 21.0 (vs), 21.7 (s), 22.3 (w), 23.4 (m), 24.0 (m), 25.6 (w), 26.2 (m);
- E) polymorph E exhibits a characteristic X-ray powder diffraction pattern with characteristic peaks expressed in  $2\theta$  at 4.4 (vw), 5.0 (s), 6.6 (s), 6.8 (s), 8.9 (s), 10.0 (m), 10.3 (s), 10.8 (m), 13.3 (s), 13.6 (m), 14.0 (s), 15.2 (vw), 15.9 (w), 16.4 (w), 16.9 (vw), 17.8 (vw), 18.3 (m), 18.9 (w), 20.2 (vs), 20.4 (m), 20.7 (m), 20.9 (m), 21.1 (vs), 21.6 (m), 21.7 (m), 22.3 (m), 23.5 (m), 23.8 (m), 24.1 (w), 24.7 (vw), 25.4 (vw), 26.6 (m), 30.2 (w), 34.0 (vw); and
- F) polymorph F exhibits a characteristic X-ray powder diffraction pattern with characteristic peaks expressed in  $2\theta$  at 5.1 (m), 5.6 (w), 7.0 (s), 8.8 (m), 9.6 (s), 10.2 (w), 10.9 (m), 11.3 (w), 11.9 (m), 12.5 (m), 13.0 (s), 13.7 (m), 14.4 (s), 14.7 (m), 15.3 (vw), 15.5 (w), 16.8 (m), 17.6 (w), 18.3 (m), 19.3 (m), 19.7 (m), 20.6 (m), 21.2 (vs), 21.8 (s), 22.8 (s), 23.1 (w), 23.8 (w, shoulder), 24.1 (s), 24.8 (s), 25.7 (m), 26.2 (vw), 26.6 (m), 26.9 (w), 28.4 (w), 29.5 (w), 29.8 (vw), 30.9 (m); wherein, for each of said polymorphs, (vs) stands for very strong intensity; (s) stands for strong intensity; (m) stands for medium intensity; (w) stands for weak intensity; (vw) stands for very weak intensity.
2. A process for preparing the crystalline polymorph or amorphous form according to claim 1, wherein: the crystalline polymorph or amorphous form being prepared is the crystalline polymorph A; and the process comprises reacting (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid sodium salt with  $\text{CaCl}_2$  in an aqueous reaction medium.
3. A process for preparing the crystalline polymorph or amorphous form according to claim 1, wherein: the crystalline polymorph or amorphous form being prepared is the crystalline polymorph B; and the process comprises suspending the crystalline polymorph A in ethanol containing water as a cosolvent.
4. The process according to claim 3, wherein the water is present in an amount of 1 to 50% by volume of the suspension of (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt.
5. A process for preparing the crystalline polymorph or amorphous form according to claim 1, wherein: the crystalline polymorph or amorphous form being prepared is the crystalline polymorph C; and the process comprises suspending the crystalline polymorph A the process comprises suspending the crystalline polymorph A in isopropanol containing water as a cosolvent.
6. The process according to claim 5, wherein the water is present in an amount of 1 to 50% by volume of the suspension of (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt.

12

7. A process for preparing the crystalline polymorph or amorphous form according to claim 1, wherein: the crystalline polymorph or amorphous form being prepared is the crystalline polymorph C; and the process comprises suspending the crystalline polymorph A in a mixture of isopropanol and a ketone solvent, containing water as a cosolvent.
8. The process according to claim 7, wherein the ketone solvent is acetone.
9. The process according to claim 7, wherein the ketone solvent is present in an amount of 1 to 30% by volume of the suspension of (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt.
10. The process according to claim 7, wherein the water is present in an amount of 1 to 20% by volume of the suspension of (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt.
11. A process for preparing the crystalline polymorph or amorphous form according to claim 1, wherein: the crystalline polymorph or amorphous form being prepared is the crystalline polymorph D; and the process comprises suspending the crystalline polymorph A in absolute ethanol.
12. A process for preparing the crystalline polymorph or amorphous form according to claim 1, wherein: the crystalline polymorph or amorphous form being prepared is the crystalline polymorph E; and the process comprises suspending the crystalline polymorph A in 1,4-dioxane containing water as a cosolvent.
13. The process according to claim 12, wherein the water is present in the amount of 1 to 50% by volume of the suspension of (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt.
14. A process for preparing the crystalline polymorph or amorphous form according to claim 1, wherein: the crystalline polymorph or amorphous form being prepared is the crystalline polymorph E; and the process comprises suspending the crystalline polymorph A in methanol containing water as a cosolvent.
15. The process according to claim 14, wherein the water is present in an amount of 1 to 50% by volume of the suspension of (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt.
16. The process according to claim 2, wherein (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt is isolated by filtration and dried in air or vacuum.
17. The process according to claim 2, wherein seeding is carried out with crystals of the desired crystalline polymorph.
18. A process preparing the crystalline polymorph or amorphous form according claim 1, wherein: the crystalline polymorph or amorphous form being prepared is the amorphous form; and the process comprises adding a non-solvent to a solution of (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt in an organic solvent.
19. The process according to claim 18, wherein the non-solvent is selected from heptane and methyl tert-butyl ether.
20. The process according to claim 18, wherein the organic solvent is selected from 1,4-dioxane, tetrahydrofuran and ethyl methyl ketone.
21. A process for preparing the crystalline polymorph or amorphous form according claim 1, wherein:



the crystalline polymorph or amorphous form being prepared is the amorphous form; and

the process comprises drying an aqueous solution of (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt by lyophilization.

22. A pharmaceutical composition comprising an effective amount of the crystalline polymorph or amorphous form according to claim 1, and a pharmaceutically acceptable carrier.

23. A crystalline polymorph A, B, C, D, E, F, or the amorphous form, of (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt of claim 1, wherein polymorph A has an X-ray powder diffraction pattern substantially as depicted in FIG. 1, polymorph B has an X-ray powder diffraction pattern substantially as depicted in FIG. 2, polymorph C has an X-ray powder diffraction pattern substantially as depicted in FIGS. 3A and 3B, polymorph D has an X-ray powder diffraction pattern substantially as depicted in FIG. 4, polymorph E has an X-ray powder diffraction pattern substantially as depicted in FIG. 5, polymorph F has an X-ray powder diffraction pattern substantially as depicted in FIG. 6, and

the amorphous form has an X-ray powder diffraction pattern substantially as depicted in FIGS. 7A and 7B.

24. A crystalline polymorph A of (3R,5S)-7[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt, which exhibits a characteristic X-ray powder diffraction pattern with characteristic peaks expressed in 2 $\theta$  at 5.0 (s), 6.8 (s), 9.1 (s), 10.0 (w), 10.5 (m), 11.0 (m), 13.3 (vw), 13.7 (s), 14.0 (w), 14.7 (w), 15.9 (vw), 16.9 (w), 17.1 (vw), 18.4 (m), 19.1 (w), 20.8 (vs), 21.1 (m), 21.6 (m), 22.9 (m), 23.7 (m), 24.2 (s), 25.2 (w), 27.1 (m), 29.6 (vw), 30.2 (w), and 34.0 (w), wherein (vs) stands for very strong intensity, (s) stands for strong intensity, (m) stands for medium intensity, (w) stands for weak intensity, and (vw) stands for very weak intensity.

25. A crystalline polymorph A of (3R,5S)-7[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt, having an X-ray powder diffraction pattern substantially as depicted in FIG. 1.

26. A crystalline polymorph B of (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt, which exhibits a characteristic X-ray powder diffraction pattern with characteristic peaks expressed in 2 $\theta$  at 4.6 (w), 5.3 (vs), 6.2 (s), 7.7 (s), 9.2 (m), 9.6 (m), 10.3 (w), 11.3 (m), 11.7 (w), 12.6 (vw), 13.0 (w), 13.9 (m), 14.7 (vw), 14.9 (w), 15.6 (w), 16.3 (m), 17.0 (vw), 17.4 (vw), 18.0 (w), 18.7 (m), 19.3 (m), 20.0 (s), 20.5 (w), 20.8 (m), 21.2 (w, shoulder), 21.5 (m), 22.4 (m), 23.2 (s), 23.8 (m), 24.4 (vw), 25.2 (w, broad), 26.0 (w), 26.4 (vw), 27.0 (w), 27.9 (vw), and 28.9 (w), wherein (vs) stands for very strong intensity, (s) stands for strong intensity, (m) stands for medium intensity, (w) stands for weak intensity, and (vw) stands for very weak intensity.

27. A crystalline polymorph B of (3R,5S)-7-[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt, having an X-ray powder diffraction pattern substantially as depicted in FIG. 2.

28. A crystalline polymorph C of (3R,5S)-7[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt, which exhibits a characteristic X-ray powder diffraction pattern with characteristic peaks expressed in 2 $\theta$  at 4.1 (m), 5.6 (s), 7.8 (m), 8.3 (m), 10.3 (m), 11.6 (w), 17.5 (w), 17.9 (w), 18.7 (m), 19.5 (s), 20.6 (m), 21.5 (vw), 21.9 (m), 23.1 (m), 24.0 (w), and 24.8 (w), wherein (vs) stands for very strong intensity, (s) stands for strong

intensity, (m) stands for medium intensity, (w) stands for weak intensity, and (vw) stands for very weak intensity.

29. A crystalline polymorph C of (3R,5S)-7[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt, having an X-ray powder diffraction pattern substantially as depicted in FIGS. 3A and 3B.

30. A crystalline polymorph D of (3R,5S)-7[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt, which exhibits a characteristic X-ray powder diffraction pattern with characteristic peaks expressed in 2 $\theta$  at 5.0 (m), 6.5 (m), 6.8 (s), 8.7 (m), 10.0 (m), 10.2 (m), 10.8 (m), 13.1 (w), 13.5 (m), 14.3 (s), 15.3 (vw), 16.1 (m), 16.8 (w), 18.2 (w), 18.5 (m), 19.0 (w), 19.9 (m), 20.5 (m), 21.0 (vs), 21.7 (s), 22.3 (w), 23.4 (m), 24.0 (m), 25.6 (w), and 26.2 (m), wherein (vs) stands for very strong intensity, (s) stands for strong intensity, (m) stands for medium intensity, (w) stands for weak intensity, and (vw) stands for very weak intensity.

31. A crystalline polymorph D of (3R,5S)-7[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt, having an X-ray powder diffraction pattern substantially as depicted in FIG. 4.

32. A crystalline polymorph E of (3R,5S)-7[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt, which exhibits a characteristic X-ray powder diffraction pattern with characteristic peaks expressed in 2 $\theta$  at 4.4 (vw), 5.0 (s), 6.6 (s), 6.8 (s), 8.9 (s), 10.0 (m), 10.3 (s), 10.8 (m), 13.3 (s), 13.6 (m), 14.0 (s), 15.2 (vw), 15.9 (w), 16.4 (w), 16.9 (vw), 17.8 (vw), 18.3 (m), 18.9 (w), 20.2 (vs), 20.4 (m), 20.7 (m), 20.9 (m), 21.1 (vs), 21.6 (m), 21.7 (m), 22.3 (m), 23.5 (m), 23.8 (m), 24.1 (w), 24.7 (vw), 25.4 (vw), 26.6 (m), 30.2 (w), and 34.0 (vw), wherein (vs) stands for very strong intensity, (s) stands for strong intensity, (m) stands for medium intensity, (w) stands for weak intensity, and (vw) stands for very weak intensity.

33. A crystalline polymorph E of (3R,5S)-7[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt, having an X-ray powder diffraction pattern substantially as depicted in FIG. 5.

34. A crystalline polymorph F of (3R,5S)-7[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt, which exhibits a characteristic X-ray powder diffraction pattern with characteristic peaks expressed in 2 $\theta$  at 5.1 (m), 5.6 (w), 7.0 (s), 8.8 (m), 9.6 (s), 10.2 (w), 10.9 (m), 11.3 (w), 11.9 (m), 12.5 (m), 13.0 (s), 13.7 (m), 14.4 (s), 14.7 (m), 15.3 (vw), 15.5 (w), 16.8 (m), 17.6 (w), 18.3 (m), 19.3 (m), 19.7 (m), 20.6 (m), 21.2 (vs), 21.8 (s), 22.8 (s), 23.1 (w), 23.8 (w, shoulder), 24.1 (s), 24.8 (s), 25.7 (m), 26.2 (vw), 26.6 (m), 26.9 (w), 28.4 (w), 29.5 (w), 29.8 (vw), and 30.9 (m), wherein (vs) stands for very strong intensity, (s) stands for strong intensity, (m) stands for medium intensity, (w) stands for weak intensity, and (vw) stands for very weak intensity.

35. A crystalline polymorph F of (3R,5S)-7[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt, having an X-ray powder diffraction pattern substantially as depicted in FIG. 6.

36. The amorphous form of (3R,5S)-7[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt.

37. The amorphous form of (3R,5S)-7[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt, having an X-ray powder diffraction pattern substantially as depicted in FIGS. 7A and 7B.

38. A process for preparing the crystalline polymorph or amorphous form according to claim 1, wherein:

the crystalline polymorph or amorphous form being prepared is the crystalline polymorph F; and the process comprises suspending the crystalline polymorph A in methanol containing water as a cosolvent.

39. The process according to claim 38, wherein the water is present in an amount of 1 to 50% by volume of the suspension of (3R,5S)-7[2-cyclopropyl-4-(4-fluorophenyl)quinolin-3-yl]-3,5-dihydroxy-6(E)-heptenoic acid hemicalcium salt.

\* \* \* \* \*