

Charles M. Lizza
William C. Baton
Saul Ewing LLP
One Riverfront Plaza, Suite 1520
Newark, NJ 07102-5426
(973) 286-6700
clizza@saul.com

*Attorneys for Plaintiff
Celgene Corporation*

**UNITED STATES DISTRICT COURT
DISTRICT OF NEW JERSEY**

CELGENE CORPORATION,

Plaintiff,

v.

**ZYDUS PHARMACEUTICALS (USA)
INC., ZYDUS INTERNATIONAL PVT.
LTD., and CADILA HEALTHCARE
LIMITED,**

Defendants.

Civil Action No. _____

**COMPLAINT FOR
PATENT INFRINGEMENT**

(Filed Electronically)

Plaintiff Celgene Corporation (“Celgene”), by its undersigned attorneys, for its Complaint against defendants Zydus Pharmaceuticals (USA) Inc. (“Zydus USA”), Zydus International Pvt. Ltd. (“Zydus International”), and Cadila Healthcare Limited (“Zydus Cadila”) (collectively with Zydus USA and Zydus International, “Zydus”) alleges as follows:

Nature of the Action

1. This is an action for patent infringement under the patent laws of the United States, 35 U.S.C. §100, *et seq.*, arising from Zydus’s filing of an Abbreviated New Drug Application (“ANDA”) No. 210154 (“Zydus’s ANDA”) with the United States Food and Drug Administration (“FDA”) seeking approval to commercially market generic versions of Celgene’s

2.5 mg, 5 mg, 10 mg, 15 mg, 20 mg, and 25 mg REVLIMID[®] drug products prior to the expiration of United States Patent Nos. 7,465,800 (the “800 patent”), 7,855,217 (the “217 patent”), 7,968,569 (the “569 patent”), 8,530,498 (the “498 patent”), 8,648,095 (the “095 patent”), 9,101,621 (the “621 patent”), and 9,101,622 (the “622 patent”), all owned by Celgene (collectively, “the patents-in-suit”).

The Parties

2. Plaintiff Celgene is a biopharmaceutical company committed to improving the lives of patients worldwide. Celgene focuses on, and invests heavily in, the discovery and development of products for the treatment of severe and life-threatening conditions, including cancer. Celgene is a world leader in the treatment of many such diseases, including cancer. Celgene is a corporation organized and existing under the laws of the State of Delaware, having a principal place of business at 86 Morris Avenue, Summit, New Jersey 07901.

3. On information and belief, Defendant Zydus Pharmaceuticals (USA) Inc. is a corporation organized and existing under the laws of the State of New Jersey, having a principal place of business at 73 Route 31 North, Pennington, New Jersey 08534.

4. On information and belief, Defendant Zydus International Pvt. Ltd. is a corporation organized and existing under the laws of Ireland, having a principal place of business at FDW House, Blackthorn Business Park, Coes Road, Dundalk, Co. Louth, Ireland.

5. On information and belief, Defendant Cadila Healthcare Limited is a corporation organized and existing under the laws of India, having a principal place of business at Zydus Tower, Satellite Cross Roads, Ahmedabad 380015, Gujarat, India.

6. On information and belief, the parent corporation of Zydus Pharmaceuticals (USA) Inc. is Zydus International Pvt. Ltd., which is a wholly owned subsidiary of Cadila Healthcare Limited (d/b/a “Zydus Cadila”).

The Patents-in-Suit

7. On December 16, 2008, the United States Patent and Trademark Office (“USPTO”) duly and lawfully issued the ’800 patent, entitled, “Polymorphic Forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione,” to Celgene as assignee of the inventors Markian S. Jaworsky, Roger Shen-Chu Chen, and George W. Muller. A copy of the ’800 patent is attached hereto as Exhibit A.

8. On December 21, 2010, the USPTO duly and lawfully issued the ’217 patent, entitled, “Polymorphic Forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione,” to Celgene as assignee of the inventors Markian S. Jaworsky, Roger Shen-Chu Chen, and George W. Muller. A copy of the ’217 patent is attached hereto as Exhibit B.

9. On June 28, 2011, the USPTO duly and lawfully issued the ’569 patent, entitled, “Methods For Treatment of Multiple Myeloma Using 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione,” to Celgene as assignee of the inventor Jerome B. Zeldis. A copy of the ’569 patent is attached hereto as Exhibit C.

10. On September 10, 2013, the USPTO duly and lawfully issued the ’498 patent, entitled, “Methods For Treating Multiple Myeloma With 3-(4-amino-1-oxo-1,3-dihydroisoindol-2-yl) piperidine-2,6-dione,” to Celgene as assignee of the inventor Jerome B. Zeldis. A copy of the ’498 patent is attached hereto as Exhibit D.

11. On February 11, 2014, the USPTO duly and lawfully issued the ’095 patent, entitled, “Methods For Treating Multiple Myeloma Using 3-(4-amino-1-oxo-1,3-dihydroisoindol-2-yl)-piperidine-2,6-dione In Combination With Proteasome Inhibitor,” to Celgene as assignee of the inventor Jerome B. Zeldis. A copy of the ’095 patent is attached hereto as Exhibit E.

12. On August 11, 2015, the USPTO duly and lawfully issued the '621 patent, entitled, "Methods For Treating Multiple Myeloma With 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione After Stem Cell Transplantation," to Celgene as assignee of the inventor Jerome B. Zeldis. A copy of the '621 patent is attached hereto as Exhibit F.

13. On August 11, 2015, the USPTO duly and lawfully issued the '622 patent, entitled, "Methods For Treating Newly-Diagnosed Multiple Myeloma 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione In Combination With Dexamethasone," to Celgene as assignee of the inventor Jerome B. Zeldis. A copy of the '622 patent is attached hereto as Exhibit G.

The REVLIMID[®] Drug Product

14. Celgene holds an approved New Drug Application ("NDA") under Section 505(a) of the Federal Food Drug and Cosmetic Act ("FFDCA"), 21 U.S.C. § 355(a), for lenalidomide capsules (NDA No. 21-880), which it sells under the trade name REVLIMID[®]. REVLIMID[®] is an FDA-approved medication used for the treatment of certain forms of cancer, including multiple myeloma (MM), in combination with dexamethasone. The claims of the patents-in-suit cover, *inter alia*, solid forms of lenalidomide, pharmaceutical compositions containing lenalidomide, and methods of use and administration of lenalidomide or pharmaceutical compositions containing lenalidomide.

15. Pursuant to 21 U.S.C. § 355(b)(1) and attendant FDA regulations, the patents-in-suit are listed in the FDA publication, "Approved Drug Products with Therapeutic Equivalence Evaluations" (the "Orange Book"), with respect to REVLIMID[®].

16. The labeling for REVLIMID[®] instructs and encourages physicians, pharmacists, and other healthcare workers and patients to administer REVLIMID[®] according to one or more of the methods claimed in the patents-in-suit.

Jurisdiction and Venue

17. This Court has jurisdiction over the subject matter of this action pursuant to 28 U.S.C. §§ 1331, 1338(a), 2201, and 2202.

18. This Court has personal jurisdiction over Zydus USA by virtue of, *inter alia*, its systematic and continuous contacts with the State of New Jersey. On information and belief, Zydus USA's principal place of business is in Pennington, New Jersey. On information and belief, Zydus USA is registered with the State of New Jersey's Division of Revenue and Enterprise Services as a business operating in New Jersey under Business Id. No. 0100915422. On information and belief, Zydus USA is registered with the State of New Jersey's Department of Health as a wholesaler under Registration No. 5003171. On information and belief, Zydus USA purposefully has conducted and continues to conduct business in this Judicial District. On information and belief, Zydus USA is a corporation organized and existing under the laws of the State of New Jersey. By virtue of its incorporation in New Jersey, this Court has personal jurisdiction over Zydus USA.

19. On information and belief, Zydus USA is in the business of, among other things, manufacturing, marketing, importing, offering for sale, and selling pharmaceutical products, including generic drug products, throughout the United States, including in this Judicial District. This Judicial District is a likely destination for the generic drug product described in Zydus's ANDA. On information and belief, Zydus USA also prepares and/or aids in the preparation and submission of ANDAs to the FDA.

20. This Court has personal jurisdiction over Cadila Healthcare Limited and Zydus International Pvt. Ltd. because, *inter alia*, they: (1) have purposely availed themselves of the privilege of doing business in New Jersey, including directly or indirectly through their subsidiary, agent, and/or alter ego, Zydus Pharmaceuticals (USA) Inc., a company incorporated

in New Jersey, with its principal place of business in New Jersey; and (2) maintain extensive and systematic contacts with the State of New Jersey, including the marketing, distribution, and/or sale of generic pharmaceutical drugs in New Jersey including through, directly or indirectly, Zydus Pharmaceuticals (USA) Inc.

21. This Court has personal jurisdiction over Zydus because, *inter alia*, it has committed an act of patent infringement under 35 U.S.C. § 271(e)(2), and has sent notice of that infringement to Celgene in the State of New Jersey. On information and belief, Zydus intends a future course of conduct that includes acts of patent infringement in New Jersey. These acts have led and will continue to lead to foreseeable harm and injury to Celgene in New Jersey and in this Judicial District.

22. Zydus Cadila's Annual Report 2015-16 states that "US is the world's largest pharmaceutical market, both for branded and generic drugs, accounting for around one third of the global market," and that "[t]he Company is present in the [US] generic pharmaceuticals market through its wholly owned subsidiary, Zydus Pharmaceuticals (USA) Inc." Cadila Healthcare Limited Annual Report 2015-16 ("Zydus Cadila Annual Report") at 9. The Zydus Cadila Annual Report further states that "the Company is ranked amongst the top 10 generics companies in the US based on prescriptions." *Id.* The Zydus Cadila Annual Report further states that "[t]he Company launched 3 new products in the US market during the year," and that "[i]n terms of ANDA filings, 30 more ANDAs were filed with the USFDA during the year, taking the cumulative number of ANDA filings to 269." *Id.* at 10. The Zydus Cadila Annual Report further states that "[g]oing forward, the Company's focus will continue to be on launching complex, difficult-to-make oral solids and formulations of other dosage forms like

injectables, nasals, creams and ointments in order to enhance its share in the US generics market.” *Id.*

23. Zydus Pharmaceuticals (USA) Inc.’s website, <http://www.zydususa.com/who-is-zydus/>, states that Zydus Pharmaceuticals (USA) Inc. “is the U.S. division of Cadila Healthcare.”

24. On information and belief, Zydus USA, Zydus International, and/or Zydus Cadila work in concert either directly or indirectly through one or more of their wholly owned subsidiaries with respect to the regulatory approval, manufacturing, marketing, sale, and distribution of generic pharmaceutical products throughout the United States, including in this Judicial District.

25. On information and belief, Zydus USA acts at the direction, and for the benefit, of Zydus International and/or Zydus Cadila, and is controlled and/or dominated by Zydus International and/or Zydus Cadila.

26. On information and belief, Zydus Pharmaceuticals (USA) Inc. and Cadila Healthcare Limited have previously been sued in this Judicial District and have not challenged personal jurisdiction. *See, e.g., Helsinn Healthcare S.A. v. Zydus Pharms. (USA) Inc. and Cadila Healthcare Ltd. (d/b/a Zydus Cadila)*, No. 16-4239 (MLC)(DEA) (D.N.J.); *AstraZeneca AB, et al. v. Zydus Pharms. (USA) Inc. and Cadila Healthcare Ltd. (dba Zydus Cadila)*, No. 15-7415 (MLC)(TJB) (D.N.J.); *Supernus Pharms., Inc. v. Zydus Pharms. (USA) Inc. and Cadila Healthcare Ltd.*, No. 14-7272 (SDW)(LDW) (D.N.J.); *Otsuka Pharm. Co., Ltd. v. Zydus Pharms. (USA) Inc. and Cadila Healthcare Ltd.*, No. 14-7252 (JBS)(KMW) (D.N.J.); *Otsuka Pharm. Co., Ltd. v. Zydus Pharms. (USA) Inc. and Cadila Healthcare Ltd.*, No. 14-3168 (JBS)(KMW) (D.N.J.).

27. Zydus Pharmaceuticals (USA) Inc. has also admitted that it is subject to personal jurisdiction in this Judicial District. *See, e.g., Takeda Pharm. Co. Ltd., et al. v. Zydus Pharms. (USA) Inc. and Cadila Healthcare Ltd.*, No. 10-1723 (Answer to Complaint, Dkt. No. 29, ¶ 12; Answer to Amended Complaint, Dkt. No. 99, ¶ 12).

28. Cadila Healthcare Limited has also admitted that it is subject to personal jurisdiction in this Judicial District. *See, e.g., Takeda Pharm. Co. Ltd., et al. v. Zydus Pharms. (USA) Inc. and Cadila Healthcare Ltd.*, No. 10-1723 (Answer to Complaint, Dkt. No. 29, ¶¶ 13, 18; Answer to Amended Complaint, Dkt. No. 99, ¶¶ 13, 18).

29. Zydus Pharmaceuticals (USA) Inc. and Cadila Healthcare Limited have further availed themselves of the jurisdiction of this Court by previously initiating litigation in this Judicial District. *See, e.g., Zydus Pharms. (USA) Inc. and Cadila Healthcare Ltd. v. Gilead Scis., Inc.*, No. 14-7080 (FLW)(LHG) (D.N.J.). Zydus Pharmaceuticals (USA) Inc. has further availed itself of the jurisdiction of this Court by previously initiating litigation in this Judicial District. *See, e.g., Zydus Pharms. USA, Inc. v. Eli Lilly and Co.*, No. 10-5584 (DMC)(JAD) (D.N.J.).

30. Venue is proper in this Judicial District pursuant to 28 U.S.C. §§ 1391 and 1400(b).

Acts Giving Rise To This Suit

31. Pursuant to Section 505 of the FFDCA, Zydus filed Zydus's ANDA seeking approval to engage in the commercial manufacture, use, sale, offer for sale, or importation into the United States of lenalidomide capsules 2.5 mg, 5 mg, 10 mg, 15 mg, 20 mg, and 25 mg ("Zydus's Proposed Products"), before the patents-in-suit expire.

32. On information and belief, following FDA approval of Zydus's ANDA, Zydus USA, Zydus International, and/or Zydus Cadila will work in concert with one another to make,

use, sell, or offer to sell Zydus's Proposed Products throughout the United States, or import such generic products into the United States.

33. On information and belief, in connection with the filing of its ANDA as described above, Zydus provided a written certification to the FDA, as called for by Section 505 of the FDCA, 21 U.S.C. § 355(j)(2)(A)(vii)(IV) ("Zydus's Paragraph IV Certification"), alleging that the claims of the patents-in-suit are invalid, unenforceable, and/or will not be infringed by the activities described in Zydus's ANDA.

34. No earlier than February 27, 2017, Zydus sent written notice of its Paragraph IV Certification to Celgene ("Zydus's Notice Letter"). Zydus's Notice Letter alleged that the claims of the patents-in-suit are invalid and/or will not be infringed by the activities described in Zydus's ANDA. Zydus's Notice Letter also informed Celgene that Zydus seeks approval to market Zydus's Proposed Products before the patents-in-suit expire. Zydus specifically directed Zydus's Notice Letter to Celgene's headquarters in Summit, New Jersey, in this Judicial District.

Count I: Infringement of the '800 Patent

35. Celgene repeats and realleges the allegations of the preceding paragraphs as if fully set forth herein.

36. Zydus's submission of its ANDA to engage in the commercial manufacture, use, sale, offer for sale, or importation into the United States of Zydus's Proposed Products, prior to the expiration of the '800 patent, constitutes infringement of one or more of the claims of that patent under 35 U.S.C. § 271(e)(2)(A).

37. There is a justiciable controversy between the parties hereto as to the infringement of the '800 patent.

38. Unless enjoined by this Court, upon FDA approval of Zydus's ANDA, Zydus will infringe one or more claims of the '800 patent under 35 U.S.C. § 271(a) by making, using, offering to sell, selling, and/or importing Zydus's Proposed Products in the United States.

39. Unless enjoined by this Court, upon FDA approval of Zydus's ANDA, Zydus will induce infringement of one or more claims of the '800 patent under 35 U.S.C. § 271(b) by making, using, offering to sell, selling, and/or importing Zydus's Proposed Products in the United States. On information and belief, upon FDA approval of Zydus's ANDA, Zydus will intentionally encourage acts of direct infringement with knowledge of the '800 patent and knowledge that its acts are encouraging infringement.

40. Unless enjoined by this Court, upon FDA approval of Zydus's ANDA, Zydus will contributorily infringe one or more claims of the '800 patent under 35 U.S.C. § 271(c) by making, using, offering to sell, selling, and/or importing Zydus's Proposed Products in the United States. On information and belief, Zydus has had and continues to have knowledge that Zydus's Proposed Products are especially adapted for a use that infringes one or more claims of the '800 patent and that there is no substantial non-infringing use for Zydus's Proposed Products.

41. Celgene will be substantially and irreparably damaged and harmed if Zydus's infringement of the '800 patent is not enjoined.

42. Celgene does not have an adequate remedy at law.

43. This case is an exceptional one, and Celgene is entitled to an award of its reasonable attorneys' fees under 35 U.S.C. § 285.

Count II: Infringement of the '217 Patent

44. Celgene repeats and realleges the allegations of the preceding paragraphs as if fully set forth herein.

45. Zydus's submission of its ANDA to engage in the commercial manufacture, use, sale, offer for sale, or importation into the United States of Zydus's Proposed Products, prior to the expiration of the '217 patent, constitutes infringement of one or more of the claims of that patent under 35 U.S.C. § 271(e)(2)(A).

46. There is a justiciable controversy between the parties hereto as to the infringement of the '217 patent.

47. Unless enjoined by this Court, upon FDA approval of Zydus's ANDA, Zydus will infringe one or more claims of the '217 patent under 35 U.S.C. § 271(a) by making, using, offering to sell, selling, and/or importing Zydus's Proposed Products in the United States.

48. Unless enjoined by this Court, upon FDA approval of Zydus's ANDA, Zydus will induce infringement of one or more claims of the '217 patent under 35 U.S.C. § 271(b) by making, using, offering to sell, selling, and/or importing Zydus's Proposed Products in the United States. On information and belief, upon FDA approval of Zydus's ANDA, Zydus will intentionally encourage acts of direct infringement with knowledge of the '217 patent and knowledge that its acts are encouraging infringement.

49. Unless enjoined by this Court, upon FDA approval of Zydus's ANDA, Zydus will contributorily infringe one or more claims of the '217 patent under 35 U.S.C. § 271(c) by making, using, offering to sell, selling, and/or importing Zydus's Proposed Products in the United States. On information and belief, Zydus has had and continues to have knowledge that Zydus's Proposed Products are especially adapted for a use that infringes one or more claims of the '217 patent and that there is no substantial non-infringing use for Zydus's Proposed Products.

50. Celgene will be substantially and irreparably damaged and harmed if Zydus's infringement of the '217 patent is not enjoined.

51. Celgene does not have an adequate remedy at law.

52. This case is an exceptional one, and Celgene is entitled to an award of its reasonable attorneys' fees under 35 U.S.C. § 285.

Count III: Infringement of the '569 Patent

53. Celgene repeats and realleges the allegations of the preceding paragraphs as if fully set forth herein.

54. Zydus's submission of its ANDA to engage in the commercial manufacture, use, sale, offer for sale, or importation into the United States of Zydus's Proposed Products, prior to the expiration of the '569 patent, constitutes infringement of one or more of the claims of that patent under 35 U.S.C. § 271(e)(2)(A).

55. There is a justiciable controversy between the parties hereto as to the infringement of the '569 patent.

56. Unless enjoined by this Court, upon FDA approval of Zydus's ANDA, Zydus will infringe one or more claims of the '569 patent under 35 U.S.C. § 271(a) by making, using, offering to sell, selling, and/or importing Zydus's Proposed Products in the United States.

57. Unless enjoined by this Court, upon FDA approval of Zydus's ANDA, Zydus will induce infringement of one or more claims of the '569 patent under 35 U.S.C. § 271(b) by making, using, offering to sell, selling, and/or importing Zydus's Proposed Products in the United States. On information and belief, upon FDA approval of Zydus's ANDA, Zydus will intentionally encourage acts of direct infringement with knowledge of the '569 patent and knowledge that its acts are encouraging infringement.

58. Unless enjoined by this Court, upon FDA approval of Zydus's ANDA, Zydus will contributorily infringe one or more claims of the '569 patent under 35 U.S.C. § 271(c) by making, using, offering to sell, selling, and/or importing Zydus's Proposed Products in the

United States. On information and belief, Zydus has had and continues to have knowledge that Zydus's Proposed Products are especially adapted for a use that infringes one or more claims of the '569 patent and that there is no substantial non-infringing use for Zydus's Proposed Products.

59. Celgene will be substantially and irreparably damaged and harmed if Zydus's infringement of the '569 patent is not enjoined.

60. Celgene does not have an adequate remedy at law.

61. This case is an exceptional one, and Celgene is entitled to an award of its reasonable attorneys' fees under 35 U.S.C. § 285.

Count IV: Infringement of the '498 Patent

62. Celgene repeats and realleges the allegations of the preceding paragraphs as if fully set forth herein.

63. Zydus's submission of its ANDA to engage in the commercial manufacture, use, sale, offer for sale, or importation into the United States of Zydus's Proposed Products, prior to the expiration of the '498 patent, constitutes infringement of one or more of the claims of that patent under 35 U.S.C. § 271(e)(2)(A).

64. There is a justiciable controversy between the parties hereto as to the infringement of the '498 patent.

65. Unless enjoined by this Court, upon FDA approval of Zydus's ANDA, Zydus will infringe one or more claims of the '498 patent under 35 U.S.C. § 271(a) by making, using, offering to sell, selling, and/or importing Zydus's Proposed Products in the United States.

66. Unless enjoined by this Court, upon FDA approval of Zydus's ANDA, Zydus will induce infringement of one or more claims of the '498 patent under 35 U.S.C. § 271(b) by making, using, offering to sell, selling, and/or importing Zydus's Proposed Products in the United States. On information and belief, upon FDA approval of Zydus's ANDA, Zydus will

intentionally encourage acts of direct infringement with knowledge of the '498 patent and knowledge that its acts are encouraging infringement.

67. Unless enjoined by this Court, upon FDA approval of Zydus's ANDA, Zydus will contributorily infringe one or more claims of the '498 patent under 35 U.S.C. § 271(c) by making, using, offering to sell, selling, and/or importing Zydus's Proposed Products in the United States. On information and belief, Zydus has had and continues to have knowledge that Zydus's Proposed Products are especially adapted for a use that infringes one or more claims of the '498 patent and that there is no substantial non-infringing use for Zydus's Proposed Products.

68. Celgene will be substantially and irreparably damaged and harmed if Zydus's infringement of the '498 patent is not enjoined.

69. Celgene does not have an adequate remedy at law.

70. This case is an exceptional one, and Celgene is entitled to an award of its reasonable attorneys' fees under 35 U.S.C. § 285.

Count V: Infringement of the '095 Patent

71. Celgene repeats and realleges the allegations of the preceding paragraphs as if fully set forth herein.

72. Zydus's submission of its ANDA to engage in the commercial manufacture, use, sale, offer for sale, or importation into the United States of Zydus's Proposed Products, prior to the expiration of the '095 patent, constitutes infringement of one or more of the claims of that patent under 35 U.S.C. § 271(e)(2)(A).

73. There is a justiciable controversy between the parties hereto as to the infringement of the '095 patent.

74. Unless enjoined by this Court, upon FDA approval of Zydus's ANDA, Zydus will infringe one or more claims of the '095 patent under 35 U.S.C. § 271(a) by making, using, offering to sell, selling, and/or importing Zydus's Proposed Products in the United States.

75. Unless enjoined by this Court, upon FDA approval of Zydus's ANDA, Zydus will induce infringement of one or more claims of the '095 patent under 35 U.S.C. § 271(b) by making, using, offering to sell, selling, and/or importing Zydus's Proposed Products in the United States. On information and belief, upon FDA approval of Zydus's ANDA, Zydus will intentionally encourage acts of direct infringement with knowledge of the '095 patent and knowledge that its acts are encouraging infringement.

76. Unless enjoined by this Court, upon FDA approval of Zydus's ANDA, Zydus will contributorily infringe one or more claims of the '095 patent under 35 U.S.C. § 271(c) by making, using, offering to sell, selling, and/or importing Zydus's Proposed Products in the United States. On information and belief, Zydus has had and continues to have knowledge that Zydus's Proposed Products are especially adapted for a use that infringes one or more claims of the '095 patent and that there is no substantial non-infringing use for Zydus's Proposed Products.

77. Celgene will be substantially and irreparably damaged and harmed if Zydus's infringement of the '095 patent is not enjoined.

78. Celgene does not have an adequate remedy at law.

79. This case is an exceptional one, and Celgene is entitled to an award of its reasonable attorneys' fees under 35 U.S.C. § 285.

Count VI: Infringement of the '621 Patent

80. Celgene repeats and realleges the allegations of the preceding paragraphs as if fully set forth herein.

81. Zydus's submission of its ANDA to engage in the commercial manufacture, use, sale, offer for sale, or importation into the United States of Zydus's Proposed Products, prior to the expiration of the '621 patent, constitutes infringement of one or more of the claims of that patent under 35 U.S.C. § 271(e)(2)(A).

82. There is a justiciable controversy between the parties hereto as to the infringement of the '621 patent.

83. Unless enjoined by this Court, upon FDA approval of Zydus's ANDA, Zydus will infringe one or more claims of the '621 patent under 35 U.S.C. § 271(a) by making, using, offering to sell, selling, and/or importing Zydus's Proposed Products in the United States.

84. Unless enjoined by this Court, upon FDA approval of Zydus's ANDA, Zydus will induce infringement of one or more claims of the '621 patent under 35 U.S.C. § 271(b) by making, using, offering to sell, selling, and/or importing Zydus's Proposed Products in the United States. On information and belief, upon FDA approval of Zydus's ANDA, Zydus will intentionally encourage acts of direct infringement with knowledge of the '621 patent and knowledge that its acts are encouraging infringement.

85. Unless enjoined by this Court, upon FDA approval of Zydus's ANDA, Zydus will contributorily infringe one or more claims of the '621 patent under 35 U.S.C. § 271(c) by making, using, offering to sell, selling, and/or importing Zydus's Proposed Products in the United States. On information and belief, Zydus has had and continues to have knowledge that Zydus's Proposed Products are especially adapted for a use that infringes one or more claims of the '621 patent and that there is no substantial non-infringing use for Zydus's Proposed Products.

86. Celgene will be substantially and irreparably damaged and harmed if Zydus's infringement of the '621 patent is not enjoined.

87. Celgene does not have an adequate remedy at law.

88. This case is an exceptional one, and Celgene is entitled to an award of its reasonable attorneys' fees under 35 U.S.C. § 285.

Count VII: Infringement of the '622 Patent

89. Celgene repeats and realleges the allegations of the preceding paragraphs as if fully set forth herein.

90. Zydus's submission of its ANDA to engage in the commercial manufacture, use, sale, offer for sale, or importation into the United States of Zydus's Proposed Products, prior to the expiration of the '622 patent, constitutes infringement of one or more of the claims of that patent under 35 U.S.C. § 271(e)(2)(A).

91. There is a justiciable controversy between the parties hereto as to the infringement of the '622 patent.

92. Unless enjoined by this Court, upon FDA approval of Zydus's ANDA, Zydus will infringe one or more claims of the '622 patent under 35 U.S.C. § 271(a) by making, using, offering to sell, selling, and/or importing Zydus's Proposed Products in the United States.

93. Unless enjoined by this Court, upon FDA approval of Zydus's ANDA, Zydus will induce infringement of one or more claims of the '622 patent under 35 U.S.C. § 271(b) by making, using, offering to sell, selling, and/or importing Zydus's Proposed Products in the United States. On information and belief, upon FDA approval of Zydus's ANDA, Zydus will intentionally encourage acts of direct infringement with knowledge of the '622 patent and knowledge that its acts are encouraging infringement.

94. Unless enjoined by this Court, upon FDA approval of Zydus's ANDA, Zydus will contributorily infringe one or more claims of the '622 patent under 35 U.S.C. § 271(c) by making, using, offering to sell, selling, and/or importing Zydus's Proposed Products in the

United States. On information and belief, Zydus has had and continues to have knowledge that Zydus's Proposed Products are especially adapted for a use that infringes one or more claims of the '622 patent and that there is no substantial non-infringing use for Zydus's Proposed Products.

95. Celgene will be substantially and irreparably damaged and harmed if Zydus's infringement of the '622 patent is not enjoined.

96. Celgene does not have an adequate remedy at law.

97. This case is an exceptional one, and Celgene is entitled to an award of its reasonable attorneys' fees under 35 U.S.C. § 285.

PRAYER FOR RELIEF

WHEREFORE, Plaintiff Celgene respectfully requests the following relief:

(A) A Judgment that Zydus has infringed the patents-in-suit by submitting ANDA No. 210154;

(B) A Judgment that Zydus has infringed, and that Zydus's making, using, selling, offering to sell, or importing Zydus's Proposed Products will infringe one or more claims of the patents-in-suit;

(C) An Order that the effective date of FDA approval of ANDA No. 210154 be a date which is not earlier than the later of the expiration of the patents-in-suit, or any later expiration of exclusivity to which Celgene is or becomes entitled;

(D) Preliminary and permanent injunctions enjoining Zydus and its officers, agents, attorneys and employees, and those acting in privity or concert with them, from making, using, selling, offering to sell, or importing Zydus's Proposed Products until after the expiration of the patents-in-suit, or any later expiration of exclusivity to which Celgene is or becomes entitled;

(E) A permanent injunction, pursuant to 35 U.S.C. § 271(e)(4)(B), restraining and enjoining Zydus, its officers, agents, attorneys and employees, and those acting in privity or

concert with them, from practicing any solid forms of lenalidomide, compositions, or methods as claimed in the patents-in-suit, or from actively inducing or contributing to the infringement of any claim of the patents-in-suit, until after the expiration of the patents-in-suit, or any later expiration of exclusivity to which Celgene is or becomes entitled;

(F) A Judgment that the commercial manufacture, use, importation into the United States, sale, and/or offer for sale of Zydus's Proposed Products will directly infringe, induce and/or contribute to infringement of the patents-in-suit;

(G) To the extent that Zydus has committed any acts with respect to the solid forms of lenalidomide, compositions, or methods claimed in the patents-in-suit, other than those acts expressly exempted by 35 U.S.C. § 271(e)(1), a Judgment awarding Celgene damages for such acts;

(H) If Zydus engages in the commercial manufacture, use, importation into the United States, sale, and/or offer for sale of Zydus's Proposed Products prior to the expiration of the patents-in-suit, a Judgment awarding damages to Celgene resulting from such infringement, together with interest;

(I) A Judgment declaring that the patents-in-suit remain valid and enforceable;

(J) A Judgment that this is an exceptional case pursuant to 35 U.S.C. § 285 and awarding Celgene its attorneys' fees incurred in this action;

(K) A Judgment awarding Celgene its costs and expenses incurred in this action; and

(L) Such further and other relief as this Court may deem just and proper.

Dated: April 12, 2017

By: s/ Charles M. Lizza

Of Counsel:

F. Dominic Cerrito
Eric C. Stops
Evangeline Shih
Andrew S. Chalson
Gabriel P. Brier
Frank C. Calvosa
Marta A. Godecki
Brian J. Forsatz
QUINN EMANUEL URQUHART & SULLIVAN, LLP
51 Madison Avenue, 22nd Floor
New York, New York 10010
(212) 849-7000

Anthony M. Insogna
J. Patrick Elsevier, Ph.D.
Steven J. Corr
Cary Miller, Ph.D.
JONES DAY
4655 Executive Drive
San Diego, CA 92121
(858) 314-1200

Charles M. Lizza
William C. Baton
SAUL EWING LLP
One Riverfront Plaza, Suite 1520
Newark, New Jersey 07102-5426
(973) 286-6700
clizza@saul.com

*Attorneys for Plaintiff
Celgene Corporation*

CERTIFICATION PURSUANT TO LOCAL CIVIL RULES 11.2 & 40.1

Pursuant to Local Civil Rules 11.2 and 40.1, I hereby certify that the matter captioned *Celgene Corporation v. Dr. Reddy's Laboratories, Ltd., et al.*, Civil Action No. 16-7704 (SDW)(LDW) (D.N.J.) is related to the matter in controversy because the matter in controversy involves the same plaintiff and the same patents, and because Defendants are seeking FDA approval to market generic versions of the same pharmaceutical product.

I further certify that, to the best of my knowledge, the matter in controversy is not the subject of any other action pending in any court or of any pending arbitration or administrative proceeding.

Dated: April 12, 2017

Of Counsel:

F. Dominic Cerrito
Eric C. Stops
Evangeline Shih
Andrew S. Chalson
Gabriel P. Brier
Frank C. Calvosa
Marta A. Godecki
Brian J. Forsatz
QUINN EMANUEL URQUHART & SULLIVAN, LLP
51 Madison Avenue, 22nd Floor
New York, New York 10010
(212) 849-7000

Anthony M. Insogna
J. Patrick Elsevier, Ph.D.
Steven J. Corr
Cary Miller, Ph.D.
JONES DAY
4655 Executive Drive
San Diego, CA 92121
(858) 314-1200

By: s/ Charles M. Lizza

Charles M. Lizza
William C. Baton
SAUL EWING LLP
One Riverfront Plaza, Suite 1520
Newark, New Jersey 07102-5426
(973) 286-6700
clizza@saul.com

*Attorneys for Plaintiff
Celgene Corporation*

EXHIBIT A



US007465800B2

(12) **United States Patent**
Jaworsky et al.

(10) **Patent No.:** **US 7,465,800 B2**
(45) **Date of Patent:** **Dec. 16, 2008**

(54) **POLYMORPHIC FORMS OF 3-(4-AMINO-1-OXO-1,3 DIHYDRO-ISOINDOL-2-YL)-PIPERIDINE-2,6-DIONE**

| | | |
|-------------|---------|-------------------|
| 5,059,595 A | 10/1991 | Le Grazie |
| 5,073,543 A | 12/1991 | Marshall et al. |
| 5,120,548 A | 6/1992 | McClelland et al. |
| 5,229,496 A | 7/1993 | Deeley et al. |
| 5,354,556 A | 10/1994 | Sparks et al. |
| 5,385,901 A | 1/1995 | Kaplan et al. |
| 5,391,485 A | 2/1995 | Deeley et al. |
| 5,393,870 A | 2/1995 | Deeley et al. |
| 5,528,823 A | 6/1996 | Rudy, Jr. et al. |
| 5,580,755 A | 12/1996 | Souza |
| 5,591,767 A | 1/1997 | Mohr et al. |
| 5,593,990 A | 1/1997 | D'Amato |
| 5,629,327 A | 5/1997 | D'Amato |
| 5,635,517 A | 6/1997 | Muller et al. |

(75) Inventors: **Markian S. Jaworsky**, Hopewell, NJ (US); **Roger Shen-Chu Chen**, Edison, NJ (US); **George W. Muller**, Bridgewater, NJ (US)

(73) Assignee: **Celgene Corporation**, Summit, NJ (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 596 days.

(21) Appl. No.: **10/934,863**

(22) Filed: **Sep. 3, 2004**

(Continued)

(65) **Prior Publication Data**

US 2005/0096351 A1 May 5, 2005

FOREIGN PATENT DOCUMENTS

WO WO 98/03502 1/1998

Related U.S. Application Data

(60) Provisional application No. 60/499,723, filed on Sep. 4, 2003.

(Continued)

OTHER PUBLICATIONS

(51) **Int. Cl.**
C07D 401/04 (2006.01)

Seddon "Pseudopolymorph: a polemic" *Crystal Growth & design* v.4(6) 1087, Web release Oct. 19, 2004.*

(52) **U.S. Cl.** **546/200; 514/323**

(Continued)

(58) **Field of Classification Search** **546/200; 514/323**

See application file for complete search history.

Primary Examiner—Celia Chang
(74) *Attorney, Agent, or Firm*—Jones Day

(56) **References Cited**


(57) **ABSTRACT**

U.S. PATENT DOCUMENTS

| | | |
|-------------|---------|------------------|
| 3,536,809 A | 10/1970 | Applezweig |
| 3,598,123 A | 8/1971 | Zaffaroni et al. |
| 3,845,770 A | 11/1974 | Theeuwes et al. |
| 3,916,899 A | 11/1975 | Theeuwes et al. |
| 4,008,719 A | 2/1977 | Theeuwes et al. |
| 4,810,643 A | 3/1989 | Souza |
| 4,999,291 A | 3/1991 | Souza |

Polymorphic forms of 3-(4-amino-1-oxo-1,3 dihydro-isoin-dol-2-yl)-piperidine-2,6-dione are disclosed. Compositions comprising the polymorphic forms, methods of making the polymorphic forms and methods of their use are also dis-closed.

14 Claims, 48 Drawing Sheets



US 7,465,800 B2

Page 2

U.S. PATENT DOCUMENTS

5,639,476 A 6/1997 Oshlack et al.
 5,674,533 A 10/1997 Santus et al.
 5,698,579 A 12/1997 Muller
 5,712,291 A 1/1998 D'Amato
 5,731,325 A 3/1998 Andrulis, Jr. et al.
 5,733,566 A 3/1998 Lewis
 5,798,368 A 8/1998 Muller et al.
 5,874,448 A 2/1999 Muller et al.
 5,877,200 A 3/1999 Muller
 5,929,117 A 7/1999 Muller et al.
 5,955,476 A 9/1999 Muller et al.
 6,020,358 A 2/2000 Muller et al.
 6,071,948 A 6/2000 D'Amato
 6,114,355 A 9/2000 D'Amato
 6,140,346 A 10/2000 Andrulis, Jr. et al.
 6,235,756 B1 5/2001 D'Amato
 6,281,230 B1 8/2001 Muller et al.
 6,316,471 B1 11/2001 Muller et al.
 6,326,388 B1 12/2001 Man et al.
 6,335,349 B1 1/2002 Muller et al.
 6,380,239 B1 4/2002 Muller et al.
 6,395,754 B1 5/2002 Muller et al.
 6,403,613 B1 6/2002 Man et al.
 6,420,414 B1 7/2002 D'Amato
 6,458,810 B1 10/2002 Muller et al.
 6,469,045 B1 10/2002 D'Amato
 6,476,052 B1 11/2002 Muller et al.
 6,518,298 B2 2/2003 Green et al.
 6,555,554 B2 4/2003 Muller et al.
 2001/0018445 A1 8/2001 Huang et al.
 2001/0056114 A1 12/2001 D'Amato
 2002/0035090 A1 3/2002 Zeldis et al.
 2002/0045643 A1 4/2002 Muller et al.
 2002/0052398 A1 5/2002 D'Amato
 2002/0054899 A1 5/2002 Zeldis
 2002/0061923 A1 5/2002 D'Amato
 2002/0128228 A1 9/2002 Hwu
 2002/0161023 A1 10/2002 D'Amato
 2002/0173658 A1 11/2002 Muller et al.
 2002/0183360 A1 12/2002 Muller et al.
 2003/0013739 A1 1/2003 Masferrer et al.
 2003/0028028 A1 2/2003 Man et al.
 2003/0045552 A1 3/2003 Robarge et al.
 2003/0069428 A1 4/2003 Muller et al.
 2003/0096841 A1 5/2003 Robarge et al.
 2003/0139451 A1 7/2003 Shah et al.
 2003/0144325 A1 7/2003 Muller et al.
 2003/0181428 A1 9/2003 Green et al.
 2003/0187024 A1 10/2003 D'Amato
 2003/0191098 A1 10/2003 D'Amato
 2003/0235909 A1 12/2003 Hariri et al.
 2004/0029832 A1 2/2004 Zeldis
 2004/0077685 A1 4/2004 Figg et al.
 2004/0077686 A1 4/2004 Dannenberg et al.
 2004/0087546 A1 5/2004 Zeldis
 2004/0091455 A1 5/2004 Zeldis
 2004/0122052 A1 6/2004 Muller et al.
 2004/0220144 A1 11/2004 Zeldis
 2004/0266809 A1 12/2004 Emanuel et al.

FOREIGN PATENT DOCUMENTS

WO WO 98/54170 12/1998
 WO WO 01/70275 9/2001
 WO WO 01/87307 11/2001
 WO WO 02/059106 8/2002
 WO WO 02/064083 8/2002
 WO PCT/US03/11578 1/2003

OTHER PUBLICATIONS

Chen et al. "Polymorphic forms of . . ." CA 142:303619 (2005).*

Haleblian, J.; McCrone, W.; *Pharmaceutical Applications of Polymorphism*, Journal of Pharmaceutical Sciences, vol. 58., No. 8, Aug. 1969.

U.S. Appl. No. 60/372,348, Hariri et al.

U.S. Appl. No. 10/732,867, D'Amato et al.

U.S. Appl. No. 09/545,654, D'Amato.

U.S. Appl. No. 09/287,377, D'Amato.

Anderson, "Moving disease biology from the laboratory to the clinic." *Seminars in Oncology*, 2002 29:17-20.

Barlogie et al., "High-dose therapy immunomodulatory drugs in multiple myeloma," *Seminars in Oncology*, 2002, 29 (6):26-33.

Barlogie et al., "Introduction: Thalidomide and the IMiDs in multiple myeloma," *Seminars in Hematology*, 2003, 40 (4):1-2.

Barlogie et al., "Total Therapy II (TII) for newly diagnosed multiple myeloma (MM): preliminary data on feasibility and efficacy in the first 231 enrolled patients; comparison with predecessor trial total therapy I (TTI) (N=231)," *Blood*, Abstract # 2857, Dec. 7-11, 2001, *American Society of Hematology*.

Barlogie, "Thalidomide and CC-5013 in Multiple Myeloma: The University of Arkansas experience," *Seminars in Hematology*, 2003, 40 (4):33-38.

Bartlett et al., "Phase I study to determine the safety, tolerability and immunostimulatory activity of thalidomide analogue CC-5013 in patients with metastatic malignant melanoma and other advanced cancers," *British Journal of Cancer*, 2004, 90:955-961.

Bartlett et al., "The evolution of thalidomide and its IMiD derivatives as anticancer agents," *Nature Reviews Cancer*, 2004, 4 (4):1-9.

Battegay, "Angiogenesis: mechanistic insights, neovascular diseases, and therapeutic prospects," *J. Mol. Med.*, 1995, 73:333-346.

Baz et al., "Doxil (D), vincristine (V), reduced frequency dexamethasone (d) and revlimid (R) (DVD-R) results in a high response rate in patients with refractory multiple myeloma (RMM)," *Blood*, Abstract # 2559, *American Society of Hematology*, Dec. 10-13, 2005.

Brennen et al., "Thalidomide and analogues: current proposed mechanisms and therapeutic usage," *Clinical Prostate Cancer*, 2004, 3 (1):54-61.

Celgene Corporation, "Additional clinical data presented on Revimid™ in myelodysplastic syndromes at the American Society of Hematology 45th annual meeting," Press Release, Dec. 2003.

Celgene Corporation, "Blood reports Revimid™ has anti-tumor activity in patients with relapsed and refractory multiple myeloma," Press Release, Nov. 1, 2002.

Celgene Corporation, "Celgene advances immunomodulatory drug (IMiD™) clinical program," Press Release, Feb. 2000.

Celgene Corporation, "Celgene announces plans to stop phase III trials in melanoma due to lack of efficacy," Press Release, Apr. 2004.

Celgene Corporation, "Celgene corporation advances ACTIMID™ (CC-4047) into phase II trial for prostate cancer," Press Release, Oct. 2003.

Celgene Corporation, "Celgene Corporation announces fourth quarter and full year results for 2002," Press Release, Jan. 2003.

Celgene Corporation, "Celgene Corporation announces third quarter results. THALOMID® (thalidomide) revenue increases 41% to \$30.5 million. Pivotal programs for THALOMID and REVIMID™ finalized. Peer-reviewed publications of THALOMID and REVIMID data. First JNK inhibitor advanced to Phase I clinical trial," Press Release, Oct. 2002.

Celgene Corporation, "Celgene Corporation announces third quarter results. Thalomid® (thalidomide) sales increase 24%. Prescriptions up 50%. Enhanced S.T.E.P.S.® launched. Pilot d-MPH data presented," Press Release, Oct. 2001.

Celgene Corporation, "Celgene Corporation receives orphan drug designation for Revimid™ for multiple myeloma," Press Release, Oct. 2001.

Celgene Corporation, "Celgene corporation reports record operating performance in first quarter with strong revenue growth and profits," Press Release, Apr. 2004.

Celgene Corporation, "Celgene corporation reports record operating performance in third quarter as total revenue increases 117% and profits rise," Press Release, Oct. 2003.

US 7,465,800 B2

Page 3

- Celgene Corporation, "Celgene corporation reports strong operating performance in second quarter as total sales increase 100 percent and profits rise," Press Release, Jul. 2003.
- Celgene Corporation, "Celgene corporation reviews 2003 achievements and announces 2004 financial outlook," Press Release, Jan. 2004.
- Celgene Corporation, "Celgene expands clinical development program for Revimid". Five additional trials of Revimid initiated in hematological and solid tumors cancers," Press Release, Jun. 2002.
- Celgene Corporation, "Celgene provides update on clinical pipeline. Celgene Announces first target indication for ACTIMID™, CC-8490. SelCID™ program to advance based on results from Phase I/II trial of CC-1088. First JNK inhibitor successfully completes phase I trial," Press Release, Jan. 2003.
- Celgene Corporation, "Celgene receives fast track status from FDA for Revimid™ in multiple myeloma," Press Release, Feb. 2003.
- Celgene Corporation, "Celgene receives fast track status from FDA for Revimid™ in myelodysplastic syndromes," Press Release, Apr. 2003.
- Celgene Corporation, "Initial Phase I solid tumor data on Celgene's lead IMiD™, Revimid™," Press Release, Jun. 2001.
- Celgene Corporation, "New Revimid™ clinical data shows potential as novel approach to treating myelodysplastic syndromes (MDS)," Press Release, May 2003.
- Celgene Corporation, "Revimid™ receives orphan drug designation from the European commission for multiple myeloma," Press Release, Feb. 2004.
- Celgene Corporation, "Revimid™ receives orphan drug designation from the European commission for myelodysplastic syndromes," Press Release, Mar. 2004.
- Chandhry, 1966, *Cancer Research*, "Effect of Prednisolone and Thalidomide on Induced Submandibular Gland Tumors in Hamster," 26(part 1)1884-86.
- Corral et al., 1999, "Differential cytokine modulation and T cell activation by two distinct classes of thalidomide analogues that are potent inhibitors of TNF-alpha," *J. Immunol.* 163(1):380-386.
- Craig et al., 1967, "Potential anticancer agents. III. 2-phthalimidoaldehydes and derivatives," *Potential Anticancer Agents III* 10:1071-1073.
- D'Amato et al., 2001, "Mechanism of action of thalidomide and 3-aminothalidomide in multiple myeloma," *Semin. Oncol.* 28:597-601.
- Dalgleish et al., "Thalidomide analogues CC-5013 and CC-4047 induce T cell activation and IL-12 production in patients with both solid tumours and relapsed and refractory multiple myeloma," *British Journal of Cancer*, 2003, 88(Suppl 1), S25-S54.
- Dalgleish, et al., "New thalidomide analogues; anti-cancer, anti-angiogenic and immunostimulatory," *British Journal of Cancer*, 2001, 85 (1)25.
- Davies et al., "Thalidomide (Thal) and immunomodulatory derivatives (IMiDs) augment natural killer (NK) cell cytotoxicity in multiple myeloma-MM," Abstract # P222, *VIIIth International Myeloma Workshop*, May 4-8, 2001.
- Davies et al., "Thalidomide (Thal) and immunomodulatory derivatives (IMiDs) augment natural killer (NK) cell cytotoxicity in multiple myeloma (MM)," Abstract # 3617, *American Society of Hematology*, Dec. 1-5, 2000.
- Davies et al., 2001, "Thalidomide and immunomodulatory derivatives augment natural killer cell cytotoxicity in multiple myeloma," *Blood* 98(1):210-216.
- Dibbs et al., "Thalidomide and thalidomide analogs suppress TNF α secretion by myocytes," Abstract # 1284, *Circulation*, 1998.
- Dimopoulos et al., "Results of thalidomide and IMiDs in multiple myeloma," Abstract # P12.1.4, *International Multiple Myeloma Workshop*, May 23-27, 2003.
- Dimopoulos et al., "Study of lenalidomide plus dexamethasone versus dexamethasone alone in relapsed or refractory multiple myeloma (MM): Results of a phase 3 Study (MM-010)," Abstract # 6, *American Society of Hematology*, Dec. 10-13, 2005.
- Dimopoulos et al., "Treatment of plasma cell dyscrasias with thalidomide and its derivatives," *Journal of Clinical Oncology*, Dec. 1, 2003, 21 (23)4444-4454.
- Dimopoulos et al., 2004, "Primary treatment with pulsed melphalan, dexamethasone, thalidomide (MDT) for symptomatic patients with multiple myeloma \geq 75 years of age," *Am. Soc. Hematol.* 46th Ann. Meeting Dec. 4-7, 2004, San Diego, CA Abstract #1482.
- Dredge et al., "A costimulatory thalidomide analog enhances the partial anti-tumor immunity of an autologous vaccination in a model of colorectal cancer, Abstract # 491, *American Association for Cancer Research*, Apr. 6-10, 2002.
- Dredge et al., "Adjuvants and the promotion of Th1-type cytokines in tumour immunotherapy," *Cancer Immunol. Immunother.*, 2002, 51:521-531.
- Dredge et al., "Angiogenesis inhibitors in cancer therapy," *Current Opinion in Investigational Drugs*, 2003, 4 (6):667-674.
- Dredge et al., "Immunological effects of thalidomide and its chemical and functional analogs," *Critical Reviews in Immunology*, 2002, 22 (5&6):425-437.
- Dredge et al., "Protective antitumor immunity induced by a costimulatory thalidomide analog in conjunction with whole tumor cell vaccination is mediated by increased Th1-type immunity¹," *The Journal of Immunology*, 2002, 168:4914-4919.
- Dredge et al., "Recent developments in antiangiogenic therapy," *Expert Opin. Biol. Ther.*, 2002, 2 (8):953-966.
- Dredge et al., "Thalidomide analogs as emerging anti-cancer drugs," *Anti-Cancer Drugs*, 2003, 14:331-335.
- Dredge et al., 2002, "Novel thalidomide analogues display anti-angiogenic activity independently of immunomodulatory effects," *Br. J. Cancer* 87(10):1166-1172.
- Fickentscher et al., "Stereochemical properties and teratogenic activity of some tetrahydrophthalimides," *Molecular Pharmacology*, 1976, 13:133-141.
- Galustian et al., "Thalidomide-derived immunomodulatory drugs as therapeutic agents," *Expert Opin. Biol. Ther.*, 2004, 4 (12):1-8.
- Glaspy et al., "The potential role of thalidomide and thalidomide analogs in melanoma," *Clinical Advances in Hematology & Oncology*, 2004, 1-7.
- Gupta et al., "Adherence of multiple myeloma cells to bone marrow stromal cells upregulates vascular endothelial growth factor secretion: therapeutic applications," *Leukemia*, 2001, 15:1950-1961.
- Haslett et al., 2003, "Thalidomide and a thalidomide analogue drug costimulate virus-specific CD8+ T cells in vitro," *J. Infect. Dis.* 187(6):946-955.
- Hayashi et al., "Mechanisms whereby immunomodulatory analogs of thalidomide augment autologous NK cell anti-myeloma immunity," *Blood*, Abstract #3219, Dec. 6-10, 2002, *American Society of Hematology*.
- He, W., et al., 1993, Abstract of papers, 206th American Chemical Society, Chicago, IL: Med. Chem., paper 216.
- Helm et al., "Comparative teratological investigation of compounds of structurally and pharmacologically related to thalidomide," *Arzneimittel Forschung/Drug Research*, 1981, 31 (I)941-949.
- Hernandez-Illizaliturri et al., "Addition of immunomodulatory drugs CC5013 or CC4047 to rituximab enhances anti-tumor activity in a severe combined immunodeficiency (SCID) mouse lymphoma model," Abstract # 235, *American Society of Hematology*, Dec. 6-9, 2003.
- Hideshima et al., "Thalidomide (Thal) and its analogs overcome drug resistance of human multiple myeloma (MM) cells to conventional therapy," Abstract 1313, *American Society of Hematology*, Dec. 1-5, 2000.
- Hideshima et al., "Thalidomide and its analogs overcome drug resistance of human multiple myeloma cells to conventional therapy," *Blood*, 2000, 96:2943-2950, *American Society of Hematology*.
- Hunt et al., "Markers of endothelial and haemostatic activation in the use of CC-4047, a structural analogue of thalidomide, in relapsed myeloma," *Blood*, Abstract # 3216, Dec. 6-10, 2002, *American Society of Hematology*.
- Hussein et al., "Doxil (D), vincristine (V), reduced frequency dexamethasone (d) and Revlimid (Dvd-R) a phase I/II trial in advanced relapsed/refractory multiple myeloma (Rmm) patients," *Blood*, Abstract #208, *American Society of Hematology*, Dec. 4-7, 2004.
- Kyle et al., "Multiple myeloma," *New England Journal of Medicine*, 2004, 351:1860-1873.

US 7,465,800 B2

Page 4

- Kyle, "Current therapy of multiple myeloma," *Internal Medicine*, 2002, 41 (3)175-180.
- Leblanc et al., "Immunomodulatory drug costimulates T cells via the B7-CD28 pathway," *Blood*, 2004, 103:1787-1790, *American Society of Hematology*.
- Lentzsch et al., "Immunomodulatory derivative of thalidomide (IMiD CC-4047) determine the lineage commitment of hematopoietic progenitors by down regulation of GATA-1 and modulation of cytokine secretion," Abstract # 3073, *American Society of Hematology*, Dec. 6-9, 2003.
- Lentzsch et al., "Immunomodulatory derivative of thalidomide (IMiD CC-4047) down regulates CAAT/enhancer-binding protein β (C/EBP β) in multiple myeloma (MM)," Abstract # 3456, *American Society of Hematology*, Dec. 6-9, 2003.
- Lentzsch et al., "In vivo activity of thalidomide and immunomodulatory drugs against multiple myeloma," *VIIIth International Myeloma Workshop*, Abstract #P225, May 4-8, 2001.
- Lentzsch et al., 2003, "Immunomodulatory analogs of thalidomide inhibit growth of Hs Sultan cells and angiogenesis in vivo," *Leukemia* 17(1):41-44.
- Liu et al., "Phase I study of CC-5013 (Revimid), a thalidomide derivative, in patients with refractory metastatic cancer," *American Society of Clinical Oncology*, Abstract #927, 2003.
- Luzzio et al., "Thalidomide analogues: derivatives of an orphan drug with diverse biological activity," *Expert Opin. Ther. Patents*, 2004, 14 (2):215-229.
- Man et al., " α - Fluoro-substituted thalidomide analogues," *Bioorganic & Medicinal Chemistry Letters* 13, 2003, 3415-3417.
- Marriott et al., "A novel subclass of thalidomide analogue with anti-solid tumor activity in which caspase-dependent apoptosis is associated with altered expression of bcl-2 family proteins," *Cancer Research*, 2003, 63:593-599.
- Marriott et al., "Immunotherapeutic and antitumor potential of thalidomide analogues," *Expert Opin. Biol. Ther.*, 2001, 1 (4):1-8.
- Marriott et al., "New thalidomide analogues; anti-cancer, anti-angiogenic and immunostimulatory," *British Journal of Cancer*, 85:25, Jul. 6, 2001.
- Marriott et al., "Thalidomide and its analogues have distinct and opposing effects on TNF- α and TNFR2 during co-stimulation of both CD4⁺ and CD8⁺ T cells," *Clin. Exp. Immunol.*, 2002, 130:75-84.
- Marriott et al., "Thalidomide derived immunomodulatory drugs (IMiDs) as potential therapeutic agents," *Current Drug Targets—Immune, Endocrine & Metabolic Disorders*, 2003, 3:181-186.
- Masellis et al., "Changes in gene expression in bone marrow mesenchymal progenitor cells as a consequence of IMiD therapy in multiple myeloma patients," *Blood*, Abstract # 1548, Dec. 7-11, 2001, *American Society of Hematology*.
- McCarty, "Thalidomide may impede cell migration in primates by down-regulating integrin β -chains: potential therapeutic utility in solid malignancies, proliferative retinopathy, inflammatory disorders, neointimal hyperplasia, and osteoporosis," *Medical Hypotheses*, 1997, 49:123-131.
- Mitsiades et al., "Apoptotic signaling induced by immunomodulatory thalidomide analogs (ImiDs) in human multiple myeloma cells: therapeutic implications," Abstract # 3224, Dec. 7-11, 2001, *American Society of Hematology*.
- Mitsiades et al., "Apoptotic signaling induced by immunomodulatory thalidomide analogs in human multiple myeloma cells: therapeutic implications," *Blood*, 2002, 99:4525-4530, *American Society of Hematology*.
- Mitsiades et al., "CC-5013 Celgene," *Current Opinion in Investigational Drugs*, 2004, 5 (6):635-647.
- Miyachi et al., 1997, "Novel biological response modifiers: phthalimides with tumor necrosis factor-alpha production-regulating activity," *J. Med. Chem.* 40:2858-2865.
- Moutouh et al., "Novel immunomodulatory drugs (IMiDs®): A potential, new therapy for β -hemoglobinopathies," Abstract # 3740, *American Society of Hematology*, Dec. 4-7, 2004.
- Patten et al., "The early use of the serum free light chain assay in patients with relapsed refractory myeloma receiving treatment with a thalidomide analogue (CC-4047)," Abstract # 1640, *American Society of Hematology*, Dec. 6-9, 2003.
- Payvandi et al., "CC-5013 inhibits the expression of adhesion molecules ICAM-1 and CD44 and prevents metastasis of B16 F10 mouse melanoma cells in an animal model," *American Society of Clinical Oncology*, Abstract # 992, 2003.
- Payvandi et al., "Effects of a thalidomide analog on binding activity of transcription factors and cell cycle progression of multiple myeloma cell lines," *Blood*, Abstract #2487, Dec. 1-5, 2000, *American Society of Hematology*.
- Payvandi et al., "Immunomodulatory drugs inhibit expression of cyclooxygenase-2 from TNF- α , IL-1 β , and LPS-stimulated human PBMC in a partially IL-10-dependent manner," *Cellular Immunology*, 2004, 81-88.
- Payvandi et al., "Thalidomide and IMiDs inhibit microvessel formation from human arterial rings in the absence of human liver microsomes," *Blood*, Abstract # 5046, Dec. 6-10, 2002, *American Society of Hematology*.
- Payvandi et al., "Thalidomide analogs IMiDs inhibit expression of cyclooxygenase-2 in multiple myeloma cell line and LPS stimulated PBMCs," *Blood*, Abstract # 2689, Dec. 7-11, 2001, *American Society of Hematology*.
- Payvandi et al., "The thalidomide analogs IMiDs enhance expression of CD69 stimulatory receptor on natural killer cells," Abstract # 1793, *American Association for Cancer Research*, Mar. 24-28, 2001.
- Raje et al., "Combination of the mTOR inhibitor rapamycin and CC-5013 has synergistic activity in multiple myeloma," *Blood*, Dec. 15, 2005, 104 (13):4188-4193.
- Rajkumar et al., "Combination therapy with lenalidomide plus dexamethasone (Rev/Dex) for newly diagnosed myeloma," *Blood*, Dec. 15, 2005, 106 (13):4050-4053.
- Richardson et al., "A multi-center, randomized, phase 2 study to evaluate the efficacy and safety of 2 CDC-5013 dose regimens when used alone or in combination with dexamethasone (Dex) for the treatment of relapsed or refractory multiple myeloma (MM)," *Blood*, Abstract # 825, *American Society of Hematology*, Dec. 6-9, 2003.
- Richardson et al., "A multicenter, single-arm, open-label study to evaluate the efficacy and safety of single-agent lenalidomide in patients with relapsed and refractory multiple myeloma: preliminary results," *10th International Myeloma Workshop*, Apr. 10-14, 2005.
- Richardson et al., "A Phase I study of oral CC5013, an immunomodulatory thalidomide (Thal) derivative, in patients with relapsed and refractory multiple myeloma (MM)," *Blood*, Abstract #3225, Dec. 7-11, 2001, *American Society of Hematology*.
- Richardson et al., "A phase I trial of lenalidomide (REVLIMID®) with bortezomib (VELCADE®) in relapsed and refractory multiple myeloma," *Blood*, Abstract # 365, *American Society of Hematology*, Dec. 10-13, 2005.
- Richardson et al., "Immunomodulatory analogs of thalidomide: an emerging new therapy in myeloma," *Journal of Clinical Oncology*, 2004, 22(16) 3212-3214.
- Richardson et al., "Immunomodulatory drug CC-5013 overcomes drug resistance and is well tolerated in patients with relapsed multiple myeloma," *Blood*, 2002, 100:3063-3067, *American Society of Hematology*.
- Richardson et al., "Novel biological therapies for the treatment of multiple myeloma," *Best Practice & Research Clinical Haematology*, 2005, 18 (4):619-634.
- Rubin et al., "Principles of cancer treatment-1," 2003, 12 ONCO IV 1.
- Schafer et al., "Enhancement of cytokine production and AP-1 transcriptional activity in T cells by thalidomide-related immunomodulatory drugs," *Journal of Pharmacology and Experimental Therapeutics*, 2003, 305(3)1222-1232.
- Schey et al., "A phase I study of an immunomodulatory thalidomide analog, CC-4047, in relapsed or refractory multiple myeloma," *Journal of Clinical Oncology*, 2004, 22 (16):1-8.
- Schey et al., "A phase I study of an immunomodulatory thalidomide analogue (CC4047) in relapse/refractory multiple myeloma," *International Society for Experimental Hematology*, Abstract #248, 2002.
- Shah et al., 1999, "Synthesis and enantiomeric separation of 2-phthalimidino-glutaric acid analogues: potent inhibitors of tumor metastasis," *J. Med. Chem.* 42:3014-3017.

US 7,465,800 B2

Page 5

- Shaughnessy et al., "Global gene expression analysis shows loss of C-MYC and IL-6 receptor gene mRNA after exposure of myeloma to thalidomide and IMiD," Abstract # 2485, *American Society of Hematology*, Dec. 1-5, 2000.
- Shire et al., "TNF- α inhibitors and rheumatoid arthritis," *Exp. Opin. Ther. Patents*, 1998, 8 (5):531-544.
- Sorbera et al., "CC-5013. Treatment of multiple myeloma. Treatment of Melanoma. Treatment of myelodysplastic syndrome. Angiogenesis inhibitor. TNF- α production inhibitor," *Drugs of the Future*, 2003, 28(5):425-431.
- Streety et al., "An update of the use and outcomes of the new immunomodulatory agent CC-4047 (Actimid) in patients with relapsed/refractory myeloma," Abstract #829, *American Society of Hematology*, Dec. 6-9, 2003.
- Streety et al., "Changes in neutrophil phenotype following the administration of CC-4047 (Actimid) to patients with multiple myeloma," Abstract # 2543, *American Society of Hematology*, Dec. 6-9, 2003.
- Streety et al., "Thalidomide analogue CC-4047 is effective in the treatment of patients with relapsed and refractory multiple myeloma (MM) and induces T-cell activation and IL-12 production," Abstract # 367, *International Multiple Myeloma Workshop*, May 23-27, 2003.
- Teo et al., "A phase I, single-blind, placebo-controlled, ascending single oral dose, safety, tolerability and pharmacokinetic study of CDC-501, a novel immunomodulatory- oncologic agent, in healthy male subjects with a comparison of fed and fasted," *Clinical Pharmacology and Therapeutics*, 2002, 71 (2)93.
- Teo et al., "Chiral inversion of the second generation IMiDTM CC-4047 (ACTIMIDTM) in human plasma and phosphate-buffered saline," *Chirality*, 2003, 15:348-351.
- Thertulien et al., "Hybrid MEL/DT Pace autotransplant regimen for Multiple Myeloma (MM)- safety and efficacy data in pilot study of 15 patients," *Blood*, Abstract # 2869, *American Society of Hematology*, Dec. 7-11, 2001.
- Tohny et al., "A phase I study of oral CC-5013 (lenalidomide, RevlimidTM), a thalidomide derivative in patients with refractory metastatic cancer," *Clinical Prostate Cancer*, 2004, 2:241-243.
- Tricot et al., "Angiochemotherapy (ACT) for multiple myeloma (MM) with DT-Pace results in a high response rate, but in contrast to tandem transplants with melphalan does not affect durable disease control," *Blood*, Abstract # 3531, *American Society of Hematology*, Dec. 7-11, 2001.
- Tsenova et al., "Use of IMiD3, a thalidomide analog, as an adjunct to therapy for experimental tuberculous meningitis," *Antimicrobial Agents and Chemotherapy*, 2002, 46 (6)1887-1895.
- Weber et al., "A multicenter, randomized, parallel-group, double-blind, placebo-controlled study of lenalidomide plus dexamethasone versus dexamethasone alone in previously treated subjects with multiple myeloma," Abstract # PO.738, *International Multiple Myeloma Workshop*, Apr. 10-14, 2005.
- Weber, "Lenalidomide (CC-5013), RevlimidTM) and other ImiDs," Abstract # PL5.02, *International Multiple Myeloma Workshop*, Apr. 10-14, 2005.
- Ye et al., "Novel IMiD drugs enhance expansion and regulate differentiation of human cord blood CD34+ cells with cytokines," *Blood*, Abstract #4099, *American Society of Hematology*, Dec. 6-10, 2002.
- Zangari et al., "Results of phase I study of CC-5013 for the treatment of multiple myeloma (MM) patients who relapse after high dose chemotherapy (HDCT)," *American Society of Hematology*, Abstract #3226, 2001.
- Zangari et al., "Revimid 25 mg (REV 25)×20 versus 50 mg (REV 50)×10 q 28 days with bridging of 5 mg×10 versus 10 mg×5 as post-transplant salvage therapy for multiple myeloma (MM)," *Blood*, Abstract # 1642, *American Society of Hematology*, Dec. 6-9, 2003.
- Zangari et al., "Risk factors for deep vein thrombosis (DVT) in a large group of myeloma patients (Pts) treated with thalidomide (Thal): The Arkansas Experience," *Blood*, Abstract # 681, *American Society of Hematology*, Dec. 7-11, 2001.
- Zeldis et al., "Potential new therapeutics for Waldenstrom's macroglobulinemia," *Seminars in Oncology*, 2003, 30 (2):275-281.
- Zeldis et al., "Update on the evolution of the IMiDTM," *International Society for Biological Therapy of Cancer*, Oral Abstract, 2003.
- Zhang et al., "CC-5079, a novel microtubule and TNF- α inhibitor with anti-angiogenic and antimetastasis activity," Abstract # B012, *International Conference on Molecular Targets and Cancer Therapeutics*, Nov. 17-21, 2003.
- U.S. Appl. No. 60/499,723, filed Sep. 4, 2003, Jaworsky.
- U.S. Appl. No. 10/693,794, filed Oct. 23, 2003, Zeldis.
- DiMartino et al., 1997, "Preparation and physical characterization of forms II and III of paracetamol," *J. Thermal Analysis* 48:447-458.
- Jonsson, 1972, "Chemical structure and teratogenic properties. 3. A review of available data on structure-activity relationships and mechanism of action of thalidomide analogues," *Acta. Pharm. Suec.* 9(6):521-542.
- Wilens et al., 1977, "Strategies in optical resolutions," *Tetrahedron* 33:2725-2736.
- Wilens, 1972, *Tables of Resolving Agents and Optical Resolutions*, Eliel, ed., U. Notre Dame Press, Notre Dame, IN pp. 268.
- Wolff, ed., 1995, *I Burger's Medicinal Chemistry and Drug Discovery*, 5th ed., pp. 172-178 and 949-982.
- Corral et al., 1999, "Immunomodulation by thalidomide and thalidomide analogs," *Ann. Rheum. Dis.* 58(Supp. 1):1107-1113.
- Grant, 1999, "Theory and Origin of Polymorphism," in *Polymorphism in Pharmaceutical Solids*, Minneapolis, MN, Ch. 1, pp. 1-10.
- He, W., et al., 1993, 206th American Chemical Society, Chicago, IL; *Med. Chem.*, paper 216.
- Lentzsch et al., 2002, "S-3-Amino-phthalimido-glutarimide inhibits angiogenesis and growth of B-cell neoplasias in mice," *Cancer Research* 62:2300-2305.
- Muller et al., 1999, "Amino-substituted thalidomide analogs: Potent inhibitors of TNF- α production," *Bioorg. Med. Chem. Lett.* 9:1625-1630.
- Muller et al., 1998, "Thalidomide analogs and PDE4 inhibition," *Bioorg. Med. Chem. Lett.* 8:2669-2674.
- Muller et al., 1996, "Structural modifications of thalidomide produce analogs with enhanced tumor necrosis factor inhibitory activity," *J. Med. Chem.* 39:3238-3240.

* cited by examiner

XRPD PATTERN OF FORM A




Fig. 1

IR SPECTRUM OF FORM A




Fig. 2

RAMAN SPECTRUM OF FORM A




Fig. 3




Fig. 4




Fig. 5




Fig. 6

IR SPECTRUM OF FORM B




Fig. 7

RAMAN SPECTRUM OF FORM B





Fig. 8



TG-IR RESULTS FOR FORM B




Fig. 10




Fig. 11




Fig. 12

IR SPECTRUM OF FORM C




Fig. 13

RAMAN SPECTRUM OF FORM C




Fig. 14

TGA (TOP) AND DSC (BOTTOM) OF FORM C




Fig. 15

TG-IR RESULTS FOR FORM C




Fig. 16




Fig. 17

XRPD PATTERN OF FORM D




Fig. 18

IR SPECTRUM OF FORM D




Fig. 19

RAMAN SPECTRUM OF FORM D




Fig. 20

TGA (TOP) AND DSC (BOTTOM) OF FORM D




Fig. 21




Fig. 22




Fig. 23




Fig. 24




Fig. 25

XRPD PATTERN OF FORM F




Fig. 26




Fig. 27

XRPD PATTERN OF FORM G




Fig. 28




Fig. 29

XRPD PATTERN OF FORM H




Fig. 30

TGA (TOP) AND DSC (BOTTOM) OF FORM H




Fig. 31

XRPD PATTERN OF POLYMORPH B




Fig. 32




Fig. 33

XRPD PATTERN OF POLYMORPH B




Fig. 34

XRPD PATTERN OF POLYMORPH E




Fig. 35

XRPD PATTERN OF POLYMORPH MIXTURE




Fig. 36




Fig. 37




Fig. 38




Fig. 39




Fig. 40




Fig. 41




Fig. 42




Fig. 43




Fig. 44




Fig. 45




Fig. 46




Fig. 47




Fig. 48




Fig. 49




Fig. 50




Fig. 51




Fig. 52




FIG. 53

INTRINSIC DISSOLUTION OF FORMS A, B, AND E (EXPERIMENT NO.2)




FIG.54

US 7,465,800 B2

1

**POLYMORPHIC FORMS OF
3-(4-AMINO-1-OXO-1,3
DIHYDRO-ISOINDOL-2-YL)-PIPERIDINE-2,6-
IONE**

This application claims the benefit of U.S. provisional application 60/499,723, filed Sep. 4, 2003, the contents of which are incorporated by reference herein their entirety.

1. FIELD OF THE INVENTION

This invention relates to polymorphic forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione, compositions comprising the polymorphic forms, methods of making the polymorphic forms and methods of their use for the treatment of diseases and conditions including, but not limited to, inflammatory diseases, autoimmune diseases, and cancer.

2. BACKGROUND OF THE INVENTION

Many compounds can exist in different crystal forms, or polymorphs, which exhibit different physical, chemical, and spectroscopic properties. For example, certain polymorphs of a compound may be more readily soluble in particular solvents, may flow more readily, or may compress more easily than others. See, e.g., P. DiMartino, et al., *J. Thermal Anal.*, 48:447-458 (1997). In the case of drugs, certain solid forms may be more bioavailable than others, while others may be more stable under certain manufacturing, storage, and biological conditions. This is particularly important from a regulatory standpoint, since drugs are approved by agencies such as the U.S. Food and Drug Administration only if they meet exacting purity and characterization standards. Indeed, the regulatory approval of one polymorph of a compound, which exhibits certain solubility and physico-chemical (including spectroscopic) properties, typically does not imply the ready approval of other polymorphs of that same compound.

Polymorphic forms of a compound are known in the pharmaceutical arts to affect, for example, the solubility, stability, flowability, fractability, and compressibility of the compound, as well as the safety and efficacy of drug products comprising it. See, e.g., Knapman, K. *Modern Drug Discoveries*, 2000, 53. Therefore, the discovery of new polymorphs of a drug can provide a variety of advantages.

U.S. Pat. Nos. 5,635,517 and 6,281,230, both to Muller et al., disclose 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione, which is useful in treating and preventing a wide range of diseases and conditions including, but not limited to, inflammatory diseases, autoimmune diseases, and cancer. New polymorphic forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione can further the development of formulations for the treatment of these chronic illnesses, and may yield numerous formulation, manufacturing and therapeutic benefits.

3. SUMMARY OF THE INVENTION

This invention encompasses polymorphs of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. In certain aspects, the invention provides polymorphs of the compound identified herein as forms A, B, C, D, E, F, G, and H. The invention also encompasses mixtures of these forms. In further embodiments, this invention provides methods of making, isolating and characterizing the polymorphs.

This invention also provides pharmaceutical compositions and single unit dosage forms comprising a polymorph of

2

3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. The invention further provides methods for the treatment or prevention of a variety of diseases and disorders, which comprise administering to a patient in need of such treatment or prevention a therapeutically effective amount of a polymorph of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione.

4. BRIEF DESCRIPTION OF THE DRAWINGS

Specific aspects of the invention can be understood with reference to the attached figures:

FIG. 1 provides a representative X-ray powder diffraction (XRPD) pattern of Form A;

FIG. 2 provides a representative IR spectrum of Form A;

FIG. 3 provides a representative Raman spectrum of Form A;

FIG. 4 provides a representative thermogravimetric analysis (TGA) curve and a representative differential scanning calorimeter (DSC) thermogram of Form A;

FIG. 5 provides a representative moisture sorption/desorption isotherm of Form A;

FIG. 6 provides a representative XRPD pattern of Form B;

FIG. 7 provides a representative IR spectrum of Form B;

FIG. 8 provides a representative Raman spectrum of Form B;

FIG. 9 provides a representative TGA curve and a representative DSC thermogram of Form B;

FIG. 10 provides representative TG-IR results of Form B;

FIG. 11 provides a representative moisture sorption/desorption isotherm of Form B;

FIG. 12 provides a representative XRPD pattern of Form C;

FIG. 13 provides a representative IR spectrum of Form C;

FIG. 14 provides a representative Raman spectrum of Form C;

FIG. 15 provides a representative TGA curve and a representative DSC thermogram of Form C;

FIG. 16 provides representative TG-IR results of Form C;

FIG. 17 provides a representative moisture sorption/desorption isotherm of Form C;

FIG. 18 provides a representative XRPD pattern of Form D;

FIG. 19 provides a representative IR spectrum of Form D;

FIG. 20 provides a representative Raman spectrum of Form D;

FIG. 21 provides a representative TGA curve and a representative DSC thermogram of Form D;

FIG. 22 provides a representative moisture sorption/desorption isotherm of Form D;

FIG. 23 provides a representative XRPD pattern of Form E;

FIG. 24 provides a representative TGA curve and a representative DSC thermogram of Form E;

FIG. 25 provides a representative moisture sorption/desorption isotherm of Form E;

FIG. 26 provides a representative XRPD pattern for a sample of Form F;

FIG. 27 provides a representative thermogram of Form F;

FIG. 28 provides a representative XRPD pattern of Form G;

FIG. 29 provides a representative DSC thermogram for a sample of Form G;

FIG. 30 provides a representative XRPD pattern of Form H;

FIG. 31 provides a representative TGA curve and a representative DSC thermogram of Form H;

US 7,465,800 B2

3

FIG. 32 provides a representative XRPD pattern of Form B;

FIG. 33 provides a representative XRPD pattern of Form B;

FIG. 34 provides a representative XRPD pattern of Form B;

FIG. 35 provides a representative XRPD pattern of Form E;
FIG. 36 provides a representative XRPD pattern of polymorph mixture;

FIG. 37 provides a representative TGA curve of Form B;

FIG. 38 provides a representative TGA curve of Form B;

FIG. 39 provides a representative TGA curve of Form B;

FIG. 40 provides a representative TGA curve of Form E;

FIG. 41 provides a representative TGA curve of polymorph mixture;

FIG. 42 provides a representative DSC thermogram of Form B;

FIG. 43 provides a representative DSC thermogram of Form B;

FIG. 44 provides a representative DSC thermogram of Form B;

FIG. 45 provides a representative DSC thermogram of Form E;

FIG. 46 provides a representative DSC thermogram of polymorph mixture;

FIG. 47 provides a UV-Vis scan of dissolution medium;

FIG. 48 provides a UV-Vis scan of 0.04 mg/ml of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione in dissolution medium;

FIG. 49 provides a UV-Vis scan of 0.008 mg/ml of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione in dissolution medium;

FIG. 50 provides a calibration curve for 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione;

FIG. 51 provides a solubility curve of Form A;

FIG. 52 provides a solubility curve of Form B;

FIG. 53 provides an intrinsic dissolution of Forms A, B and E; and

FIG. 54 provides an intrinsic dissolution of Forms A, B and E.

5. DETAILED DESCRIPTION OF THE INVENTION

5.1 Definitions

As used herein and unless otherwise indicated, the terms “treat,” “treating” and “treatment” refer to the alleviation of a disease or disorder and/or at least one of its attendant symptoms.

As used herein and unless otherwise indicated, the terms “prevent,” “preventing” and “prevention” refer to the inhibition of a symptom of a disease or disorder or the disease itself.

As used herein and unless otherwise indicated, the terms “polymorph” and “polymorphic form” refer to solid crystalline forms of a compound or complex. Different polymorphs of the same compound can exhibit different physical, chemical and/or spectroscopic properties. Different physical properties include, but are not limited to stability (e.g., to heat or light), compressibility and density (important in formulation and product manufacturing), and dissolution rates (which can affect bioavailability). Differences in stability can result from changes in chemical reactivity (e.g., differential oxidation, such that a dosage form discolors more rapidly when comprised of one polymorph than when comprised of another polymorph) or mechanical characteristics (e.g., tablets crumble on storage as a kinetically favored polymorph converts to thermodynamically more stable polymorph) or both

4

(e.g., tablets of one polymorph are more susceptible to breakdown at high humidity). Different physical properties of polymorphs can affect their processing. For example, one polymorph might be more likely to form solvates or might be more difficult to filter or wash free of impurities than another due to, for example, the shape or size distribution of particles of it.


Polymorphs of a molecule can be obtained by a number of methods known in the art. Such methods include, but are not limited to, melt recrystallization, melt cooling, solvent recrystallization, desolvation, rapid evaporation, rapid cooling, slow cooling, vapor diffusion and sublimation. Polymorphs can be detected, identified, classified and characterized using well-known techniques such as, but not limited to, differential scanning calorimetry (DSC), thermogravimetry (TGA), X-ray powder diffractometry (XRPD), single crystal X-ray diffractometry, vibrational spectroscopy, solution calorimetry, solid state nuclear magnetic resonance (NMR), infrared (IR) spectroscopy, Raman spectroscopy, hot stage optical microscopy, scanning electron microscopy (SEM), electron crystallography and quantitative analysis, particle size analysis (PSA), surface area analysis, solubility, and rate of dissolution.

As used herein to refer to the spectra or data presented in graphical form (e.g., XRPD, IR, Raman and NMR spectra), and unless otherwise indicated, the term “peak” refers to a peak or other special feature that one skilled in the art would recognize as not attributable to background noise. The term “significant peaks” refers to peaks at least the median size (e.g., height) of other peaks in the spectrum or data, or at least 1.5, 2, or 2.5 times the median size of other peaks in the spectrum or data.

As used herein and unless otherwise indicated, the term “substantially pure” when used to describe a polymorph of a compound means a solid form of the compound that comprises that polymorph and is substantially free of other polymorphs of the compound. A representative substantially pure polymorph comprises greater than about 80% by weight of one polymorphic form of the compound and less than about 20% by weight of other polymorphic forms of the compound, more preferably greater than about 90% by weight of one polymorphic form of the compound and less than about 10% by weight of the other polymorphic forms of the compound, even more preferably greater than about 95% by weight of one polymorphic form of the compound and less than about 5% by weight of the other polymorphic forms of the compound, and most preferably greater than about 97% by weight of one polymorphic forms of the compound and less than about 3% by weight of the other polymorphic forms of the compound.

5.2 Polymorphic Forms

This invention is directed to polymorphic forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione, which has the structure shown below:



This compound can be prepared according to the methods described in U.S. Pat. Nos. 6,281,230 and 5,635,517, the

US 7,465,800 B2

5

entireties of which are incorporated herein by reference. For example, the compound can be prepared through catalytic hydrogenation of 3-(4-nitro-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. 3-(4-Nitro-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione can be obtained by allowing 2,6-dioxopiperidin-3-ammonium chloride to react with methyl 2-bromomethyl-4-nitrobenzoate in dimethylformamide in the presence of triethylamine. The methyl 2-bromomethyl-4-nitrobenzoate in turn is obtained from the corresponding methyl ester of nitro-ortho-toluic acid by conventional bromination with N-bromosuccinimide under the influence of light.

Polymorphs of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione can be obtained by techniques known in the art, including solvent recrystallization, desolvation, vapor diffusion, rapid evaporation, slow evaporation, rapid cooling and slow cooling. Polymorphs can be made by dissolving a weighed quantity of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione in various solvents at elevated temperatures. The solutions of the compound can then be filtered and allowed to evaporate either in an open vial (for fast hot evaporation) or in a vial covered with aluminum foil containing pinholes (hot slow evaporation). Polymorphs can also be obtained from slurries. Polymorphs can be crystallized from solutions or slurries using several methods. For example, a solution created at an elevated temperature (e.g., 60° C.) can be filtered quickly then allowed to cool to room temperature. Once at room temperature, the sample that did not crystallize can be moved to a refrigerator then filtered. Alternatively, the solutions can be crash cooled by dissolving the solid in a solvent at an increased temperature (e.g., 45-65° C.) followed by cooling in a dry ice/solvent bath.

One embodiment of the invention encompasses Form A of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form A is an unsolvated, crystalline material that can be obtained from non-aqueous solvent systems. Another embodiment of the invention encompasses Form B of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form B is a hemihydrated, crystalline material that can be obtained from various solvent systems. Another embodiment of the invention encompasses Form C of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form C is a hemisolvated crystalline material that can be obtained from solvents such as, but not limited to, acetone. Another embodiment of the invention encompasses Form D of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form D is a crystalline, solvated polymorph prepared from a mixture of acetonitrile and water. Another embodiment of the invention encompasses Form E of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form E is a dihydrated, crystalline material. Another embodiment of the invention encompasses Form F of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form F is an unsolvated, crystalline material that can be obtained from the dehydration of Form E. Another embodiment of the invention encompasses Form G of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form G is an unsolvated, crystalline material that can be obtained from slurrying forms B and E in a solvent such as, but not limited to, tetrahydrofuran (THF). Another embodiment of the invention encompasses Form H of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form H is a partially hydrated crystalline material that can be obtained by exposing Form E to 0% relative humidity. Each of these forms is discussed in detail below.

Another embodiment of the invention encompasses a composition comprising amorphous 3-(4-amino-1-oxo-1,3 dihydro-

6

dro-isoindol-2-yl)-piperidine-2,6-dione and crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione of form A, B, C, D, E, F, G or H. Specific compositions can comprise greater than about 50, 75, 90 or 95 weight percent crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione.

Another embodiment of the invention encompasses a composition comprising at least two crystalline forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione (e.g., a mixture of polymorph forms B and E).

5.2.1 Form A

The data described herein for Form A, as well as for Forms B-H, were obtained using the experimental methods described in Examples 6.3-6.7, provided below.

Form A can be obtained from various solvents, including, but not limited to 1-butanol, butyl acetate, ethanol, ethyl acetate, methanol, methyl ethyl ketone, and THF. FIG. 1 shows a representative XRPD pattern of Form A. The pattern is characterized by peaks, preferably significant peaks, at approximately 8, 14.5, 16, 17.5, 20.5, 24, and 26 degrees 2 θ . Representative IR and Raman spectra data are provided in FIGS. 2 and 3.

Representative thermal characteristics of Form A are shown in FIG. 4. TGA data show a small weight increase up to about 150° C., indicating an unsolvated material. Weight loss above 150° C. is attributed to decomposition. The DSC curve of Form A exhibits an endotherm at about 270° C.

Representative moisture sorption and desorption data are plotted in FIG. 5. Form A does not exhibit a significant weight gain from 5 to 95% relative humidity. Equilibrium can be obtained at each relative humidity step. As the form dries from 95% back down to 5% relative humidity, it tends to maintain its weight such that at 5% relative humidity it has typically lost only about 0.003% by weight from start to finish. Form A is capable of remaining a crystalline solid for about 11 days when stored at about 22, 45, 58, and 84% relative humidity.

Interconversion studies show that Form A can convert to Form B in aqueous solvent systems and can convert to Form C in acetone solvent systems. Form A tends to be stable in anhydrous solvent systems. In water systems and in the presence of Form E, Form A tends to convert to Form E.

When stored for a period of about 85 days under two different temperature/relative humidity stress conditions (room temperature/0% relative humidity (RH) and 40° C./93% RH), Form A typically does not convert to a different form.

In sum, Form A is a crystalline, unsolvated solid that melts at approximately 270° C. Form A is weakly or not hygroscopic and appears to be the most thermodynamically stable anhydrous polymorph of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione discovered thus far.

5.2.2 Form B

Form B can be obtained from many solvents, including, but not limited to, hexane, toluene, and water. FIG. 6 shows a representative XRPD pattern of Form B, characterized by peaks at approximately 16, 18, 22 and 27 degrees 2 θ .

Solution proton NMR confirm that Form B is a form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Representative IR and Raman spectra are shown in FIGS. 7 and 8, respectively. Compared to Form A, the IR spectrum for Form B has peaks at approximately 3513 and 1960 cm⁻¹.

Representative DSC and TGA data for Form B are shown in FIG. 9. The DSC curve exhibits endotherms at about 146 and 268° C. These events are identified as dehydration and melting by hot stage microscopy experiments. Form B typi-

US 7,465,800 B2

7

cally loses about 3.1% volatiles up to about 175° C. (per approximately 0.46 moles of water). Comparison of the IR spectrum of the volatiles with that of water indicates that they are water (See FIG. 10). Calculations from TGA data indicate that Form B is a hemihydrate. Karl Fischer water analysis also supports this conclusion.

Representative moisture sorption and desorption data are shown in FIG. 11. Form B typically does not exhibit a significant weight gain from 5% to 95% relative humidity, when equilibrium is obtained at each relative humidity step. As Form B dries from 95% back down to 5% relative humidity, it tends to maintain its weight such that at 5% relative humidity it typically has gained only about 0.022% by weight (about 0.003 mg) from start to finish. Form B does not convert to a different form upon exposure to about 84% relative humidity for about ten days.

Interconversion studies show that Form B typically converts to Form A in a THF solvent system, and typically converts to Form C in an acetone solvent system. In aqueous solvent systems such as pure water and 10% water solutions, Form B is the most stable of the polymorphic forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. However, it can convert to Form E in the presence of water. Desolvation experiments show that upon heating at about 175° C. for about five minutes, Form B typically converts to Form A.

When stored for a period of about 85 days under two different temperature/relative humidity stress conditions (room temperature/0% RH and 40° C./93% RH), Form B does not convert to a different form.

In sum, Form B is a hemihydrated, crystalline solid which has a DSC thermogram exhibiting endotherms at about 146 and about 268° C. Interconversion studies show that Form B converts to Form E in aqueous solvent systems, and converts to other forms in acetone and other anhydrous systems.

5.2.3 Form C

Form C can be obtained from evaporations, slurries and slow cools in acetone solvent systems. A representative XRPD pattern of this form is shown in FIG. 12. The data are characterized by peaks at approximately 15.5 and 25 degrees 2θ.

Solution proton NMR indicates that the 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione molecule is intact. Representative IR and Raman spectra are shown in FIGS. 13 and 14, respectively. The IR spectrum of Form C is characterized by peaks at approximately 3466, 3373, and 3318 cm⁻¹. The Raman spectrum of Form C is characterized by peaks at about 3366, 3321, 1101, and 595 cm⁻¹.

Representative thermal characteristics for Form C are plotted in FIG. 15. Form C loses about 10.02% volatiles up to about 175° C., indicating it is a solvated material. Weight loss above about 175° C. is attributed to decomposition. Identification of volatiles in Form C can be accomplished with TG-IR experiments. The representative IR spectrum captured after several minutes of heating, as depicted in FIG. 13, when compared with a spectral library, shows acetone to be the best match. Calculations from TGA data show that Form C is a hemisolvate (approximately 0.497 moles of acetone). The DSC curve for Form C, shown in FIG. 15, exhibits endotherms at about 150 and about 269° C. The endotherm at about 150° C. is attributed to solvent loss based on observations made during hot stage microscopy experiments. The endotherm at about 269° C. is attributed to the melt based on hot stage experiments.

Representative moisture sorption and desorption balance data are shown in FIG. 17. Form C does not exhibit a significant weight gain from 5 to 85% relative humidity, when

8

equilibrium is obtained at each relative humidity step up to 85% relative humidity. At 95% relative humidity, Form C experiences a significant weight loss of about 6.03%. As the sample dries from 95% back down to 5% relative humidity, the sample maintains the weight achieved at the end of the adsorption phase at each step down to 5% relative humidity. Form C is capable of converting to Form B when stored at about 84% relative humidity for approximately ten days.

Interconversion studies show that Form C typically converts to Form A in a THF solvent system and typically converts to Form E in an aqueous solvent system. In an acetone solvent system, Form C is the most stable form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Desolvation experiments performed on Form C show that upon heating at about 150° C. for about five minutes, Form C will typically convert to Form A.

In sum, Form C is a crystalline, hemisolvated solid, which has a DSC thermogram exhibiting endotherms at about 150 and about 269° C. Form C is not hygroscopic below about 85% RH, but can convert to Form B at higher relative humidities.

5.2.4 Form D

Form D can be obtained from evaporation in acetonitrile solvent systems. A representative XRPD pattern of the form is shown in FIG. 18. The pattern is characterized by peaks at approximately 27 and 28 degrees 2θ.

Solution proton NMR indicates that the 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione molecule is intact. Representative IR and Raman spectra are shown in FIGS. 19 and 20, respectively. The IR spectrum of Form D is characterized by peaks at approximately 3509, 2299, and 2256 cm⁻¹. The Raman spectrum of Form D is characterized by peaks at approximately 2943, 2889, 2297, 2260, 1646, and 1150 cm⁻¹.

Representative thermal characteristics for Form D are plotted in FIG. 21. Form D loses about 6.75% volatiles up to about 175° C., indicating a solvated material. Weight loss above about 175° C. is attributed to decomposition. TG-IR experiments indicate that the volatiles are water and acetonitrile. Calculations from TG data show that about one mole of water is present in the sample. A representative DSC curve for Form D exhibits endotherms at about 122 and about 270° C. The endotherm at about 122° C. is attributed to loss of volatiles based on observations made during hot stage microscopy experiments. The endotherm at about 270° C. is attributed to the melt based on hot stage experiments.

Representative moisture sorption and desorption data are plotted in FIG. 22. Form D does not exhibit a significant weight gain from 5 to 95% relative humidity when equilibrium is obtained at each relative humidity step. As the form dries from 95% back down to 5% relative humidity, it maintains its weight such that at 5% relative humidity the form has typically gained only about 0.39% by weight (about 0.012 mg) from start to finish. Form A is capable of converting to Form B when stored at about 84% relative humidity for approximately ten days.

Interconversion studies show that Form D is capable of converting to Form A in a THF solvent system, to Form E in an aqueous solvent system, and to Form C in an acetone solvent system. Desolvation experiments performed on Form D show that upon heating at about 150° C. for about five minutes Form D will typically convert to Form A.

In sum, Form D is a crystalline solid, solvated with both water and acetonitrile, which has a DSC thermogram exhibiting endotherms at about 122 and about 270° C. Form D is either weakly or not hygroscopic, but will typically convert to Form B when stressed at higher relative humidities.

5.2.5 Form E

Form E can be obtained by slurring 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione in water and by a slow evaporation of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione in a solvent system with a ratio of about 9:1 acetone:water. A representative XRPD pattern is shown in FIG. 23. The data are characterized by peaks at approximately 20, 24.5 and 29 degrees 2 θ .

Representative thermal characteristics of Form E are plotted in FIG. 24. Form E typically loses about 10.58% volatiles up to about 125° C., indicating that it is a solvated material. A second weight loss of an additional about 1.38% was observed between about 125° C. and about 175° C. Weight loss above about 175° C. is attributed to decomposition. Karl Fischer and TG-IR experiments support the conclusion that the volatile weight loss in Form E is due to water. The representative DSC curve for Form E exhibits endotherms at about 99, 161 and 269° C. Based on observations made during hot stage microscopy experiments, the endotherms at about 99 and about 161° C. are attributed to loss of volatiles. The endotherm at about 269° C. is attributed to the melt based on hot stage experiments.

Representative moisture sorption and desorption data are plotted in FIG. 25. Form E typically does not exhibit a significant weight change from 5 to 95% relative humidity when equilibrium is obtained at each relative humidity step. As the sample dried from 95% back down to 5% relative humidity, the sample continues to maintain weight such that at 5% relative humidity the sample has lost only about 0.0528% by weight from start to finish.

Interconversion studies show that Form E can convert to Form C in an acetone solvent system and to Form G in a THF solvent system. In aqueous solvent systems, Form E appears to be the most stable form. Desolvation experiments performed on Form E show that upon heating at about 125° C. for about five minutes, Form E can convert to Form B. Upon heating at 175° C. for about five minutes, Form B can convert to Form F.

When stored for a period of 85 days under two different temperature/relative humidity stress conditions (room temperature/0% RH and 40° C./93% RH) Form E typically does not convert to a different form. When stored for seven days at room temperature/0% RH, Form E can convert to a new form, Form H.

5.2.6 Form F

Form F can be obtained by complete dehydration of Form E. A representative XRPD pattern of Form F, shown in FIG. 26, is characterized by peaks at approximately 19, 19.5 and 25 degrees 2 θ .

Representative thermal characteristics of Form F are shown in FIG. 27. The representative DSC curve for Form F exhibits an endotherm at about 269° C. preceded directly by two smaller endotherms indicative of a crystallized form of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. The DSC thermogram does not show any thermal events prior to the melt, suggesting that it is an unsolvated material.

5.2.7 Form G

Form G can be obtained by slurring forms B and E in THF. A representative XRPD pattern of this form, shown in FIG. 28, is characterized by a peak at approximately 23 degrees 2 θ . Two other peaks unique to Form G appear at approximately 21 and 24.5 degrees 2 θ .

Representative thermal characteristics of Form G are plotted in FIG. 29. A representative DSC curve for Form G exhibits an endotherm at about 248° C. followed by a small, broad exotherm at about 267° C. No thermal events are seen

in the DSC thermogram at lower temperatures, suggesting that it is an unsolvated material.

5.2.8 Form H

Form H can be obtained by storing Form E at room temperature and 0% RH for about 7 days. A representative XRPD pattern is shown in FIG. 30. The pattern is characterized by a peak at 15 degrees 2 θ , and two other peaks at 26 and 31 degrees 2 θ .

Representative thermal characteristics are shown in FIG. 31. Form H loses about 1.67% volatiles up to about 150° C. Weight loss above about 150° C. is attributed to decomposition. Karl Fischer data shows that Form H typically contains about 1.77% water (about 0.26 moles), suggesting that the weight loss seen in the TG is due to dehydration. The DSC thermogram shows a broad endotherm between about 50° C. and about 125° C., corresponding to the dehydration of Form H and a sharp endotherm at about 269° C., which is likely due to a melt.

When slurried in water with either Forms A or B, after about 14 days Form H can convert to Form E. When slurried in THF, Form H can convert to Form A. When slurried in acetone, Form H can convert to Form C.

In sum, Form H is a crystalline solid, hydrated with about 0.25 moles of water, which has a DSC thermogram exhibiting an endotherm between about 50 and 125° C. and an endotherm at about 269° C.

5.3 Methods of Use and Pharmaceutical Compositions

Polymorphs of the invention exhibit physical characteristics that are beneficial for drug manufacture, storage or use. All polymorphs of the invention have utility as pharmaceutically active ingredients or intermediates thereof.

This invention encompasses methods of treating and preventing a wide variety of diseases and conditions using polymorphs of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. In each of the methods, a therapeutically or prophylactically effective amount of the compound is administered to a patient in need of such treatment or prevention. Examples of such disease and conditions include, but are not limited to, diseases associated with undesired angiogenesis, cancer (e.g., solid and blood borne tumors), inflammatory diseases, autoimmune diseases, and immune diseases. Examples of cancers and pre-cancerous conditions include those described in U.S. Pat. Nos. 6,281,230 and 5,635,517 to Muller et al. and in various U.S. patent applications to Zeldis, including application Ser. No. 10/411,649, filed Apr. 11, 2003 (Treatment of Myelodysplastic Syndrome); Ser. No. 10/438,213 filed May 15, 2003 (Treatment of Various Types of Cancer); Ser. No. 10/411,656, filed Apr. 11, 2003 (Treatment of Myeloproliferative Diseases). Examples of other diseases and disorders that can be treated or prevented using compositions of the invention are described in U.S. Pat. Nos. 6,235,756 and 6,114,335 to D'Amato and in other U.S. patent applications to Zeldis, including Ser. No. 10/693,794, filed Oct. 23, 2003 (Treatment of Pain Syndrome) and Ser. No. 10/699,154, filed Oct. 30, 2003 (Treatment of Macular Degeneration). The entirety of each of the patents and patent applications cited herein is incorporated herein by reference.

Depending on the disease to be treated and the subject's condition, polymorphs of the invention can be administered by oral, parenteral (e.g., intramuscular, intraperitoneal, intravenous, ICV, intracisternal injection or infusion, subcutaneous injection, or implantation), inhalation spray, nasal, vaginal, rectal, sublingual, or topical routes of administration and may be formulated, alone or together, in suitable dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles appropriate for each route of administration. Because individual poly-

US 7,465,800 B2

11

morphs have different dissolution, stability, and other properties, the optimal polymorph used in methods of treatment may depend on the route of administration. For example, forms that are readily soluble in aqueous solutions are preferably used to provide liquid dosage forms, whereas forms that exhibit great thermal stability may be preferred in the manufacture of solid dosage forms (e.g., tablets and capsules).

Although the physical characteristics of polymorphs can, in some cases, affect their bioavailability, amounts of the polymorphs that are therapeutically or prophylactically effective in the treatment of various disease and conditions can be readily determined by those of ordinary skill in the pharmacy or medical arts. In certain embodiments of the invention, a polymorph is administered orally and in a single or divided daily doses in an amount of from about 0.10 to about 150 mg/day, or from about 5 to about 25 mg/day. In other embodiments, a polymorph is administered every other day in an amount of from about 0.10 to about 150 mg/day, or from about 5 to about 25 mg/day.

The invention encompasses pharmaceutical compositions and single unit dosage forms that can be used in methods of treatment and prevention, which comprise one or more polymorphs of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione and optionally one or more excipients or diluents. Specific compositions and dosage forms are disclosed in the various patents and patent applications incorporated herein by reference. In one embodiment, a single dosage form comprises a polymorph (e.g., Form B) in an amount of about 5, 10, 25 or 50 mg.

6. EXAMPLES

6.1 Polymorph Screen

A polymorph screen to generate the different solid forms of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione was carried out as follows.

A weighed sample of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (usually about 10 mg) was treated with aliquots of the test solvent. Solvents were either reagent or HPLC grade. The aliquots were usually about 200 μ L. Between additions, the mixture was usually shaken or sonicated. When the solids dissolved, as judged by visual inspection, estimated solubilities were calculated. Solubilities were estimated from these experiments based on the total solvent used to provide a solution. Actual solubilities may have been greater than those calculated due to the use of too-large solvent aliquots or to a slow rate of dissolution.

Samples were created by generating solutions (usually about 30 mg in 20 mL) at elevated temperatures, filtering, and allowing the solution to evaporate whether in an open vial (hot fast evaporation) or in a vial covered with aluminum foil containing pinholes (hot slow evaporation).

Slurry experiments were also performed. Usually about 25 mg of solid was placed in either 3 or 5 mL of solvent. The samples were then placed on orbital shakers at either ambient temperature or 40° C. for 4-10 days.

Crystallizations were performed using various cooling methods. Solid was dissolved in a solvent at an elevated temperature (e.g., about 60° C.), filtered quickly and allowed to cool to room temperature. Once at room temperature, samples that did not crystallize were moved to a refrigerator. Solids were removed by filtration or decantation and allowed to dry in the air. Crash cools were performed by dissolving solid in a solvent at an increased temperature (e.g., about 45-65° C.) followed by cooling in a dry ice/acetone bath.

12

Hygroscopicity studies were performed by placing portions of each polymorph in an 84% relative humidity chamber for approximately one week.

Desolvation studies were carried out by heating each polymorph in a 70° C. oven for approximately one week.

Interconversion experiments were carried out by making slurries containing two forms in a saturated solvent. The slurries were agitated for approximately 7-20 days at ambient temperature. The insoluble solids were recovered by filtration and analyzed using XRPD.

6.2 Preparation of Polymorphic Forms

Eight solid forms of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione were prepared as described below.

Form A was obtained by crystallization from various non-aqueous solvents including 1-butanol, butyl acetate, ethanol, ethyl acetate, methanol, methyl ethyl ketone, and tetrahydrofuran. Form B was also obtained by crystallization from the solvents hexane, toluene and water. Form C was obtained from evaporations, slurries, and slow cools in acetone solvent systems. Form D was obtained from evaporations in acetonitrile solvent systems. Form E was obtained most readily by slurrying 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione in water. Form F was obtained by complete desolvation of Form E. It is found to be an unsolvated, crystalline material that melts at about 269° C. Form G was obtained by slurrying forms B and E in THF. Form H was obtained by stressing Form E at room temperature and 0% RH for 7 days.

6.2.1 Synthesis of Polymorphs B and E

Form B is the desired polymorph for the active pharmaceutical ingredient (API) of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. This form has been used in the formulation of API into drug product for clinical studies. Three batches were produced as apparent mixtures of polymorphs in the non-micronized API of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione.

Development work was carried out to define a process that would generate polymorph B from this mixture of polymorphs and could be implemented for strict polymorphic controls in the validation batches and future manufacturing of API of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. Characterization of polymorphic forms produced during the work was performed by XRPD, DSC, TGA and KF.

A process was also developed for the large-scale preparation of Form E. Polymorph E material was prepared in order to carry out a comparison with polymorph B drug product in capsule dissolution testing of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. 150 g of a mixture of polymorphs in 3 L of water was stirred at room temperature for 48 hours. The product was collected by filtration and dried at 25° C. for 24 hours under vacuum. XRPD, DSC, TGA, KF and HPLC analyses confirmed that the material isolated was polymorph E.

In a preliminary work, it was demonstrated that stirring a suspension of a mixture of polymorphs of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione with water at high temperature (75° C.) for an extended period of time converted this mixture of polymorphs exclusively to form B. Several specific parameters were identified including temperature, solvent volume and drying parameters (temperature and vacuum). XRPD, DSC, TGA, KF and HPLC analyses were used to characterize all of the batches. After completing the optimization work, the optimized process was scaled-up to 100-200 g on three lots of API. Drying studies were carried

US 7,465,800 B2

13

out at 20° C., 30° C. and 40° C., and 65° C. with a vacuum of 150 mm of Hg. The results are shown in Tables 1-5.

The cooling and holding periods of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione slurry were studied. The experimental laboratory data suggests that polymorph B seems to be forming first, and overtime equilibration to polymorph E at RT conditions occurs, therefore generating a mixture of polymorphs B and E. This result supports the fact that polymorph B seems to be a kinetic product, and that prolonged processing time converts the material to polymorph E resulting in a mixture of polymorphs B and E.

A laboratory procedure was developed to exclusively produce polymorph B of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. The procedure includes a stirred 10 volume water slurry at ~75° C. for 6-24 hours. The following preferred process parameters have been identified:

1. Hot slurry temperature of 70-75° C.
2. Product filtration of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione at 65-75° C.
3. Drying under vacuum at 60-70° C. is preferred for an efficient removal of unbound water in 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione wet cake.
4. The filtration step of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione may be a time sensitive operation. The use of efficient solid-liquid separation equipment is preferred.
5. Holding periods of water-wet cake of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione at KF higher than 5% may cause the kinetic equilibrations of polymorph B to mixed polymorphs of E and B.

Drying to KF <4.0% water was achieved in ~3 hours (30-70° C., 152 mm Hg). Polymorphs B and E were distinguished by the water levels as measured by KF and TGA. The reference sample of polymorph B is micronized API. In order to make accurate comparison by XRPD samples were gently grinded before submission for analysis. This increases the clarity of the identification of the polymorphic form. All samples were analyzed for XRPD, DSC, TGA, KF and HPLC.

TABLE 1

| Preliminary Studies | | | |
|---------------------|--|--------------------|--------------------|
| Amount | Reaction conditions | Analysis | Results/conclusion |
| 2 g | Water, rt, 48 h | XRPD, DSC, TGA, KF | Polymorph E |
| 25 g | Water, rt, 48 h | XRPD, DSC, TGA, KF | Polymorph E |
| 5 g | Water, 70-75° C., 24 h then rt 24 h | XRPD, DSC, TGA, KF | Polymorph B |
| 1 g | 9:1 Acetone - water, Slow evpo. 175° C. 1 h in an oven | XRPD, DSC, TGA, KF | Polymorph Mixture |
| 1 g | Water, rt, 24 h | XRPD, DSC, TGA, KF | Polymorph A |
| 0.5 g (polymorph A) | Water, rt, 24 h | XRPD, DSC, TGA, KF | Polymorph E |
| 1 g polymorph B | Water, rt, 48 h | XRPD, DSC, TGA, KF | Polymorph E |
| 1 g polymorph E | Water, 70-75° C., 24 h | XRPD, DSC, TGA, KF | Polymorph B |
| 1 g | Slurry in heptane | XRPD, DSC, TGA, KF | No change |

14

TABLE 2

| Optimization of Temperature, Time and Solvent Volume | | | | |
|--|-------------------|-------------|----------|--------------------|
| Amount | Amount Water (mL) | Temp (° C.) | Time (h) | Results/conclusion |
| 10 g | 50 | 75 | 6 | Mix |
| 10 g | 50 | 75 | 24 | Polymorph B |
| 10 g | 100 | 70 | 6 | Polymorph B |
| 10 g | 100 | 70 | 14 | Polymorph B |
| 10 g | 100 | 70 | 21 | Polymorph B |
| 10 g | 100 | 75 | 6 | Polymorph B |
| 10 g | 100 | 75 | 24 | Polymorph B |
| 10 g | 100 | 75 | 6 | Polymorph B |
| 10 g | 100 | 75 | 19 | Polymorph B |
| 10 g | 100 | 75 | 14 | Polymorph B |
| 10 g | 100 | 75 | 24 | Polymorph B |
| 5 g | 100 | 75 | 18 | Polymorph B |
| 10 g | 100 | 80 | 6 | Polymorph B |
| 10 g | 100 | 80 | 20 | Polymorph B |
| 10 g | 200 | 45 | 6 | Polymorph B + E |
| 10 g | 200 | 45 | 24 | Polymorph E |
| 10 g | 200 | 60 | 48 | Polymorph B |
| 10 g | 200 | 75 | 6 | Mix |
| 10 g | 200 | 75 | 24 | Polymorph B |
| 10 g | 200 | 75 | 13 | Polymorph B |
| 10 g | 200 | 75 | 24 | Polymorph B |

Optimum conditions were determined to be 10 volumes of solvent (H₂O), 70-80° C. for 6-24 hours.

TABLE 3

| Holding Time | | | | |
|--------------|-----------------------------|------------------|---------------------|--------------------|
| Amount | Reaction Conditions | Holding Time (h) | Holding Temp (° C.) | Results/Conclusion |
| 5 g | Water, 70-75° C., 24 h | 24 | 23-25 | Polymorph B |
| 1 g | Water, 70-75° C., 24 h | 48 | 23-25 | Polymorph E |
| Polymorph B | | | | |
| 2 g | Water, 40 mL | 16 | 23-25 | Polymorph E |
| 150 g | Water, 3.0 L | 24 | 23-25 | Polymorph E |
| 150 g | Water, 3.0 L | 48 | 23-25 | Polymorph E |
| 10 g | Water, 100 mL, 24 h, 75° C. | 18 | 23-25 | Polymorph B |
| 10 g | Water, 100 mL, 24 h, 75° C. | 18 | 40 | Polymorph B |
| 10 g | Water, 200 mL, 24 h, 75° C. | 14 | -5 | Mix |
| 10 g | Water, 200 mL, 24 h, 75° C. | 14 | 23-25 | Polymorph E |
| 10 g | Water, 200 mL, 24 h, 75° C. | 14 | 40 | Mix |
| 10 g | Water, 100 mL, 24 h, 75° C. | 21 | 23-25 | Polymorph E |
| 10 g | Water, 100 mL, 24 h, 75° C. | 21 | 40 | Mix |
| 10 g | Water, 100 mL, 14 h, 75° C. | 2 | 23-25 | Mix |

Holding time gave mixed results and it was determined that the material should be filtered at 60-65° C. and the material washed with 0.5 volume of warm (50-60° C.) water.

TABLE 4

| Scale-up Experiments | | | | |
|----------------------|------------------|-------------|----------|--------------------|
| Amount | Amount Water (L) | Temp (° C.) | Time (h) | Results/Conclusion |
| 100 g | 1.0 | 75 | 6 | Polymorph B |
| 100 g | 1.0 | 75 | 22 | Polymorph B |

US 7,465,800 B2

15

TABLE 4-continued

| Scale-up Experiments | | | | |
|----------------------|------------------|-------------|----------|---------------------|
| Amount | Amount Water (L) | Temp (° C.) | Time (h) | Results/ Conclusion |
| 100 g | 1.0 | 75 | 6 | Polymorph B |
| 100 g | 1.0 | 75 | 24 | Polymorph B |
| 100 g | 1.0 | 75 | 6 | Polymorph B |
| 100 g | 1.0 | 75 | 22 | Polymorph B |

TABLE 5

| Drying Studies | | | | | |
|----------------|-----------------|--------------------|----------------|---------|---------------------|
| Amount | Drying Time (h) | Drying Temp (° C.) | Vacuum (mm Hg) | KF§ (%) | Results/ Conclusion |
| 100 g | 0 | — | — | 3.690 | Polymorph B |
| 100 g | 3 | 30 | 152 | 3.452 | Polymorph B |
| 100 g | 8 | 30 | 152 | 3.599 | Polymorph B |
| 100 g | 0 | — | — | 3.917 | Polymorph B |
| 100 g | 5 | 40 | 152 | 3.482 | Polymorph B |
| 100 g | 22 | 40 | 152 | 3.516 | Polymorph B |
| 100 g | 3 | 40 | 152 | 3.67 | Polymorph B |
| 100 g | 22 | 40 | 152 | 3.55 | Polymorph B |

* Reaction Conditions: Water 1 L, 75° C., 22-24 h;
 § Average of 2 runs.

Drying studies determined that the material should be dried at 35-40° C., 125-152 mm Hg for 3 to 22 h or until the water content reaches $\leq 4\%$ w/w.

For a large scale preparation of polymorph E (5222-152-B), a 5-L round bottom flask was charged with 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione (150 g, 0.579 mol) and water (3000 mL, 20 vol). The mixture was mechanically stirred at room temperature (23-25° C.) for 48 h under nitrogen atmosphere.

Samples were taken after 24 h and 48 h before the mixture was filtered and air-dried on the filter for 1 h. The material was transferred to a drying tray and dried at room temperature (23-25° C.) for 24 h. KF analysis on the dried material showed water content of 11.9%. The material was submitted for XRPD, TGA, DSC and HPLC analysis. Analysis showed the material was pure polymorph E.

For a large scale preparation of polymorph B (5274-104), a 2 L-3-necked round bottom flask was charged with 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione (polymorph mixture, 100 g, 0.386 mol) and water (1000 mL, 10.0 vol). The mixture was heated to 75° C. over approximately 30 minutes with mechanical stirring under nitrogen atmosphere.

Samples were taken after 6 h and 24 h before the mixture was allowed to cool to 60-65° C., filtered and the material washed with warm (50-60° C.) water (50 mL, 0.5 vol). The material was transferred to a drying tray and dried at 30° C., 152 mm Hg for 8 h. KF analysis on the dried material showed water content of 3.6%. After grinding the material was submitted for XRPD, TGA, DSC and HPLC analysis. Analysis showed the material was pure polymorph B. The results of the analyses are shown in FIGS. 32-46.

6.3 X-Ray Powder Diffraction Measurements

X-ray powder diffraction analyses were carried out on a Shimadzu XRD-6000 X-ray powder diffractometer using Cu K α radiation. The instrument is equipped with a fine-focus X-ray tube. The tube voltage and amperage were set at 40 kV and 40 mA, respectively. The divergence and scattering slits were set at 1° and the receiving slit was set at 0.15 mm.

16

Diffraction radiation was detected by a NaI scintillation detector. A theta-two theta continuous scan at 3°/min (0.4 sec/0.02° step) from 2.5 degrees 2 θ to 40 degrees 2 θ was used. A silicon standard was analyzed each day to check the instrument alignment.

X-ray powder diffraction analyses were also carried out using Cu K α radiation on an Inel XRG-3000 diffractometer equipped with a curved position-sensitive detector. Data were collected in real time over a theta-two theta range of 120° at a resolution of 0.030°. The tube voltage and current were 40 kV and 30 mA, respectively. A silicon standard was analyzed each day to check for instrument alignment. Only the region between 2.5 and 40 degrees 2 θ is shown in the figures.

6.4 Thermal Analysis

TG analyses were carried out on a TA Instrument TGA 2050 or 2950. The calibration standards were nickel and alumel. Approximately 5 mg of sample was placed on a pan, accurately weighed, and inserted into the TG furnace. The samples were heated in nitrogen at a rate of 10° C./min, up to a final temperature of 300 or 350° C.

DSC data were obtained on a TA 2920 instrument. The calibration standard was indium. Approximately 2-5 mg samples were placed into a DSC pan and the weight accurately recorded. Crimped pans with one pinhole were used for analysis and the samples were heated under nitrogen at a rate of 10° C./min, up to a final temperature of 350° C.

Hot-stage microscopy was carried out using a Kofler hot stage mounted on a Leica Microscope. The instrument was calibrated using USP standards.

A TA Instruments TGA 2050 interfaced with a Nicolet model 560 Fourier transform IR spectrophotometer, equipped with a globar source, XT/KBr beamsplitter, and deuterated triglycine sulfate (DTGS) detector, was utilized for TG-IR experiments. The IR spectrometer was wavelength calibrated with polystyrene on the day of use while the TG was temperature and weight calibrated biweekly, using indium for the temperature calibration. A sample of approximately 10 mg of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione was weighed into an aluminum pan and heated from 25 to 30° C. to 200° C. at a rate of 20° C./min with a helium purge. IR spectra were obtained in series, with each spectrum representing 32 co-added scans at a resolution of 4 cm⁻¹. Spectra were collected with a 17-second repeat time. TG/IR analysis data are presented as Gram-Schmidt plots and IR spectra linked to the time. Gram-Schmidt plots show total IR intensity vs. time; hence, the volatiles can be identified at each time point. They also show when the volatiles are detected. From the Gram-Schmidt plots, time points were selected and the IR spectra of these time points are presented in the stacked linked spectra. Each spectrum identifies volatiles evolving at that time point. Volatiles were identified from a search of the HR Nicolet TGA vapor phase spectral library. The library match results are also presented to show the identified vapor.

6.5 Spectroscopy Measurements

Raman spectra were acquired on a Nicolet model 750 Fourier transform Raman spectrometer utilizing an excitation wavelength of 1064 nm and approximately 0.5 W of Nd:YAG laser power. The spectra represent 128 to 256 co-added scans acquired at 4 cm⁻¹ resolution. The samples were prepared for analysis by placing the material in a sample holder and positioning this in the spectrometer. The spectrometer was wavelength calibrated using sulfur and cyclohexane at the time of use.

The mid-IR spectra were acquired on a Nicolet model 860 Fourier transform IR spectrophotometer equipped with a glo-

bar source XT/KBr beamsplitter and a deuterated triglycine sulfate (DTGS) detector. A Spectra-Tech, Inc. diffuse reflectance accessory was utilized for sampling. Each spectrum represents 128 co-added scans at a spectral resolution of 4 cm^{-1} . A background data set was acquired with an alignment mirror in place. A single beam sample data set was then acquired. Subsequently, a $\log 1/R$ (where R =reflectance) spectrum was acquired by rationing the two data sets against each other. The spectrophotometer was calibrated (wavelength) with polystyrene at the time of use.

6.6 Moisture Sorption/Desorption Measurements

Moisture sorption/desorption data were collected on a VTI SGA-100 moisture balance system. For sorption isotherms, a sorption range of 5 to 95% relative humidity (RH) and a desorption range of 95 to 5% RH in 10% RH increments was used for analysis. The sample was not dried prior to analysis. Equilibrium criteria used for analysis were less than 0.0100 weight percent change in 5 minutes with a maximum equilibration time of 3 hours if the weight criterion was not met. Data were not corrected for the initial moisture content of the samples.

6.7 Solution Proton NMR Measurements

NMR spectra not previously reported were collected at SSCI, Inc, 3065 Kent Avenue, West Lafayette, Ind. Solution phase ^1H NMR spectra were acquired at ambient temperature on a Bruker model AM spectrometer. The ^1H NMR spectrum represents 128 co-added transients collected with a 4 μsec pulse and a relaxation delay time of 5 seconds. The free induction decay (FID) was exponentially multiplied with a 0.1 Hz Lorentzian line broadening factor to improve the signal-to-noise ratio. The NMR spectrum was processed utilizing GRAMS software, version 5.24. Samples were dissolved in dimethyl sulfoxide- d_6 .

The scope of this invention can be understood with reference to the appended claims.

6.8 Intrinsic Dissolution and Solubility Studies

Intrinsic dissolution experiments were conducted on Form A (anhydrous), Form B (hemihydrate), and Form E (dihydrate) of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Equilibrium solubility experiments were conducted on Forms A and B. Aliquots were analyzed by ultraviolet-visible spectrophotometry, and the solids remaining from each experiment were analyzed by X-ray powder diffraction (XRPD).

6.8.1 Experimental

6.8.1.1 Dissolution

Dissolution experiments were carried out in a VanKel VK6010-8 dissolution apparatus equipped with a VK650A heater/circulator. An intrinsic dissolution apparatus (Woods apparatus) was used. Samples were compressed at 1.5 metric tons (1000 psi) for 1 min using the Woods apparatus in a hydraulic press, giving a sample surface of 0.50 cm^2 . A dissolution medium consisting of 900 mL HCl buffer, pH 1.8, with 1% sodium lauryl sulfate, was used for each experiment. The medium was degassed by vacuum filtration through a $0.22\text{-}\mu\text{m}$ nylon filter disk and maintained at 37°C . The apparatus was rotated at 50 rpm for each experiment. Aliquots were filtered immediately using $0.2\text{-}\mu\text{m}$ nylon syringe filters. In some cases, the undissolved solids were recovered and analyzed by X-ray powder diffraction (XRPD).

6.8.1.2 Solubility

Equilibrium solubility experiments were conducted in a 100-mL, three-neck, round-bottom flask immersed in a constant temperature oil bath maintained at 25°C . A solid sample of 400-450 mg was stirred in 50 mL of dissolution medium (HCl buffer, pH 1.8, with 1% sodium lauryl sulfate) using a mechanical stir rod. Aliquots were filtered using $0.2\text{-}\mu\text{m}$

nylon syringe filters and immediately diluted $1\text{ mL}\rightarrow 50\text{ mL}$, then $5\text{ mL}\rightarrow 25\text{ mL}$ with dissolution medium in Class A glassware, a final dilution factor of 250.

6.8.1.3 UV-Vis Spectrophotometry

Dissolution and solubility samples solutions were analyzed by a Beckman DU 640 single-beam spectrophotometer. A 1.000-cm quartz cuvette and an analysis wavelength of 228.40 nm were utilized. The detector was zeroed with a cuvette filled with dissolution medium.

6.8.1.4 X-Ray Powder Diffraction

XRPD analyses were carried out on a Shimadzu XRD-6000 X-ray powder diffractometer using $\text{Cu K}\alpha$ radiation. The instrument is equipped with a fine focus X-ray tube. The tube power and amperage were set at 40 kV and 40 mA, respectively. The divergence and scattering slits were set at 1° and the receiving slit was set at 0.15 mm. Diffracted radiation was detected by a NaI scintillation detector. A theta-two theta continuous scan at $3^\circ/\text{min}$ ($0.4\text{ sec}/0.02^\circ$ step) from 2.5 to $40^\circ 2\theta$ was used. A silicon standard was analyzed each day to check the instrument alignment. Samples were packed in an aluminum holder with silicon insert.

6.8.2 Results

The results of these solubility and intrinsic studies are summarized in Table 6. Both the solubility and dissolution experiments were conducted in a medium of HCl buffer, pH 1.8, containing 1% sodium lauryl sulfate. Form A was found to be unstable in the medium, converting to Form E. The solubilities of Forms A, B, and E were estimated to be 6.2, 5.8, and 4.7 mg/mL, respectively. The dissolution rates of Forms A, B, and E were estimated to be 0.35, 0.34, and 0.23 mg/mL, respectively.

6.8.2.1 UV-Vis Spectrophotometry Method Development

A UV-Vis scan of the dissolution medium (blanked with an empty cuvette) was done to identify any interfering peaks. A small peak at 225 nm was present as shown in FIG. 47.

Solutions of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione at varying concentrations were analyzed by UV-Vis spectrophotometry. A preliminary scan of a 1.0 mg/mL solution was done, with the instrument blanked with dissolution medium. The solution was highly absorbing and noisy from 200-280 nm, making dilution necessary.

A 0.04 mg/mL solution of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione was then scanned from 200-300 nm. The plot was still noisy between 200 and 230 nm as shown in FIG. 48. The sample was further diluted to 0.008 mg/mL. A wavelength scan of 200-350 nm for this sample showed a peak at 228.4 nm with no interference, as shown in FIG. 49. Therefore, a wavelength of 228.4 was chosen for analysis of the solubility and dissolution samples.

A six-point calibration curve was generated with standards of the following concentrations: 0.001 mg/mL, 0.002 mg/mL, 0.005 mg/mL, 0.010 mg/mL, 0.015 mg/mL, and 0.020 mg/mL (Notebook 569-90). A linearity coefficient of $R^2=0.9999$ was obtained as shown in FIG. 50.

6.8.2.2 Solubility

A sample consisting of 449.4 mg Form A was slurried in dissolution medium. Particle size was not controlled. Aliquots were taken at 7, 15, 30, 60, 90, and 150 min. The concentration reached 6.0 mg/mL by the first time point. The highest concentration reached was 6.2 mg/mL, at 30 min. From that point the concentration decreased, reaching 4.7 mg/mL at 150 min as in FIG. 51. The solids remaining at the final time point were analyzed by XRPD and found to be Form E as shown in Table 7. No peaks attributed to Form A can be seen in the pattern. Since the concentration did not plateau at 4.7 mg/mL, the solubility of Form E may be lower than that.

US 7,465,800 B2

19

A sample consisting of 401.4 mg Form B was slurried in dissolution medium. Particle size was not controlled. Aliquots were taken at 7, 15, 30, 60, 90, 180, 420, and 650 min. Form B dissolved much more slowly than Form A, reaching 3.3 mg/mL in 90 min. The concentration stabilized at 5.6-5.7 mg/mL at the final three time points as in FIG. 52. The remaining solids were shown to be Form B as in Table 7, suggesting Form B has good stability in water.

A summary of the solubilities is given in Table 6. The amounts dissolved at each time point are shown in Tables 8 and 9.

TABLE 6

| <u>Summary of Results</u> | | | | |
|---------------------------|------------|--------------------------|--------------------------|------------------------------------|
| Form | Solubility | Intrinsic Dissolution #1 | Intrinsic Dissolution #2 | Average Intrinsic Dissolution Rate |
| Form A | 6.2 mg/mL | 0.35 | 0.22 ^a | 0.29 ^a |
| Form B | 5.8 mg/mL | 0.35 | 0.32 | 0.34 |
| Form E | 4.7 mg/mL | 0.21 | 0.25 | 0.23 |

^aThe Form A dissolution experiment #2 may have converted to Form E on the surface of the disk, skewing the average rate lower.

TABLE 7

| <u>Experimental Details</u> | |
|-----------------------------|------------|
| Experiment | Final Form |
| Pressed Form A | A |
| Pressed Form B | B |
| Form A Solubility | E |
| Form B Solubility | B |
| Form A Dissolution | — |
| Form A Dissolution | A |
| Form B Dissolution | — |
| Form B Dissolution | B |
| Form E Dissolution | E |
| Form E Dissolution | — |

TABLE 8

| <u>Form A Solubility</u> | |
|--------------------------|-----------------------|
| Time Point (min) | Concentration (mg/mL) |
| 7 | 6.00 |
| 15 | 6.11 |
| 30 | 6.16 |
| 60 | 6.10 |
| 90 | 5.46 |
| 150 | 4.73 |

TABLE 9

| <u>Form B Solubility</u> | |
|--------------------------|-----------------------|
| Time Point (min) | Concentration (mg/mL) |
| 7 | 1.63 |
| 15 | 2.14 |
| 30 | 2.33 |
| 60 | 2.94 |
| 90 | 3.34 |
| 180 | 5.67 |
| 420 | 5.76 |
| 650 | 5.61 |

20

6.8.2.3 Intrinsic Dissolution

Approximately 200 mg each of Forms A and B were compressed into disks in the Woods apparatus using 2 metric tons of pressure. The samples were subsequently scraped out, ground gently, and analyzed by XRPD. The study showed that compression and grinding does not cause a form change in either case. (See Table 7).

Two preliminary dissolution runs were performed. The disks fractured to some extent in both experiments, compromising the requirement of constant surface area.

The first experiment of intrinsic dissolution that strictly followed the USP chapter on intrinsic dissolution utilized approximately 150 mg each of Forms A and B. Seven aliquots, beginning at 5 min and ending at 90 min, were taken to maintain sink conditions. The experiment resulted in linear dissolution profiles, with a rate of 0.35 mg per cm² per minute for both forms. The Form E experiment was done later under the same conditions and added to the graph for comparison. (See FIG. 53). The Form E dissolution rate was 0.21 mg per cm² per minute, significantly lower than the dissolution rate of Forms A and B. This is in line with expectations based on the solubility data. The crystal form of the remaining solids did not change in any case.

The second experiment utilized approximately 250 mg each of Forms A and B. The Form E experiment (135 mg) was done later and added to the graph for comparison. (See FIG. 54). Nine aliquots were taken, beginning at 5 min and ending at 150 min. The dissolution rates were 0.22, 0.32, and 0.25 mg per cm² per minute, respectively, for Forms A, B, and E. The dissolution rate for Form A in this experiment was low, while the rates for Forms B and E were similar to those found in the first experiment. It is believed that in this case, a thin layer of the Form A sample disk may have converted to Form E upon exposure to water. This is supported by the evidence of rapid conversion of Form A to Form E in the solubility experiment. The diffraction pattern of the undissolved solids does not indicate a form change. However, the bulk of the sample disk is not exposed to water. Therefore, the true intrinsic dissolution rate of Form A is believed to be close to 0.35 mg per cm² per minute. An insufficient quantity of Form A was available to repeat the experiment.

A summary of the intrinsic dissolution rates is given in Table 6. The amounts dissolved at each time point are summarized in Tables 10 and 11.

TABLE 10

| <u>Intrinsic Dissolution Experiment #1 Results</u> | | | |
|--|---------------------|---------------------|---------------------|
| Time Point | Form A ^a | Form B ^a | Form E ^a |
| 5 min | 5.76 | 10.80 ^b | 2.70 |
| 10 min | 7.73 | 6.85 | 4.13 |
| 20 min | 11.31 | 10.25 | 6.96 |
| 30 min | 15.59 | 14.35 | 9.60 |
| 45 min | 21.98 | 20.57 | 12.57 |
| 60 min | 27.11 | 25.70 | 15.16 |
| 90 min | 34.17 | 34.34 | 20.82 |

^aResults are reported as Cumulative Amount Dissolved per Unit Area (mg/cm²)

^bThis date point not included in graph since the value is higher than the next two data points.

US 7,465,800 B2

21

TABLE 11

| Intrinsic Dissolution Experiment #2 Results | | | |
|---|---------------------|---------------------|---------------------|
| Time Point | Form A ^a | Form B ^a | Form E ^a |
| 5 min | 4.50 | 5.04 | 3.06 |
| 10 min | 5.22 | 6.12 | 4.31 |
| 20 min | 7.54 | 7.73 | 11.40 |
| 30 min | 11.46 | 12.72 | 11.93 |
| 45 min | 15.01 | 17.33 | 14.72 |
| 60 min | 18.38 | 21.93 | 18.52 |
| 90 min | 24.38 | 31.64 | 26.24 |
| 120 min | 30.35 | 41.31 | 33.56 |
| 150 min | 35.26 | 49.54 | 40.82 |

^aResults are reported as Cumulative Amount Dissolved per Unit Area (mg/cm²)

6.9 Analyses of Mixtures of Polymorphs

This invention encompasses mixtures of different polymorphs. For example, an X-ray diffraction analysis of one production sample yielded a pattern that contained two small peaks seen at approximately 12.6° and 25.8° 2θ in addition to those representative of Form B. In order to determine the composition of that sample, the following steps were performed:

- 1) Matching of the new production pattern to known forms along with common pharmaceutical excipients and contaminants;
- 2) Cluster analysis of the additional peaks to identify if any unknown phase is mixed with the original Form B;
- 3) Harmonic analysis of the additional peaks to identify if any preferred orientation may be present or if any changes in the crystal habit may have occurred; and
- 4) Indexing of the unit cells for both Form B and the new production sample to identify any possible crystallographic relationships.

Based on these tests, which can be adapted for the analysis of any mixture of polymorphs, it was determined that the sample contained a mixture of polymorph forms B and E.

6.10 Dosage Form

Table 12 illustrates a batch formulation and single dosage formulation for a 25 mg single dose unit of a polymorphic form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione.

TABLE 12

| Formulation for a 25 mg capsule | | | |
|--|-------------------|----------------------|---------------------|
| Material | Percent By Weight | Quantity (mg/tablet) | Quantity (kg/batch) |
| Polymorphic Form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione | 40.0% | 25 mg | 16.80 kg |
| Pregelatinized Corn Starch, NF | 59.5% | 37.2 mg | 24.99 kg |
| Magnesium Stearate | 0.5% | 0.31 mg | 0.21 kg |
| Total | 100.0% | 62.5 mg | 42.00 kg |

22

The pregelatinized corn starch (SPRESS B-820) and polymorphic form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione components are passed through a screen (i.e., a 710 μm screen) and then loaded into a Diffusion Mixer with a baffle insert and blended for about 15 minutes. The magnesium stearate is passed through a screen (i.e., a 210 μm screen) and added to the Diffusion Mixer. The blend is then encapsulated in capsules using a Dosator type capsule filling machine.

The entire scope of this invention is not limited by the specific examples described herein, but is more readily understood with reference to the appended claims.

What is claimed is:

1. Crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione hemihydrate.

2. The hemihydrate of claim 1 having an X-ray powder diffraction pattern comprising peaks at approximately 16, 22 and 27 degrees 2θ.

3. The hemihydrate of claim 2 wherein the pattern further comprises a peak at approximately 18 degrees 2θ.

4. The hemihydrate of claim 1 having an X-ray powder diffraction pattern comprising peaks at 15.8, 22.2 and 26.7 degrees 2θ.

5. The hemihydrate of claim 4 wherein the pattern further comprises a peak at 18.2 degrees 2θ.

6. The hemihydrate of claim 1 having an X-ray powder diffraction pattern corresponding to the representative X-ray powder diffraction patterns depicted in FIG. 6, FIG. 32, FIG. 33 and FIG. 34.

7. The hemihydrate of claim 1 having a differential scanning calorimetry thermogram comprising an endotherm with a maximum at about 268° C.

8. The hemihydrate of claim 7 wherein the thermogram further comprises an endotherm corresponding to dehydration.

9. The hemihydrate of claim 1 having a differential scanning calorimetry thermogram corresponding to the representative differential scanning calorimetry thermograms depicted in FIG. 9, FIG. 42, FIG. 43 and FIG. 44.

10. The hemihydrate of claim 1 having between approximately 0.46 and approximately 0.59 moles of water per mole of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione.

11. The hemihydrate of claim 1 having a thermogravimetric analysis thermogram comprising a weight loss of between about 3.1% and about 4.0% when heated from about 30° C. to about 175° C.

12. The hemihydrate of claim 1 having an infrared spectrum comprising peaks at approximately 3513 and 1960 cm⁻¹.

13. The hemihydrate of claim 1 having an infrared spectrum corresponding to the representative infrared spectrum depicted in FIG. 7.

14. The hemihydrate of claim 1 having a Raman spectrum corresponding to the representative Raman spectrum depicted in FIG. 8.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 7,465,800 B2
APPLICATION NO. : 10/934863
DATED : December 16, 2008
INVENTOR(S) : Jaworsky et al.

Page 1 of 11

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Please replace Drawing Sheets 15, 26, 31, 32, 33, 35, 36, 38, 39, and 40 (of 48) with the Replacement Drawing Sheets on the following pages of this Certificate of Correction:

Page 2 of 11: Replacement Drawing Sheet 15 of 48 (Fig. 15);

Page 3 of 11: Replacement Drawing Sheet 26 of 48 (Fig. 26);

Page 4 of 11: Replacement Drawing Sheet 31 of 48 (Fig. 31);

Page 5 of 11: Replacement Drawing Sheet 32 of 48 (Figs. 32 and 33);

Page 6 of 11: Replacement Drawing Sheet 33 of 48 (Figs. 34 and 35);

Page 7 of 11: Replacement Drawing Sheet 35 of 48 (Figs. 37 and 38);


Page 8 of 11: Replacement Drawing Sheet 36 of 48 (Figs. 39 and 40);

Page 9 of 11: Replacement Drawing Sheet 38 of 48 (Figs. 42 and 43);

Page 10 of 11: Replacement Drawing Sheet 39 of 48 (Figs. 44 and 45);

Page 11 of 11: Replacement Drawing Sheet 40 of 48 (Fig. 46).

Signed and Sealed this
Nineteenth Day of April, 2011



David J. Kappos
Director of the United States Patent and Trademark Office




FIG.15




FIG.26




FIG.31

XRPD PATTERN OF POLYMORPH B

File: Process 5274-104-B

Date: 06-04-04 16:10 Step: 0.020° Cnt Time: 0.500 Sec.

Range: 1.50-40.00 (Deg) Step Scan Rate: 2.40 Deg/min.




FIG.32

XRPD PATTERN OF POLYMORPH B

File: Process 5274-100-C

Date: 06-02-04 16:11 Step: 0.020° Cnt Time: 0.500 Sec.

Range: 1.50-40.00 (Deg) Step Scan Rate: 2.40 Deg/min.




FIG.33

XRPD PATTERN OF POLYMORPH B

File: Process 5222-157-C

Date: 06/04/04 15:07 Step: 0.020° Cnt Time: 0.500 Sec.

Range: 1.50-40.00 (Deg) Step Scan Rate: 2.40 Deg/min.




FIG.34

XRPD PATTERN OF POLYMORPH E

File: Process 5222-152-B Form E

Date: 05/21/04 10:46 Step: 0.020° Cnt Time: 0.500 Sec.

Range: 1.50-40.00 (Deg) Step Scan Rate: 2.40 Deg/min.




FIG.35

TGA CURVE OF POLYMORPH B
Sample: 5274-104-B
Size 7.8320mg
Method: Ramp
TGA
Run Date: 09-Jun-04 17:04
Instrument: TGA 0.500 V5.3 Build 171




FIG.37

TGA CURVE OF POLYMORPH B
Sample: 5274-100-C
Size 10.9430mg
Method: Ramp
Comment: Evotec Batch 1675C H2O
TGA
Run Date: 03-Jun-04 11:20
Instrument: TGA 0.500 V5.3 Build 171




FIG.38

TGA CURVE OF POLYMORPH B
TGA
Sample: 5222-157-C
Size: 24.4610mg
Method: Ramp
Comment: Evotec Batch 17
Run Date: 07-Jun-04 09:46
Instrument: TGA 0.500 V5.3 Build 171




FIG.39

TGA CURVE OF POLYMORPH E
TGA
Sample: 5222-152-B
Size: 11.3850mg
Method: Ramp
Comment: Form E
Run Date: 21-May-04 09:34
Instrument: TGA 0.500 V5.3 Build 171




FIG.40

DSC OF POLYMORPH B
DSC

Sample: 5274-104-B
 Size: 0.0000mg
 Method: Cell constant calibration
 Comment: OAI batch 15 22b

Run Date: 07-Jun-04 11:30
 Instrument: DSC Q 1000 V7.3 Build 249




FIG.42

DSC OF POLYMORPH B
DSC

Sample: 5274-100-C
 Size: 0.0000mg
 Method: Cell constant calibration
 Comment: Evotec Batch 1675C 10vol H2O 24h 3h dry

Run Date: 02-Jun-04 17:01
 Instrument: DSC Q 1000 V7.3 Build 249




FIG.43

DSC OF POLYMORPH B
DSC

Sample: 5222-157-C
 Size: 0.0000mg
 Method: Cell constant calibration
 Comment: OAI batch 17

Run Date: 07-Jun-04 09:45
 Instrument: DSC Q 1000 V7.3 Build 249




FIG.44

DSC OF POLYMORPH E
DSC

Sample: 5222-152-B
 Size: 3.4500mg
 Method: Cell constant calibration
 Comment: Evotec batch 15 Form E

Run Date: 21-May-04 09:32
 Instrument: DSC Q 1000 V7.3 Build 249




FIG.45

DSC OF POLYMORPH MIXTURE

DSC

Sample: 5222-161-A

Size: 0.0000mg

Method: Cell constant calibration

Comment: OAI batch 15

Run Date: 11-Jun-04 12:44

Instrument: DSC Q 1000 V7.3 Build 249




FIG.46

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 7,465,800 B2
APPLICATION NO. : 10/934863
DATED : December 16, 2008
INVENTOR(S) : Jaworsky et al.

Page 1 of 1


It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

On the Title Page:

The first or sole Notice should read --

Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b)
by 966 days.

Signed and Sealed this
Fifteenth Day of May, 2012

A handwritten signature in black ink that reads "David J. Kappos". The signature is written in a cursive, slightly slanted style.

David J. Kappos
Director of the United States Patent and Trademark Office

EXHIBIT B



US007855217B2

(12) **United States Patent**
Jaworsky et al.

(10) **Patent No.:** **US 7,855,217 B2**
(45) **Date of Patent:** ***Dec. 21, 2010**

(54) **POLYMORPHIC FORMS OF 3-(4-AMINO-1-OXO-1,3 DIHYDRO-ISOINDOL-2-YL)-PIPERIDINE-2,6-DIONE**

(75) Inventors: **Markian S. Jaworsky**, Hopewell, NJ (US); **Roger Shen-Chu Chen**, Edison, NJ (US); **George W. Muller**, Bridgewater, NJ (US)

(73) Assignee: **Celgene Corporation**, Summit, NJ (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 82 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **12/335,395**
(22) Filed: **Dec. 15, 2008**

(65) **Prior Publication Data**
US 2009/0149500 A1 Jun. 11, 2009

Related U.S. Application Data
(62) Division of application No. 10/934,863, filed on Sep. 3, 2004, now Pat. No. 7,465,800.
(60) Provisional application No. 60/499,723, filed on Sep. 4, 2003.

(51) **Int. Cl.**
A61K 31/454 (2006.01)
C07D 401/04 (2006.01)

(52) **U.S. Cl.** **514/323; 546/200**
(58) **Field of Classification Search** **514/323; 546/200**
See application file for complete search history.

(56) **References Cited**
U.S. PATENT DOCUMENTS
3,536,809 A 10/1970 Applezweig
3,598,123 A 8/1971 Zaffaroni
3,845,770 A 11/1974 Theeuwes et al.
3,916,899 A 11/1975 Theeuwes et al.
4,008,719 A 2/1977 Theeuwes et al.
4,810,643 A 3/1989 Souza
4,999,291 A 3/1991 Souza
5,059,595 A 10/1991 Le Grazie
5,073,543 A 12/1991 Marshall et al.
5,120,548 A 6/1992 McClelland et al.
5,229,496 A 7/1993 Deeley et al.
5,354,556 A 10/1994 Sparks et al.
5,385,901 A 1/1995 Kaplan et al.
5,391,485 A 2/1995 Deeley et al.
5,393,870 A 2/1995 Deeley et al.
5,528,823 A 6/1996 Rudy, Jr. et al.
5,580,755 A 12/1996 Souza
5,591,767 A 1/1997 Mohr et al.
5,593,990 A 1/1997 D'Amato
5,629,327 A 5/1997 D'Amato
5,635,517 A 6/1997 Muller et al.

5,639,476 A 6/1997 Oshlack et al.
5,674,533 A 10/1997 Santus et al.
5,698,579 A 12/1997 Muller
5,712,291 A 1/1998 D'Amato
5,731,325 A 3/1998 Andrulis, Jr. et al.
5,733,566 A 3/1998 Lewis
5,798,368 A 8/1998 Muller et al.
5,874,448 A 2/1999 Muller et al.
5,877,200 A 3/1999 Muller
5,929,117 A 7/1999 Muller et al.
5,955,476 A 9/1999 Muller et al.
6,020,358 A 2/2000 Muller et al.
6,071,948 A 6/2000 D'Amato
6,114,335 A 9/2000 D'Amato
6,114,355 A 9/2000 D'Amato
6,140,346 A 10/2000 Andrulis, Jr. et al.
6,235,756 B1 5/2001 D'Amato
6,281,230 B1 8/2001 Muller et al.
6,316,471 B1 11/2001 Muller et al.
6,326,388 B1 12/2001 Man et al.
6,335,349 B1 1/2002 Muller et al.
6,380,239 B1 4/2002 Muller et al.
6,395,754 B1 5/2002 Muller et al.
6,403,613 B1 6/2002 Man et al.
6,420,414 B1 7/2002 D'Amato
6,458,810 B1 10/2002 Muller et al.
6,469,045 B1 10/2002 D'Amato
6,476,052 B1 11/2002 Muller et al.
6,518,298 B2 2/2003 Green et al.
6,555,554 B2 4/2003 Muller et al.

(Continued)
FOREIGN PATENT DOCUMENTS

JP A H10-53576 2/1998

(Continued)
OTHER PUBLICATIONS

Dimartino et al., 1997, "Preparation and physical characterization of forms II and III of paracetamol," J. Thermal Analysis 48:447-458.

(Continued)

Primary Examiner—Celia Chang
(74) Attorney, Agent, or Firm—Jones Day

(57) **ABSTRACT**

Polymorphic forms of 3-(4-amino-1-oxo-1,3 dihydro-isoin-dol-2-yl)-piperidine-2,6-dione are disclosed. Compositions comprising the polymorphic forms, methods of making the polymorphic forms and methods of their use are also dis-closed.

10 Claims, 48 Drawing Sheets

US 7,855,217 B2

Page 2

U.S. PATENT DOCUMENTS

| | | | |
|--------------|----|---------|-------------------|
| 7,112,602 | B2 | 9/2006 | D'Amato et al. |
| 7,465,800 | B2 | 12/2008 | Jaworsky et al. |
| 2001/0018445 | A1 | 8/2001 | Huang et al. |
| 2001/0056114 | A1 | 12/2001 | D'Amato |
| 2002/0035090 | A1 | 3/2002 | Zeldis et al. |
| 2002/0045643 | A1 | 4/2002 | Muller et al. |
| 2002/0052398 | A1 | 5/2002 | D'Amato |
| 2002/0054899 | A1 | 5/2002 | Zeldis |
| 2002/0061923 | A1 | 5/2002 | D'Amato |
| 2002/0128228 | A1 | 9/2002 | Hwu |
| 2002/0161023 | A1 | 10/2002 | D'Amato |
| 2002/0173658 | A1 | 11/2002 | Muller et al. |
| 2002/0183360 | A1 | 12/2002 | Muller et al. |
| 2003/0013739 | A1 | 1/2003 | Masferrer et al. |
| 2003/0028028 | A1 | 2/2003 | Man et al. |
| 2003/0045552 | A1 | 3/2003 | Robarge et al. |
| 2003/0069428 | A1 | 4/2003 | Muller et al. |
| 2003/0096841 | A1 | 5/2003 | Robarge et al. |
| 2003/0139451 | A1 | 7/2003 | Shah et al. |
| 2003/0144325 | A1 | 7/2003 | Muller et al. |
| 2003/0181428 | A1 | 9/2003 | Green et al. |
| 2003/0187024 | A1 | 10/2003 | D'Amato |
| 2003/0191098 | A1 | 10/2003 | D'Amato |
| 2003/0235909 | A1 | 12/2003 | Hariri et al. |
| 2004/0029832 | A1 | 2/2004 | Zeldis |
| 2004/0077685 | A1 | 4/2004 | Figg et al. |
| 2004/0077686 | A1 | 4/2004 | Dannenberg et al. |
| 2004/0087546 | A1 | 5/2004 | Zeldis |
| 2004/0091455 | A1 | 5/2004 | Zeldis |
| 2004/0122052 | A1 | 6/2004 | Muller et al. |
| 2004/0220144 | A1 | 11/2004 | Zeldis |
| 2004/0266809 | A1 | 12/2004 | Emanuel et al. |
| 2005/0203142 | A1 | 9/2005 | Zeldis |
| 2006/0052609 | A1 | 3/2006 | Muller et al. |
| 2008/0064876 | A1 | 3/2008 | Muller et al. |
| 2008/0132541 | A1 | 6/2008 | Zeldis et al. |
| 2009/0062343 | A1 | 3/2009 | Jaworsky et al. |
| 2009/0149449 | A1 | 6/2009 | Liu et al. |
| 2009/0149499 | A1 | 6/2009 | Jaworsky et al. |
| 2009/0176832 | A1 | 7/2009 | Jaworsky et al. |
| 2009/0187023 | A1 | 7/2009 | Jaworsky et al. |

FOREIGN PATENT DOCUMENTS

| | | | |
|----|----------------|----------------|------------|
| JP | A | 2001-503384 | 3/2001 |
| WO | | WO97/46526 | A1 12/1997 |
| WO | | WO 98/03502 | 1/1998 |
| WO | | WO98/03502 | A1 1/1998 |
| WO | | WO 98/54170 | 12/1998 |
| WO | | WO 01/70275 | 9/2001 |
| WO | | WO 01/87307 | 11/2001 |
| WO | | WO 02/059106 | 8/2002 |
| WO | | WO 02/064083 | 8/2002 |
| WO | PCT/US03/11578 | | 1/2003 |
| WO | | WO 03097052 | 11/2003 |
| WO | | WO02/26737 | A1 4/2004 |
| WO | | WO 2004103274 | 12/2004 |
| WO | | WO 2005023192 | 3/2005 |
| WO | | WO 2006/028964 | A1 3/2006 |
| WO | | WO 2007/136640 | A2 11/2007 |
| WO | | WO 2009/111948 | A1 9/2009 |
| WO | | WO 2009/114601 | A2 9/2009 |
| WO | | WO 2010/054833 | 5/2010 |
| WO | | WO 2010/056384 | 5/2010 |
| WO | | WO 2010/061209 | 6/2010 |

WO WO 2010/100476 9/2010

OTHER PUBLICATIONS

Jonsson, 1972, "Chemical structure and teratogenic properties. 3. A review of available data on structure-activity relationships and mechanism of action of thalidomide analogues," *Acta. Pharm. Suec.* 9(6):521-542.

Wilens et al., 1977, "Strategies in optical resolutions," *Tetrahedron* 33:2725-2736.

Wilens, 1972, *Tables of Resolving Agents and Optical Resolutions*, Eliel, ed., U. Notre Dame Press, Notre Dame, IN pp. 268.

Wolff, ed., 1995, *I Burger's Medicinal Chemistry and Drug Discovery*, 5th ed., pp. 172-178 and 949-982.

Corral et al., 1999, "Immunomodulation by thalidomide and thalidomide analogs," *Ann. Rheum. Dis.* 58(Supp. 1):1107-1113.

Grant, 1999, "Theory and Origin of Polymorphism," in *Polymorphism in Pharmaceutical Solids*, Minneapolis, MN, Ch. 1, pp. 1-10.

He, W., et al., 1993, 206th American Chemical Society, Chicago, IL; *Med. Chem.*, paper 216.

Lentzsch et al., 2002, "S-3-Amino-phthalimido-glutarimide inhibits angiogenesis and growth of B-cell neoplasias in mice," *Cancer Research* 62:2300-2305.

Muller et al., 1999, "Amino-substituted thalidomide analogs: Potent inhibitors of TNF- γ production," *Bioorg. Med. Chem. Lett.* 9:1625-1630.

Muller et al., 1998, "Thalidomide analogs and PDE4 inhibition," *Bioorg. Med. Chem. Lett.* 8:2669-2674.

Muller et al., 1996, "Structural modifications of thalidomide produce analogs with enhanced tumor necrosis factor inhibitory activity," *J. Med. Chem.* 39:3238-3240.

Anderson, "Moving disease biology from the laboratory to the clinic," *Seminars in Oncology*, 2002 29:17-20.

Barlogie et al., "High-dose therapy immunomodulatory drugs in multiple myeloma," *Seminars in Oncology*, 2002, 29 (6):26-33.

Barlogie et al., "Introduction: Thalidomide and the IMiDs in multiple myeloma," *Seminars in Hematology*, 2003, 40 (4):1-2.

Barlogie et al., "Total Therapy II (TTII) for newly diagnosed multiple myeloma (MM): preliminary data on feasibility and efficacy in the first 231 enrolled patients; comparison with predecessor trial total therapy I (TTI) (N=231)," *Blood, Abstract # 2857*, Dec. 7-11, 2001, *American Society of Hematology*.

Barlogie, "Thalidomide and CC-5013 in Multiple Myeloma: The University of Arkansas experience," *Seminars in Hematology*, 2003, 40 (4):33-38.

Bartlett et al., "Phase I study to determine the safety, tolerability and immunostimulatory activity of thalidomide analogue CC-5013 in patients with metastatic malignant melanoma and other advanced cancers," *British Journal of Cancer*, 2004, 90:955-961.

Bartlett et al., "The evolution of thalidomide and its IMiD derivatives as anticancer agents," *Nature Reviews Cancer*, 2004, 4 (4):1-9.

Battegay, "Angiogenesis: mechanistic insights, neovascular diseases, and therapeutic prospects," *J. Mol. Med.*, 1995, 73:333-346.

Baz et al., "Doxil (D), vincristine (V), reduced frequency dexamethasone (d) and revlimid (R) (DVd-R) results in a high response rate in patients with refractory multiple myeloma (RMM)," *Blood, Abstract # 2559*, *American Society of Hematology*, Dec. 10-13, 2005.

Brennen et al., "Thalidomide and analogues: current proposed mechanisms and therapeutic usage," *Clinical Prostate Cancer*, 2004, 3 (1):54-61.

Celgene Corporation, "Additional clinical data presented on Revimid™ in myelodysplastic syndromes at the American Society of Hematology 45th annual meeting," Press Release, Dec. 2003.

Celgene Corporation, "Blood reports Revimid™ has anti-tumor activity in patients with relapsed and refractory multiple myeloma," Press Release, Nov. 1, 2002.

Celgene Corporation, "Celgene advances immunomodulatory drug (IMiD™) clinical program," Press Release, Feb. 2000.

Celgene Corporation, "Celgene announces plans to stop phase III trials in melanoma due to lack of efficacy," Press Release, Apr. 2004.

US 7,855,217 B2

Page 3

- Celgene Corporation, "Celgene corporation advances ACTIMID™ (CC-4047) into phase II trial for prostate cancer," Press Release, Oct. 2003.
- Celgene Corporation, "Celgene Corporation announces fourth quarter and full year results for 2002," Press Release, Jan. 2003.
- Celgene Corporation, "Celgene Corporation announces third quarter results. THALOMID® (thalidomide) revenue increases 41% to \$30.5 million. Pivotal programs for THALOMID and REVIMID™ finalized. Peer-reviewed publications of THALOMID and REVIMID data. First JNK inhibitor advanced to Phase I clinical trial," Press Release, Oct. 2002.
- Celgene Corporation, "Celgene Corporation announces third quarter results. Thalomid® (thalidomide) sales increase 24%. Prescriptions up 50%. Enhanced S.T.E.P.S.® launched. Pilot d-MPH data presented," Press Release, Oct. 2001.
- Celgene Corporation, "Celgene Corporation receives orphan drug designation for Revimid™ for multiple myeloma," Press Release, Oct. 2001.
- Celgene Corporation, "Celgene corporation reports record operating performance in first quarter with strong revenue growth and profits," Press Release, Apr. 2004.
- Celgene Corporation, "Celgene corporation reports record operating performance in third quarter as total revenue increases 117% and profits rise," Press Release, Oct. 2003.
- Celgene Corporation, "Celgene corporation reports strong operating performance in second quarter as total sales increase 100 percent and profits rise," Press Release, Jul. 2003.
- Celgene Corporation, "Celgene corporation reviews 2003 achievements and announces 2004 financial outlook," Press Release, Jan. 2004.
- Celgene Corporation, "Celgene expands clinical development program for Revimid™. Five additional trials of Revimid initiated in hematological and solid tumor cancers," Press Release, Jun. 2002.
- Celgene Corporation, "Celgene provides update on clinical pipeline. Celgene Announces first target indication for ACTIMID™, CC-8490. SeICID™ program to advance based on results from Phase I/II trial of CC-1088. First JNK inhibitor successfully completes phase I trial," Press Release, Jan. 2003.
- Celgene Corporation, "Celgene receives fast track status from FDA for Revimid™ in multiple myeloma," Press Release, Feb. 2003.
- Celgene Corporation, "Celgene receives fast track status from FDA for Revimid™ in myelodysplastic syndromes," Press Release, Apr. 2003.
- Celgene Corporation, "Initial Phase I solid tumor data on Celgene's lead IMiD™, Revimid™," Press Release, Jun. 2001.
- Celgene Corporation, "New Revimid™ clinical data shows potential as novel approach to treating myelodysplastic syndromes (MDS)," Press Release, May 2003.
- Celgene Corporation, "Revlimid™ receives orphan drug designation from the European commission for multiple myeloma," Press Release, Feb. 2004.
- Celgene Corporation, "Revlimid™ receives orphan drug designation from the European commission for myelodysplastic syndromes," Press Release, Mar. 2004.
- Chaudhry, 1966, *Cancer Research*, "Effect of Prednisolone and Thalidomide on Induced Submandibular Gland Tumors in Hamster," 26(part 1):1884-86.
- Corral et al., 1999, "Differential cytokine modulation and T cell activation by two distinct classes of thalidomide analogues that are potent inhibitors of TNF-alpha," *J. Immunol.* 163(1):380-386.
- Craig et al., 1967, "Potential anticancer agents. III. 2-phthalimidoaldehydes and derivatives," *Potential Anticancer Agents III* 10:1071-1073.
- D'Amato et al., 2001, "Mechanism of action of thalidomide and 3-aminothalidomide in multiple myeloma," *Semin. Oncol.* 28:597-601.
- Dalgleish et al., "Thalidomide analogues CC-5013 and CC-4047 induce T cell activation and IL-12 production in patients with both solid tumours and relapsed and refractory multiple myeloma," *British Journal of Cancer*, 2003, 88(Suppl 1). S25-S54.
- Dalgleish, et al., "New thalidomide analogues; anti-cancer, anti-angiogenic and immunostimulatory," *British Journal of Cancer*, 2001, 85 (1)25.
- Davies et al., "Thalidomide (Thal) and immunomodulatory derivatives (IMiDs) augment natural killer (NK) cell cytotoxicity in multiple myeloma (MM)," Abstract # P222, *VIIIth International Myeloma Workshop*, May 4-8, 2001.
- Davies et al., "Thalidomide (Thal) and immunomodulatory derivatives (IMiDs) augment natural killer (NK) cell cytotoxicity in multiple myeloma(MM)," Abstract # 3617, *American Society of Hematology*, Dec. 1-5, 2000.
- Davies et al., 2001, "Thalidomide and immunomodulatory derivatives augment natural killer cell cytotoxicity in multiple myeloma," *Blood* 98(1):210-216.
- Dibbs et al., "Thalidomide and thalidomide analogs suppress TNF α secretion by myocytes," Abstract # 1284, *Circulation*, 1998.
- Dimopoulos et al., "Results of thalidomide and IMiDs in multiple myeloma," Abstract # P12.1.4, *International Multiple Myeloma Workshop*, May 23-27, 2003.
- Dimopoulos et al., "Study of lenalidomide plus dexamethasone versus dexamethasone alone in relapsed or refractory multiple myeloma (MM): Results of a phase 3 Study (MM-010)," Abstract # 6, *American Society of Hematology*, Dec. 10-13, 2005.
- Dimopoulos et al., "Treatment of plasma cell dyscrasias with thalidomide and its derivatives," *Journal of Clinical Oncology*, Dec. 1, 2003, 21 (23):4444-4454.
- Dimopoulos et al., 2004, "Primary treatment with pulsed melphalan, dexamethasone, thalidomide (MDT) for symptomatic patients with multiple myeloma ≤ 75 years of age," *Am. Soc. Hematol.* 46th Ann. Meeting Dec. 4-7, 2004, San Diego, CA Abstract # 1482.
- Dredge et al., "A costimulatory thalidomide analog enhances the partial anti-tumor immunity of an autologous vaccination in a model of colorectal cancer," Abstract # 491, *American Association for Cancer Research*, Apr. 6-10, 2002.
- Dredge et al., "Adjuvants and the promotion of Th1-type cytokines in tumour immunotherapy," *Cancer Immunol Immunother.*, 2002, 51:521-531.
- Dredge et al., "Angiogenesis inhibitors in cancer therapy," *Current Opinion in Investigational Drugs*, 2003, 4 (6):667-674.
- Dredge et al., "Immunological effects of thalidomide and its chemical and functional analogs," *Critical Reviews in Immunology*, 2002, 22 (5&6):425-437.
- Dredge et al., "Protective antitumor immunity induced by a costimulatory thalidomide analog in conjunction with whole tumor cell vaccination is mediated by increased Th1-type immunity¹," *The Journal of Immunology*, 2002, 168:4914-4919.
- Dredge et al., "Recent developments in antiangiogenic therapy," *Expert Opin. Biol. Ther.*, 2002, 2 (8):953-966.
- Dredge et al., "Thalidomide analogs as emerging anti-cancer drugs," *Anti-Cancer Drugs*, 2003, 14:331-335.
- Dredge et al., 2002, "Novel thalidomide analogues display anti-angiogenic activity independently of immunomodulatory effects," *Br. J. Cancer* 87(10):1166-1172.
- Fickentscher et al., "Stereochemical properties and teratogenic activity of some tetrahydrophthalimides," *Molecular Pharmacology*, 1976, 13:133-141.
- Galustian et al., "Thalidomide-derived immunomodulatory drugs as therapeutic agents," *Expert Opin. Biol. Ther.*, 2004, 4 (12):1-8.
- Glaspay et al., "The potential role of thalidomide and thalidomide analogs in melanoma," *Clinical Advances in Hematology & Oncology*, 2004, 1-7.
- Gupta et al., "Adherence of multiple myeloma cells to bone marrow stromal cells upregulates vascular endothelial growth factor secretion: therapeutic applications," *Leukemia*, 2001, 15:1950-1961.
- Haslett et al., 2003, "Thalidomide and a thalidomide analogue drug costimulate virus-specific CD8 + T cells in vitro," *J. Infect. Dis.* 187(6):946-955.
- Hayashi et al., "Mechanisms whereby immunomodulatory analogs of thalidomide augment autologous NK cell anti-myeloma immunity," *Blood*, Abstract # 3219, Dec. 6-10, 2002, *American Society of Hematology*.
- He, W., et al., 1993, Abstract of papers. 206th American Chemical Society, Chicago, IL; *Med. Chem.*, paper 216.
- Helm et al., "Comparative teratological investigation of compounds of structurally arid pharmacologically related to thalidomide," *Arzneimittel Forschung Drug Research*, 1981, 31 (1)941-949.

US 7,855,217 B2

Page 4

- Hernandez-Illizaliturri et al., "Addition of immunomodulatory drugs CC5013 or CC4047 to rituximab enhances anti-tumor activity in a severe combined immunodeficiency (SCID) mouse lymphoma model," *Abstract # 235, American Society of Hematology*, Dec. 6-9, 2003.
- Hideshima et al., "Thalidomide (Thal) and its analogs overcome drug resistance of human multiple myeloma (MM) cells to conventional therapy," *Abstract 1313, American Society of Hematology*, Dec. 1-5, 2000.
- Hideshima et al., "Thalidomide and its analogs overcome drug resistance of human multiple myeloma cells to conventional therapy," *Blood*, 2000, 96:2943-2950, *American Society of Hematology*.
- Hunt et al., "Markers of endothelial and haemostatic activation in the use of CC-4047, a structural analogue of thalidomide, in relapsed myeloma," *Blood*, *Abstract # 3216*, Dec. 6-10, 2002, *American Society of Hematology*.
- Hussein et al., "Doxil (D), vincristine (V), reduced frequency dexamethasone (d) and Revlimid (DVd-R) a phase I/II trial in advanced relapsed/refractory multiple myeloma (Rmm) patients," *Blood*, *Abstract #208, American Society of Hematology*, Dec. 4-7, 2004.
- Kyle et al., "Multiple myeloma," *New England Journal of Medicine*, 2004, 351:1860-1873.
- Kyle, "Current therapy of multiple myeloma," *Internal Medicine*, 2002, 41 (3)175-180.
- Leblanc et al., "Immunomodulatory drug costimulates T cells via the B7-CD28 pathway," *Blood*, 2004, 103:1787-1790, *American Society of Hematology*.
- Lentzsch et al., "Immunomodulatory derivative of thalidomide (IMiD CC-4047) determine the lineage commitment of hematopoietic progenitors by down regulation of GATA-1 and modulation of cytokine secretion," *Abstract # 3073, American Society of Hematology*, Dec. 6-9, 2003.
- Lentzsch et al., "Immunomodulatory derivative of thalidomide (IMiD CC-4047) down regulates CAAT/enhancer-binding protein β (C/EBP β) in multiple myeloma (MM)," *Abstract # 3456, American Society of Hematology*, Dec. 6-9, 2003.
- Lentzsch et al., "In vivo activity of thalidomide and immunomodulatory drugs against multiple myeloma," *VIIIth International Myeloma Workshop*, *Abstract #P225*, May 4-8, 2001.
- Lentzsch et al., 2003, "Immunomodulatory analogs of thalidomide inhibit growth of Hs Sultan cells and angiogenesis in vivo," *Leukemia* 17(1):41-44.
- Liu et al., "Phase I study of CC-5013 (Revimid), a thalidomide derivative, in patients with refractory metastatic cancer," *American Society of Oncology*, *Abstract #927*, 2003.
- Luzzio et al., "Thalidomide analogues: derivatives of an orphan drug with diverse biological activity," *Expert Opin. Ther. Patents*, 2004, 14 (2):215-229.
- Man et al., " α -Fluoro-substituted thalidomide analogues," *Bioorganic & Medicinal Chemistry Letters* 13, 2003, 3415-3417.
- Marriott et al., "A novel subclass of thalidomide analogue with anti-solid tumor activity in which caspase-dependent apoptosis is associated with altered expression of bcl-2 family proteins," *Cancer Research*, 2003, 63:593-599.
- Marriott et al., "Immunotherapeutic and antitumour potential of thalidomide analogues," *Expert Opin. Biol. Ther.*, 2001, 1 (4):1-8.
- Marriott et al., "New thalidomide analogues; anti-cancer, anti-angiogenic and immunostimulatory," *British Journal of Cancer*, 85:25, Jul. 6, 2001.
- Marriott et al., "Thalidomide and its analogues have distinct and opposing effects on TNF- α and TNFR2 during co-stimulation of both CD4⁺ and CD8⁺ T cells," *Clin. Exp. Immunol.*, 2002, 130:75-84.
- Marriott et al., "Thalidomide derived immunomodulatory drugs (IMiDs) as potential therapeutic agents," *Current Drug Targets—Immune, Endocrine & Metabolic Disorders*, 2003, 3:181-186.
- Masellis et al., "Changes in gene expression in bone marrow mesenchymal progenitor cells as a consequence of IMiD therapy in multiple myeloma patients," *Blood*, *Abstract # 1548*, Dec. 7-11, 2001, *American Society of Hematology*.
- McCarty, "Thalidomide may impede cell migration in primates by down-regulating integrin β -chains: potential therapeutic utility in solid malignancies, proliferative retinopathy, inflammatory disorders, neointimal hyperplasia, and osteoporosis," *Medical Hypotheses*, 1997, 49:123-131.
- Mitsiades et al., "Apoptotic signaling induced by immunomodulatory thalidomide analogs (Imids) in human multiple myeloma cells: therapeutic implications," *Abstract # 3224*, Dec. 7-11, 2001, *American Society of Hematology*.
- Mitsiades et al., "Apoptotic signaling induced by immunomodulatory thalidomide analogs in human multiple myeloma cells: therapeutic implications," *Blood*, 2002, 99:4525-4530, *American Society of Hematology*.
- Mitsiades et al., "CC-5013 Celgene," *Current Opinion in Investigational Drugs*, 2004, 5 (6):635-647.
- Miyachi et al., 1997, "Novel biological response modifiers: phthalimides with tumor necrosis factor-alpha production-regulating activity," *J. Med. Chem.* 40:2858-2865.
- Moutouh et al., "Novel immunomodulatory drugs (IMiDs®): A potential, new therapy for β -hemoglobinopathies," *Abstract # 3740, American Society of Hematology*, Dec. 4-7, 2004.
- Patten et al., "The early use of the serum free light chain assay in patients with relapsed refractory myeloma receiving treatment with a thalidomide analogue (CC-4047)," *Abstract # 1640, American Society of Hematology*, Dec. 6-9, 2003.
- Payvandi et al., "CC-5013 inhibits the expression of adhesion molecules ICAM-1 and CD44 and prevents metastasis of B16 F10 mouse melanoma cells in an animal model," *American Society of Clinical Oncology*, *Abstract # 992*, 2003.
- Payvandi et al., "Effects of a thalidomide analog on binding activity of transcription factors and cell cycle progression of multiple myeloma cell lines," *Blood*, *Abstract #2487*, Dec. 1-5, 2000, *American Society of Hematology*.
- Payvandi et al., "Immunomodulatory drugs inhibit expression of cyclooxygenase-2 from TNF- α , IL-1 β , and LPS-stimulated human PBMC in a partially IL-10-dependent manner," *Cellular Immunology*, 2004, 81-88.
- Payvandi et al., "Thalidomide and IMiDs inhibit microvessel formation from human arterial rings in the absence of human liver microsomes," *Blood*, *Abstract # 5046*, Dec. 6-10, 2002, *American Society of Hematology*.
- Payvandi et al., "Thalidomide analogs IMiDs inhibit expression of cyclooxygenase-2 in multiple myeloma cell line and LPS stimulated PBMCs," *Blood*, *Abstract # 2689*, Dec. 7-11, 2001, *American Society of Hematology*.
- Payvandi Yvandi et al., "The thalidomide analogs IMiDs enhance expression of CD69 stimulatory receptor on natural killer cells," *Abstract # 1793, American Association for Cancer Research*, Mar. 24-28, 2001.
- Raje et al., "Combination of the mTOR inhibitor rapamycin and CC-5013 has synergistic activity in multiple myeloma," *Blood*, Dec. 15, 2004, 104 (13)4188-4193.
- Rajkumar et al., "Combination therapy with lenalidomide plus dexamethasone (Rev/Dex) for newly diagnosed myeloma," *Blood*, Dec. 15, 2005, 106 (13)4050-4053.
- Richardson et al., "A multi-center, randomized, phase 2 study to evaluate the efficacy and safety of 2 CDC-5013 dose regimens when used alone or in combination with dexamethasone (Dex) for the treatment of relapsed or refractory multiple myeloma (MM)," *Blood*, *Abstract # 825, American Society of Hematology*, Dec. 6-9, 2003.
- Richardson et al., "A multicenter, single-arm, open-label study to evaluate the efficacy and safety of single-agent lenalidomide in patients with relapsed and refractory multiple myeloma; preliminary results," *10th International Myeloma Workshop*, Apr. 10-14, 2005.
- Richardson et al., "A Phase I study of oral CC5013, an immunomodulatory thalidomide (Thal) derivative, in patients with relapsed and refractory multiple myeloma (MM)," *Blood*, *Abstract #3225*, Dec. 7-11, 2001, *American Society of Hematology*.
- Richardson et al., "A phase I trial of lenalidomide (REVLIMID®) with bortezomib (VELCADE®) in relapsed and refractory multiple myeloma," *Blood*, *Abstract # 365, American Society of Hematology*, Dec. 10-13, 2005.
- Richardson et al., "Immunomodulatory analogs of thalidomide: an emerging new therapy in myeloma," *Journal of Clinical Oncology*, 2004, 22(16) 3212-3214.

US 7,855,217 B2

Page 5

- Richardson et al., "Immunomodulatory drug CC-55013 overcomes drug resistance and is well tolerated in patients with relapsed multiple myeloma," *Blood*, 2002 100:3063-3067, *American Society of Hematology*.
- Richardson et al., "Novel biological therapies for the treatment of multiple myeloma," *Best Practice & Research Clinical Haematology*, 2005, 18 (4):619-634.
- Rubin et al., "Principles of cancer treatment-I," 2003, 12 ONCO IV 1.
- Schafer et al., "Enhancement of cytokine production and AP-1 transcriptional activity in T cells by thalidomide-related immunomodulatory drugs," *Journal of Pharmacology and Experimental Therapeutics*, 2003, 305(3):1222-1232.
- Schey et al., "A phase I study of an immunomodulatory thalidomide analog, CC-4047, in relapsed or refractory multiple myeloma," *Journal of Clinical Oncology*, 2004, 22 (16):1-8.
- Schey et al., "A phase I study of an immunomodulatory thalidomide analogue (CC4047) in relapse/refractory multiple myeloma," *International Society for Experimental Hematology, Abstract #248*, 2002.
- Shah et al., 1999, "Synthesis and enantiomeric separation of 2-phthalimidino-glutaric acid analogues: potent inhibitors of tumor metastasis," *J. Med. Chem.* 42:3014-3017.
- Shaughnessy et al., "Global gene expression analysis shows loss of C-MYC and IL-6 receptor gene mRNA after exposure of myeloma to thalidomide and IMiD," *Abstract # 2485, The American Society of Hematology*, Dec. 1-5, 2000.
- Shire et al., "TNF- α inhibitors and rheumatoid arthritis," *Exp. Opin. Ther. Patents*, 1998, 8 (5):531-544.
- Sorbera et al., "CC-5013. Treatment of multiple myeloma. Treatment of Melanoma. Treatment of myelodysplastic syndrome. Angiogenesis inhibitor. TNF- α production inhibitor," *Drugs of the Future*, 2003, 28(5):425-431.
- Streetly et al., "An update of the use and outcomes of the new immunomodulatory agent CC-4047 (Actimid) in patients with relapsed/refractory myeloma," *Abstract #829, American Society of Hematology*, Dec. 6-9, 2003.
- Streetly et al., "Changes in neutrophil phenotype following the administration of CC-4047 (Actimid) to patients with multiple myeloma," *Abstract # 2543, American Society of Hematology*, Dec. 6-9, 2003.
- Streetly et al., "Thalidomide analogue CC-4047 is effective in the treatment of patients with relapsed and refractory multiple myeloma (MM) and induces T-cell activation and IL-12 production," *Abstract # 367, International Multiple Myeloma Workshop*, May 23-27, 2003.
- Teo et al., "A phase I, single-blind, placebo-controlled, ascending single oral dose, safety, tolerability and pharmacokinetic study of CDC-501, a novel immunomodulatory- oncologic agent, in healthy male subjects with a comparison of fed and fasted," *Clinical Pharmacology and Therapeutics*, 2002, 71 (2):93.
- Teo et al., "Chiral inversion of the second generation IMiDTM CC-4047 (ACTIMIDTM) in human plasma and phosphate-buffered saline," *Chirality*, 2003, 15:348-351.
- Thertulien et al., "Hybrid MEL/DT PACE autotransplant regimen for Multiple Myeloma (MM)- safety and efficacy data in pilot study of 15 patients," *Blood, Abstract # 2869, American Society of Hematology*, Dec. 7-11, 2001.
- Tohny et al., "A phase I study of oral CC-5013 (lenalidomide, RevlimidTM), a thalidomide derivative, in patients with refractory metastatic cancer," *Clinical Prostate Cancer*, 2004, 2:241-243.
- Tricot et al., "Angiochemotherapy (ACT) for multiple myeloma (MM) with DT-PACE results in a high response rate, but in contrast to tandem transplants with melphalan does not affect durable disease control," *Blood, Abstract # 3531, American Society of Hematology*, Dec. 7-11, 2001.
- Tsenova et al., "Use of IMiD3, a thalidomide analog, as an adjunct to therapy for experimental tuberculous meningitis," *Antimicrobial Agents and Chemotherapy*, 2002, 46 (6):1887-1895.
- Weber et al., "A multicenter, randomized, parallel-group, double-blind, placebo-controlled study of lenalidomide plus dexamethasone versus dexamethasone alone in previously treated subjects with multiple myeloma," *Abstract # PO.738, International Multiple Myeloma Workshop*, Apr. 10-14, 2005.
- Weber, "Lenalidomide (CC-5013, RevlimidTM) and other IMiDs," *Abstract # PL5.02, International Multiple Myeloma Workshop*, Apr. 10-14, 2005.
- Ye et al., "Novel IMiD drugs enhance expansion and regulate differentiation of human cord blood CD34+ cells with cytokines," *Blood, Abstract #4099, American Society of Hematology*, Dec. 6-10, 2002.
- Zangari et al., "Results of phase I study of CC-5013 for the treatment of multiple myeloma (MM) patients who relapse after high dose chemotherapy (HDCT)," *American Society of Hematology, Abstract # 3226*, 2001.
- Zangari et al., "Revimid 25 mg (Rev 25) \times 20 versus 50 mg (REV 50) \times 10 q 28 days with bridging of 5 mg \times 10 versus 10 mg \times 5 as post-transplant salvage therapy for multiple myeloma (MM)," *Blood, Abstract # 1642, American Society of Hematology*, Dec. 6-9, 2003.
- Zangari et al., "Risk factors for deep vein thrombosis (DVT) in a large group of myeloma patients (PTS) treated with thalidomide (Thal): The Arkansas Experience," *Blood, Abstract # 681, American Society of Hematology*, Dec. 7-11, 2001.
- Zeldis et al., "Potential new therapeutics for Waldenstrom's macroglobulinemia," *Seminars in Oncology*, 2003, 30 (2):275-281.
- Zeldis et al., "Update on the evolution of the IMiDTM," *International Society for Biological Therapy of Cancer. Oral Abstract*, 2003.
- Zhang et al., "CC-5079, a novel microtubule and TNF- α inhibitor with anti-angiogenic and antimetastasis activity," *Abstract # B012, International Conference on Molecular Targets and Cancer Therapeutics*, Nov. 17-21, 2003.
- Haleblian, J.; McCrone, W.; "Pharmaceutical Applications of Polymorphism", *Journal of Pharmaceutical Sciences*, Aug. 1969, 58(8):911-929.
- Davies, F.E.; et al., "Thalidomide and immunomodulatory derivatives augment natural killer cell cytotoxicity in multiple myeloma," *Blood*, Jul. 2001, 98(1):210-216.
- Kyle, R.A.; Rajkumar, S.V., "Therapeutic Application of Thalidomide in Multiple Myeloma," *Seminars in Oncology*, Dec. 2001, 28(6):583-587.
- USPTO Office Action dated Dec. 31, 2007 in currently pending U.S. Appl. No. 10/438,213.
- Office Action dated Apr. 16, 2010 in corresponding Japanese Patent Application No. 2006-525471 (with summary).
- USPTO Office Action dated Oct. 10, 2007 in U.S. Appl. No. 7,465,800.
- Bernstein, 2002, *Polymorphism in Molecular Crystals*, Clarendon Press, 115-18.
- Bernstein, 2004, "Crystal Structure Prediction and Polymorphism," *ACA Transactions*, 39, 14-23.
- Bond et al., 2002, "Controlling Crystal Architecture in Molecular Solids: the Supramolecular Approach," in *Supramolecular Organization and Materials Design*, Jones & Rao eds., Cambridge University Press, Chapter 12, pp. 391-443.
- Brittain, 1999, "Methods for the Characterization of Polymorphs and Solvates," in *Polymorphism in Pharmaceutical Solids*, Brittain ed., Chapter 6, pp. 227-278.
- Brittain, 2002, "Polymorphism: Pharmaceutical Aspects," in *Encyclopedia of Pharmaceutical Technology*, 2nd Edition, vol. 3, Swarbrick & Boylan eds., pp. 2239-2249.
- U.S. Appl. No. 60/499,723, filed Sep. 4, 2003, Jaworsky.
- U.S. Appl. No. 60/372,348, filed Apr. 12, 2002, Hariiri et al.
- U.S. Appl. No. 09/545,654, filed Apr. 10, 2000, D'Amato.
- U.S. Appl. No. 09/287,377, filed Apr. 7, 1999, D'Amato.
- Brittain, 2007, "Polymorphism and Solvatomorphism 2005," *Journal of Pharmaceutical Sciences*, 96(4), 705-28.
- Bruns et al., 1984, "Thermochemical Investigation of Theophylline, Theophylline Hydrate and Their Aqueous Solutions," *Thermochimica Acta*, 72, 31-40.
- Byrn et al., 1999, "Drugs as Molecular Solids," in *Solid-State Chemistry of Drugs*, 2nd Edition, Chapter 1, pp. 3-43.
- Cruz Cabeza et al., 2006, "Prediction and Observation of Isostructurality Induced by Solvent Incorporation in Multicomponent Crystals," *J. Am. Chem. Soc.*, 128, 14466-67.
- Cruz Cabeza et al., 2007, "Importance of Molecular Shape for the Overall Stability of Hydrogen Bond Motifs in the Crystal Structures of Various Carbamazepine-Type Drug Molecules," *Crystal Growth & Design*, 7(1), 100-07.

US 7,855,217 B2

Page 6

- Day et al., 2004, "An Assessment of Lattice Energy Minimization for the Prediction of Molecular Organic Crystal Structures," *Crystal Growth & Design*, 4(6), 1327-40.
- Day et al., 2006, "Investigating the Latent Polymorphism of Maleic Acid," *Chemical Communications*, 54-56.
- Dean, 1993, *Analytical Chemistry Handbook*, 10.24-10.26.
- Gadamasetti et al., 2007, "Process Chemistry in the Pharmaceutical Industry," in *Encyclopedia of Pharmaceutical Technology*, 3rd Edition, vol. 5, Swarbrick ed., pp. 2993-3007.
- Jones et al., 2006, "Pharmaceutical Cocrystals: An Emerging Approach to Physical Property Enhancement," *MRS Bulletin*, 31, 875-79.
- Kirk-Othmer, 2002, "Crystallization," in *Encyclopedia of Chemical Technology*, vol. 8, pp. 95-147.
- Knapman, 2000, "Polymorphic Predictions," *Modern Drug Discovery*, 3(2), 53-57.
- Office Action dated Jun. 18, 2010 in U.S. Appl. No. 12/220,336.
- Richardson et al., "A Multi-Center, Randomized, Phase II Study To Evaluate the Efficacy and Safety of Two CDC-5013 Dose Regimens When Used Alone or in Combination with Dexamethasone (Dex) for the Treatment of Relapsed or Refractory Multiple Myeloma (MM)," *Blood*, Journal of the American Society of Hematology, Dec. 6-10, 2002, 100(11):Abstract #386.
- Seddon, 2004, "Pseudopolymorph: A Polemic," *Crystal Growth & Design*, 4(6), 1087.
- Sharma et al., 2006, "Toxicity Profile of the Immunomodulatory Thalidomide Analogue, Lenalidomide: Phase I Clinical Trial of Three Dosing Schedules in Patients with Solid Malignancies," *European Journal of Cancer*, 42, 2318-25.
- Vippagunta et al., 2001, "Crystalline Solids," *Advanced Drug Delivery Reviews*, 48, 3-26.
- Caira, 1998, "Crystalline Polymorphism of Organic Compounds," *Topics in Current Chemistry*, 198, 163-208.
- Fauci et al, 1998, "Harrison's Principles of Internal Medicine, Chronic Visual Loss," in *Harrison's Principles of Internal Medicine*, 168.
- Notice of Allowability mailed Nov. 3, 2008 in U.S. Appl. No. 10/934,863.
- Office Action mailed Jun. 17, 2010 in U.S. Appl. No. 10/557,302.




Fig. 1

IR SPECTRUM OF FORM A




Fig. 2

RAMAN SPECTRUM OF FORM A




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8

TGA (TOP) AND DSC (BOTTOM) OF FORM B




Fig. 9

TG-IR RESULTS FOR FORM B





Fig. 10






Fig. 12

IR SPECTRUM OF FORM C




Fig. 13





Fig. 14

TGA (TOP) AND DSC (BOTTOM) OF FORM C






Fig. 16




Fig. 17

XRPD PATTERN OF FORM D




Fig. 18

IR SPECTRUM OF FORM D




Fig. 19

RAMAN SPECTRUM OF FORM D




Fig. 20

TGA (TOP) AND DSC (BOTTOM) OF FORM D




Fig. 21




Fig. 22

XRPD PATTERN OF FORM E




Fig. 23

TGA (TOP) AND DSC (BOTTOM) OF FORM E




Fig. 24




Fig. 25

XRPD PATTERN OF FORM F




Fig. 26




Fig. 27

XRPD PATTERN OF FORM G




Fig. 28




Fig. 29




Fig. 30

TGA (TOP) AND DSC (BOTTOM) OF FORM H





Fig. 31

XRPD PATTERN OF POLYMORPH B




Fig. 32




Fig. 33

XRPD PATTERN OF POLYMORPH B




Fig. 34

XRPD PATTERN OF POLYMORPH E




Fig. 35

XRPD PATTERN OF POLYMORPH MIXTURE




Fig. 36




Fig. 37




Fig. 38




Fig. 39




Fig. 40




Fig. 41




Fig. 42




Fig. 43




Fig. 44




Fig. 45




Fig. 46




Fig. 47




Fig. 48




Fig. 49

CALIBRATION CURVE FOR
3-(4-AMINO-1-OXO-1,3 DIHYDROISOINDOL-2-YL)-PIPERIDINE-2,6-DIONE




Fig. 50




Fig. 51




Fig. 52




FIG. 53

INTRINSIC DISSOLUTION OF FORMS A, B, AND E (EXPERIMENT NO.2)




FIG.54

US 7,855,217 B2

1

**POLYMORPHIC FORMS OF
3-(4-AMINO-1-OXO-1,3
DIHYDRO-ISOINDOL-2-YL)-
PIPERIDINE-2,6-DIONE**

This application is a divisional application of U.S. patent application Ser. No. 10/934,863, filed Sep. 3, 2004, now U.S. Pat. No. 7,465,800 presently pending, which claims the benefit of U.S. provisional application 60/499,723, filed Sep. 4, 2003, the contents of each of which are incorporated by reference herein in their entireties.

1. FIELD OF THE INVENTION

This invention relates to polymorphic forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione, compositions comprising the polymorphic forms, methods of making the polymorphic forms and methods of their use for the treatment of diseases and conditions including, but not limited to, inflammatory diseases, autoimmune diseases, and cancer.

2. BACKGROUND OF THE INVENTION

Many compounds can exist in different crystal forms, or polymorphs, which exhibit different physical, chemical, and spectroscopic properties. For example, certain polymorphs of a compound may be more readily soluble in particular solvents, may flow more readily, or may compress more easily than others. See, e.g., P. DiMartino, et al., *J. Thermal Anal.*, 48:447-458 (1997). In the case of drugs, certain solid forms may be more bioavailable than others, while others may be more stable under certain manufacturing, storage, and biological conditions. This is particularly important from a regulatory standpoint, since drugs are approved by agencies such as the U.S. Food and Drug Administration only if they meet exacting purity and characterization standards. Indeed, the regulatory approval of one polymorph of a compound, which exhibits certain solubility and physico-chemical (including spectroscopic) properties, typically does not imply the ready approval of other polymorphs of that same compound.

Polymorphic forms of a compound are known in the pharmaceutical arts to affect, for example, the solubility, stability, flowability, fractability, and compressibility of the compound, as well as the safety and efficacy of drug products comprising it. See, e.g., Knapman, K. *Modern Drug Discoveries*, 2000, 53. Therefore, the discovery of new polymorphs of a drug can provide a variety of advantages.

U.S. Pat. Nos. 5,635,517 and 6,281,230, both to Muller et al., disclose 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione, which is useful in treating and preventing a wide range of diseases and conditions including, but not limited to, inflammatory diseases, autoimmune diseases, and cancer. New polymorphic forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione can further the development of formulations for the treatment of these chronic illnesses, and may yield numerous formulation, manufacturing and therapeutic benefits.

3. SUMMARY OF THE INVENTION

This invention encompasses polymorphs of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. In certain aspects, the invention provides polymorphs of the compound identified herein as forms A, B, C, D, E, F, G, and H. The invention also encompasses mixtures of these forms. In further embodiments, this invention provides methods of making, isolating and characterizing the polymorphs.

2

This invention also provides pharmaceutical compositions and single unit dosage forms comprising a polymorph of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. The invention further provides methods for the treatment or prevention of a variety of diseases and disorders, which comprise administering to a patient in need of such treatment or prevention a therapeutically effective amount of a polymorph of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione.

4. BRIEF DESCRIPTION OF THE DRAWINGS

Specific aspects of the invention can be understood with reference to the attached figures:

FIG. 1 provides a representative X-ray powder diffraction (XRPD) pattern of Form A;

FIG. 2 provides a representative IR spectrum of Form A;

FIG. 3 provides a representative Raman spectrum of Form A;

FIG. 4 provides a representative thermogravimetric analysis (TGA) curve and a representative differential scanning calorimeter (DSC) thermogram of Form A;

FIG. 5 provides a representative moisture sorption/desorption isotherm of Form A;

FIG. 6 provides a representative XRPD pattern of Form B;

FIG. 7 provides a representative IR spectrum of Form B;

FIG. 8 provides a representative Raman spectrum of Form B;

FIG. 9 provides a representative TGA curve and a representative DSC thermogram of Form B;

FIG. 10 provides representative TG-IR results of Form B;

FIG. 11 provides a representative moisture sorption/desorption isotherm of Form B;

FIG. 12 provides a representative XRPD pattern of Form C;

FIG. 13 provides a representative IR spectrum of Form C;

FIG. 14 provides a representative Raman spectrum of Form C;

FIG. 15 provides a representative TGA curve and a representative DSC thermogram of Form C;

FIG. 16 provides representative TG-IR results of Form C;

FIG. 17 provides a representative moisture sorption/desorption isotherm of Form C;

FIG. 18 provides a representative XRPD pattern of Form D;

FIG. 19 provides a representative IR spectrum of Form D;

FIG. 20 provides a representative Raman spectrum of Form D;

FIG. 21 provides a representative TGA curve and a representative DSC thermogram of Form D;

FIG. 22 provides a representative moisture sorption/desorption isotherm of Form D;

FIG. 23 provides a representative XRPD pattern of Form E;

FIG. 24 provides a representative TGA curve and a representative DSC thermogram of Form E;

FIG. 25 provides a representative moisture sorption/desorption isotherm of Form E;

FIG. 26 provides a representative XRPD pattern for a sample of Form F;

FIG. 27 provides a representative thermogram of Form F;

FIG. 28 provides a representative XRPD pattern of Form G;

FIG. 29 provides a representative DSC thermogram for a sample of Form G;

FIG. 30 provides a representative XRPD pattern of Form H;

US 7,855,217 B2

3

FIG. 31 provides a representative TGA curve and a representative DSC thermogram of Form H;

FIG. 32 provides a representative XRPD pattern of Form B;

FIG. 33 provides a representative XRPD pattern of Form B;

FIG. 34 provides a representative XRPD pattern of Form B;

FIG. 35 provides a representative XRPD pattern of Form E;

FIG. 36 provides a representative XRPD pattern of polymorph mixture;

FIG. 37 provides a representative TGA curve of Form B;

FIG. 38 provides a representative TGA curve of Form B;

FIG. 39 provides a representative TGA curve of Form B;

FIG. 40 provides a representative TGA curve of Form E;

FIG. 41 provides a representative TGA curve of polymorph mixture;

FIG. 42 provides a representative DSC thermogram of Form B;

FIG. 43 provides a representative DSC thermogram of Form B;

FIG. 44 provides a representative DSC thermogram of Form B;

FIG. 45 provides a representative DSC thermogram of Form E;

FIG. 46 provides a representative DSC thermogram of polymorph mixture;

FIG. 47 provides a UV-Vis scan of dissolution medium;

FIG. 48 provides a UV-Vis scan of 0.04 mg/ml of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione in dissolution medium;

FIG. 49 provides a UV-Vis scan of 0.008 mg/ml of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione in dissolution medium;

FIG. 50 provides a calibration curve for 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione;

FIG. 51 provides a solubility curve of Form A;

FIG. 52 provides a solubility curve of Form B;

FIG. 53 provides an intrinsic dissolution of Forms A, B and E; and

FIG. 54 provides an intrinsic dissolution of Forms A, B and E.

5. DETAILED DESCRIPTION OF THE INVENTION

5.1 Definitions

As used herein and unless otherwise indicated, the terms “treat,” “treating” and “treatment” refer to the alleviation of a disease or disorder and/or at least one of its attendant symptoms.

As used herein and unless otherwise indicated, the terms “prevent,” “preventing” and “prevention” refer to the inhibition of a symptom of a disease or disorder or the disease itself.

As used herein and unless otherwise indicated, the terms “polymorph” and “polymorphic form” refer to solid crystalline forms of a compound or complex. Different polymorphs of the same compound can exhibit different physical, chemical and/or spectroscopic properties. Different physical properties include, but are not limited to stability (e.g., to heat or light), compressibility and density (important in formulation and product manufacturing), and dissolution rates (which can affect bioavailability). Differences in stability can result from changes in chemical reactivity (e.g., differential oxidation, such that a dosage form discolors more rapidly when comprised of one polymorph than when comprised of another polymorph) or mechanical characteristics (e.g., tablets crumble on storage as a kinetically favored polymorph converts to thermodynamically more stable polymorph) or both

4

(e.g., tablets of one polymorph are more susceptible to breakdown at high humidity). Different physical properties of polymorphs can affect their processing. For example, one polymorph might be more likely to form solvates or might be more difficult to filter or wash free of impurities than another due to, for example, the shape or size distribution of particles of it.


Polymorphs of a molecule can be obtained by a number of methods known in the art. Such methods include, but are not limited to, melt recrystallization, melt cooling, solvent recrystallization, desolvation, rapid evaporation, rapid cooling, slow cooling, vapor diffusion and sublimation. Polymorphs can be detected, identified, classified and characterized using well-known techniques such as, but not limited to, differential scanning calorimetry (DSC), thermogravimetry (TGA), X-ray powder diffractometry (XRPD), single crystal X-ray diffractometry, vibrational spectroscopy, solution calorimetry, solid state nuclear magnetic resonance (NMR), infrared (IR) spectroscopy, Raman spectroscopy, hot stage optical microscopy, scanning electron microscopy (SEM), electron crystallography and quantitative analysis, particle size analysis (PSA), surface area analysis, solubility, and rate of dissolution.

As used herein to refer to the spectra or data presented in graphical form (e.g., XRPD, IR, Raman and NMR spectra), and unless otherwise indicated, the term “peak” refers to a peak or other special feature that one skilled in the art would recognize as not attributable to background noise. The term “significant peaks” refers to peaks at least the median size (e.g., height) of other peaks in the spectrum or data, or at least 1.5, 2, or 2.5 times the median size of other peaks in the spectrum or data.

As used herein and unless otherwise indicated, the term “substantially pure” when used to describe a polymorph of a compound means a solid form of the compound that comprises that polymorph and is substantially free of other polymorphs of the compound. A representative substantially pure polymorph comprises greater than about 80% by weight of one polymorphic form of the compound and less than about 20% by weight of other polymorphic forms of the compound, more preferably greater than about 90% by weight of one polymorphic form of the compound and less than about 10% by weight of the other polymorphic forms of the compound, even more preferably greater than about 95% by weight of one polymorphic form of the compound and less than about 5% by weight of the other polymorphic forms of the compound, and most preferably greater than about 97% by weight of one polymorphic forms of the compound and less than about 3% by weight of the other polymorphic forms of the compound.

5.2 Polymorphic Forms

This invention is directed to polymorphic forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione, which has the structure shown below:



This compound can be prepared according to the methods described in U.S. Pat. Nos. 6,281,230 and 5,635,517, the entireties of which are incorporated herein by reference. For

US 7,855,217 B2

5

example, the compound can be prepared through catalytic hydrogenation of 3-(4-nitro-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. 3-(4-Nitro-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione can be obtained by allowing 2,6-dioxopiperidin-3-ammonium chloride to react with methyl 2-bromomethyl-4-nitrobenzoate in dimethylformamide in the presence of triethylamine. The methyl 2-bromomethyl-4-nitrobenzoate in turn is obtained from the corresponding methyl ester of nitro-ortho-toluic acid by conventional bromination with N-bromosuccinimide under the influence of light.

Polymorphs of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione can be obtained by techniques known in the art, including solvent recrystallization, desolvation, vapor diffusion, rapid evaporation, slow evaporation, rapid cooling and slow cooling. Polymorphs can be made by dissolving a weighed quantity of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione in various solvents at elevated temperatures. The solutions of the compound can then be filtered and allowed to evaporate either in an open vial (for fast hot evaporation) or in a vial covered with aluminum foil containing pinholes (hot slow evaporation). Polymorphs can also be obtained from slurries. Polymorphs can be crystallized from solutions or slurries using several methods. For example, a solution created at an elevated temperature (e.g., 60° C.) can be filtered quickly then allowed to cool to room temperature. Once at room temperature, the sample that did not crystallize can be moved to a refrigerator then filtered. Alternatively, the solutions can be crash cooled by dissolving the solid in a solvent at an increased temperature (e.g., 45-65° C.) followed by cooling in a dry ice/solvent bath.

One embodiment of the invention encompasses Form A of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form A is an unsolvated, crystalline material that can be obtained from non-aqueous solvent systems. Another embodiment of the invention encompasses Form B of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form B is a hemihydrated, crystalline material that can be obtained from various solvent systems. Another embodiment of the invention encompasses Form C of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form C is a hemisolvated crystalline material that can be obtained from solvents such as, but not limited to, acetone. Another embodiment of the invention encompasses Form D of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form D is a crystalline, solvated polymorph prepared from a mixture of acetonitrile and water. Another embodiment of the invention encompasses Form E of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form E is a dihydrated, crystalline material. Another embodiment of the invention encompasses Form F of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form F is an unsolvated, crystalline material that can be obtained from the dehydration of Form E. Another embodiment of the invention encompasses Form G of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form G is an unsolvated, crystalline material that can be obtained from slurrying forms B and E in a solvent such as, but not limited to, tetrahydrofuran (THF). Another embodiment of the invention encompasses Form H of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Form H is a partially hydrated crystalline material that can be obtained by exposing Form E to 0% relative humidity. Each of these forms is discussed in detail below.

Another embodiment of the invention encompasses a composition comprising amorphous 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione and crystalline 3-(4-

6

amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione of form A, B, C, D, E, F, G or H. Specific compositions can comprise greater than about 50, 75, 90 or 95 weight percent crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione.

Another embodiment of the invention encompasses a composition comprising at least two crystalline forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione (e.g., a mixture of polymorph forms B and E).

5.2.1 Form A

The data described herein for Form A, as well as for Forms B-H, were obtained using the experimental methods described in Examples 6.3-6.7, provided below.

Form A can be obtained from various solvents, including, but not limited to 1-butanol, butyl acetate, ethanol, ethyl acetate, methanol, methyl ethyl ketone, and THF. FIG. 1 shows a representative XRPD pattern of Form A. The pattern is characterized by peaks, preferably significant peaks, at approximately 8, 14.5, 16, 17.5, 20.5, 24, and 26 degrees 2 θ . Representative IR and Raman spectra data are provided in FIGS. 2 and 3.

Representative thermal characteristics of Form A are shown in FIG. 4. TGA data show a small weight increase up to about 150° C., indicating an unsolvated material. Weight loss above 150° C. is attributed to decomposition. The DSC curve of Form A exhibits an endotherm at about 270° C.

Representative moisture sorption and desorption data are plotted in FIG. 5. Form A does not exhibit a significant weight gain from 5 to 95% relative humidity. Equilibrium can be obtained at each relative humidity step. As the form dries from 95% back down to 5% relative humidity, it tends to maintain its weight such that at 5% relative humidity it has typically lost only about 0.003% by weight from start to finish. Form A is capable of remaining a crystalline solid for about 11 days when stored at about 22, 45, 58, and 84% relative humidity.

Interconversion studies show that Form A can convert to Form B in aqueous solvent systems and can convert to Form C in acetone solvent systems. Form A tends to be stable in anhydrous solvent systems. In water systems and in the presence of Form E, Form A tends to convert to Form E.

When stored for a period of about 85 days under two different temperature/relative humidity stress conditions (room temperature/0% relative humidity (RH) and 40° C./93% RH), Form A typically does not convert to a different form.

In sum, Form A is a crystalline, unsolvated solid that melts at approximately 270° C. Form A is weakly or not hygroscopic and appears to be the most thermodynamically stable anhydrous polymorph of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione discovered thus far.

5.2.2 Form B

Form B can be obtained from many solvents, including, but not limited to, hexane, toluene, and water. FIG. 6 shows a representative XRPD pattern of Form B, characterized by peaks at approximately 16, 18, 22 and 27 degrees 2 θ .

Solution proton NMR confirm that Form B is a form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Representative IR and Raman spectra are shown in FIGS. 7 and 8, respectively. Compared to Form A, the IR spectrum for Form B has peaks at approximately 3513 and 1960 cm⁻¹.

Representative DSC and TGA data for Form B are shown in FIG. 9. The DSC curve exhibits endotherms at about 146 and 268° C. These events are identified as dehydration and melting by hot stage microscopy experiments. Form B typically loses about 3.1% volatiles up to about 175° C. (per

US 7,855,217 B2

7

approximately 0.46 moles of water). Comparison of the IR spectrum of the volatiles with that of water indicates that they are water (See FIG. 10). Calculations from TGA data indicate that Form B is a hemihydrate. Karl Fischer water analysis also supports this conclusion.

Representative moisture sorption and desorption data are shown in FIG. 11. Form B typically does not exhibit a significant weight gain from 5% to 95% relative humidity, when equilibrium is obtained at each relative humidity step. As Form B dries from 95% back down to 5% relative humidity, it tends to maintain its weight such that at 5% relative humidity it typically has gained only about 0.022% by weight (about 0.003 mg) from start to finish. Form B does not convert to a different form upon exposure to about 84% relative humidity for about ten days.

Interconversion studies show that Form B typically converts to Form A in a THF solvent system, and typically converts to Form C in an acetone solvent system. In aqueous solvent systems such as pure water and 10% water solutions, Form B is the most stable of the polymorphic forms of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. However, it can convert to Form E in the presence of water. Desolvation experiments show that upon heating at about 175° C. for about five minutes, Form B typically converts to Form A.

When stored for a period of about 85 days under two different temperature/relative humidity stress conditions (room temperature/0% RH and 40° C./93% RH), Form B does not convert to a different form.

In sum, Form B is a hemihydrated, crystalline solid which has a DSC thermogram exhibiting endotherms at about 146 and about 268° C. Interconversion studies show that Form B converts to Form E in aqueous solvent systems, and converts to other forms in acetone and other anhydrous systems.

5.2.3 Form C

Form C can be obtained from evaporations, slurries and slow cools in acetone solvent systems. A representative XRPD pattern of this form is shown in FIG. 12. The data are characterized by peaks at approximately 15.5 and 25 degrees 2θ.

Solution proton NMR indicates that the 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione molecule is intact. Representative IR and Raman spectra are shown in FIGS. 13 and 14, respectively. The IR spectrum of Form C is characterized by peaks at approximately 3466, 3373, and 3318 cm⁻¹. The Raman spectrum of Form C is characterized by peaks at about 3366, 3321, 1101, and 595 cm⁻¹.

Representative thermal characteristics for Form C are plotted in FIG. 15. Form C loses about 10.02% volatiles up to about 175° C., indicating it is a solvated material. Weight loss above about 175° C. is attributed to decomposition. Identification of volatiles in Form C can be accomplished with TG-IR experiments. The representative IR spectrum captured after several minutes of heating, as depicted in FIG. 13, when compared with a spectral library, shows acetone to be the best match. Calculations from TGA data show that Form C is a hemisolvate (approximately 0.497 moles of acetone). The DSC curve for Form C, shown in FIG. 15, exhibits endotherms at about 150 and about 269° C. The endotherm at about 150° C. is attributed to solvent loss based on observations made during hot stage microscopy experiments. The endotherm at about 269° C. is attributed to the melt based on hot stage experiments.

Representative moisture sorption and desorption balance data are shown in FIG. 17. Form C does not exhibit a significant weight gain from 5 to 85% relative humidity, when equilibrium is obtained at each relative humidity step up to

8

85% relative humidity. At 95% relative humidity, Form C experiences a significant weight loss of about 6.03%. As the sample dries from 95% back down to 5% relative humidity, the sample maintains the weight achieved at the end of the adsorption phase at each step down to 5% relative humidity. Form C is capable of converting to Form B when stored at about 84% relative humidity for approximately ten days.

Interconversion studies show that Form C typically converts to Form A in a THF solvent system and typically converts to Form E in an aqueous solvent system. In an acetone solvent system, Form C is the most stable form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Desolvation experiments performed on Form C show that upon heating at about 150° C. for about five minutes, Form C will typically convert to Form A.

In sum, Form C is a crystalline, hemisolvated solid, which has a DSC thermogram exhibiting endotherms at about 150 and about 269° C. Form C is not hygroscopic below about 85% RH, but can convert to Form B at higher relative humidities.

5.2.4 Form D

Form D can be obtained from evaporation in acetonitrile solvent systems. A representative XRPD pattern of the form is shown in FIG. 18. The pattern is characterized by peaks at approximately 27 and 28 degrees 2θ.

Solution proton NMR indicates that the 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione molecule is intact. Representative IR and Raman spectra are shown in FIGS. 19 and 20, respectively. The IR spectrum of Form D is characterized by peaks at approximately 3509, 2299, and 2256 cm⁻¹. The Raman spectrum of Form D is characterized by peaks at approximately 2943, 2889, 2297, 2260, 1646, and 1150 cm⁻¹.

Representative thermal characteristics for Form D are plotted in FIG. 21. Form D loses about 6.75% volatiles up to about 175° C., indicating a solvated material. Weight loss above about 175° C. is attributed to decomposition. TG-IR experiments indicate that the volatiles are water and acetonitrile. Calculations from TG data show that about one mole of water is present in the sample. A representative DSC curve for Form D exhibits endotherms at about 122 and about 270° C. The endotherm at about 122° C. is attributed to loss of volatiles based on observations made during hot stage microscopy experiments. The endotherm at about 270° C. is attributed to the melt based on hot stage experiments.

Representative moisture sorption and desorption data are plotted in FIG. 22. Form D does not exhibit a significant weight gain from 5 to 95% relative humidity when equilibrium is obtained at each relative humidity step. As the form dries from 95% back down to 5% relative humidity, it maintains its weight such that at 5% relative humidity the form has typically gained only about 0.39% by weight (about 0.012 mg) from start to finish. Form A is capable of converting to Form B when stored at about 84% relative humidity for approximately ten days.

Interconversion studies show that Form D is capable of converting to Form A in a THF solvent system, to Form E in an aqueous solvent system, and to Form C in an acetone solvent system. Desolvation experiments performed on Form D show that upon heating at about 150° C. for about five minutes Form D will typically convert to Form A.

In sum, Form D is a crystalline solid, solvated with both water and acetonitrile, which has a DSC thermogram exhibiting endotherms at about 122 and about 270° C. Form D is either weakly or not hygroscopic, but will typically convert to Form B when stressed at higher relative humidities.

5.2.5 Form E

Form E can be obtained by slurring 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione in water and by a slow evaporation of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione in a solvent system with a ratio of about 9:1 acetone:water. A representative XRPD pattern is shown in FIG. 23. The data are characterized by peaks at approximately 20, 24.5 and 29 degrees 2 θ .

Representative thermal characteristics of Form E are plotted in FIG. 24. Form E typically loses about 10.58% volatiles up to about 125° C., indicating that it is a solvated material. A second weight loss of an additional about 1.38% was observed between about 125° C. and about 175° C. Weight loss above about 175° C. is attributed to decomposition. Karl Fischer and TG-IR experiments support the conclusion that the volatile weight loss in Form E is due to water. The representative DSC curve for Form E exhibits endotherms at about 99, 161 and 269° C. Based on observations made during hot stage microscopy experiments, the endotherms at about 99 and about 161° C. are attributed to loss of volatiles. The endotherm at about 269° C. is attributed to the melt based on hot stage experiments.

Representative moisture sorption and desorption data are plotted in FIG. 25. Form E typically does not exhibit a significant weight change from 5 to 95% relative humidity when equilibrium is obtained at each relative humidity step. As the sample dried from 95% back down to 5% relative humidity, the sample continues to maintain weight such that at 5% relative humidity the sample has lost only about 0.0528% by weight from start to finish.

Interconversion studies show that Form E can convert to Form C in an acetone solvent system and to Form G in a THF solvent system. In aqueous solvent systems, Form E appears to be the most stable form. Desolvation experiments performed on Form E show that upon heating at about 125° C. for about five minutes, Form E can convert to Form B. Upon heating at 175° C. for about five minutes, Form B can convert to Form F.

When stored for a period of 85 days under two different temperature/relative humidity stress conditions (room temperature/0% RH and 40° C./93% RH) Form E typically does not convert to a different form. When stored for seven days at room temperature/0% RH, Form E can convert to a new form, Form H.

5.2.6 Form F

Form F can be obtained by complete dehydration of Form E. A representative XRPD pattern of Form F, shown in FIG. 26, is characterized by peaks at approximately 19, 19.5 and 25 degrees 2 θ .

Representative thermal characteristics of Form F are shown in FIG. 27. The representative DSC curve for Form F exhibits an endotherm at about 269° C. preceded directly by two smaller endotherms indicative of a crystallized form of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. The DSC thermogram does not show any thermal events prior to the melt, suggesting that it is an unsolvated material.

5.2.7 Form G

Form G can be obtained by slurring forms B and E in THF. A representative XRPD pattern of this form, shown in FIG. 28, is characterized by a peak at approximately 23 degrees 2 θ . Two other peaks unique to Form G appear at approximately 21 and 24.5 degrees 2 θ .

Representative thermal characteristics of Form G are plotted in FIG. 29. A representative DSC curve for Form G exhibits an endotherm at about 248° C. followed by a small, broad exotherm at about 267° C. No thermal events are seen

in the DSC thermogram at lower temperatures, suggesting that it is an unsolvated material.

5.2.8 Form H

Form H can be obtained by storing Form E at room temperature and 0% RH for about 7 days. A representative XRPD pattern is shown in FIG. 30. The pattern is characterized by a peak at 15 degrees 2 θ , and two other peaks at 26 and 31 degrees 2 θ .

Representative thermal characteristics are shown in FIG. 31. Form H loses about 1.67% volatiles up to about 150° C. Weight loss above about 150° C. is attributed to decomposition. Karl Fischer data shows that Form H typically contains about 1.77% water (about 0.26 moles), suggesting that the weight loss seen in the TG is due to dehydration. The DSC thermogram shows a broad endotherm between about 50° C. and about 125° C., corresponding to the dehydration of Form H and a sharp endotherm at about 269° C., which is likely due to a melt.

When slurried in water with either Forms A or B, after about 14 days Form H can convert to Form E. When slurried in THF, Form H can convert to Form A. When slurried in acetone, Form H can convert to Form C.

In sum, Form H is a crystalline solid, hydrated with about 0.25 moles of water, which has a DSC thermogram exhibiting an endotherm between about 50 and 125° C. and an endotherm at about 269° C.

5.3 Methods of Use and Pharmaceutical Compositions

Polymorphs of the invention exhibit physical characteristics that are beneficial for drug manufacture, storage or use. All polymorphs of the invention have utility as pharmaceutically active ingredients or intermediates thereof.

This invention encompasses methods of treating and preventing a wide variety of diseases and conditions using polymorphs of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. In each of the methods, a therapeutically or prophylactically effective amount of the compound is administered to a patient in need of such treatment or prevention. Examples of such disease and conditions include, but are not limited to, diseases associated with undesired angiogenesis, cancer (e.g., solid and blood borne tumors), inflammatory diseases, autoimmune diseases, and immune diseases. Examples of cancers and pre-cancerous conditions include those described in U.S. Pat. Nos. 6,281,230 and 5,635,517 to Muller et al. and in various U.S. patent applications to Zeldis, including application Ser. Nos. 10/411,649, filed Apr. 11, 2003 (Treatment of Myelodysplastic Syndrome); 10/438,213 filed May 15, 2003 (Treatment of Various Types of Cancer); 10/411,656, filed Apr. 11, 2003 (Treatment of Myeloproliferative Diseases). Examples of other diseases and disorders that can be treated or prevented using compositions of the invention are described in U.S. Pat. Nos. 6,235,756 and 6,114,335 to D'Amato and in other U.S. patent applications to Zeldis, including Ser. Nos. 10/693,794, filed Oct. 23, 2003 (Treatment of Pain Syndrome) and 10/699,154, filed Oct. 30, 2003 (Treatment of Macular Degeneration). The entirety of each of the patents and patent applications cited herein is incorporated herein by reference.

Depending on the disease to be treated and the subject's condition, polymorphs of the invention can be administered by oral, parenteral (e.g., intramuscular, intraperitoneal, intravenous, ICV, intracisternal injection or infusion, subcutaneous injection, or implantation), inhalation spray, nasal, vaginal, rectal, sublingual, or topical routes of administration and may be formulated, alone or together, in suitable dosage unit

US 7,855,217 B2

11

formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles appropriate for each route of administration. Because individual polymorphs have different dissolution, stability, and other properties, the optimal polymorph used in methods of treatment may depend on the route of administration. For example, forms that are readily soluble in aqueous solutions are preferably used to provide liquid dosage forms, whereas forms that exhibit great thermal stability may be preferred in the manufacture of solid dosage forms (e.g., tablets and capsules).

Although the physical characteristics of polymorphs can, in some cases, affect their bioavailability, amounts of the polymorphs that are therapeutically or prophylactically effective in the treatment of various disease and conditions can be readily determined by those of ordinary skill in the pharmacy or medical arts. In certain embodiments of the invention, a polymorph is administered orally and in a single or divided daily doses in an amount of from about 0.10 to about 150 mg/day, or from about 5 to about 25 mg/day. In other embodiments, a polymorph is administered every other day in an amount of from about 0.10 to about 150 mg/day, or from about 5 to about 25 mg/day.

The invention encompasses pharmaceutical compositions and single unit dosage forms that can be used in methods of treatment and prevention, which comprise one or more polymorphs of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione and optionally one or more excipients or diluents. Specific compositions and dosage forms are disclosed in the various patents and patent applications incorporated herein by reference. In one embodiment, a single dosage form comprises a polymorph (e.g., Form B) in an amount of about 5, 10, 25 or 50 mg.

6. EXAMPLES

6.1 Polymorph Screen

A polymorph screen to generate the different solid forms of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione was carried out as follows.

A weighed sample of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (usually about 10 mg) was treated with aliquots of the test solvent. Solvents were either reagent or HPLC grade. The aliquots were usually about 200 μ L. Between additions, the mixture was usually shaken or sonicated. When the solids dissolved, as judged by visual inspection, estimated solubilities were calculated. Solubilities were estimated from these experiments based on the total solvent used to provide a solution. Actual solubilities may have been greater than those calculated due to the use of too-large solvent aliquots or to a slow rate of dissolution.

Samples were created by generating solutions (usually about 30 mg in 20 mL) at elevated temperatures, filtering, and allowing the solution to evaporate whether in an open vial (hot fast evaporation) or in a vial covered with aluminum foil containing pinholes (hot slow evaporation).

Slurry experiments were also performed. Usually about 25 mg of solid was placed in either 3 or 5 mL of solvent. The samples were then placed on orbital shakers at either ambient temperature or 40° C. for 410 days.

Crystallizations were performed using various cooling methods. Solid was dissolved in a solvent at an elevated temperature (e.g., about 60° C.), filtered quickly and allowed to cool to room temperature. Once at room temperature, samples that did not crystallize were moved to a refrigerator. Solids were removed by filtration or decantation and allowed

12

to dry in the air. Crash cools were performed by dissolving solid in a solvent at an increased temperature (e.g., about 45-65° C.) followed by cooling in a dry ice/acetone bath.

Hygroscopicity studies were performed by placing portions of each polymorph in an 84% relative humidity chamber for approximately one week.

Desolvation studies were carried out by heating each polymorph in a 70° C. oven for approximately one week.

Interconversion experiments were carried out by making slurries containing two forms in a saturated solvent. The slurries were agitated for approximately 7-20 days at ambient temperature. The insoluble solids were recovered by filtration and analyzed using XRPD.

6.2 Preparation of Polymorphic Forms

Eight solid forms of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione were prepared as described below.

Form A was obtained by crystallization from various non-aqueous solvents including 1-butanol, butyl acetate, ethanol, ethyl acetate, methanol, methyl ethyl ketone, and tetrahydrofuran. Form B was also obtained by crystallization from the solvents hexane, toluene and water. Form C was obtained from evaporations, slurries, and slow cools in acetone solvent systems. Form D was obtained from evaporations in acetonitrile solvent systems. Form E was obtained most readily by slurrying 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione in water. Form F was obtained by complete desolvation of Form E. It is found to be an unsolvated, crystalline material that melts at about 269° C. Form G was obtained by slurrying forms B and E in THF. Form H was obtained by stressing Form E at room temperature and 0% RH for 7 days.

6.2.1 Synthesis of Polymorphs B and E

Form B is the desired polymorph for the active pharmaceutical ingredient (API) of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. This form has been used in the formulation of API into drug product for clinical studies. Three batches were produced as apparent mixtures of polymorphs in the non-micronized API of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione.

Development work was carried out to define a process that would generate polymorph B from this mixture of polymorphs and could be implemented for strict polymorphic controls in the validation batches and future manufacturing of API of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. Characterization of polymorphic forms produced during the work was performed by XRPD, DSC, TGA and KF.

A process was also developed for the large-scale preparation of Form E. Polymorph E material was prepared in order to carry out a comparison with polymorph B drug product in capsule dissolution testing of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. 150 g of a mixture of polymorphs in 3 L of water was stirred at room temperature for 48 hours. The product was collected by filtration and dried at 25° C. for 24 hours under vacuum. XRPD, DSC, TGA, KF and HPLC analyses confirmed that the material isolated was polymorph E.

In a preliminary work, it was demonstrated that stirring a suspension of a mixture of polymorphs of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione with water at high temperature (75° C.) for an extended period of time converted this mixture of polymorphs exclusively to form B. Several specific parameters were identified including temperature, solvent volume and drying parameters (temperature

US 7,855,217 B2

13

and vacuum). XRPD, DSC, TGA, KF and HPLC analyses were used to characterize all of the batches. After completing the optimization work, the optimized process was scaled-up to 100-200 g on three lots of API. Drying studies were carried out at 20° C., 30° C. and 40° C., and 65° C. with a vacuum of 150 mm of Hg. The results are shown in Tables 1-5.

The cooling and holding periods of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione slurry were studied. The experimental laboratory data suggests that polymorph B seems to be forming first, and overtime equilibration to polymorph E at RT conditions occurs, therefore generating a mixture of polymorphs B and E. This result supports the fact that polymorph B seems to be a kinetic product, and that prolonged processing time converts the material to polymorph E resulting in a mixture of polymorphs B and E.

A laboratory procedure was developed to exclusively produce polymorph B of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. The procedure includes a stirred 10 volume water slurry at ~75° C. for 6-24 hours. The following preferred process parameters have been identified:

1. Hot slurry temperature of 70-75° C.
2. Product filtration of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)piperidine-2,6-dione at 65-75° C.
3. Drying under vacuum at 60-70° C. is preferred for an efficient removal of unbound water in 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione wet cake.
4. The filtration step of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione may be a time sensitive operation. The use of efficient solid-liquid separation equipment is preferred.
5. Holding periods of water-wet cake of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione at KF higher than 5% may cause the kinetic equilibrations of polymorph B to mixed polymorphs of E and B.

Drying to KF<4.0% water was achieved in ~3 hours (30-70° C., 152 mm Hg). Polymorphs B and E were distinguished by the water levels as measured by KF and TGA. The reference sample of polymorph B is micronized API. In order to make accurate comparison by XRPD samples were gently grinded before submission for analysis. This increases the clarity of the identification of the polymorphic form. All samples were analyzed for XRPD, DSC, TGA, KF and HPLC.

TABLE 1

| Preliminary Studies | | | |
|---------------------|-------------------------------------|--------------------|--------------------|
| Amount | Reaction conditions | Analysis | Results/conclusion |
| 2 g | Water, rt, 48 h | XRPD, DSC, TGA, KF | Polymorph E |
| 25 g | Water, rt, 48 h | XRPD, DSC, TGA, KF | Polymorph E |
| 5 g | Water, 70-75° C., 24 h then rt 24 h | XRPD, DSC, TGA, KF | Polymorph B |
| 1 g | 9:1 Acetone - water, Slow evpo. | XRPD, DSC, TGA, KF | Polymorph Mixture |
| 1 g | 175° C. 1 h in an oven | XRPD, DSC, TGA, KF | Polymorph A |
| 0.5 g (polymorph A) | Water, rt, 24 h | XRPD, DSC, TGA, KF | Polymorph E |
| 1 g polymorph B | Water, rt, 48 h | XRPD, DSC, TGA, KF | Polymorph E |
| 1 g polymorph E | Water, 70-75° C., 24 h | XRPD, DSC, TGA, KF | Polymorph B |
| 1 g | Slurry in heptane | XRPD, DSC, TGA, KF | No change |

14

TABLE 2

| Optimization of Temperature, Time and Solvent Volume | | | | |
|--|-------------------|-------------|----------|--------------------|
| Amount | Amount Water (mL) | Temp (° C.) | Time (h) | Results/conclusion |
| 10 g | 50 | 75 | 6 | Mix |
| 10 g | 50 | 75 | 24 | Polymorph B |
| 10 g | 100 | 70 | 6 | Polymorph B |
| 10 g | 100 | 70 | 14 | Polymorph B |
| 10 g | 100 | 70 | 21 | Polymorph B |
| 10 g | 100 | 75 | 6 | Polymorph B |
| 10 g | 100 | 75 | 24 | Polymorph B |
| 10 g | 100 | 75 | 6 | Polymorph B |
| 10 g | 100 | 75 | 19 | Polymorph B |
| 10 g | 100 | 75 | 14 | Polymorph B |
| 10 g | 100 | 75 | 24 | Polymorph B |
| 5 g | 100 | 75 | 18 | Polymorph B |
| 10 g | 100 | 80 | 6 | Polymorph B |
| 10 g | 100 | 80 | 20 | Polymorph B |
| 10 g | 200 | 45 | 6 | Polymorph B + E |
| 10 g | 200 | 45 | 24 | Polymorph E |
| 10 g | 200 | 60 | 48 | Polymorph B |
| 10 g | 200 | 75 | 6 | Mix |
| 10 g | 200 | 75 | 24 | Polymorph B |
| 10 g | 200 | 75 | 13 | Polymorph B |
| 10 g | 200 | 75 | 24 | Polymorph B |

Optimum conditions were determined to be 10 volumes of solvent (H₂O), 70-80° C. for 6-24 hours.

TABLE 3

| Holding Time | | | | |
|--------------|-----------------------------|------------------|---------------------|--------------------|
| Amount | Reaction Conditions | Holding Time (h) | Holding Temp (° C.) | Results/Conclusion |
| 5 g | Water, 70-75° C., 24 h | 24 | 23-25 | Polymorph B |
| 1 g | Water, 70-75° C., 24 h | 48 | 23-25 | Polymorph E |
| Polymorph B | | | | |
| 2 g | Water, 40 mL | 16 | 23-25 | Polymorph E |
| 150 g | Water, 3.0 L | 24 | 23-25 | Polymorph E |
| 150 g | Water, 3.0 L | 48 | 23-25 | Polymorph E |
| 10 g | Water, 100 mL, 24 h, 75° C. | 18 | 23-25 | Polymorph B |
| 10 g | Water, 100 mL, 24 h, 75° C. | 18 | 40 | Polymorph B |
| 10 g | Water, 200 mL, 24 h, 75° C. | 14 | -5 | Mix |
| 10 g | Water, 200 mL, 24 h, 75° C. | 14 | 23-25 | Polymorph E |
| 10 g | Water, 200 mL, 24 h, 75° C. | 14 | 40 | Mix |
| 10 g | Water, 100 mL, 24 h, 75° C. | 21 | 23-25 | Polymorph E |
| 10 g | Water, 100 mL, 24 h, 75° C. | 21 | 40 | Mix |
| 10 g | Water, 100 mL, 14 h, 75° C. | 2 | 23-25 | Mix |

Holding time gave mixed results and it was determined that the material should be filtered at 60-65° C. and the material washed with 0.5 volume of warm (50-60° C.) water.

TABLE 4

| Scale-up Experiments | | | | |
|----------------------|------------------|-------------|----------|--------------------|
| Amount | Amount Water (L) | Temp (° C.) | Time (h) | Results/Conclusion |
| 100 g | 1.0 | 75 | 6 | Polymorph B |
| 100 g | 1.0 | 75 | 22 | Polymorph B |

US 7,855,217 B2

15

TABLE 4-continued

| Scale-up Experiments | | | | |
|----------------------|------------------|-------------|----------|--------------------|
| Amount | Amount Water (L) | Temp (° C.) | Time (h) | Results/Conclusion |
| 100 g | 1.0 | 75 | 6 | Polymorph B |
| 100 g | 1.0 | 75 | 24 | Polymorph B |
| 100 g | 1.0 | 75 | 6 | Polymorph B |
| 100 g | 1.0 | 75 | 22 | Polymorph B |

TABLE 5

| Drying Studies | | | | | |
|----------------|-----------------|--------------------|----------------|---------|--------------------|
| Amount | Drying Time (h) | Drying Temp (° C.) | Vacuum (mm Hg) | KF§ (%) | Results/Conclusion |
| 100 g | 0 | — | — | 3.690 | Polymorph B |
| 100 g | 3 | 30 | 152 | 3.452 | Polymorph B |
| 100 g | 8 | 30 | 152 | 3.599 | Polymorph B |
| 100 g | 0 | — | — | 3.917 | Polymorph B |
| 100 g | 5 | 40 | 152 | 3.482 | Polymorph B |
| 100 g | 22 | 40 | 152 | 3.516 | Polymorph B |
| 100 g | 3 | 40 | 152 | 3.67 | Polymorph B |
| 100 g | 22 | 40 | 152 | 3.55 | Polymorph B |

*Reaction Conditions: Water 1 L, 75° C., 22-24 h;

§Average of 2 runs.

Drying studies determined that the material should be dried at 35-40° C., 125-152 mm Hg for 3 to 22 h or until the water content reaches $\leq 4\%$ w/w.

For a large scale preparation of polymorph E (5222-152-B), a 5-L round bottom flask was charged with 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione (150 g, 0.579 mol) and water (3000 mL, 20 vol). The mixture was mechanically stirred at room temperature (23-25° C.) for 48 h under nitrogen atmosphere.

Samples were taken after 24 h and 48 h before the mixture was filtered and air-dried on the filter for 1 h. The material was transferred to a drying tray and dried at room temperature (23-25° C.) for 24 h. KF analysis on the dried material showed water content of 11.9%. The material was submitted for XRPD, TGA, DSC and HPLC analysis. Analysis showed the material was pure polymorph E.

For a large scale preparation of polymorph B (5274-104), a 2 L-3-necked round bottom flask was charged with 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione (polymorph mixture, 100 g, 0.386 mol) and water (1000 mL, 10.0 vol). The mixture was heated to 75° C. over approximately 30 minutes with mechanical stirring under nitrogen atmosphere.

Samples were taken after 6 h and 24 h before the mixture was allowed to cool to 60-65° C., filtered and the material washed with warm (50-60° C.) water (50 mL, 0.5 vol). The material was transferred to a drying tray and dried at 30° C., 152 mm Hg for 8 h. KF analysis on the dried material showed water content of 3.6%. After grinding the material was submitted for XRPD, TGA, DSC and HPLC analysis. Analysis showed the material was pure polymorph B. The results of the analyses are shown in FIGS. 32-46.

6.3 X-Ray Powder Diffraction Measurements

X-ray powder diffraction analyses were carried out on a Shimadzu XRD 6000 X-ray powder diffractometer using Cu K α radiation. The instrument is equipped with a fine-focus X-ray tube. The tube voltage and amperage were set at 40 kV

16

and 40 mA, respectively. The divergence and scattering slits were set at 1° and the receiving slit was set at 0.15 mm. Diffracted radiation was detected by a NaI scintillation detector. A theta-two theta continuous scan at 3°/min (0.4 sec/0.02° step) from 2.5 degrees 2 θ to 40 degrees 2 θ was used. A silicon standard was analyzed each day to check the instrument alignment.

X-ray powder diffraction analyses were also carried out using Cu K α radiation on an Inel XRG-3000 diffractometer equipped with a curved position-sensitive detector. Data were collected in real time over a theta-two theta range of 120° at a resolution of 0.03°. The tube voltage and current were 40 kV and 30 mA, respectively. A silicon standard was analyzed each day to check for instrument alignment. Only the region between 2.5 and 40 degrees 2 θ is shown in the figures.

6.4 Thermal Analysis

TG analyses were carried out on a TA Instrument TGA 2050 or 2950. The calibration standards were nickel and alumel. Approximately 5 mg of sample was placed on a pan, accurately weighed, and inserted into the TG furnace. The samples were heated in nitrogen at a rate of 10° C./min, up to a final temperature of 300 or 350° C.

DSC data were obtained on a TA 2920 instrument. The calibration standard was indium. Approximately 2-5 mg samples were placed into a DSC pan and the weight accurately recorded. Crimped pans with one pinhole were used for analysis and the samples were heated under nitrogen at a rate of 10° C./min, up to a final temperature of 350° C.

Hot-stage microscopy was carried out using a Kofler hot stage mounted on a Leica Microscope. The instrument was calibrated using USP standards.

A TA Instruments TGA 2050 interfaced with a Nicolet model 560 Fourier transform IR spectrophotometer, equipped with a globar source, XT/KBr beamsplitter, and deuterated triglycine sulfate (DTGS) detector, was utilized for TG-IR experiments. The IR spectrometer was wavelength calibrated with polystyrene on the day of use while the TG was temperature and weight calibrated biweekly, using indium for the temperature calibration. A sample of approximately 10 mg of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione was weighed into an aluminum pan and heated from 25 to 30° C. to 200° C. at a rate of 20° C./min with a helium purge. IR spectra were obtained in series, with each spectrum representing 32 co-added scans at a resolution of 4 cm⁻¹. Spectra were collected with a 17-second repeat time. TG/IR analysis data are presented as Gram-Schmidt plots and IR spectra linked to the time. Gram-Schmidt plots show total IR intensity vs. time; hence, the volatiles can be identified at each time point. They also show when the volatiles are detected. From the Gram-Schmidt plots, time points were selected and the IR spectra of these time points are presented in the stacked linked spectra. Each spectrum identifies volatiles evolving at that time point. Volatiles were identified from a search of the HR Nicolet TGA vapor phase spectral library. The library match results are also presented to show the identified vapor.

6.5 Spectroscopy Measurements

Raman spectra were acquired on a Nicolet model 750 Fourier transform Raman spectrometer utilizing an excitation wavelength of 1064 nm and approximately 0.5 W of Nd:YAG laser power. The spectra represent 128 to 256 co-added scans acquired at 4 cm⁻¹ resolution. The samples were prepared for analysis by placing the material in a sample holder and posi-

tioning this in the spectrometer. The spectrometer was wavelength calibrated using sulfur and cyclohexane at the time of use.

The mid-IR spectra were acquired on a Nicolet model 860 Fourier transform IR spectrophotometer equipped with a globar source XT/KBr beamsplitter and a deuterated triglycine sulfate (DTGS) detector. A Spectra-Tech, Inc. diffuse reflectance accessory was utilized for sampling. Each spectrum represents 128 co-added scans at a spectral resolution of 4 cm^{-1} . A background data set was acquired with an alignment mirror in place. A single beam sample data set was then acquired. Subsequently, a log 1/R (where R=reflectance) spectrum was acquired by rationing the two data sets against each other. The spectrophotometer was calibrated (wavelength) with polystyrene at the time of use.

6.6 Moisture Sorption/Desorption Measurements

Moisture sorption/desorption data were collected on a VTI SGA-100 moisture balance system. For sorption isotherms, a sorption range of 5 to 95% relative humidity (RH) and a desorption range of 95 to 5% RH in 10% RH increments was used for analysis. The sample was not dried prior to analysis. Equilibrium criteria used for analysis were less than 0.0100 weight percent change in 5 minutes with a maximum equilibration time of 3 hours if the weight criterion was not met. Data were not corrected for the initial moisture content of the samples.

6.7 Solution Proton NMR Measurements

NMR spectra not previously reported were collected at SSCI, Inc, 3065 Kent Avenue, West Lafayette, Ind. Solution phase ^1H NMR spectra were acquired at ambient temperature on a Bruker model AM spectrometer. The ^1H NMR spectrum represents 128 co added transients collected with a 4 μsec pulse and a relaxation delay time of 5 seconds. The free induction decay (FID) was exponentially multiplied with a 0.1 Hz Lorentzian line broadening factor to improve the signal-to-noise ratio. The NMR spectrum was processed utilizing GRAMS software, version 5.24. Samples were dissolved in dimethyl sulfoxide- d_6 .

The scope of this invention can be understood with reference to the appended claims.

6.8 Intrinsic Dissolution and Solubility Studies

Intrinsic dissolution experiments were conducted on Form A (anhydrous), Form B (hemihydrate), and Form E (dihydrate) of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione. Equilibrium solubility experiments were conducted on Forms A and B. Aliquots were analyzed by ultraviolet-visible spectrophotometry, and the solids remaining from each experiment were analyzed by X-ray powder diffraction (XRPD).

6.8.1 Experimental

6.8.1.1 Dissolution

Dissolution experiments were carried out in a VanKel VK6010-8 dissolution apparatus equipped with a VK650A heater/circulator. An intrinsic dissolution apparatus (Woods apparatus) was used. Samples were compressed at 1.5 metric tons (1000 psi) for 1 min using the Woods apparatus in a hydraulic press, giving a sample surface of 0.50 cm^2 . A dissolution medium consisting of 900 mL HCl buffer, pH 1.8, with 1% sodium lauryl sulfate, was used for each experiment. The medium was degassed by vacuum filtration through a 0.22- μm nylon filter disk and maintained at 37° C. The appa-

ratus was rotated at 50 rpm for each experiment. Aliquots were filtered immediately using 0.2- μm nylon syringe filters. In some cases, the undissolved solids were recovered and analyzed by X-ray powder diffraction (XRPD).

6.8.1.2 Solubility

Equilibrium solubility experiments were conducted in a 100-mL, three-neck, round-bottom flask immersed in a constant temperature oil bath maintained at 25° C. A solid sample of 400-450 mg was stirred in 50 mL of dissolution medium (HCl buffer, pH 1.8, with 1% sodium lauryl sulfate) using a mechanical stir rod. Aliquots were filtered using 0.2- μm nylon syringe filters and immediately diluted 1 mL→50 mL, then 5 mL→25 mL with dissolution medium in Class A glassware, a final dilution factor of 250.

6.8.1.3 UV-Vis Spectrophotometry

Dissolution and solubility samples solutions were analyzed by a Beckman DU 640 single-beam spectrophotometer. A 1.000-cm quartz cuvette and an analysis wavelength of 228.40 nm were utilized. The detector was zeroed with a cuvette filled with dissolution medium.

6.8.1.4 X-Ray Powder Diffraction

XRPD analyses were carried out on a Shimadzu XRD-6000 X-ray powder diffractometer using Cu $K\alpha$ radiation. The instrument is equipped with a fine focus X-ray tube. The tube power and amperage were set at 40 kV and 40 mA, respectively. The divergence and scattering slits were set at 1° and the receiving slit was set at 0.15 mm. Diffracted radiation was detected by a NaI scintillation detector. A theta-two theta continuous scan at 3°/min (0.4 sec/0.02° step) from 2.5 to 40° 2 θ was used. A silicon standard was analyzed each day to check the instrument alignment. Samples were packed in an aluminum holder with silicon insert.

6.8.2 Results

The results of these solubility and intrinsic studies are summarized in Table 6. Both the solubility and dissolution experiments were conducted in a medium of HCl buffer, pH 1.8, containing 1% sodium lauryl sulfate. Form A was found to be unstable in the medium, converting to Form E. The solubilities of Forms A, B, and E were estimated to be 6.2, 5.8, and 4.7 mg/mL, respectively. The dissolution rates of Forms A, B, and E were estimated to be 0.35, 0.34, and 0.23 mg/mL, respectively.

6.8.2.1 UV-Vis Spectrophotometry Method Development

A UV-V is scan of the dissolution medium (blanked with an empty cuvette) was done to identify any interfering peaks. A small peak at 225 nm was present as shown in FIG. 47.

Solutions of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione at varying concentrations were analyzed by UV-V is spectrophotometry. A preliminary scan of a 1.0 mg/mL solution was done, with the instrument blanked with dissolution medium. The solution was highly absorbing and noisy from 200-280 nm, making dilution necessary.

A 0.04 mg/mL solution of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione was then scanned from 200-300 nm. The plot was still noisy between 200 and 230 nm as shown in FIG. 48. The sample was further diluted to 0.008 mg/mL. A wavelength scan of 200-350 nm for this sample showed a peak a 228.4 nm with no interference, as shown in FIG. 49. Therefore, a wavelength of 228.4 was chosen for analysis of the solubility and dissolution samples.

A six-point calibration curve was generated with standards of the following concentrations: 0.001 mg/mL, 0.002 mg/mL, 0.005 mg/mL, 0.010 mg/mL, 0.015 mg/mL, and 0.020 mg/mL (Notebook 569-90). A linearity coefficient of $R^2=0.9999$ was obtained as shown in FIG. 50.

6.8.2.2 Solubility

A sample consisting of 449.4 mg Form A was slurried in dissolution medium. Particle size was not controlled. Aliquots were taken at 7, 15, 30, 60, 90, and 150 min. The concentration reached 6.0 mg/mL by the first time point The highest concentration reached was 6.2 mg/mL, at 30 min. From that point the concentration decreased, reaching 4.7 mg/mL at 150 min as in FIG. 51. The solids remaining at the final time point were analyzed by XRPD and found to be Form E as shown in Table 7. No peaks attributed to Form A can be seen in the pattern. Since the concentration did not plateau at 4.7 mg/mL, the solubility of Form E may be lower than that.

A sample consisting of 401.4 mg Form B was slurried in dissolution medium. Particle size was not controlled. Aliquots were taken at 7, 15, 30, 60, 90, 180, 420, and 650 min. Form B dissolved much more slowly than Form A, reaching 3.3 mg/mL in 90 min. The concentration stabilized at 5.6-5.7 mg/mL at the final three time points as in FIG. 52. The remaining solids were shown to be Form B as in Table 7, suggesting Form B has good stability in water.

A summary of the solubilities is given in Table 6. The amounts dissolved at each time point are shown in Tables 8 and 9.

TABLE 6

| Summary of Results | | | | |
|--------------------|------------|--------------------------|--------------------------|------------------------------------|
| Form | Solubility | Intrinsic Dissolution #1 | Intrinsic Dissolution #2 | Average Intrinsic Dissolution Rate |
| Form A | 6.2 mg/mL | 0.35 | 0.22 ^a | 0.29 ^a |
| Form B | 5.8 mg/mL | 0.35 | 0.32 | 0.34 |
| Form E | 4.7 mg/mL | 0.21 | 0.25 | 0.23 |

^aThe Form A dissolution experiment #2 may have converted to Form E on the surface of the disk, skewing the average rate lower.

TABLE 7

| Experimental Details | |
|----------------------|------------|
| Experiment | Final Form |
| Pressed Form A | A |
| Pressed Form B | B |
| Form A Solubility | E |
| Form B Solubility | B |
| Form A Dissolution | — |
| Form A Dissolution | A |
| Form B Dissolution | — |
| Form B Dissolution | B |
| Form E Dissolution | E |
| Form E Dissolution | — |

TABLE 8

| Form A Solubility | |
|-------------------|-----------------------|
| Time Point (min) | Concentration (mg/mL) |
| 7 | 6.00 |
| 15 | 6.11 |
| 30 | 6.16 |
| 60 | 6.10 |
| 90 | 5.46 |
| 150 | 4.73 |

TABLE 9

| Form B Solubility | |
|-------------------|-----------------------|
| Time Point (min) | Concentration (mg/mL) |
| 7 | 1.63 |
| 15 | 2.14 |
| 30 | 2.33 |
| 60 | 2.94 |
| 90 | 3.34 |
| 180 | 5.67 |
| 420 | 5.76 |
| 650 | 5.61 |

6.8.2.3 Intrinsic Dissolution

Approximately 200 mg each of Forms A and B were compressed into disks in the Woods apparatus using 2 metric tons of pressure. The samples were subsequently scraped out, ground gently, and analyzed by XRPD. The study showed that compression and grinding does not cause a form change in either case. (See Table 7).

Two preliminary dissolution runs were performed. The disks fractured to some extent in both experiments, compromising the requirement of constant surface area.

The first experiment of intrinsic dissolution that strictly followed the USP chapter on intrinsic dissolution utilized approximately 150 mg each of Forms A and B. Seven aliquots, beginning at 5 min and ending at 90 min, were taken to maintain sink conditions. The experiment resulted in linear dissolution profiles, with a rate of 0.35 mg per cm² per minute for both forms. The Form E experiment was done later under the same conditions and added to the graph for comparison. (See FIG. 53). The Form E dissolution rate was 0.21 mg per cm² per minute, significantly lower than the dissolution rate of Forms A and B. This is in line with expectations based on the solubility data. The crystal form of the remaining solids did not change in any case.

The second experiment utilized approximately 250 mg each of Forms A and B. The Form E experiment (135 mg) was done later and added to the graph for comparison. (See FIG. 54). Nine aliquots were taken, beginning at 5 min and ending at 150 min. The dissolution rates were 0.22, 0.32, and 0.25 mg per cm² per minute, respectively, for Forms A, B, and E. The dissolution rate for Form A in this experiment was low, while the rates for Forms B and E were similar to those found in the first experiment. It is believed that in this case, a thin layer of the Form A sample disk may have converted to Form E upon exposure to water. This is supported by the evidence of rapid conversion of Form A to Form E in the solubility experiment. The diffraction pattern of the undissolved solids does not indicate a form change. However, the bulk of the sample disk is not exposed to water. Therefore, the true intrinsic dissolution rate of Form A is believed to be close to 0.35 mg per cm² per minute. An insufficient quantity of Form A was available to repeat the experiment.

A summary of the intrinsic dissolution rates is given in Table 6. The amounts dissolved at each time point are summarized in Tables 10 and 11.

TABLE 10

| Intrinsic Dissolution Experiment #1 Results | | | |
|---|---------------------|---------------------|---------------------|
| Time Point | Form A ^a | Form B ^a | Form E ^a |
| 5 min | 5.76 | 10.80 ^b | 2.70 |
| 10 min | 7.73 | 6.85 | 4.13 |

US 7,855,217 B2

21

TABLE 10-continued

| Intrinsic Dissolution Experiment #1 Results | | | |
|---|---------------------|---------------------|---------------------|
| Time Point | Form A ^a | Form B ^a | Form E ^a |
| 20 min | 11.31 | 10.25 | 6.96 |
| 30 min | 15.59 | 14.35 | 9.60 |
| 45 min | 21.98 | 20.57 | 12.57 |
| 60 min | 27.11 | 25.70 | 15.16 |
| 90 min | 34.17 | 34.34 | 20.82 |

^aResults are reported as Cumulative Amount Dissolved per Unit Area (mg/cm²)^bThis date point not included in graph since the value is higher than the next two data points.

TABLE 11

| Intrinsic Dissolution Experiment #2 Results | | | |
|---|---------------------|---------------------|---------------------|
| Time Point | Form A ^a | Form B ^a | Form E ^a |
| 5 min | 4.50 | 5.04 | 3.06 |
| 10 min | 5.22 | 6.12 | 4.31 |
| 20 min | 7.54 | 7.73 | 11.40 |
| 30 min | 11.46 | 12.72 | 11.93 |
| 45 min | 15.01 | 17.33 | 14.72 |
| 60 min | 18.38 | 21.93 | 18.52 |
| 90 min | 24.38 | 31.64 | 26.24 |
| 120 min | 30.35 | 41.31 | 33.56 |
| 150 min | 35.26 | 49.54 | 40.82 |

^aResults are reported as Cumulative Amount Dissolved per Unit Area (mg/cm²)

6.9 Analyses of Mixtures of Polymorphs

This invention encompasses mixtures of different polymorphs. For example, an X-ray diffraction analysis of one production sample yielded a pattern that contained two small peaks seen at approximately 12.6° and 25.8° 2θ in addition to those representative of Form B. In order to determine the composition of that sample, the following steps were performed.

- 1) Matching of the new production pattern to known forms along with common pharmaceutical excipients and contaminants;
- 2) Cluster analysis of the additional peaks to identify if any unknown phase is mixed with the original Form B;
- 3) Harmonic analysis of the additional peaks to identify if any preferred orientation may be present or if any changes in the crystal habit may have occurred; and
- 4) Indexing of the unit cells for both Form B and the new production sample to identify any possible crystallographic relationships.

Based on these tests, which can be adapted for the analysis of any mixture of polymorphs, it was determined that the sample contained a mixture of polymorph forms B and E.

6.10 Dosage Form

Table 12 illustrates a batch formulation and single dosage formulation for a 25 mg single dose unit of a polymorphic form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione.

22

TABLE 12

| Formulation for a 25 mg capsule | | | |
|--|-------------------|----------------------|---------------------|
| Material | Percent By Weight | Quantity (mg/tablet) | Quantity (kg/batch) |
| Polymorphic Form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione | 40.0% | 25 mg | 16.80 kg |
| Pregelatinized Corn Starch, NF | 59.5% | 37.2 mg | 24.99 kg |
| Magnesium Stearate | 0.5% | 0.31 mg | 0.21 kg |
| Total | 100.0% | 62.5 mg | 42.00 kg |

The pregelatinized corn starch (SPRESS B-820) and polymorphic form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione components are passed through a screen (i.e., a 710 μm screen) and then loaded into a Diffusion Mixer with a baffle insert and blended for about 15 minutes. The magnesium stearate is passed through a screen (i.e., a 210 μm screen) and added to the Diffusion Mixer. The blend is then encapsulated in capsules using a Dosator type capsule filling machine.

The entire scope of this invention is not limited by the specific examples described herein, but is more readily understood with reference to the appended claims.

What is claimed is:

1. A solid form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione comprising crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione hemihydrate, wherein the crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione hemihydrate is present at greater than about 80% by weight of the solid form.

2. A solid form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione comprising crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione hemihydrate, wherein the crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione hemihydrate is present at greater than about 90% by weight of the solid form.

3. A solid form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione comprising crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione hemihydrate, wherein the crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione hemihydrate is present at greater than about 95% by weight of the solid form.

4. A solid form of 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione comprising crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione hemihydrate, wherein the crystalline 3-(4-amino-1-oxo-1,3 dihydro-isoindol-2-yl)-piperidine-2,6-dione hemihydrate is present at greater than about 97% by weight of the solid form.

5. A pharmaceutical composition comprising a therapeutically effective amount of the solid form of claim 1, 2, 3, or 4, and a pharmaceutically acceptable excipient or carrier.

6. The pharmaceutical composition of claim 5, which is a single unit dosage form.

7. The pharmaceutical composition of claim 5, wherein the therapeutically effective amount is about 5 mg.

8. The pharmaceutical composition of claim 5, wherein the therapeutically effective amount is about 10 mg.

9. The pharmaceutical composition of claim 5, wherein the therapeutically effective amount is about 25 mg.

10. The pharmaceutical composition of claim 5, wherein the therapeutically effective amount is about 50 mg.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 7,855,217 B2
APPLICATION NO. : 12/335395
DATED : December 21, 2010
INVENTOR(S) : Jaworsky et al.

Page 1 of 11

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Please replace Drawing Sheets 15, 26, 31, 32, 33, 35, 36, 38, 39, and 40 (of 48) with the Replacement Drawing Sheets on the following pages of this Certificate of Correction:

Page 2 of 11: Replacement Drawing Sheet 15 of 48 (Fig. 15);

Page 3 of 11: Replacement Drawing Sheet 26 of 48 (Fig. 26);

Page 4 of 11: Replacement Drawing Sheet 31 of 48 (Fig. 31);

Page 5 of 11: Replacement Drawing Sheet 32 of 48 (Figs. 32 and 33);

Page 6 of 11: Replacement Drawing Sheet 33 of 48 (Figs. 34 and 35);

Page 7 of 11: Replacement Drawing Sheet 35 of 48 (Figs. 37 and 38);

Page 8 of 11: Replacement Drawing Sheet 36 of 48 (Figs. 39 and 40);

Page 9 of 11: Replacement Drawing Sheet 38 of 48 (Figs. 42 and 43);

Page 10 of 11: Replacement Drawing Sheet 39 of 48 (Figs. 44 and 45);

Page 11 of 11: Replacement Drawing Sheet 40 of 48 (Fig. 46).

Signed and Sealed this
Third Day of May, 2011

A handwritten signature in black ink that reads "David J. Kappos". The signature is written in a cursive, slightly slanted style.

David J. Kappos
Director of the United States Patent and Trademark Office

TGA (TOP) AND DSC (BOTTOM) OF FORM C




FIG.15




FIG.26

TGA (TOP) AND DSC (BOTTOM) OF FORM H




FIG.31

XRPD PATTERN OF POLYMORPH B




FIG.32

XRPD PATTERN OF POLYMORPH B




FIG.33

XRPD PATTERN OF POLYMORPH B

File: Process 5222-157-C

Date: 06/04/04 15:07 Step: 0.020° Cnt Time: 0.500 Sec.

Range: 1.50-40.00 (Deg) Step Scan Rate: 2.40 Deg/min.




FIG.34

XRPD PATTERN OF POLYMORPH E

File: Process 5222-152-B Form E

Date: 05/21/04 10:46 Step: 0.020° Cnt Time: 0.500 Sec.

Range: 1.50-40.00 (Deg) Step Scan Rate: 2.40 Deg/min.




FIG.35

TGA CURVE OF POLYMORPH B
Sample: 5274-104-B
Size 7.8320mg
Method: Ramp
TGA
Run Date: 09-Jun-04 17:04
Instrument: TGA 0.500 V5.3 Build 171




FIG.37

TGA CURVE OF POLYMORPH B
Sample: 5274-100-C
Size 10.9430mg
Method: Ramp
Comment: Evotec Batch 1675C H2O
TGA
Run Date: 03-Jun-04 11:20
Instrument: TGA 0.500 V5.3 Build 171




FIG.38

TGA CURVE OF POLYMORPH B
TGA
Sample: 5222-157-C
Size: 24.4610mg
Method: Ramp
Comment: Evotec Batch 17
Run Date: 07-Jun-04 09:46
Instrument: TGA 0.500 V5.3 Build 171




FIG.39

TGA CURVE OF POLYMORPH E
TGA
Sample: 5222-152-B
Size: 11.3850mg
Method: Ramp
Comment: Form E
Run Date: 21-May-04 09:34
Instrument: TGA 0.500 V5.3 Build 171




FIG.40

DSC OF POLYMORPH B
DSC

Sample: 5274-104-B
Size: 0.0000mg
Method: Cell constant calibration
Comment: OAI batch 15 22b

Run Date: 07-Jun-04 11:30
Instrument: DSC Q 1000 V7.3 Build 249




FIG.42

DSC OF POLYMORPH B
DSC

Sample: 5274-100-C
Size: 0.0000mg
Method: Cell constant calibration
Comment: Evotec Batch 1675C 10vol H2O 24h 3h dry

Run Date: 02-Jun-04 17:01
Instrument: DSC Q 1000 V7.3 Build 249




FIG.43

DSC OF POLYMORPH B
DSC

Sample: 5222-157-C
 Size: 0.0000mg
 Method: Cell constant calibration
 Comment: OAI batch 17

Run Date: 07-Jun-04 09:45
 Instrument: DSC Q 1000 V7.3 Build 249




FIG.44

DSC OF POLYMORPH E
DSC

Sample: 5222-152-B
 Size: 3.4500mg
 Method: Cell constant calibration
 Comment: Evotec batch 15 Form E

Run Date: 21-May-04 09:32
 Instrument: DSC Q 1000 V7.3 Build 249




FIG.45

Sample: 5222-161-A
Size: 0.0000mg
Method: Cell constant calibration
Comment: OAI batch 15

DSC OF POLYMORPH MIXTURE
DSC

Run Date: 11-Jun-04 12:44
Instrument: DSC Q 1000 V7.3 Build 249




FIG.46

EXHIBIT C



US007968569B2

(12) **United States Patent**
Zeldis

(10) **Patent No.:** **US 7,968,569 B2**
(45) **Date of Patent:** **Jun. 28, 2011**

(54) **METHODS FOR TREATMENT OF MULTIPLE MYELOMA USING 3-(4-AMINO-1-OXO-1,3-DIHYDRO-ISOINDOL-2-YL)-PIPERIDINE-2,6-DIONE**

(75) Inventor: **Jerome B. Zeldis**, Princeton, NJ (US)

(73) Assignee: **Celgene Corporation**, Summit, NJ (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 145 days.

(21) Appl. No.: **10/438,213**

(22) Filed: **May 15, 2003**

(65) **Prior Publication Data**

US 2004/0029832 A1 Feb. 12, 2004

Related U.S. Application Data

(60) Provisional application No. 60/380,842, filed on May 17, 2002, provisional application No. 60/424,600, filed on Nov. 6, 2002.

(51) **Int. Cl.**
A61K 31/445 (2006.01)

(52) **U.S. Cl.** **514/323**

(58) **Field of Classification Search** 514/321,
514/323

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

| | | | | |
|-----------|-----|---------|----------------------|---------|
| 3,536,809 | A | 10/1970 | Applezweig | |
| 3,598,123 | A | 8/1971 | Zaffaroni et al. | |
| 3,845,770 | A | 11/1974 | Theeuwes et al. | |
| 3,916,899 | A | 11/1975 | Theeuwes et al. | |
| 4,008,719 | A | 2/1977 | Theeuwes et al. | |
| 4,810,643 | A | 3/1989 | Souza | |
| 4,999,291 | A | 3/1991 | Souza | |
| 5,059,595 | A | 10/1991 | Le Grazie | |
| 5,073,543 | A | 12/1991 | Marshall et al. | |
| 5,120,548 | A | 6/1992 | McClelland et al. | |
| 5,134,127 | A | 7/1992 | Stella et al. | |
| 5,229,496 | A | 7/1993 | Deeley et al. | |
| 5,354,556 | A | 10/1994 | Sparks et al. | |
| 5,385,901 | A | 1/1995 | Kaplan et al. | |
| 5,391,485 | A | 2/1995 | Deeley et al. | |
| 5,393,870 | A | 2/1995 | Deeley et al. | |
| 5,528,823 | A | 6/1996 | Rudy, Jr. et al. | |
| 5,580,755 | A | 12/1996 | Souza | |
| 5,591,767 | A | 1/1997 | Mohr et al. | |
| 5,593,990 | A | 1/1997 | D'Amato | |
| 5,629,327 | A | 5/1997 | D'Amato | |
| 5,635,517 | A * | 6/1997 | Muller et al. | 514/323 |
| 5,639,476 | A | 6/1997 | Oshlack et al. | |
| 5,674,533 | A | 10/1997 | Santus et al. | |
| 5,698,579 | A | 12/1997 | Muller | |
| 5,712,291 | A | 1/1998 | D'Amato | |
| 5,731,325 | A | 3/1998 | Andrulis, Jr. et al. | |
| 5,733,566 | A | 3/1998 | Lewis | |
| 5,798,368 | A | 8/1998 | Muller et al. | |
| 5,874,448 | A | 2/1999 | Muller et al. | |
| 5,877,200 | A | 3/1999 | Muller | |
| 5,929,117 | A | 7/1999 | Muller et al. | |
| 5,955,476 | A | 9/1999 | Muller et al. | |

| | | | | |
|-----------|------|---------|----------------------|---------|
| 6,020,358 | A | 2/2000 | Muller et al. | |
| 6,071,948 | A | 6/2000 | D'Amato | |
| 6,077,822 | A * | 6/2000 | Dyrsting et al. | 514/8 |
| 6,114,355 | A | 9/2000 | D'Amato | |
| 6,140,346 | A | 10/2000 | Andrulis, Jr. et al. | |
| 6,228,879 | B1 | 5/2001 | Green et al. | |
| 6,235,756 | B1 | 5/2001 | D'Amato | |
| 6,281,230 | B1 * | 8/2001 | Muller et al. | 514/323 |
| 6,316,471 | B1 | 11/2001 | Muller et al. | |
| 6,326,388 | B1 | 12/2001 | Man et al. | |
| 6,335,349 | B1 | 1/2002 | Muller et al. | |
| 6,380,239 | B1 | 4/2002 | Muller et al. | |
| 6,395,754 | B1 | 5/2002 | Muller et al. | |
| 6,403,613 | B1 | 6/2002 | Man et al. | |
| 6,420,414 | B1 | 7/2002 | D'Amato | |
| 6,458,810 | B1 | 10/2002 | Muller et al. | |
| 6,469,045 | B1 | 10/2002 | D'Amato | |
| 6,476,052 | B1 | 11/2002 | Muller et al. | |
| 6,518,298 | B2 | 2/2003 | Green et al. | |
| 6,555,554 | B2 * | 4/2003 | Muller et al. | 514/323 |
| 6,673,828 | B1 | 1/2004 | Green et al. | |
| 7,119,106 | B2 * | 10/2006 | Muller et al. | 514/323 |
| 7,189,740 | B2 * | 3/2007 | Zeldis | 514/323 |
| 7,393,862 | B2 * | 7/2008 | Zeldis | 514/320 |
| 7,435,745 | B2 | 10/2008 | D'Amato | |

(Continued)

FOREIGN PATENT DOCUMENTS

WO WO 92/14455 9/1992

(Continued)

OTHER PUBLICATIONS

Corral et al. Immunomodulation by thalidomide and thalidomide analogues . . . Ann. Rheum. 1999; 58; 107-13.*
 Kyle et al. ("The Application of Thalidomide in Multiple Myeloma", Semin Oncol. Dec. 2001; 28(6):583-7).*
 Davies et al. ("Thalidomide and immunomodulatory derivatives augment natural killer cell cytotoxicity in multiple myeloma", Blood. Jul. 1, 2001;98(1):210-6).*
 Broder et al. ("Dideoxycytidine: current clinical experience and future prospects. A summary.", Am J Med. May 21, 1990;88(5B):31S-33S).*
 Filella et al. (Cancer Detect Prey. 1996;20(1):52-6).*
 Raza et al., 2001, "Thalidomide produces transfusion independence in long-standing refractory anemias of patients with myelodysplastic syndromes," Blood 98(4):958-965.

(Continued)

Primary Examiner — Frederick Krass
Assistant Examiner — Chris E Simmons
 (74) *Attorney, Agent, or Firm* — Jones Day

(57) **ABSTRACT**

Methods of treating, preventing and/or managing cancer as well as and diseases and disorders associated with, or characterized by, undesired angiogenesis are disclosed. Specific methods encompass the administration of an immunomodulatory compound alone or in combination with a second active ingredient. The invention further relates to methods of reducing or avoiding adverse side effects associated with chemotherapy, radiation therapy, hormonal therapy, biological therapy or immunotherapy which comprise the administration of an immunomodulatory compound. Pharmaceutical compositions, single unit dosage forms, and kits suitable for use in methods of the invention are also disclosed.

15 Claims, 1 Drawing Sheet

US 7,968,569 B2

Page 2

U.S. PATENT DOCUMENTS

| | | | | |
|--------------|-----|---------|-------------------|---------|
| 2001/0018445 | A1 | 8/2001 | Huang et al. | |
| 2001/0022973 | A1* | 9/2001 | Ortyl et al. | 424/452 |
| 2001/0056114 | A1 | 12/2001 | D'Amato | |
| 2002/0035090 | A1 | 3/2002 | Zeldis et al. | |
| 2002/0045643 | A1 | 4/2002 | Muller et al. | |
| 2002/0052398 | A1 | 5/2002 | D'Amato | |
| 2002/0054899 | A1 | 5/2002 | Zeldis | |
| 2002/0061923 | A1 | 5/2002 | D'Amato | |
| 2002/0128228 | A1 | 9/2002 | Hwu | |
| 2002/0161023 | A1 | 10/2002 | D'Amato | |
| 2002/0173658 | A1 | 11/2002 | Muller et al. | |
| 2002/0183360 | A1 | 12/2002 | Muller et al. | |
| 2003/0013739 | A1 | 1/2003 | Masferrer et al. | |
| 2003/0028028 | A1 | 2/2003 | Man et al. | |
| 2003/0045552 | A1 | 3/2003 | Robarge et al. | |
| 2003/0069428 | A1 | 4/2003 | Muller et al. | |
| 2003/0096841 | A1 | 5/2003 | Robarge et al. | |
| 2003/0139451 | A1 | 7/2003 | Shah et al. | |
| 2003/0144325 | A1 | 7/2003 | Muller et al. | |
| 2003/0181428 | A1 | 9/2003 | Green et al. | |
| 2003/0187024 | A1 | 10/2003 | D'Amato | |
| 2003/0191098 | A1 | 10/2003 | D'Amato | |
| 2003/0235909 | A1 | 12/2003 | Hariri et al. | |
| 2004/0029832 | A1 | 2/2004 | Zeldis | |
| 2004/0067953 | A1* | 4/2004 | Stein et al. | 514/251 |
| 2004/0077685 | A1 | 4/2004 | Figg et al. | |
| 2004/0077686 | A1 | 4/2004 | Dannenberg et al. | |
| 2004/0087546 | A1 | 5/2004 | Zeldis | |
| 2004/0091455 | A1 | 5/2004 | Zeldis | |
| 2004/0122052 | A1 | 6/2004 | Muller et al. | |
| 2004/0147558 | A1* | 7/2004 | Treston et al. | 514/323 |
| 2004/0266809 | A1 | 12/2004 | Emanuel et al. | |
| 2005/0049265 | A1* | 3/2005 | Adams | 514/267 |

FOREIGN PATENT DOCUMENTS

| | | |
|----|--------------|---------|
| WO | WO 94/20085 | 9/1994 |
| WO | WO 98/03502 | 1/1998 |
| WO | WO 98/54170 | 12/1998 |
| WO | WO 01/70275 | 9/2001 |
| WO | WO 01/87307 | 11/2001 |
| WO | WO 02/015926 | 2/2002 |
| WO | WO 02/059106 | 8/2002 |
| WO | WO 02/064083 | 8/2002 |
| WO | WO03086373 | 10/2003 |

OTHER PUBLICATIONS

Shah et al., 1999, "Synthesis and enantiomeric separation of 2-phthalimidino-glutaric acid analogues: potent inhibitors of tumor metastasis," *J. Med. Chem.* 42:3014-3017.

Shibata et al., 1995, "N-alkylphthalimides: structural requirement of thalidomidal action on 12-O-tetradecanoylphorbol-13-acetate-induced tumor necrosis factor production by human leukemia HL-60 cells," *Chem. Pharm. Bull.* 43(1):177-179.

Shimazawa et al., 1999, "Antiangiogenic activity of tumor necrosis factor-alpha production regulators derived from thalidomide," *Biol. Pharm. Bull.* 22(2):224-226.

Rubin et al., "Principles of Cancer Treatment-I", 12 ONCO IV I, May 2003.

Wilen et al., 1977, *Tetrahedron* 33:2725.

Wilen, 1972, *Tables of Resolving Agents and Optical Resolutions*, E.L. Eliel, ed., Univ. of Notre Dame Press, Notre Dame, IN pp. 268.

Wolff ed., 1995, *Burger's Medicinal Chemistry and Drug Discovery*, 5th ed., pp. 172-178, 949-982.

Bach, 1963, "Thalidomide in Cancer Chemotherapy," *The Lancet*, No. 1271, p. 71.

Bach, 1963, "Studies on the Possible Anti-Neoplastic Effect of Thalidomide," *Acta Pathologica Et Microbiologica Scandinavica* 59:491-499.

Chaundhry, 1966, *Cancer Research*, "Effect of Prednisolone and Thalidomide on Induced Submandibular Gland Tumors in Hamster," 26(part 1):1884-86.

DiPaolo, 1963, "Effect of Thalidomide on a Variety of Transplantable Tumors," *Cancer Chemotherapy Reports* No. 29, p. 99-102.

DiPaolo, 1963, "In vitro Test Systems for Cancer Chemotherapy, II. Correlation of in vitro Inhibition of Dehydrogenase and Growth with

in vivo Inhibition of Ehrlich Ascites Tumor," *Proceedings of the Society for Experimental Biology & Medicine*, 114:384-387.

DiPaolo, 1964, "Thalidomide: Effects on Ehrlich Ascites Tumor Cells in vitro" *Science* 144:1583.

Mauad, 1963, "Clinical Improvements Obtained in Advanced Cancer Patients with Treatment with Thalidomide Associated with Hormones," *Anais Paulistas de Medicina e Cirurgia* 86:13-40.

Roe and Mitchley, 1963, "Thalidomide and Neoplasia" *Nature* 200:1016-1017.

Alexanian et al., 2004, "VTD (Velcade, thalidomide, dexamethasone) as primary therapy for newly-diagnosed multiple myeloma," *Am. Soc. Hematol. 46th Ann. Meeting Dec. 4-7, 2004*, San Diego, CA Abstract #210.

Anderson, 2000, "Thalidomide: Therapeutic potential in hematologic malignancies," *Seminars in Hematology* 37(1 Supp 3): 1-4.

Attal et al., 2004, "Maintenance treatment with thalidomide after autologous transplantation for myeloma: First analysis of a prospective randomized study of the Intergroupe Francophone du Myelome (IFM99 02)," *Am. Soc. Hematol. 46th Ann. Meeting Dec. 4-7, 2004*, San Diego, CA Abstract #535.

Bemardeschi et al., 2003, *J. Exp. Clin. Cancer Res.* 22(4):129-133.

Corral et al., 1999, "Differential cytokine modulation and T cell activation by two distinct classes of thalidomide analogues that are potent inhibitors of TNF-alpha," *J. Immunol.* 163(1):380-386.

Davies et al., 2001, "Thalidomide and immunomodulatory derivatives augment natural killer cell cytotoxicity in multiple myeloma," *Blood* 98(1):210-216.

Dimopoulos et al., 2004, "Primary treatment with pulsed melphalan, dexamethasone, thalidomide (MDT) for symptomatic patients with multiple myeloma ≥ 75 years of age," *Am. Soc. Hematol. 46th Ann. Meeting Dec. 4-7, 2004*, San Diego, CA Abstract #1482.

Eisen et al., 2000, "Continuous low dose Thalidomide: a phase II study in advanced melanoma, renal cell, ovarian and breast cancer," *Br. J. Cancer* 82(4):812-817.

Fakhouri et al., 2004, "Thalidomide in patients with multiple myeloma and renal failure," *Br. J. Haematol.* 125:90-102.

Fenk et al., 2005, "Single-agent thalidomide for treatment of first relapse following high-dose chemotherapy in patients with multiple myeloma," *Leukemia* 19(1):156-159.

Gupta et al., 2001, "Adherence of multiple myeloma cells to bone marrow stromal cells upregulates vascular endothelial growth factor secretion: therapeutic applications," *Leukemia* 15(12):1950-1961.

Haslett et al., 2003, "Thalidomide and a thalidomide analogue drug costimulate virus-specific CD8+ T cells in vitro," *J. Infect. Dis.* 187(6):946-955.

Hideshima et al., 2000, "Thalidomide and its analogs overcome drug resistance of human multiple myeloma cells to conventional therapy," *Blood* 96(9):2943-2950.

Offidani et al., 2003, "Thalidomide plus oral melphalan for advanced multiple myeloma: a phase II study," *Haematologica*, Dec. 2003;88(12):1432-1433.

Palumbo et al., 2004, "A prospective randomized trial of oral melphalan prednisone, thalidomide (MPT) vs. oral melphalan, prednisone (MP): An interim analysis," *Am. Soc. Hematol. 46th Ann. Meeting Dec. 4-7, 2004*, San Diego, CA Abstract #207.

Raje et al., 1999, "Thalidomide—a revival story," *N. Engl. J. Med.* 341(21):1606-1609.

Rajkumar et al., 2004, "Thalidomide plus dexamethasone versus dexamethasone alone in newly diagnosed multiple myeloma (E1A00): Results of a phase III trial coordinated by the Eastern Cooperative Oncology Group," *Am. Soc. Hematol. 46th Ann. Meeting Dec. 4-7, 2004*, San Diego, CA Abstract #205.

Rajkumar et al., 2000, "Prognostic value of bone marrow angiogenesis in multiple myeloma," *Clin. Cancer Res.* 6(8):3111-3116.

Ribatti et al., 1999, "Bone marrow angiogenesis and mast cell density increase simultaneously with progression of human multiple myeloma," *Br. J. Cancer* 79(3-4):451-455.

Singhal et al., 1999, "Antitumor activity of thalidomide in refractory multiple myeloma," *N. Engl. J. Med.* 341(21):1565-1571.

Steins et al., 2002, "Efficacy and safety of thalidomide in patients with acute myeloid leukemia," *Blood* 99(3):834-839.

US 7,968,569 B2

Page 3

- Vacca et al., 1999, "Bone marrow neovascularization, plasma cell angiogenic potential, and matrix metalloproteinase-2 secretion parallel progression of human multiple myeloma," *Blood* 93(9):3064-3073.
- Wohrer et al., 2004, "Effective treatment of primary plasma cell leukemia with thalidomide and dexamethasone—a case report," *Hematol. J.* 5(4):361-363.
- N. Ake Jonnson, 1972, "Chemical Structure and Teratogenic Properties," *Acta Pharm.*, pp. 521-542.
- Anderson, "Moving disease biology from the laboratory to the clinic," *Seminars in Oncology*, 2002 29:17-20.
- Barlogie et al., "Total Therapy II (TTII) for newly diagnosed multiple myeloma (MM): preliminary data on feasibility and efficacy in the first 231 enrolled patients; comparison with predecessor trial total therapy I (TTI) (N=231)," *Blood, Abstract # 2857*, Dec. 7-11, 2001, *American Society of Hematology*.
- Barlogie et al., "High-dose therapy immunomodulatory drugs in multiple myeloma," *Seminars in Oncology*, 2002, 29 (6):26-33.
- Barlogie et al., "Introduction: Thalidomide and the IMiDs in multiple myeloma," *Seminars in Hematology*, 2003, 40 (4):1-2.
- Barlogie, "Thalidomide and CC-5013 in Multiple Myeloma: The University of Arkansas experience," *Seminars in Hematology*, 2003, 40 (4):33-38.
- Bartlett et al., "The evolution of thalidomide and its IMiD derivatives as anticancer agents," *Nature Reviews Cancer*, 2004, 4 (4):1-9.
- Bartlett et al., "Phase I study to determine the safety, tolerability and immunostimulatory activity of thalidomide analogue CC-5013 in patients with metastatic malignant melanoma and other advanced cancers," *British Journal of Cancer*, 2004, 90:955-961.
- Battagay, "Angiogenesis: mechanistic insights, neovascular diseases, and therapeutic prospects," *J. Mol. Med.*, 1995, 73:333-346.
- Baz et al., "Doxil (D), vincristine (V), reduced frequency dexamethasone (d) and revlimid (R) (DVd-R) results in a high response rate in patients with refractory multiple myeloma (RMM)," *Blood, Abstract # 2559*, *American Society of Hematology*, Dec. 10-13, 2005.
- Brennen et al., "Thalidomide and analogues: current proposed mechanisms and therapeutic usage," *Clinical Prostate Cancer*, 2004, 3 (1):54-61.
- Celgene Corporation, "Celgene advances immunomodulatory drug (IMiD™) clinical program," Press Release, Feb. 2000.
- Celgene Corporation, "Initial Phase I solid tumor data on Celgene's lead ImiD™, Revimid™," Press Release, Jun. 2001.
- Celgene Corporation, "Celgene Corporation receives orphan drug designation for Revimid™ for multiple myeloma," Press Release, Oct. 2001.
- Celgene Corporation, "Celgene Corporation announces third quarter results. Thalomid® (thalidomide) sales increase 24%. Prescriptions up 50%. Enhanced S.T.E.P.S.® launched. Pilot d-MPH data presented," Press Release, Oct. 2001.
- Celgene Corporation, "Celgene expands clinical development program for Revimid™—Five additional trials of Revimid initiated in hematological and solid tumor cancers," Press Release, Jun. 2002.
- Celgene Corporation, "Celgene Corporation announces third quarter results. THALOMID® (thalidomide) revenue increases 41% to \$30.5 million. Pivotal programs for THALOMID and REVIMID™ finalized. Peer-reviewed publications of THALOMID and REVIMID data. First JNK inhibitor advanced to Phase I clinical trial," Press Release, Oct. 2002.
- Celgene Corporation, "Blood reports Revimid™ has anti-tumor activity in patients with relapsed and refractory multiple myeloma," Press Release, Nov. 1, 2002.
- Celgene Corporation, "Celgene provides update on clinical pipeline. Celgene Announces first target indication for ACTIMID™, CC-8490. SelCID™ program to advance based on results from Phase I/II trial of CC-1088. First JNK inhibitor successfully completes phase I trial," Press Release, Jan. 2003.
- Celgene Corporation, "Celgene Corporation announces fourth quarter and full year results for 2002," Press Release, Jan. 2003.
- Celgene Corporation, "Celgene receives fast track status from FDA for Revimid™ in multiple myeloma," Press Release, Feb. 2003.
- Celgene Corporation, "Celgene receives fast track status from FDA for Revimid™ in myelodysplastic syndromes," Press Release, Apr. 2003.
- Celgene Corporation, "New Revimid™ clinical data shows potential as novel approach to treating myelodysplastic syndromes (MDS)," Press Release, May 2003.
- Celgene Corporation, "Celgene corporation reports strong operating performance in second quarter as total sales increase 100 percent and profits rise," Press Release, Jul. 2003.
- Celgene Corporation, "Celgene corporation reports record operating performance in third quarter as total revenue increases 117% and profits rise," Press Release, Oct. 2003.
- Celgene Corporation, "Celgene corporation advances ACTIMID™ (CC-4047) into phase II trial for prostate cancer," Press Release, Oct. 2003.
- Celgene Corporation, "Additional clinical data presented on Revimid™ in myelodysplastic syndromes at the American Society of Hematology 45th annual meeting," Press Release, Dec. 2003.
- Celgene Corporation, "Celgene corporation reviews 2003 achievements and announces 2004 financial outlook," Press Release, Jan. 2004.
- Celgene Corporation, "Revlimid™ receives orphan drug designation from the European commission for multiple myeloma," Press Release, Feb. 2004.
- Celgene Corporation, "Revlimid™ receives orphan drug designation from the European commission for myelodysplastic syndromes," Press Release, Mar. 2004.
- Celgene Corporation, "Celgene corporation reports record operating performance in first quarter with strong revenue growth and profits," Press Release, Apr. 2004.
- Celgene Corporation, "Celgene announces plans to stop phase III trials in melanoma due to lack of efficacy," Press Release, Apr. 2004.
- Dalgleish, et al., "New thalidomide analogues; anti-cancer, anti-angiogenic and immunostimulatory," *British Journal of Cancer*, 2001, 85 (1)25.
- Dalgleish et al., "Thalidomide analogues CC-5013 and CC-4047 induce T cell activation and IL-12 production in patients with both solid tumours and relapsed and refractory multiple myeloma," *British Journal of Cancer*, 2003, 88(Suppl 1), S25-S54.
- Database Pharmaml XP002369094 retrieved from STN. Database accession No. 1659300, & Marketletter, Oct. 9, 2001.
- Database NLDB XP002369095 retrieved from STN. Database accession No. 2002:35280, & Marketletter, Jun. 18, 2001.
- Davies et al., "Thalidomide (Thal) and immunomodulatory derivatives (IMiDs) augment natural killer (NK) cell cytotoxicity in multiple myeloma (MM)," Abstract # 3617, *American Society of Hematology*, Dec. 1-5, 2000.
- Davies et al., "Thalidomide (Thal) and immunomodulatory derivatives (IMiDs) augment natural killer (NK) cell cytotoxicity in multiple myeloma ~MM," Abstract # P222, *VIIIth International Myeloma Workshop*, May 4-8, 2001.
- Dibbs et al., "Thalidomide and thalidomide analogs suppress TNF α secretion by myocytes," Abstract # 1284, *Circulation*, 1998.
- Dimopoulos et al., "Results of thalidomide and IMiDs in multiple myeloma," Abstract # P12.1.4, *International Multiple Myeloma Workshop*, May 23-27, 2003.
- Dimopoulos et al., "Treatment of plasma cell dyscrasias with thalidomide and its derivatives," *Journal of Clinical Oncology*, Dec. 1, 2003, 21 (23)4444-4454.
- Dimopoulos et al., "Study of lenalidomide plus dexamethasone versus dexamethasone alone in relapsed or refractory multiple myeloma (MM): Results of a phase 3 Study (MM-010)," Abstract # 6, *American Society of Hematology*, Dec. 10-13, 2005.
- Dredge et al., A costimulatory thalidomide analog enhances the partial anti-tumor immunity of an autologous vaccination in a model of colorectal cancer, Abstract # 491, *American Association for Cancer Research*, Apr. 6-10, 2002.
- Dredge et al., "Adjuvants and the promotion of Th1-type cytokines in tumour immunotherapy," *Cancer Immunol. Immunother.*, 2002, 51:521-531.
- Dredge et al., "Immunological effects of thalidomide and its chemical and functional analogs," *Critical Reviews in Immunology*, 2002, 21 (5&6):425-437.

US 7,968,569 B2

Page 4

- Dredge et al., "Protective antitumor immunity induced by a costimulatory thalidomide analog in conjunction with whole tumor cell vaccination is mediated by increased Th1 -type immunity¹," *The Journal of Immunology*, 2002, 168:4914-4919.
- Dredge et al., "Recent developments in antiangiogenic therapy," *Expert Opin. Biol. Ther.*, 2002, 2 (8):953-966.
- Dredge et al., "Angiogenesis inhibitors in cancer therapy," *Current Opinion in Investigational Drugs*, 2003, 4 (6):667-674.
- Dredge et al., "Thalidomide analogs as emerging anti-cancer drugs," *Anti-Cancer Drugs*, 2003, 14:331-335.
- Fickentscher et al., "Stereochemical properties and teratogenic activity of some tetrahydrophthalimides," *Molecular Pharmacology*, 1976, 13:133-141.
- Figg et al., "Inhibition of angiogenesis: treatment options for patients with metastatic prostate cancer," *Investigational New Drugs*, 2002, 20(2):183-194.
- Galustian et al., "Thalidomide-derived immunomodulatory drugs as therapeutic agents," *Expert Opin. Biol. Ther.*, 2004, 4 (12):1-8.
- Glaspy et al., "The potential role of thalidomide and thalidomide analogs in melanoma," *Clinical Advances in Hematology & Oncology*, 2004, 1-7.
- Gupta et al., "Adherence of multiple myeloma cells to bone marrow stromal cells upregulates vascular endothelial growth factor secretion: therapeutic applications," *Leukemia*, 2001, 15:1950-1961.
- Hayashi et al., "Mechanisms whereby immunomodulatory analogs of thalidomide augment autologous NK cell anti-myeloma immunity," *Blood Abstract #3219*, Dec. 6-10, 2002, *American Society of Hematology*.
- He, W., et al., 1993, Abstract of papers, 206th American Chemical Society, Chicago, IL; Med. Chem., paper 216.
- Helm et al., "Comparative teratological investigation of compounds of structurally and pharmacologically related to thalidomide," *Arzneimittel Forschung/Drug Research*, 1981, 31 (1)941-949.
- Hernandez-Illizaliturri et al., "Addition of immunomodulatory drugs CC5013 or CC4047 to rituximab enhances anti-tumor activity in a severe combined immunodeficiency (SCID) mouse lymphoma model," Abstract # 235, *American Society of Hematology*, Dec. 6-9, 2003.
- Hideshima et al., "Thalidomide and its analogs overcome drug resistance of human multiple myeloma cells to conventional therapy," *Blood*, 2000, 96:2943-2950, *American Society of Hematology*.
- Hideshima et al., "Thalidomide (Thal) and its analogs overcome drug resistance of human multiple myeloma (MM) cells to conventional therapy," Abstract 1313, *American Society of Hematology*, Dec. 1-5, 2000.
- Hunt et al., "Markers of endothelial and haemostatic activation in the use of CC-4047, a structural analogue of thalidomide, in relapsed myeloma," *Blood Abstract # 3216*, Dec. 6-10, 2002, *American Society of Hematology*.
- Hussein et al., "Doxil (D), vincristine (V), reduced frequency dexamethasone (d) and Revlimid (DVd-R) a phase I/II trial in advanced relapsed/refractory multiple myeloma (Rmm) patients," *Blood, Abstract #208*, *American Society of Hematology*, Dec. 4-7, 2004.
- Hwu et al., "Thalidomide and its analogues in the treatment of metastatic melanoma," *Chemotherapy Foundation Symposium*, Abstract #44, 2002.
- Kyle, "Current therapy of multiple myeloma," *Internal Medicine*, 2002, 41 (3)175-180.
- Kyle et al., "Multiple myeloma," *New England Journal of Medicine*, 2004, 351:1860-1873.
- LeBlanc et al., "Immunomodulatory drug costimulates T cells via the B7-CD28 pathway," *Blood*, 2004, 103:1787-1790, *American Society of Hematology*.
- Lentzsch et al., "In vivo activity of thalidomide and immunomodulatory drugs against multiple myeloma," *VIIIth International Myeloma Workshop. Abstract #P225*, May 4-8, 2001.
- Lentzsch et al., "Immunomodulatory derivative of thalidomide (IMiD CC-4047) determine the lineage commitment of hematopoietic progenitors by down regulation of GATA-1 and modulation of cytokine secretion," Abstract # 3073, *American Society of Hematology*, Dec. 6-9, 2003.
- Lentzsch et al., "Immunomodulatory derivative of thalidomide (IMiD CC-4047) down regulates CAAT/enhancer-binding protein β (C/EBP β) in multiple myeloma (MM)," Abstract # 3456, *American Society of Hematology*, Dec. 6-9, 2003.
- Luzzio et al., "Thalidomide analogues: derivatives of an orphan drug with diverse biological activity," *Expert Opin. Ther. Patents*, 2004, 14 (2):215-229.
- Man et al., " α - Fluoro-substituted thalidomide analogues," *Bioorganic & Medicinal Chemistry Letters* 13, 2003, 3415-3417.
- Marriott et al., "Immunotherapeutic and antitumor potential of thalidomide analogues," *Expert Opin. Biol. Ther.*, 2001, 1 (4):1-8.
- Marriott et al., "New thalidomide analogues; anti-cancer, anti-angiogenic and immunostimulatory," *British Journal of Cancer*, 85:25, Jul. 6, 2001.
- Marriott et al., "Thalidomide and its analogues have distinct and opposing effects on TNF- α and TNFR2 during co-stimulation of both CD4⁺ and CD8⁺T cells," *Clin. Exp. Immunol.*, 2002, 130:75-84.
- Marriott et al., "A novel subclass of thalidomide analogue with anti-solid tumor activity in which caspase-dependent apoptosis is associated with altered expression of bcl-2 family proteins¹," *Cancer Research*, 2003, 63:593-599.
- Marriott et al., "Thalidomide derived immunomodulatory drugs (IMiDs) as potential therapeutic agents," *Current Drug Targets—Immune. Endocrine & Metabolic Disorders*, 2003, 3:181-186.
- Masellis et al., "Changes in gene expression in bone marrow mesenchymal progenitor cells as a consequence of IMiD therapy in multiple myeloma patients," *Blood, Abstract #1548*, Dec. 7-11, 2001, *American Society of Hematology*.
- McCarty, "Thalidomide may impede cell migration in primates by down-regulating integrin β -chains: potential therapeutic utility in solid malignancies, proliferative retinopathy, inflammatory disorders, neointimal hyperplasia, and osteoporosis," *Medical Hypotheses*, 1997, 49:123-131.
- Mitsiades et al., "Apoptotic signaling induced by immunomodulatory thalidomide analogs (Imids) in human multiple myeloma cells: therapeutic implications," Abstract # 3224, Dec. 7-11, 2001, *American Society of Hematology*.
- Mitsiades et al., "Apoptotic signaling induced by immunomodulatory thalidomide analogs in human multiple myeloma cells: therapeutic implications," *Blood*, 2002, 99:4525-4530, *American Society of Hematology*.
- Mitsiades et al., "CC-5013 Celgene," *Current Opinion in Investigational Drugs*, 2004, 5 (6):635-647.
- Moutouh et al., "Novel immunomodulatory drugs (IMiD®): A potential, new therapy for β - hemoglobinopathies," Abstract # 3740, *American Society of Hematology*, Dec. 4-7, 2004.
- Patten et al., "The early use of the serum free light chain assay in patients with relapsed refractory myeloma receiving treatment with a thalidomide analogue (CC-4047)," Abstract # 1640, *American Society of Hematology*, Dec. 6-9, 2003.
- Payvandi et al., "Effects of a thalidomide analog on binding activity of transcription factors and cell cycle progression of multiple myeloma cell lines," *Blood Abstract #2487*, Dec. 1-5, 200, *American Society of Hematology*.
- Payvandi et al., "The thalidomide analogs IMiDs enhance expression of CD69 stimulatory receptor on natural killer cells," Abstract # 1793, *American Association for Cancer Research*, Mar. 24-28, 2001.
- Payvandi et al., "Thalidomide analogs IMiDs inhibit expression of cyclooxygenase-2 in multiple myeloma cell line and LPS stimulated PBMCs," *Blood, Abstract # 2689*, Dec. 7-11, 2001, *American Society of Hematology*.
- Payvandi et al., "Thalidomide and IMiDs inhibit microvessel formation from human arterial rings in the absence of human liver microsomes," *Blood, Abstract # 5046*, Dec. 6-10, 2002, *American Society of Hematology*.
- Payvandi et al., "CC-5013 inhibits the expression of adhesion molecules ICAM-1 and CD44 and prevents metastasis of B16 F10 mouse melanoma cells in an animal model," *American Society of Clinical Oncology, Abstract # 992*, 2003.
- Payvandi et al., "Immunomodulatory drugs inhibit expression of cyclooxygenase-2 from TNF- α , IL-1 β , and LPS-stimulated stimulated human PMBC in a partially IL-10-dependent manner," *Cellular Immunology*, 2004, 81-88.

US 7,968,569 B2

Page 5

- Raje et al., "Combination of the mTOR inhibitor rapamycin and CC-5013 has synergistic activity in multiple myeloma," *Blood*, Dec. 15, 2004, 104 (13):4188-4193.
- Rajkumar et al., "Combination therapy with lenalidomide plus dexamethasone (Rev/Dex) for newly diagnosed myeloma," *Blood*, Dec. 15, 2005, 106 (13):4050-4053.
- Richardson et al., "A Phase 1 study of oral CC5013, an immunomodulatory thalidomide (Thal) derivative, in patients with relapsed and refractory multiple myeloma (MM)," *Blood Abstract #3225*, Dec. 7-11, 2001, *American Society of Hematology*.
- Richardson et al., "Immunomodulatory drug CC-5013 overcomes drug resistance and is well tolerated in patients with relapsed multiple myeloma," *Blood*, 2002 100:3063-3067, *American Society of Hematology*.
- Richardson et al., "A multi-center, randomized, phase 2 study to evaluate the efficacy and safety of 2 CDC-5013 dose regimens when used alone or in combination with dexamethasone (Dex) for the treatment of relapsed or refractory multiple myeloma (MM)," *Blood Abstract # 825*, *American Society of Hematology*, Dec. 6-9, 2003.
- Richardson et al., "Immunomodulatory analogs of thalidomide: an emerging new therapy in myeloma," *Journal of Clinical Oncology*, 2004, 22(16):3212-3214.
- Richardson et al., "A multicenter, single-arm, open-label study to evaluate the efficacy and safety of single-agent lenalidomide in patients with relapsed and refractory multiple myeloma; preliminary results," *10th International Myeloma Workshop*, Apr. 10-14, 2005.
- Richardson et al., "Novel biological therapies for the treatment of multiple myeloma," *Best Practice & Research Clinical Haematology*, 2005, 18 (4):619-634.
- Richardson et al., "A phase 1 trial of lenalidomide (REVLIMID®) with bortezomib (VELCADE®) in relapsed and refractory multiple myeloma," *Blood Abstract # 365*, *American Society of Hematology*, Dec. 10-13, 2005.
- Rubin et al., "Principles of cancer treatment-1," 2003, 12 ONCO IV 1.
- Schafer et al., "Enhancement of cytokine production and AP-1 transcriptional activity in T cells by thalidomide-related immunomodulatory drugs," *Journal of Pharmacology and Experimental Therapeutics*, 2003, 305(3):1222-1232.
- Schey et al., "A phase I study of an immunomodulatory thalidomide analog, CC-4047, in relapsed or refractory multiple myeloma," *Journal of Clinical Oncology*, 2004, 22 (16):1-8.
- Schey et al., "A phase I study of an immunomodulatory thalidomide analogue (CC4047) in relapse/refractory multiple myeloma," *International Society for Experimental Hematology Abstract #248*, 2002.
- Shaughnessy et al., "Global gene expression analysis shows loss of C-MYC and IL-6 receptor gene mRNA after exposure of myeloma to thalidomide and ImiD," Abstract # 2485, *The American Society of Hematology*, Dec. 1-5, 2000.
- Shire et al., "TNF- α inhibitors and rheumatoid arthritis," *Exp. Opin. Ther. Patents*, 1998, 8 (5):531-544.
- Sorbera et al., "CC-5013. Treatment of multiple myeloma. Treatment of Melanoma. Treatment of myelodysplastic syndrome. Angiogenesis inhibitor. TNF- α production inhibitor," *Drugs of the Future*, 2003, 28(5):425-431.
- Streetly et al., "Thalidomide analogue CC-4047 is effective in the treatment of patients with relapsed and refractory multiple myeloma (MM) and induces T-cell activation and IL-12 production," Abstract # 367, *International Multiple Myeloma Workshop*, May 23-27, 2003.
- Streetly et al., "Changes in neutrophil phenotype following the administration of CC-4047 (Actimid) to patients with multiple myeloma," Abstract # 2543, *American Society of Hematology*, Dec. 6-9, 2003.
- Streetly et al., "An update of the use and outcomes of the new immunomodulatory agent CC-4047 (Actimid) in patients with relapsed/refractory myeloma," Abstract #829, *American Society of Hematology*, Dec. 6-9, 2003.
- Teo et al., "A phase I, single-blind, placebo-controlled, ascending single oral dose, safety, tolerability and pharmacokinetic study of CDC-501, a novel immunomodulatory—oncologic agent, in healthy male subjects with a comparison of fed and fasted," *Clinical Pharmacology and Therapeutics*, 2002, 71 (2):93.
- Teo et al., "Chiral inversion of the second generation ImiD™ CC-4047 (ACTIMID™) in human plasma and phosphate-buffered saline," *Chirality*, 2003, 15:348-351.
- Thertulien et al., "Hybrid MEL/DT PACE autotransplant regimen for Multiple Myeloma (MM)—safety and efficacy data in pilot study of 15 patients," *Blood Abstract # 2869*, *American Society of Hematology*, Dec. 7-11, 2001.
- Tohny et al., "A phase I study of oral CC-5013 (lenalidomide, Revlimid™), a thalidomide derivative, in patients with refractory metastatic cancer," *Clinical Prostate Cancer*, 2004, 2:241-243.
- Tricot et al., "Angiochemotherapy (ACT) for multiple myeloma (MM) with DT-PACE results in a high response rate, but in contrast to tandem transplants with melphalan does not affect durable disease control," *Blood Abstract # 3531*, *American Society of Hematology*, Dec. 7-11, 2001.
- Tsenova et al., "Use of ImiD3, a thalidomide analog, as an adjunct to therapy for experimental tuberculous meningitis," *Antimicrobial Agents and Chemotherapy*, 2002, 46 (6):1887-1895.
- Weber, "Lenalidomide (CC-5013, Revlimid™) and other ImiDs," Abstract # PL5.02, *International Multiple Myeloma Workshop*, Apr. 10-14, 2005.
- Weber et al., "A multicenter, randomized, parallel-group, double-blind, placebo-controlled study of lenalidomide plus dexamethasone versus dexamethasone alone in previously treated subjects with multiple myeloma," Abstract # PO.738, *International Multiple Myeloma Workshop*, Apr. 10-14, 2005.
- Ye et al., "Novel ImiD drugs enhance expansion and regulate differentiation of human cord blood CD34+ cells with cytokines," *Blood Abstract #4099*, *American Society of Hematology*, Dec. 6-10, 2002.
- Zangari et al., "Risk factors for deep vein thrombosis (DVT) in a large group of myeloma patients (Pts) treated with thalidomide (Thal): The Arkansas Experience," *Blood Abstract # 681*, *American Society of Hematology*, Dec. 7-11, 2001.
- Zangari et al., "Revimid 25 mg (REV 25) \times 20 versus 50 mg (REV 50) \times 10 q 28 days with bridging of 5 mg \times 10 versus 10 mg \times 5 as post-transplant salvage therapy for multiple myeloma (MM)," *Blood Abstract # 1642*, *American Society of Hematology*, Dec. 6-9, 2003.
- Zeldis et al., "Potential new therapeutics for Waldenstrom's macroglobulinemia," *Seminars in Oncology*, 2003, 30 (2):275-281.
- Zhang et al., "CC-5079, a novel microtubule and TNF- α inhibitor with anti-angiogenic and antimetastasis activity," Abstract # B012, *International Conference on Molecular Targets and Cancer Therapeutics*, Nov. 17-21, 2003.
- Anderson, "The Role of Immunomodulatory Drugs in Multiple Myeloma," *Seminars in Hematology*, vol. 40, No. 4, Suppl 4, 2003: pp. 23-32.
- Weber, "Thalidomide and Its Derivatives: New Promise for Multiple Myeloma," *Cancer Control*, vol. 10, No. 5, 375-383, 2003.
- U.S. Appl. No. 60/499,723, filed Sep. 4, 2003, Markian.
- U.S. Appl. No. 60/372,348, filed Apr. 12, 2002, Hariri et al.
- U.S. Appl. No. 10/732,867, filed Dec. 9, 2003, D'Amato et al.
- U.S. Appl. No. 09/545,654, filed Apr. 10, 2000, D'Amato.
- U.S. Appl. No. 09/287,377, filed Apr. 7, 1999, D'Amato.
- Carstensen, 1995, *Drug Stability: Principles & Practice*, 2nd ed., Marcel Dekker, New York, NY pp. 379-380.
- Corral et al., 1999, "Immunomodulation by thalidomide and thalidomide analogues," *Ann. Rheum. Dis.* 58(Suppl 1):1107-1113.
- Craig et al., 1967, "Potential anticancer agents. III. 2-phthalimidoaldehydes and derivatives," *Potential Anticancer Agents III* 10:1071-1073.
- D'Amato et al., 2001, "Mechanism of action of thalidomide and 3-aminothalidomide in multiple myeloma," *Semin. Oncol.* 28:597-601.
- D'Amato et al., 1994, "Thalidomide is an Inhibitor of Angiogenesis", *Proc. Natl. Acad. Sci.* 91:4082-4085.
- De et al., 1976, "Hansch analysis for some antineoplastic glutarimides," *J. Indian Chem. Soc. I.III:* 825-826.
- De et al., 1976, "Possible antineoplastic agents: III. Synthesis of 6-alkyl-2-[4'-methoxyphthalimido] and 6-alkyl-3-[3'-4'-dimethoxyphenyl] glutarimides," *J. Indian Chem. Soc. I.III:*1122-1125.

US 7,968,569 B2

Page 6

- Dredge et al., 2002, "Novel thalidomide analogues display anti-angiogenic activity independently of immunomodulatory effects," *Br. J. Cancer* 87(10):1166-1172.
- Folkman et al., 1983, "Angiogenesis inhibition and tumor regression caused by heparin or a heparin fragment in the presence of cortisone," *Science* 221(4612):719-725.
- Gershbein, 1991, "The thalidomide analog, EM 12, enhances 1,2-dimethylhydrazine-induction of rat colon adenocarcinomas," *Cancer Letters* 60: 129-133.
- Grabstald et al., 1965, "Clinical experiences with thalidomide in patients with cancer," *Clinical Pharmacology and Therapeutics* 6:298-302.
- Lentzsch et al., 2003, "Immunomodulatory analogs of thalidomide inhibit growth of Hs Sultan cells and angiogenesis in vivo," *Leukemia* 17(1):41-44.
- Lentzsch et al., 2002, "S-3-amino-phthalimido-glutarimide inhibits angiogenesis and growth of B-cell neoplasias in mice", *Cancer Research* 62:2300-2305.
- Miyachi et al., 1997, "Novel biological response modifiers: phthalimides with tumor necrosis factor-alpha production-regulating activity," *J. Med. Chem.* 40:2858-2865.
- Muller et al., 1999, "Amino-substituted thalidomide analogs: potent inhibitors of TNF-alpha production," *Bioorg. Med. Chem. Lett.* 9(11):1625-1630.
- Muller et al., 1998, "Thalidomide analogs and PDE4 inhibition," *Bioorg. Med. Chem. Lett.* 8(19):2669-2674.
- Muller et al., 1996, "Structural modifications of thalidomide produce analogs with enhanced tumor necrosis factor inhibitory activity," *J. Med. Chem.* 39(17):3238-3240.
- Olson et al., 1965, "Thalidomide (N-phthaloylglutaramide) in the treatment of advanced cancer," *Clinical Pharmacology and Therapeutics* 6(3):292-297.
- Penichet et al., 2001, "Antibody-cytokine fusion proteins for the therapy of cancer," *J. Immunol. Methods* 248(1-2):91-101.
- Physician's Desk Reference*, 2002, 56th ed., pp. 1755-1760.
- Attal et al., 2004, "Maintenance treatment with thalidomide after autologous transplantation for myeloma: First analysis of a prospective randomized study of the Intergroupe Francophone du Myelome (IFM 99 02)," *Am. Soc. Hematol. 46th Ann. Meeting Dec. 4-7, 2004, San Diego, CA Abstract #535.*
- Bernardeschi et al., 2003, *J. Exp. Clin. Cancer Res.* 22(4):129-133.
- Dimopoulos et al., 2004, "Primary treatment with pulsed melphalan, dexamethasone, thalidomide (MDT) for symptomatic patients with multiple myeloma \geq 75 years of age," *Am. Soc. Hematol. 46th Ann. Meeting Dec. 4-7, 2004, San Diego, CA Abstract #1482.*
- Rajkumar et al., 2004, "Thalidomide plus dexamethasone versus dexamethasone alone in newly diagnosed multiple myeloma (E1A00): Results of a phase III trial coordinated by the Eastern Cooperative Oncology Group," *Am. Soc. Hematol. 46th Ann. Meeting Dec. 4-7, 2004, San Diego, CA Abstract #205.*
- Bach, 1963, "Thalidomide in Cancer Chemotherapy," *The Lancet*, No. 1271, p. 71.
- Bach, 1963, "Studies on the Possible Anti-Neoplastic Effect of Thalidomide," *Acta Pathologica Et Microbiologica Scandinavica* 59:491-499.
- Chandhry, 1966, *Cancer Research*, "Effect of Prednisolone and Thalidomide on Induced Submandibular Gland Tumors in Hamster," 26(part 1)1884-86.
- DiPaolo, 1963, "Effect of Thalidomide on a Variety of Transplantable Tumors," *Cancer Chemotherapy Reports* No. 29, p. 99-102.
- DiPaolo, 1963, "In vitro Test Systems for Cancer Chemotherapy. II. Correlation of in vitro Inhibition of Dehydrogenase and Growth with in vivo Inhibition of Ehrlich Asoites Tumor," *Proceedings of the Society for Experimental Biology & Medicine*, 114:384-387.
- Mauad, 1963, "Clinical Improvements Obtained in Advanced Cancer Patients with Treatment with Thalidomide Associated with Hormones," *Anais Paulistas de Medicina e Cirurgia* 86:13-40.
- Roe and Mitchley, 1963, "Thalidomide and Neoplasia" *Nature* 200:1016-1017.
- Liu et al., "Phase I study of CC-5013 (Revimid), a thalidomide derivative, in patients with refractory metastatic cancer," *American Society of Clinical Oncology, Abstract #927, 2003.*
- Zangari et al., "Results of phase I study of CC-5013 for the treatment of multiple myeloma (MM) patients who relapse after high dose chemotherapy (HDCT)," *American Society of Hematology, Abstract #3226, 2001.*
- Zeldis et al., "Update on the evolution of the IMiD™," *International Society for Biological Therapy of Cancer, Oral Abstract, 2003.*
- Anderson, "Moving disease biology from the laboratory to the clinic," *Seminars in Oncology, 2002 29:17-20.*
- Barlogie et al., "Total Therapy II (TTII) for newly diagnosed multiple myeloma (MM): preliminary data on feasibility and efficacy in the first 231 enrolled patients; comparison with predecessor trial total therapy I (TTI) (N=231)," *Blood, Abstract # 2857, Dec. 7-11, 2001, American Society of Hematology.*
- Barlogie et al., "High-dose therapy immunomodulatory drugs in multiple myeloma," *Seminars in Oncology, 2002, 29 (6):26-33.*
- Barlogie et al., "Introduction: Thalidomide and the IMiDs in multiple myeloma," *Seminars in Hematology, 2003, 40 (4):1-2.*
- Barlogie, "Thalidomide and CC-5013 in Multiple Myeloma: The University of Arkansas experience," *Seminars in Hematology, 2003, 40 (4):33-38.*
- Bartlett et al., "The evolution of thalidomide and its IMiD derivatives as anticancer agents," *Nature Reviews Cancer, 2004, 4 (4): 1 -9.*
- Bartlett et al., "Phase I study to determine the safety, tolerability and immunostimulatory activity of thalidomide analogue CC-5013 in patients with metastatic malignant melanoma and other advanced cancers," *British Journal of Cancer, 2004, 90:955-961.*
- Battegay, "Angiogenesis: mechanistic insights, neovascular diseases, and therapeutic prospects," *J. Mol. Med., 1995, 73:333-346.*
- Baz et al., "Doxil (D), vincristine (V), reduced frequency dexamethasone (d) and revlimid (R) (DvD-R) results in a high response rate in patients with refractory multiple myeloma (RMM)," *Blood, Abstract # 2559, American Society of Hematology, Dec. 10-13, 2005.*
- Brennen et al., "Thalidomide and analogues: current proposed mechanisms and therapeutic usage," *Clinical Prostate Cancer, 2004, 3 (1):54-61.*
- Celgene Corporation, "Celgene advances immunomodulatory drug (IMiD™) clinical program," *Press Release, Feb. 2000.*
- Celgene Corporation, "Additional clinical data presented on Revimid™ in myelodysplastic syndromes at the American Society of Hematology 45th annual meeting," *Press Release, Dec. 2003.*
- Dalgleish, et al., "New thalidomide analogues; anti-cancer, anti-angiogenic and immunostimulatory," *British Journal of Cancer, 2001, 85 (1)25.*
- Dalgleish et al., "Thalidomide analogues CC-5013 and CC-4047 induce T cell activation and IL-12 production in patients with both solid tumours and relapsed and refractory multiple myeloma," *British Journal of Cancer, 2003, 88(Suppl 1), S25-S54.*
- Davies et al., "Thalidomide (Thal) and immunomodulatory derivatives (IMiDs) augment natural killer (NK) cell cytotoxicity in multiple myeloma(MM)," *Abstract # 3617, American Society of Hematology, Dec. 1-5, 2000.*
- Davies et al., "Thalidomide (Thal) and immunomodulatory derivatives (IMiDs) augment natural killer (NK) cell cytotoxicity in multiple myeloma ~MM)," *Abstract # P222, VIIIth International Myeloma Workshop, May 4-8, 2001.*
- Dibbs et al., "Thalidomide and thalidomide analogs suppress TNF γ secretion by myocytes," *Abstract # 1284, Circulation, 1998.*
- Dimopoulos et al., "Results of thalidomide and IMiDs in multiple myeloma," *Abstract # P12.1.4, International Multiple Myeloma Workshop, May 23-27, 2003.*
- Dimopoulos et al., "Treatment of plasma cell dyscrasias with thalidomide and its derivatives," *Journal of Clinical Oncology, Dec. 1, 2003, 21 (23)4444-4454.*
- Dimopoulos et al., "Study of lenalidomide plus dexamethasone versus dexamethasone alone in relapsed or refractory multiple myeloma (MM): Results of a phase 3 Study (MM-010)," *Abstract # 6, American Society of Hematology, Dec. 10-13, 2005.*
- Dredge et al., A costimulatory thalidomide analog enhances the partial anti-tumor immunity of an autologous vaccination in a model of colorectal cancer, *Abstract # 491, American Association for Cancer Research, Apr. 6-10, 2002.*

- Dredge et al., "Adjuvants and the promotion of Th 1-type cytokines in tumour immunotherapy," *Cancer Immunol. Immunother.*, 2002, 51:521-531.
- Dredge et al., "Immunological effects of thalidomide and its chemical and functional analogs," *Critical Reviews in Immunology*, 2002, 22 (5&6):425-437.
- Dredge et al., "Protective antitumor immunity induced by a costimulatory thalidomide analog in conjunction with whole tumor cell vaccination is mediated by increased Th 1-type immunity1," *The Journal of Immunology*, 2002, 168:4914-4919.
- Dredge et al., "Recent developments in antiangiogenic therapy," *Expert Opin. Biol. Ther.*, 2002, 2 (8):953-966.
- Dredge et al., "Angiogenesis inhibitors in cancer therapy," *Current Opinion in Investigational Drugs*, 2003, 4 (6):667-674.
- Dredge et al., "Thalidomide analogs as emerging anti-cancer drugs," *Anti-Cancer Drugs*, 2003, 14:331-335.
- Fickentscher et al., "Stereochemical properties and teratogenic activity of some tetrahydrophthalimides," *Molecular Pharmacology*, 1976, 13:133-141.
- Figg et al., "Inhibition of angiogenesis: treatment options for patients with metastatic prostate cancer," *Investigational New Drugs*, 2002, 20(2):183-194.
- Galustian et al., "Thalidomide-derived immunomodulatory drugs as therapeutic agents," *Expert Opin. Biol. Ther.*, 2004,4(12):1-8.
- Glaspy et al., "The potential role of thalidomide and thalidomide analogs in melanoma," *Clinical Advances in Hematology & Oncology*, 2004, 1-7.
- Gupta et al., "Adherence of multiple myeloma cells to bone marrow stromal cells upregulates vascular endothelial growth factor secretion: therapeutic applications," *Leukemia*, 2001, 15:1950-1961.
- Hayashi et al., "Mechanisms whereby immunomodulatory analogs of thalidomide augment autologous NK cell anti-myeloma immunity," *Blood*, Abstract #3219, Dec. 6-10, 2002, American Society of Hematology.
- Helm et al., "Comparative teratological investigation of compounds of structurally and pharmacologically related to thalidomide," *Arzneimittel Forschung/Drug Research*, 1981, 31 (1)941-949.
- Hernandez-ILlizarurr et al., "Addition of immunomodulatory drugs CC5013 or CC4047 to rituximab enhances anti-tumor activity in a severe combined immunodeficiency (SCID) mouse lymphoma model," Abstract # 235, American Society of Hematology, Dec. 6-9, 2003.
- Hideshima et al., "Thalidomide and its analogs overcome drug resistance of human multiple myeloma cells to conventional therapy," *Blood*, 2000, 96:2943-2950, American Society of Hematology.
- Hideshima et al., "Thalidomide (Thal) and its analogs overcome drug resistance of human multiple myeloma (MM) cells to conventional therapy," Abstract 1313, American Society of Hematology, Dec. 1-5, 2000.
- Hunt et al., "Markers of endothelial and haemostatic activation in the use of CC-4047, a structural analogue of thalidamide, in relapsed myeloma," *Blood*, Abstract # 3216, Dec. 6-10, 2002, American Society of Hematology.
- Hussein et al., "Doxil (D), vincristine (V), reduced frequency dexamethasone (d) and Revlimid (DVd-R) a phase I/II trial in advanced relapsed/refractory multiple myeloma (Rmm) patients," *Blood*, Abstract #208, American Society of Hematology, Dec. 4-7, 2004.
- Hwu et al., "Thalidomide and its analogues in the treatment of metastatic melanoma," *Chemotherapy Foundation Symposium*, Abstract #44, 2002.
- Kyle, "Current therapy of multiple myeloma," *Internal Medicine*, 2002, 41 (3)175-180.
- Kyle et al., "Multiple myeloma," *New England Journal of Medicine*, 2004, 351:1860-1873.
- LeBlanc et al., "Immunomodulatory drug costimulates T cells via the B7-CD28 pathway," *Blood*, 2004, 103:1787-1790, American Society of Hematology.
- Lentzsch et al., "In vivo activity of thalidomide and immunomodulatory drugs against multiple myeloma," VIIIth International Myeloma Workshop, Abstract #P225, May 4-8, 2001.
- Lentzsch et al., "Immunomodulatory derivative of thalidomide (IMiD CC-4047) determine the lineage commitment of hematopoietic progenitors by down regulation of GATA-1 and modulation of cytokine secretion," Abstract # 3073, American Society of Hematology, Dec. 6-9, 2003.
- Lentzsch et al., "Immunomodulatory derivative of thalidomide (IMiD CC-4047) down regulates CAAT/enhancer-binding protein γ (C/EBP γ) in multiple myeloma (MM)," Abstract # 3456, American Society of Hematology, Dec. 6-9, 2003.
- Luzzio et al., "Thalidomide analogues: derivatives of an orphan drug with diverse biological activity," *Expert Opin. Ther. Patents*, 2004, 14 (2):215-229.
- Man et al., " γ -Fluoro-substituted thalidomide analogues," *Bioorganic & Medicinal Chemistry Letters* 13, 2003, 3415-3417.
- Marriott et al., "Immunotherapeutic and antitumor potential of thalidomide analogues," *Expert Opin. Biol. Ther.*, 2001, 1 (4):1-8.
- Marriott et al., "New thalidomide analogues; anti-cancer, anti-angiogenic and immunostimulatory," *British Journal of Cancer*, 85:25, Jul. 6, 2001.
- Marriott et al., "Thalidomide and its analogues have distinct and opposing effects on TNF- α and TNFR2 during co-stimulation of both CD4+ and CD8+ T cells," *Clin. Exp. Immunol.*, 2002, 130:75-84.
- Marriott et al., "A novel subclass of thalidomide analogue with anti-solid tumor activity in which caspase-dependent apoptosis is associated with altered expression of bcl-2 family proteins 1," *Cancer Research*, 2003, 63:593-599.
- Marriott et al., "Thalidomide derived immunomodulatory drugs (IMiDs) as potential therapeutic agents," *Current Drug Targets—Immunologic, Endocrine & Metabolic Disorders*, 2003, 3:181-186.
- Masellis et al., "Changes in gene expression in bone marrow mesenchymal progenitor cells as a consequence of IMiD therapy in multiple myeloma patients," *Blood*, Abstract # 1548, Dec. 7-11, 2001, American Society of Hematology.
- McCarty, "Thalidomide may impede cell migration in primates by down-regulating integrin γ -chains: potential therapeutic utility in solid malignancies, proliferative retinopathy, inflammatory disorders, neointimal hyperplasia, and osteoporosis," *Medical Hypotheses*, 1997, 49:123-131.
- Mitsiades et al., "Apoptotic signaling induced by immunomodulatory thalidomide analogs (Imids) in human multiple myeloma cells: therapeutic implications," Abstract # 3224, Dec. 7-11, 2001, American Society of Hematology.
- Mitsiades et al., "Apoptotic signaling induced by immunomodulatory thalidomide analogs in human multiple myeloma cells: therapeutic implications," *Blood*, 2002, 99:4525-4530, American Society of Hematology.
- Mitsiades et al., "CC-5013 Celgene," *Current Opinion in Investigational Drugs*, 2004, 5 (6):635-647.
- Moutouh et al., "Novel immunomodulatory drugs (IMiDs®): A potential, new therapy for γ -hemoglobinopathies," Abstract # 3740, American Society of Hematology, Dec. 4-7, 2004.
- Patten et al., "The early use of the serum free light chain assay in patients with relapsed refractory myeloma receiving treatment with a thalidomide analogue (CC-4047)," Abstract # 1640, American Society of Hematology, Dec. 6-9, 2003.
- Payvandi et al., "Effects of a thalidomide analog on binding activity of transcription factors and cell cycle progression of multiple myeloma cell lines," *Blood*, Abstract #2487, Dec. 1-5, 2000, American Society of Hematology.
- Payvandi et al., "The thalidomide analogs IMiDs enhance expression of CD69 stimulatory receptor on natural killer cells," Abstract # 1793, American Association for Cancer Research, Mar. 24-28, 2001.
- Payvandi et al., "Thalidomide analogs IMiDs inhibit expression of cyclooxygenase-2 in multiple myeloma cell line and LPS stimulated PBMCs," *Blood*, Abstract # 2689, Dec. 7-11, 2001, American Society of Hematology.
- Payvandi et al., "Thalidomide and IMiDs inhibit microvessel formation from human arterial rings in the absence of human liver microsomes," *Blood*, Abstract # 5046, Dec. 6-10, 2002, American Society of Hematology.
- Payvandi et al., "CC-5013 inhibits the expression of adhesion molecules ICAM-1 and CD44 and prevents metastasis of B16 F10 mouse melanoma cells in an animal model," *American Society of Clinical Oncology*, Abstract # 992, 2003.

US 7,968,569 B2

Page 8

- Payvandi et al., "Immunomodulatory drugs inhibit expression of cyclooxygenase-2 from TNF- α , IL-1 α , and LPS-stimulated human PBMC in a partially IL-10-dependent manner," *Cellular Immunology*, 2004, 81-88.
- Raje et al., "Combination of the mTOR inhibitor rapamycin and CC-5013 has synergistic activity in multiple myeloma," *Blood*, Dec. 15, 2004, 104 (13):4188-4193.
- Rajkumar et al., "Combination therapy with lenalidomide plus dexamethasone (Rev/Dex) for newly diagnosed myeloma," *Blood*, Dec. 15, 2005, 106 (13):4050-4053.
- Richardson et al., "A Phase I study of oral CC5013, an immunomodulatory thalidomide (Thal) derivative, in patients with relapsed and refractory multiple myeloma (MM)," *Blood*, Abstract #3225, Dec. 7-11, 2001, American Society of Hematology.
- Richardson et al., "Immunomodulatory drug CC-5013 overcomes drug resistance and is well tolerated in patients with relapsed multiple myeloma," *Blood*, 2002 100:3063-3067, American Society of Hematology.
- Richardson et al., "A multi-center, randomized, phase 2 study to evaluate the efficacy and safety of 2 CDC-5013 dose regimens when used alone or in combination with dexamethasone (Dex) for the treatment of relapsed or refractory multiple myeloma (MM)," *Blood*, Abstract # 825, American Society of Hematology, Dec. 6-9, 2003.
- Richardson et al., "Immunomodulatory analogs of thalidomide: an emerging new therapy in myeloma," *Journal of Clinical Oncology*, 2004, 22(16) 3212-3214.
- Richardson et al., "A multicenter, single-arm, open-label study to evaluate the efficacy and safety of single-agent lenalidomide in patients with relapsed and refractory multiple myeloma; preliminary results," 10th International Myeloma Workshop, Apr. 10-14, 2005.
- Richardson et al., "Novel biological therapies for the treatment of multiple myeloma," *Best Practice & Research Clinical Haematology*, 2005, 18 (4):619-634.
- Richardson et al., "A phase I trial of lenalidomide (REVLIMID®) with bortezomib (VELCADE®) in relapsed and refractory multiple myeloma," *Blood*, Abstract # 365, American Society of Hematology, Dec. 10-13, 2005.
- Schafer et al., "Enhancement of cytokine production and AP-1 transcriptional activity in T cells by thalidomide-related immunomodulatory drugs," *Journal of Pharmacology and Experimental Therapeutics*, 2003, 305(3):1222-1232.
- Schey et al., "A phase I study of an immunomodulatory thalidomide analog, CC-4047, in relapsed or refractory multiple myeloma," *Journal of Clinical Oncology*, 2004, 22 (16):1-8.
- Schey et al., "A phase I study of an immunomodulatory thalidomide analogue (CC4047) in relapse/refractory multiple myeloma," *International Society for Experimental Hematology*, Abstract #248, 2002.
- Shaughnessy et al., "Global gene expression analysis shows loss of C-MYC and IL-6 receptor gene mRNA after exposure of myeloma to thalidomide and ImiD," Abstract # 2485, The American Society of Hematology, Dec. 1-5, 2000.
- Shire et al., "TNF- β inhibitors and rheumatoid arthritis," *Exp. Opin. Ther. Patents*, 1998, 8 (5):531-544.
- Sorbera et al., "CC-5013. Treatment of multiple myeloma. Treatment of Melanoma. Treatment of myelodysplastic syndrome. Angiogenesis inhibitor. TNF- β production inhibitor," *Drugs of the Future*, 2003, 28(5):425-431.
- Streety et al., "Thalidomide analogue CC-4047 is effective in the treatment of patients with relapsed and refractory multiple myeloma (MM) and induces T-cell activation and IL-12 production," Abstract # 367, International Multiple Myeloma Workshop, May 23-27, 2003.
- Streety et al., "Changes in neutrophil phenotype following the administration of CC-4047 (Actimid) to patients with multiple myeloma," Abstract # 2543, American Society of Hematology, Dec. 6-9, 2003.
- Streety et al., "An update of the use and outcomes of the new immunomodulatory agent CC-4047 (Actimid) in patients with relapsed/refractory myeloma," Abstract #829, American Society of Hematology, Dec. 6-9, 2003.
- Teo et al., "A phase I, single-blind, placebo-controlled, ascending single oral dose, safety, tolerability and pharmacokinetic study of CDC-501, a novel immunomodulatory- oncologic agent, in healthy male subjects with a comparison of fed and fasted," *Clinical Pharmacology and Therapeutics*, 2002, 71 (2):93.
- Teo et al., "Chiral inversion of the second generation IMiD™ CC-4047 (ACTIMID™) in human plasma and phosphate-buffered saline," *Chirality*, 2003, 15:348-351.
- Thertulien et al., "Hybrid MEL/DT PACE autotransplant regimen for Multiple Myeloma (MM)—safety and efficacy data in pilot study of 15 patients," *Blood*, Abstract # 2869, American Society of Hematology, Dec. 7-11, 2001.
- Tohnya et al., "A phase I study of oral CC-5013 (lenalidomide, Revlimid™), a thalidomide derivative, in patients with refractory metastatic cancer," *Clinical Prostate Cancer*, 2004, 2:241-243.
- Tricot et al., "Angiochemotherapy (ACT) for multiple myeloma (MM) with DT-PACE results in a high response rate, but in contrast to tandem transplants with melphalan does not affect durable disease control," *Blood*, Abstract # 3531, American Society of Hematology, Dec. 7-11, 2001.
- Tsenova et al., "Use of IMiD3, a thalidomide analog, as an adjunct to therapy for experimental tuberculous meningitis," *Antimicrobial Agents and Chemotherapy*, 2002, 46 (6):1887-1895.
- Weber, "Lenalidomide (CC-5013, Revlimid™) and other ImiDs," Abstract # PL5.02, International Multiple Myeloma Workshop, Apr. 10-14, 2005.
- Weber et al., "A multicenter, randomized, parallel-group, double-blind, placebo-controlled study of lenalidomide plus dexamethasone versus dexamethasone alone in previously treated subjects with multiple myeloma," Abstract # PO.738, International Multiple Myeloma Workshop, Apr. 10-14, 2005.
- Ye et al., "Novel IMiD drugs enhance expansion and regulate differentiation of human cord blood CD34+ cells with cytokines," *Blood*, Abstract #4099, American Society of Hematology, Dec. 6-10, 2002.
- Zangari et al., "Risk factors for deep vein thrombosis (DVT) in a large group of myeloma patients (Pts) treated with thalidomide (Thal): The Arkansas Experience," *Blood*, Abstract # 681, American Society of Hematology, Dec. 7-11, 2001.
- Zangari et al., "Revimid 25 mg (REV 25) \times 20 versus 50 mg (REV 50) \times 10 q 28 days with bridging of 5 mg \times 10 versus 10 mg \times 5 as post-transplant salvage therapy for multiple myeloma (MM)," *Blood*, Abstract # 1642, American Society of Hematology, Dec. 6-9, 2003.
- Zeldis et al., "Potential new therapeutics for Waldenstrom's macroglobulinemia," *Seminars in Oncology*, 2003, 30 (2):275-281.
- Zhang et al., "CC-5079, a novel microtubule and TNF- α inhibitor with anti-angiogenic and antimetastasis activity," Abstract # B012, International Conference on Molecular Targets and Cancer Therapeutics, Nov. 17-21, 2003.
- Anderson, "The Role of Immunomodulatory Drugs in Multiple Myeloma," *Seminars in Hematology*, vol. 40, No. 4, Suppl 4, 2003: pp. 23-32.
- Weber, "Thalidomide and Its Derivatives: New Promise for Multiple Myeloma," *Cancer Control*, vol. 10, No. 5, 375-383, 2003.
- Patt, Yehuda A.; Hassan, Manal M.; Lozano, Richard D.; Ellis, Lee M.; Peterson, J. Andrew; Waugh, Kimberly A.; Durable Clinical Response of Refractory Hepatocellular Carcinoma to Orally Administered Thalidomide. *American Journal of Clinical Oncology*, 2000.
- Richardson, Paul; Hideshima, Teru; Anderson, Kenneth; Thalidomide: The Revival of a Drug with Therapeutic Promise in the Treatment of Cancer; *Principles & Practice of Oncology*, vol. 15, No. 2, 2001.
- Thomas, Melodie; DOSS, Deborah, Thalidomide Nursing Roundtable Update, Monograph, Sep. 2002.
- Richardson, Paul; Hideshima, Teru; Anderson, Kenneth; Thalidomide: Emerging Role in Cancer Medicine; *Annual Review of Medicine*, 2002.
- Berenson, J.R.; Bergsagel, P. L.; Munshi, N.; Initiation and Maintenance of Multiple Myeloma; *Seminars in Hematology*, vol. 36, No. 1, Supp. 3, Jan. 1999, pp. 9-13.
- Gollob, J.A.; Schinpper, C.P.; Orsini, E.; Murphy, E.; Daley, J.F.; Lazo, S.B.; Frank, D.A.; Characterization of a Novel Subset of CD8 T Cells That Expands in patients Receiving Interleukin-12, 02, *Am. Soc. For Clin. Investigation, Inc.*, vol. 102, No. 3, Aug. 1998, pp. 561-575.
- Cavanagh, L.L.; Barnetson, R.S.; Basten, A.; Halliday, G.M.; Dendritic Epidermal T-Cell Involvement in Induction of CD8+

US 7,968,569 B2

Page 9

- T-Cell-Mediated Immunity Against an Ultraviolet Radiation-Induced Skin Tumor Int. J. Cancer: 70, 98-105, 1997.
- Thomas, D.A., Aguayo, A., Estey, E., Albitar, M., O'Brien, S., Giles, F.J., Beran, M., Cortes, J., Zeldis, J., Keating, M.J., Barlogie, B., Kantarjian, H.M., Thalidomide as anti-angiogenesis therapy (rx) in refractory or relapsed leukemia. Abstract #2269, American Society of Hematology, Dec. 3-7, 1999.
- Barlogie, B., Desikan, R., Munshi, N., Siegel, D., Mehta, J., Singhal, S., Anaissie, E., Single Course D.T. Pace Anti-Angiochemotherapy Effects CR in Plasma Cell Leukemia and Fulminant Multiple Myeloma (MM). Abstract #4180. American Society of Hematology, Dec. 4-9, 1998.
- Hideshima, T., Chauhan, D., Shima, Y., Noopur, R., Davies, F.E., Tai, Y., Treon, S.P., Lin, B.K., Schlossman, R.L., Richardson, P.G., Gupta, D., Muller, G.W., Stirling, D.I., Anderson, K.C., Thalidomide (THAL) and its Analogs Overcome Drug Resistance of Human Multiple Myeloma (MM) Cells to Conventional Therapy. Abstract #1313. American Society of Hematology, Dec. 1-5, 2000.
- Payvandi, F., Wu, L., Gupta, D., Hideshima, T., Haley, M., Muller, G., Chen, R., Anderson, K.C., Stirling, D., Effects of a Thalidomide Analog on Binding Activity of Transcription Factors and Cell Cycle Progression of Multiple Myeloma Cell Lines. Abstract #2487. American Society of Hematology, Dec. 1-5, 2000.
- Davies, F.E., Raje, N., Hideshima, T., Lentzsch, S., Young, G., Tai, Y., Lin, B.K., Podar, K., Chauhan, D., Treon, S.P., Gupta, D., Mitsiades, C., Mitsiades, N., Hayashi, T., Richardson, P.G., Schlossman, R.L., Muller, G.W., Stirling, D. I., Anderson, K.C., Thalidomide (THAL) and Immunomodulatory Derivatives (IMiDS) Augment Natural Killer (NK) Cell Cytotoxicity in Multiple Myeloma (MM). Abstract #3617. American Society of Hematology, Dec. 1-5, 2000.
- Hideshima, T., Chauhan, D., Castro, A., Hayashi, T., Mitsiades, C., Mitsiades, N., Akiyama, M., Richardson, P.G., Schlossman, R.L., Adams, J., Anderson, K.C., NF- γ B as a Therapeutic Target in Multiple Myeloma (MM). Abstract #1581. American Society of Hematology, Dec. 7-11, 2001.
- Lentsch, S., Rogers, M., LeBlanc, R., Birsner, A., Shah, J., Anderson K., D'Amato R., 3-Amino-Phthalimido-Glutarimide (S-3APG) Inhibits Angiogenesis and Growth in Drug Resistant Multiple Myeloma (MM) in vivo. Abstract #1976, American Society of Hematology, Dec. 7-11, 2001.
- Park, Y., Kim, S.A., Kim, C.J., Chung, J.H., Mechanism of the Effect of Thalidomide on Human Multiple Myeloma Cells. Abstract #2685. American Society of Clinical Oncology, May 12-17, 2001.
- Payvandi, F., Wu, L., Haley M., Gupta, D., Zhang, L., Schafer, P., Muller, G.W., Chen, R., Anderson, K.C., Stirling, D., Thalidomide Analogs IMiDS Inhibit Expression of Cyclooxygenase-2 in Multiple Myeloma Cell Line and LPS Stimulated PBMCs. Abstract #2689. American Society of Hematology, Dec. 7-11, 2001.
- Mitsiades, N., Mitsiades, C., Poulaki, V., Akiyama, M., Tai, Y., Lin, B., Hayashi, T., Catley, L., Hideshima, T., Chauhan, D., Treon, S.P., Anderson, K.C., Apoptotic Signaling Induced by Immunomodulatory Thalidomide Analogs (Imids) in Human Multiple Myeloma Cells; Therapeutic Implications. Abstract #3224. American Society of Hematology, Dec. 7-11, 2001.
- "Celgene drug promises activity in solid tumors," *Marketletter*, Jun. 18, 2001.
- Marriott et al., "Thalidomide analogue CDC-501 is safe and well tolerated by patients with end stage cancer and shows evidence of clinical responses and extensive immune activation," *Br. J. Cancer*, 2002, 86(Supp. 1):Abst 6.4.
- Kast, R.E., "Evidence of a mechanism by which etanercept increased TNF-alpha in multiple myeloma: New insights into the biology of TNF-alpha giving new treatment opportunities—the role of burproion," *Leukemia Research*, 2005, 29:1459-1463.
- Tsimberidou, A. et al., "Pilot study of recombinant human soluble tumor necrosis factor (TNF) receptor (p75) fusion protein (TNFR:Fc:Enbrel) in patients with refractory multiple myeloma: increase in plasma TNF α levels during treatment," *Leukemia Research*, 2003, 27:375-380.
- Dimopoulos, et al., "Long-term follow-up on overall survival from the MM-009 and MM-010 phase III trials of lenalidomide plus dexamethasone in patients with relapsed or refractory multiple myeloma," *Leukemia*, 2009, 1-6.
- Hideshima, T., et al., "A review of lenalidomide in combination with dexamethasone for the treatment of multiple myeloma," *Therapeutics and Clinical Risk Management*, 2008, 4(1):129-136.
- Wang, M., et al., "Lenalidomide plus dexamethasone is more effective than dexamethasone alone in patients with relapsed or refractory multiple myeloma regardless of prior thalidomide exposure," *Blood*, 2008, 112(12):4445-4451.
- Gandhi, A., et al., "Dexamethasone Synergizes with Lenalidomide to Inhibit Multiple Myeloma Tumor Growth, But Reduces Lenalidomide-Induced Immunomodulation of T and NK Cell Function," *Current Cancer Drug Targets*, 2010, 10(1):1-13.
- Gay, F. et al., "Lenalidomide plus dexamethasone versus thalidomide plus dexamethasone in newly diagnosed multiple myeloma: a comparative analysis of 411 patients," *Blood*, 2010, 115(97):1343-150.
- Notification letter dated Aug. 30, 2010 from Natco Pharma Limited to Celgene Corporation re: Notification pursuant to § 505(j)(2)(B) of the Federal Food, Drug and Cosmetic Act.
- Complaint for Patent Infringement filed on Oct. 8, 2010 by Celgene Corporation in the U.S. District Court, District of New Jersey against Natco Pharma Limited.
- Answer to Complaint filed on Nov. 18, 2010 by Natco Pharma Limited in the U.S. District Court, District of New Jersey.
- Grosshans, E. and Illy, G., "Thalidomide Therapy for Inflammatory Dermatoses," *International Journal of Dermatology*, 1984, 23(9):598-602.
- Krenn, M. et al., "Improvements in Solubility and Stability of Thalidomide upon Complexation with Hydropropyl- β -Cyclodextrin," *Journal of Pharmaceutical Sciences*, 1992, 81(7):685-689.
- Schmahl, H. J. et al., "Pharmacokinetics of the Teratogenic and Nonteratogenic Thalidomide Analogs EM 12 and Supidimide in the Rat and Marmoset Monkey", in *Pharmacokinetics in Teratogenesis*, CRC Press, 1987, vol. I, Ch. 12, pp. 181-192.
- Schumacher, H. et al., "The Teratogenic Activity of a Thalidomide Analogue, EM₁₂, in Rabbits, Rats, and Monkeys," *Teratology*, 1971, 5:233-240.
- Smith, R. et al., "Studies on the Relationship Between the Chemical Structure and Embryotoxic Activity of Thalidomide and Related Compounds," in *A Symposium on Embryopathic Activity of Drugs*, J. & A. Churchill Ltd., 1965, Session 6, pp. 194-209.
- Sheskin, J. and Sagher, F., "Trials with Thalidomide Derivatives in Leprosy Reactions," *Leprosy Review*, 1968, 39(4):203-205.
- Sheskin, J., "Study with Nine Thalidomide Derivatives in the Leprosy Reaction," *Pharmacology and Therapeutics*, 1978, 17:82-84.
- Raje, N. and Anderson, K., "thalidomide and immunomodulatory drugs as cancer therapy," *Current Opinions in Oncology*, 2002, 14:635-640.
- Kumar, S. et al., "Thalidomide as an anti-cancer agent," *J. Cell. Mod. Med.*, 2002, 6(2):160-174.
- Singhal, S. and Mehta, J., "Thalidomide in Cancer," *BioDrugs*, 2001, 15(3):163-172.
- List, A., "New Approaches to the Treatment of Myelodysplasia," *The Oncologist*, 2002, 7(suppl. 1):39-49.
- Kurzrock, R., "Myelodysplastic syndrome overview," *Seminars in Hematology* (Abstract only), 2002, 39(3)suppl. 2):18-25 Abstract only.
- Goerner, et al., "Morbidity and mortality of chronic GVHD after hematopoietic stem cell transplantation from HLA-identical siblings for patients with aplastic or refractory anemias," *Biology of Blood and Marrow Transplantation* (Abstract only), 2002, 8(1):47-56.
- Thomas, D., "Pilot studies of Thalidomide in Acute Myelogenous Leukemia, Myelodysplastic Syndromes, and Myeloproliferative Disorders," *Seminars in Hematology*, 2000, 37(1)(suppl. 3):26-34.
- Zorat, F. et al., "The clinical and biological effects of thalidomide in patients with myelodysplastic syndromes," *British Journal of Haematology*, 2001, 115:881-894.
- Official Action dated Feb. 10, 2009 in JP Application No. 2004-545192. (English translation provided).
- Teramura, M., Men-ekiyokusei Ryouhou, *Current Therapy*, 2000, 18(5):140-144 (in Japanese).
- Kon-nichi no Chiryuu Shishin, 1997 [Pocket Edition], Igaku Shoin, 1997, 513-514 (in Japanese).



US 7,968,569 B2

Page 10

- Okamoto, T., Kotsuzuikeisei Shoukougun to Men-eki Ijo, Bessatsu Nihon Rinsho. Syndrome Series for each area, No. 22, Blood Syndromes III, Nihon Rinsho, 213-216 (in Japanese), (Oct. 1998).
- Merck Manual, 17th ed. Japanese version, 1999, 951-952.
- Notice of Allowance from U.S. Appl. No. 11/096,155 dated Jan. 12, 2010.
- Rajkumar et al., "Combination therapy with thalidomide plus dexamethasone for newly diagnosed multiple myeloma," *American Society of Hematology*, 43rd Annual Meeting, Dec. 7-11, 2001, Abstract #3525.
- Scheffler et al., "Safety and pharmacokinetics of CDC-501, a novel immunomodulatory-oncologic agent, after single then multiple, oral 100 mg twice daily dOSes," *American Society for Clinical Pharmacology and Therapeutics*, Mar. 24-27, 2002, Abstract #WPIII-63.
- Meregalli et al., "High-dose dexamethasone as first line therapy of multiple myeloma?," *Recenti Progressi in Medicina*, 1998, 89(1):18-20.
- Official Action in corresponding Canadian Application No. 2,476,983, (dated Aug. 21, 2009).
- Liu et al., "Phase I study of CC-5013 (Revimid), a thalidomide derivative, in patients with refractory metastatic cancer," *American Society of Clinical Oncology*, Abstract #927, 2003.
- Zeldis et al. "Update on the evolution of the IMiD™," *International Society for Biological Therapy of Cancer, Oral Abstract*, 2003.
- Patt, Yehuda A.; Hassan, Manal M.; Lozano, Richard D.; Ellis, Lee M.; Peterson, J. Andrew; Waugh, Kimberly A.; *Durable Clinical Response of Refractory Hepatocellular Carcinoma to Orally Administered Thalidomide*, American Journal of Clinical Oncology, 2000.
- Richardson, Paul; Hideshima, Teru; Anderson, Kenneth; *Thalidomide: The Revival of a Drug with Therapeutic Promise in the Treatment of Cancer*, Principles & Practice of Oncology, vol. 15, No. 2, 2001.
- Thomas, Melodie; Doss, Deborah, *Thalidomide Nursing Roundtable Update*, Monograph, Sep. 2002.
- Richardson, Paul; Hideshima, Teru; Anderson, Kenneth; *Thalidomide: Emerging Role in Cancer Medicine*; Annual Review of Medicine, 2002.
- Berenson, J.R.; Bergsagel, P. L.; Munshi, N., *Initiation and Maintenance of Multiple Myeloma*; Seminars in Hematology, vol. 36 No. 1, Supp.3, Jan. 1999, pp. 9-13.
- Gollob, J.A.; Schinipper, C.P.; Orsini, E.; Murphy, E.; Daley, J.F.; Lazo, S.B.; Frank, D.A.; *Characterization of a Novel Subset of CD8 T Cells That Expands in patients Receiving interleukin-12*, 02, Am. Soc. For Clin. Investigation, Inc., vol. 102. No. 3 Aug. 1998, pp. 561-575.
- Cavanagh, L.L.; Barnetson, R.S.; Basten, A.; Halliday, G.M.; *Dendritic Epidermal T-Cell Involvement in Induction of CD8+ T-Cell-Mediated Immunity Against an Ultraviolet Radiation-Induced Skin Tumor* Int. J. Cancer: 70, 98-105, 1997.
- Thomas, D.A., Aguayo, A., Estey, E., Albitar, M., O'Brien, S., Giles, F. J., Beran, M., Cortes, J., Zeldis, I., Keating, M.J., Barlogie, B., Kantarjian, H.M., *Thalidomide as anti-angiogenesis therapy (rx) in refractory or relapsed leukemia*. Abstract #2269, American Society of Hematology, Dec. 3-7, 1999.
- Barlogie, B., Desikan, R., Munshi, N., Siegel, D., Meheas, J., Singhal, S., Anaissie, E., *Single Course D.T. Pace Anti-Angiochemotherapy Effects CR in Plasma Cell Leukemia and Fulminant Multiple Myeloma (MM)*, Abstract #4180. American Society of Hematology, Dec. 1-9, 1998.
- Davies, F. E., Raje, N., Hideshima, T., Lentzsch, S., Young G., Tai, Y., Lin, B.K., Podar, K., Chauhan, D., Treon, S.P., Gupta, D., Mitsiades, C., Mitsiades, N., Hayashi, T., Richardson, P.G., Schlossman, R.L., Muller, G.W., Stirling, D.L., Anderson, K.C., *Thalidomide (THAL) and Immunomodulatory Derivatives (IMiDS) Augment Natural Killer (NK) Cell Cytotoxicity in Multiple Myeloma (MM)*. Abstract #3617. American Society of Hematology, Dec. 1-5, 2000.
- Hideshima, T., Chauhan, D., Castro, A., Hayashi, T Mitsiades, C., Mitsiades, N., Akiyama, M., Richardson, P.G., Schlossman, R.L., Adams, J., Anderson, K.C., *NF-KB as a Therapeutic Target in Multiple Myeloma (MM)*. Abstract # 1581. American Society of Hematology. Dec. 7-11, 2001.
- Lentsch, S., Rogers, M., LeBlanc, R., Birsner, A., Shah, J., Anderson K., D'Amato R., *3-Amino-Phthalimido-Glutarimide (S-3APG) Inhibits Angiogenesis and Growth in Drug Resistant Multiple Myeloma (MM) in vivo*. Abstract # 1976. American Society of Hematology, Dec. 7-11, 2001.
- Park, Y., Kim, S.A., Kim, C.J., Chung, J.H., *Mechanism of the Effect of Thalidomide on Human Multiple Myeloma Cells*. Abstract #2685. American Society of Clinical Oncology, May 12-17, 2001.
- Payvandi, F., Wu, I., Haley M., Gupta, D., Zhang, L., Schafer, P., Muller, G.W., Chen, R., Anderson, K.C., Stirling, D., *Thalidomide Analogs IMiDS Inhibit Expression of Cyclooxygenase-2 in Multiple Myeloma Cell Line and LPS Stimulated PBMCs*. Abstract #2689. American Society of Hematology, Dec. 7-11, 2001.
- Mitslades, N., Mitslades, C., Poulaki, V., Akiyama, M., Tai, Y., Lin, B., Hayashi, T., Cati, F.Y.L., Hideshima, T., Chauhan, D., Treon, S.P., Anderson, K.C., *Apoptotic Signaling Induced By Immunomodulatory Thalidomide Analogs (Imids) in Human Multiple Myeloma Cells; Therapeutic Implications*, Abstract #3224. American Society of Hematology, Dec. 7-11, 2001.
- Richardson, P.G., Schlossman, R.L., Hideshima, T., Davies, F., LeBlanc, R., Catley, L., Doss, D., Kelly, K.A., McKenney, M., Mechlowicz, J., Freeman, A., DeOcampo, Il, R., Rich, R., Ryoo, J., Chauhan, D., Munshi, N., Weller, E., Zeldis, J., Anderson, K.C., *A Phase I Study of Oral CC5013, an Immunomodulatory Thalidomide (Thal) Derivative, in Patients With Relapsed and Refractory Multiple Myeloma (MM)*. Abstract #3225. American Society of Hematology, Dec. 7-11, 2001.
- Zangari, M. Tricot, G., Zeldis, J., Eddlemon, P., Saghafifar, F., Barlogie, B., *Results of Phase I Study of CC5013, for the Treatment of Multiple Myeloma (MM) Patients Who Relapse After High Dose Chemotherapy (HDCT)*. Abstract #3226. American Society of Hematology, Dec. 7-11, 2001.
- Richardson, P. et al., "Thalidomide in multiple myeloma," *Biomed Pharmacother*, 2002, 56:115-28.
- Swartz, G. et al., "Pre-clinical evaluation of ENMD-0995: A thalidomide analog with activity against multiple myeloma and solid tumors," *Cell and Tumor Biology*, 2002, 43:181-182, Abstract# 910.
- Mazucco, R., "Angiogenesis and Anti-angiogenesis Therapeutics," *IDrugs*, 2002, 5(4): 320-322.
- Worker, C., "JP Morgan Hambrecl & Quist—20th Annual Healthcare Conference," *IDrugs*, 5(2):113-116, (2002).
- Treston, A. et al., "Pre-Clinical Evaluation of a Thalidomide Analog with Activity Against Multiple Myeloma and Solid Tumors—ENMD-0995 (S-(-)-3-(3-amino-phthalimido)-glutarimide)," *Blood*, 100(11):816a, Abstract #3225.
- Mazucco, R. and Williams, L., "Immunotherapy, chemoprevention and angiogenesis," *IDrugs*, 2002, 5(5):408-411.
- Fernandes, P., "Anti-Cancer Drug Discovery and Development Summit," *IDrugs*, 2002, 5(8):757-764.
- Notice of Opposition to EP 1 505 973 filed by Synthon B.V. on Nov. 30, 2010.
- Notice of Opposition to EP 1 505 973 filed by Strawman Limited on Dec. 1, 2010.
- Samson, D. et al., "Infusion of Vincristine and Doxorubicin with Oral Dexamethasone as First-Line Therapy for Multiple Myeloma," *The Lancet*, 1989, 334(8668):882-885.
- Barlogie, B. et al., "Effective Treatment of Advanced Multiple Myeloma Refractory to Alkylating Agents," *N. Engl. J. Med.*, 1984, 310(21):1353-1356.
- Dimopoulos, M. et al., "Thalidomide and dexamethasone combination for refractory multiple myeloma," *Annals of Oncology*, 2001, 12:991-995.
- Zangari, M., et al., "Thrombogenic activity of doxorubicin in myeloma patients receiving thalidomide: implications for therapy," *Blood*, 2002, 100:1168-1171.
- List, A. et al., "High Erythropoietic Remitting Activity of the Immunomodulatory Thalidomide Analog, CC5013, in Patients with Myelodysplastic Syndrome (MDS)," Abstract #353, *Blood*, 2002, 100(11):96a.
- Mufti, G. et al., "Myelodysplastic Syndrome," *American Society of Hematology*, 2003, pp. 176-199.
- Extracts from drug databases: retrieved from <http://www.nextbio.com/b/search/ov/IMiD3%20cpd> on Nov. 26, 2010 and <http://pubchem.ncbi.nlm.nih.gov/summary/summary.cgi?cid=216326> on Nov. 26, 2010.

* cited by examiner

Figure 1
Effects of Revimid™ and Thalidomide on MM Cell Proliferation



US 7,968,569 B2

1

**METHODS FOR TREATMENT OF MULTIPLE
MYELOMA USING 3-(4-AMINO-1-OXO-1,3-
DIHYDRO-ISOINDOL-2-YL)-PIPERIDINE-
2,6-DIONE**

This application claims the benefit of U.S. provisional application No. 60/380,842, filed May 17, 2002, and No. 60/424,600, filed Nov. 6, 2002, the entireties of which are incorporated herein by reference.

1. FIELD OF THE INVENTION

This invention relates to methods of treating, preventing and/or managing specific cancers, and other diseases including, but not limited to, those associated with, or characterized by, undesired angiogenesis, by the administration of one or more immunomodulatory compounds alone or in combination with other therapeutics. In particular, the invention encompasses the use of specific combinations, or “cocktails,” of drugs and other therapy, e.g., radiation to treat these specific cancers, including those refractory to conventional therapy. The invention also relates to pharmaceutical compositions and dosing regimens.

2. BACKGROUND OF THE INVENTION

2.1 Pathobiology of Cancer and Other Diseases

Cancer is characterized primarily by an increase in the number of abnormal cells derived from a given normal tissue, invasion of adjacent tissues by these abnormal cells, or lymphatic or blood-borne spread of malignant cells to regional lymph nodes and to distant sites (metastasis). Clinical data and molecular biologic studies indicate that cancer is a multistep process that begins with minor preneoplastic changes, which may under certain conditions progress to neoplasia. The neoplastic lesion may evolve clonally and develop an increasing capacity for invasion, growth, metastasis, and heterogeneity, especially under conditions in which the neoplastic cells escape the host’s immune surveillance. Roitt, I., Brostoff, J and Kale, D., *Immunology*, 17.1-17.12 (3rd ed., Mosby, St. Louis, Mo., 1993).

There is an enormous variety of cancers which are described in detail in the medical literature. Examples includes cancer of the lung, colon, rectum, prostate, breast, brain, and intestine. The incidence of cancer continues to climb as the general population ages, as new cancers develop, and as susceptible populations (e.g., people infected with AIDS or excessively exposed to sunlight) grow. A tremendous demand therefore exists for new methods and compositions that can be used to treat patients with cancer.

Many types of cancers are associated with new blood vessel formation, a process known as angiogenesis. Several of the mechanisms involved in tumor-induced angiogenesis have been elucidated. The most direct of these mechanisms is the secretion by the tumor cells of cytokines with angiogenic properties. Examples of these cytokines include acidic and basic fibroblastic growth factor (a,b-FGF), angiogenin, vascular endothelial growth factor (VEGF), and TNF- α . Alternatively, tumor cells can release angiogenic peptides through the production of proteases and the subsequent breakdown of the extracellular matrix where some cytokines are stored (e.g., b-FGF). Angiogenesis can also be induced indirectly through the recruitment of inflammatory cells (particularly macrophages) and their subsequent release of angiogenic cytokines (e.g., TNF- α , bFGF).

A variety of other diseases and disorders are also associated with, or characterized by, undesired angiogenesis. For

2

example, enhanced or unregulated angiogenesis has been implicated in a number of diseases and medical conditions including, but not limited to, ocular neovascular diseases, choroidal neovascular diseases, retina neovascular diseases, rubeosis (neovascularization of the angle), viral diseases, genetic diseases, inflammatory diseases, allergic diseases, and autoimmune diseases. Examples of such diseases and conditions include, but are not limited to: diabetic retinopathy; retinopathy of prematurity; corneal graft rejection; neovascular glaucoma; retrolental fibroplasia; and proliferative vitreoretinopathy.

Accordingly, compounds that can control angiogenesis or inhibit the production of certain cytokines, including TNF- α , may be useful in the treatment and prevention of various diseases and conditions.

2.2 Methods of Treating Cancer

Current cancer therapy may involve surgery, chemotherapy, hormonal therapy and/or radiation treatment to eradicate neoplastic cells in a patient (see, for example, Stockdale, 1998, *Medicine*, vol. 3, Rubenstein and Federman, eds., Chapter 12, Section IV). Recently, cancer therapy could also involve biological therapy or immunotherapy. All of these approaches pose significant drawbacks for the patient. Surgery, for example, may be contraindicated due to the health of a patient or may be unacceptable to the patient. Additionally, surgery may not completely remove neoplastic tissue. Radiation therapy is only effective when the neoplastic tissue exhibits a higher sensitivity to radiation than normal tissue. Radiation therapy can also often elicit serious side effects. Hormonal therapy is rarely given as a single agent. Although hormonal therapy can be effective, it is often used to prevent or delay recurrence of cancer after other treatments have removed the majority of cancer cells. Biological therapies and immunotherapies are limited in number and may produce side effects such as rashes or swellings, flu-like symptoms, including fever, chills and fatigue, digestive tract problems or allergic reactions.

With respect to chemotherapy, there are a variety of chemotherapeutic agents available for treatment of cancer. A majority of cancer chemotherapeutics act by inhibiting DNA synthesis, either directly, or indirectly by inhibiting the biosynthesis of deoxyribonucleotide triphosphate precursors, to prevent DNA replication and concomitant cell division. Gilman et al., *Goodman and Gilman’s: The Pharmacological Basis of Therapeutics*, Tenth Ed. (McGraw Hill, New York).

Despite availability of a variety of chemotherapeutic agents, chemotherapy has many drawbacks. Stockdale, *Medicine*, vol. 3, Rubenstein and Federman, eds., ch. 12, sect. 10, 1998. Almost all chemotherapeutic agents are toxic, and chemotherapy causes significant, and often dangerous side effects including severe nausea, bone marrow depression, and immunosuppression. Additionally, even with administration of combinations of chemotherapeutic agents, many tumor cells are resistant or develop resistance to the chemotherapeutic agents. In fact, those cells resistant to the particular chemotherapeutic agents used in the treatment protocol often prove to be resistant to other drugs, even if those agents act by different mechanism from those of the drugs used in the specific treatment. This phenomenon is referred to as pleiotropic drug or multidrug resistance. Because of the drug resistance, many cancers prove refractory to standard chemotherapeutic treatment protocols.

Other diseases or conditions associated with, or characterized by, undesired angiogenesis are also difficult to treat. However, some compounds such as protamine, heparin and steroids have been proposed to be useful in the treatment of certain specific diseases. Taylor et al., *Nature* 297:307 (1982);

US 7,968,569 B2

3

Folkman et al., *Science* 221:719 (1983); and U.S. Pat. Nos. 5,001,116 and 4,994,443. Thalidomide and certain derivatives of it have also been proposed for the treatment of such diseases and conditions. U.S. Pat. Nos. 5,593,990, 5,629,327, 5,712,291, 6,071,948 and 6,114,355 to D'Amato.

Still, there is a significant need for safe and effective methods of treating, preventing and managing cancer and other diseases and conditions, particularly for diseases that are refractory to standard treatments, such as surgery, radiation therapy, chemotherapy and hormonal therapy, while reducing or avoiding the toxicities and/or side effects associated with the conventional therapies.

2.3 IMiDS™

A number of studies have been conducted with the aim of providing compounds that can safely and effectively be used to treat diseases associated with abnormal production of TNF- α . See, e.g., Marriott, J. B., et al., *Expert Opin. Biol. Ther.* 1(4):1-8 (2001); G. W. Muller, et al., *Journal of Medicinal Chemistry* 39(17): 3238-3240 (1996); and G. W. Muller, et al, *Bioorganic & Medicinal Chemistry Letters* 8: 2669-2674 (1998). Some studies have focused on a group of compounds selected for their capacity to potently inhibit TNF- α production by LPS stimulated PBMC. L. G. Corral, et al., *Ann. Rheum. Dis.* 58:(Suppl I) 1107-1113 (1999). These compounds, which are referred to as IMiDS™ (Celgene Corporation) or Immunomodulatory Drugs, show not only potent inhibition of TNF- α but also marked inhibition of LPS induced monocyte IL1 β and IL12 production. LPS induced IL6 is also inhibited by immunomodulatory compounds, albeit partially. These compounds are potent stimulators of LPS induced IL10. Id. Particular examples of IMiD™s include, but are not limited to, the substituted 2-(2,6-dioxopiperidin-3-yl) phthalimides and substituted 2-(2,6-dioxopiperidin-3-yl)-1-oxoisindoles described in U.S. Pat. Nos. 6,281,230 and 6,316,471, both to G. W. Muller, et al.

3. SUMMARY OF THE INVENTION

This invention encompasses methods of treating and preventing certain types of cancer, including primary and metastatic cancer, as well as cancers that are refractory or resistant to conventional chemotherapy. The methods comprise administering to a patient in need of such treatment or prevention a therapeutically or prophylactically effective amount of an immunomodulatory compound, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof. The invention also encompasses methods of managing certain cancers (e.g., preventing or prolonging their recurrence, or lengthening the time of remission) which comprise administering to a patient in need of such management a prophylactically effective amount of an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

In particular methods of the invention, an immunomodulatory compound is administered in combination with a therapy conventionally used to treat, prevent or manage cancer. Examples of such conventional therapies include, but are not limited to, surgery, chemotherapy, radiation therapy, hormonal therapy, biological therapy and immunotherapy.

This invention also encompasses methods of treating, managing or preventing diseases and disorders other than cancer that are associated with, or characterized by, undesired angiogenesis, which comprise administering to a patient in need of such treatment, management or prevention a therapeutically or prophylactically effective amount of an immunomodula-

4

tory compound, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

In other methods of the invention, an immunomodulatory compound is administered in combination with a therapy conventionally used to treat, prevent or manage diseases or disorders associated with, or characterized by, undesired angiogenesis. Examples of such conventional therapies include, but are not limited to, surgery, chemotherapy, radiation therapy, hormonal therapy, biological therapy and immunotherapy.

This invention encompasses pharmaceutical compositions, single unit dosage forms, dosing regimens and kits which comprise an immunomodulatory compound, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, and a second, or additional, active agent. Second active agents include specific combinations, or "cocktails," of drugs.

4. BRIEF DESCRIPTION OF FIGURE

FIG. 1 shows a comparison of the effects of 3-(4-amino-1-oxo-1,3-dihydro-isindol-2-yl)-piperidine-2,6-dione (Revimid™) and thalidomide in inhibiting the proliferation of multiple myeloma (MM) cell lines in an in vitro study. The uptake of [³H]-thymidine by different MM cell lines (MM. 1S, Hs Sultan, U266 and RPMI-8226) was measured as an indicator of the cell proliferation.

5. DETAILED DESCRIPTION OF THE INVENTION

A first embodiment of the invention encompasses methods of treating, managing, or preventing cancer which comprises administering to a patient in need of such treatment or prevention a therapeutically or prophylactically effective amount of an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

In particular methods encompassed by this embodiment, the immunomodulatory compound is administered in combination with another drug ("second active agent") or method of treating, managing, or preventing cancer. Second active agents include small molecules and large molecules (e.g., proteins and antibodies), examples of which are provided herein, as well as stem cells. Methods, or therapies, that can be used in combination with the administration of the immunomodulatory compound include, but are not limited to, surgery, blood transfusions, immunotherapy, biological therapy, radiation therapy, and other non-drug based therapies presently used to treat, prevent or manage cancer.

Another embodiment of the invention encompasses methods of treating, managing or preventing diseases and disorders other than cancer that are characterized by undesired angiogenesis. These methods comprise the administration of a therapeutically or prophylactically effective amount of an immunomodulatory compound, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

Examples of diseases and disorders associated with, or characterized by, undesired angiogenesis include, but are not limited to, inflammatory diseases, autoimmune diseases, viral diseases, genetic diseases, allergic diseases, bacterial diseases, ocular neovascular diseases, choroidal neovascular diseases, retina neovascular diseases, and rubeosis (neovascularization of the angle).

In particular methods encompassed by this embodiment, the immunomodulatory compound is administered in combina-

US 7,968,569 B2

5

tion with a second active agent or method of treating, managing, or preventing the disease or condition. Second active agents include small molecules and large molecules (e.g., proteins and antibodies), examples of which are provided herein, as well as stem cells. Methods, or therapies, that can be used in combination with the administration of the immunomodulatory compound include, but are not limited to, surgery, blood transfusions, immunotherapy, biological therapy, radiation therapy, and other non-drug based therapies presently used to treat, prevent or manage disease and conditions associated with, or characterized by, undesired angiogenesis.

The invention also encompasses pharmaceutical compositions (e.g., single unit dosage forms) that can be used in methods disclosed herein. Particular pharmaceutical compositions comprise an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, and a second active agent.

5.1 Immunomodulatory Compounds

Compounds used in the invention include immunomodulatory compounds that are racemic, stereomerically enriched or stereomerically pure, and pharmaceutically acceptable salts, solvates, hydrates, stereoisomers, clathrates, and prodrugs thereof. Preferred compounds used in the invention are small organic molecules having a molecular weight less than about 1,000 g/mol, and are not proteins, peptides, oligonucleotides, oligosaccharides or other macromolecules.

As used herein and unless otherwise indicated, the terms "immunomodulatory compounds" and "IMiDs™" (Celgene Corporation) encompasses small organic molecules that markedly inhibit TNF- α , LPS induced monocyte IL1 β and IL12, and partially inhibit IL6 production. Specific immunomodulatory compounds are discussed below.

TNF- α is an inflammatory cytokine produced by macrophages and monocytes during acute inflammation. TNF- α is responsible for a diverse range of signaling events within cells. TNF- α may play a pathological role in cancer. Without being limited by theory, one of the biological effects exerted by the immunomodulatory compounds of the invention is the reduction of synthesis of TNF- α . Immunomodulatory compounds of the invention enhance the degradation of TNF- α mRNA.


Further, without being limited by theory, immunomodulatory compounds used in the invention may also be potent co-stimulators of T cells and increase cell proliferation dramatically in a dose dependent manner. Immunomodulatory compounds of the invention may also have a greater co-stimulatory effect on the CD8+ T cell subset than on the CD4+ T cell subset. In addition, the compounds preferably have anti-inflammatory properties, and efficiently co-stimulate T cells.

Specific examples of immunomodulatory compounds of the invention, include, but are not limited to, cyano and carboxy derivatives of substituted styrenes such as those disclosed in U.S. Pat. No. 5,929,117; 1-oxo-2-(2,6-dioxo-3-fluoropiperidin-3-yl) isoindolines and 1,3-dioxo-2-(2,6-dioxo-3-fluoropiperidine-3-yl) isoindolines such as those described in U.S. Pat. No. 5,874,448; the tetra substituted 2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolines described in U.S. Pat. No. 5,798,368; 1-oxo and 1,3-dioxo-2-(2,6-dioxopiperidin-3-yl) isoindolines (e.g., 4-methyl derivatives of thalidomide and EM-12), including, but not limited to, those disclosed in U.S. Pat. No. 5,635,517; and a class of non-polypeptide cyclic amides disclosed in U.S. Pat. Nos. 5,698,579 and 5,877,200; analogs and derivatives of thalidomide, including hydrolysis products, metabolites, derivatives and precursors of thalidomide, such as those described in U.S.

6

Pat. Nos. 5,593,990, 5,629,327, and 6,071,948 to D'Amato; aminothalidomide, as well as analogs, hydrolysis products, metabolites, derivatives and precursors of aminothalidomide, and substituted 2-(2,6-dioxopiperidin-3-yl) phthalimides and substituted 2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindoles such as those described in U.S. Pat. Nos. 6,281,230 and 6,316,471; isoindole-imide compounds such as those described in U.S. patent application Ser. No. 09/972,487 filed on Oct. 5, 2001, U.S. patent application Ser. No. 10/032,286 filed on Dec. 21, 2001, and International Application No. PCT/US01/50401 (International Publication No. WO 02/059106). The entireties of each of the patents and patent applications identified herein are incorporated herein by reference. Immunomodulatory compounds of the invention do not include thalidomide.


Other specific immunomodulatory compounds of the invention include, but are not limited to, 1-oxo- and 1,3-dioxo-2-(2,6-dioxopiperidin-3-yl) isoindolines substituted with amino in the benzo ring as described in U.S. Pat. No. 5,635,517 which is incorporated herein by reference. These compounds have the structure I:



in which one of X and Y is C=O, the other of X and Y is C=O or CH₂, and R² is hydrogen or lower alkyl, in particular methyl. Specific immunomodulatory compounds include, but are not limited to:


- 1-oxo-2-(2,6-dioxopiperidin-3-yl)-4-aminoisoindoline;
- 1-oxo-2-(2,6-dioxopiperidin-3-yl)-5-aminoisoindoline;
- 1-oxo-2-(2,6-dioxopiperidin-3-yl)-6-aminoisoindoline;
- 1-oxo-2-(2,6-dioxopiperidin-3-yl)-7-aminoisoindoline;
- 1,3-dioxo-2-(2,6-dioxopiperidin-3-yl)-4-aminoisoindoline;
- and
- 1,3-dioxo-2-(2,6-dioxopiperidin-3-yl)-5-aminoisoindoline.

Other specific immunomodulatory compounds of the invention belong to a class of substituted 2-(2,6-dioxopiperidin-3-yl) phthalimides and substituted 2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindoles, such as those described in U.S. Pat. Nos. 6,281,230; 6,316,471; 6,335,349; and 6,476,052, and International Patent Application No. PCT/US97/13375 (International Publication No. WO 98/03502), each of which is incorporated herein by reference. Compounds representative of this class are of the formulas:




US 7,968,569 B2

7



wherein R^1 is hydrogen or methyl. In a separate embodiment, the invention encompasses the use of enantiomerically pure forms (e.g. optically pure (R) or (S) enantiomers) of these compounds.

Still other specific immunomodulatory compounds of the invention belong to a class of isoindole-imides disclosed in U.S. patent application Ser. Nos. 10/032,286 and 09/972,487, and International Application No. PCT/US01/50401 (International Publication No. WO 02/059106), each of which are incorporated herein by reference. Representative compounds are of formula II:



and pharmaceutically acceptable salts, hydrates, solvates, clathrates, enantiomers, diastereomers, racemates, and mixtures of stereoisomers thereof, wherein:

one of X and Y is C=O and the other is CH₂ or C≡O;

R^1 is H, (C₁-C₈)alkyl, (C₃-C₇)cycloalkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, benzyl, aryl, (C₀-C₄)alkyl-(C₁-C₆)heterocycloalkyl, (C₀-C₄)alkyl-(C₂-C₅)heteroaryl, C(O)R³, C(S)R³, C(O)OR⁴, (C₁-C₈)alkyl-N(R⁶)₂, (C₁-C₈)alkyl-OR⁵, (C₁-C₈)alkyl-C(O)OR⁵, C(O)NHR³, C(S)NHR³, C(O)NR³R^{3'}, C(S)NR³R^{3'} or (C₁-C₈)alkyl-O(CO)R⁵;

R^2 is H, F, benzyl, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, or (C₂-C₈)alkynyl;

R^3 and $R^{3'}$ are independently (C₁-C₈)alkyl, (C₃-C₇)cycloalkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, benzyl, aryl, (C₀-C₄)alkyl-(C₁-C₆)heterocycloalkyl, (C₀-C₄)alkyl-(C₂-C₅)heteroaryl, (C₀-C₈)alkyl-N(R⁶)₂, (C₁-C₈)alkyl-OR⁵, (C₁-C₈)alkyl-C(O)OR⁵, (C₁-C₈)alkyl-O(CO)R⁵, or C(O)OR⁵;

R^4 is (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, (C₁-C₄)alkyl-OR⁵, benzyl, aryl, (C₀-C₄)alkyl-(C₁-C₆)heterocycloalkyl, or (C₀-C₄)alkyl-(C₂-C₅)heteroaryl;

8

R^5 is (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, benzyl, aryl, or (C₂-C₅)heteroaryl;

each occurrence of R^6 is independently H, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, benzyl, aryl, (C₂-C₅)heteroaryl, or (C₀-C₈)alkyl-C(O)O—R⁵ or the R^6 groups can join to form a heterocycloalkyl group;

n is 0 or 1; and

* represents a chiral-carbon center.

In specific compounds of formula II, when n is 0 then R^1 is (C₃-C₇)cycloalkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, benzyl, aryl, (C₀-C₄)alkyl-(C₁-C₆)heterocycloalkyl, (C₀-C₄)alkyl-(C₂-C₅)heteroaryl, C(O)R³, C(O)OR⁴, (C₁-C₈)alkyl-N(R⁶)₂, (C₁-C₈)alkyl-OR⁵, (C₁-C₈)alkyl-C(O)OR⁵, C(S)NHR³, or (C₁-C₈)alkyl O(CO)R⁵;


R^2 is H or (C₁-C₈)alkyl; and

R^3 is (C₁-C₈)alkyl, (C₃-C₇)cycloalkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, benzyl, aryl, (C₀-C₄)alkyl-(C₁-C₆)heterocycloalkyl, (C₀-C₄)alkyl-(C₂-C₅)heteroaryl, (C₅-C₈)alkyl-N(R⁶)₂; (C₀-C₈)alkyl-NH—C(O)O—R⁵; (C₁-C₈)alkyl-OR⁵, (C₁-C₈)alkyl-C(O)OR⁵, (C₁-C₈)alkyl-O(CO)R⁵, or C(O)OR⁵; and the other variables have the same definitions.


In other specific compounds of formula II, R^2 is H or (C₁-C₄)alkyl.

In other specific compounds of formula II, R^1 is (C₁-C₈)alkyl or benzyl.

In other specific compounds of formula II, R^1 is H, (C₁-C₈)alkyl, benzyl, CH₂OCH₃, CH₂CH₂OCH₃, or



In another embodiment of the compounds of formula II, R^1 is



wherein Q is O or S, and each occurrence of R^7 is independently H, (C₁-C₈)alkyl, benzyl, CH₂OCH₃, or CH₂CH₂OCH₃.

In other specific compounds of formula II, R^1 is C(O)R³.

In other specific compounds of formula II, R^3 is (C₀-C₄)alkyl-(C₂-C₅)heteroaryl, (C₁-C₅)alkyl, aryl, or (C₀-C₄)alkyl-OR⁵.

In other specific compounds of formula II, heteroaryl is pyridyl, furyl, or thienyl.


In other specific compounds of formula II, R^1 is C(O)OR⁴.

In other specific compounds of formula II, the H of C(O)NHC(O) can be replaced with (C₁-C₄)alkyl, aryl, or benzyl.

Still other specific immunomodulatory compounds of the invention belong to a class of isoindole-imides disclosed in U.S. patent application Ser. No. 09/781,179, International Publication No. WO 98/54170, and U.S. Pat. No. 6,395,754, each of which are incorporated herein by reference. Representative compounds are of formula III:

US 7,968,569 B2

9



and pharmaceutically acceptable salts, hydrates, solvates, clathrates, enantiomers, diastereomers, racemates, and mixtures of stereoisomers thereof, wherein:

one of X and Y is C=O and the other is CH₂ or C=O;

R is H or CH₂OCOR⁵;

(i) each of R¹, R², R³, or R⁴, independently of the others, is halo, alkyl of 1 to 4 carbon atoms, or alkoxy of 1 to 4 carbon atoms or (ii) one of R¹, R², R³, or R⁴ is nitro or —NHR⁵ and the remaining of R¹, R², R³, or R⁴ are hydrogen;

R⁵ is hydrogen or alkyl of 1 to 8 carbons

R⁶ hydrogen, alkyl of 1 to 8 carbon atoms, benzo, chloro, or fluoro;

R⁷ is R⁷—CHR¹⁰—N(R⁸R⁹);


R⁷ is m-phenylene or p-phenylene or —(C_nH_{2n})— in which n has a value of 0 to 4;

each of R⁸ and R⁹ taken independently of the other is hydrogen or alkyl of 1 to 8 carbon atoms, or R⁸ and R⁹ taken together are tetramethylene, pentamethylene, hexamethylene, or —CH₂CH₂[X]₁CH₂CH₂— in which [X]₁ is —O—, —S—, or —NH—;


R¹⁰ is hydrogen, alkyl of 1 to 8 carbon atoms, or phenyl; and

* represents a chiral-carbon center.

The most preferred immunomodulatory compounds of the invention are 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione and 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. The compounds can be obtained via standard, synthetic methods (see e.g., U.S. Pat. No. 5,635, 517, incorporated herein by reference). The compounds are available from Celgene Corporation, Warren, N.J. 4-(Amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (ACTIMID™) has the following chemical structure:



The compound 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (REVIMID™) has the following chemical structure:



10

III
Compounds of the invention can either be commercially purchased or prepared according to the methods described in the patents or patent publications disclosed herein. Further, optically pure compounds can be asymmetrically synthesized or resolved using known resolving agents or chiral columns as well as other standard synthetic organic chemistry techniques.

As used herein and unless otherwise indicated, the term “pharmaceutically acceptable salt” encompasses non-toxic acid and base addition salts of the compound to which the term refers. Acceptable non-toxic acid addition salts include those derived from organic and inorganic acids or bases known in the art, which include, for example, hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid, methanesulphonic acid, acetic acid, tartaric acid, lactic acid, succinic acid, citric acid, malic acid, maleic acid, sorbic acid, aconitic acid, salicylic acid, phthalic acid, embolic acid, enanthic acid, and the like.

Compounds that are acidic in nature are capable of forming salts with various pharmaceutically acceptable bases. The bases that can be used to prepare pharmaceutically acceptable base addition salts of such acidic compounds are those that form non-toxic base addition salts, i.e., salts containing pharmacologically acceptable cations such as, but not limited to, alkali metal or alkaline earth metal salts and the calcium, magnesium, sodium or potassium salts in particular. Suitable organic bases include, but are not limited to, N,N-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine), lysine, and procaine.

As used herein and unless otherwise indicated, the term “prodrug” means a derivative of a compound that can hydrolyze, oxidize, or otherwise react under biological conditions (in vitro or in vivo) to provide the compound. Examples of prodrugs include, but are not limited to, derivatives of immunomodulatory compounds of the invention that comprise biodegradable moieties such as biodegradable amides, biodegradable esters, biodegradable carbamates, biodegradable carbonates, biodegradable ureides, and biodegradable phosphate analogues. Other examples of prodrugs include derivatives of immunomodulatory compounds of the invention that comprise —NO, —NO₂, —ONO, or —ONO₂ moieties. Prodrugs can typically be prepared using well-known methods, such as those described in 1 *Burger's Medicinal Chemistry and Drug Discovery*, 172-178, 949-982 (Manfred E. Wolff ed., 5th ed. 1995), and *Design of Prodrugs* (H. Bundgaard ed., Elsevier, N.Y. 1985).

As used herein and unless otherwise indicated, the terms “biohydrolyzable amide,” “biohydrolyzable ester,” “biohydrolyzable carbamate,” “biohydrolyzable carbonate,” “biohydrolyzable ureide,” “biohydrolyzable phosphate” mean an amide, ester, carbamate, carbonate, ureide, or phosphate, respectively, of a compound that either: 1) does not interfere with the biological activity of the compound but can confer upon that compound advantageous properties in vivo, such as uptake, duration of action, or onset of action; or 2) is biologically inactive but is converted in vivo to the biologically active compound. Examples of biohydrolyzable esters include, but are not limited to, lower alkyl esters, lower acyloxyalkyl esters (such as acetoxymethyl, acetoxylethyl, aminocarbonyloxymethyl, pivaloyloxymethyl, and pivaloyloxyethyl esters), lactonyl esters (such as phthalidyl and thiophthalidyl esters), lower alkoxyacyloxyalkyl esters (such as methoxycarbonyl-oxyethyl, ethoxycarbonyloxyethyl and isopropoxycarbonyloxyethyl esters), alkoxyalkyl esters, choline esters, and acylamino alkyl esters (such as acetamidomethyl esters). Examples of biohydrolyzable amides

US 7,968,569 B2

11

include, but are not limited to, lower alkyl amides, α -amino acid amides, alkoxyacyl amides, and alkylaminoalkylcarbonyl amides. Examples of biohydrolyzable carbamates include, but are not limited to, lower alkylamines, substituted ethylenediamines, amino acids, hydroxyalkylamines, heterocyclic and heteroaromatic amines, and polyether amines.

Various immunomodulatory compounds of the invention contain one or more chiral centers, and can exist as racemic mixtures of enantiomers or mixtures of diastereomers. This invention encompasses the use of stereomerically pure forms of such compounds, as well as the use of mixtures of those forms. For example, mixtures comprising equal or unequal amounts of the enantiomers of a particular immunomodulatory compounds of the invention may be used in methods and compositions of the invention. These isomers may be asymmetrically synthesized or resolved using standard techniques such as chiral columns or chiral resolving agents. See, e.g., Jacques, J., et al., *Enantiomers, Racemates and Resolutions* (Wiley-Interscience, New York, 1981); Wilen, S. H., et al., *Tetrahedron* 33:2725 (1977); Eliel, E. L., *Stereochemistry of Carbon Compounds* (McGraw-Hill, N.Y., 1962); and Wilen, S. H., *Tables of Resolving Agents and Optical Resolutions* p. 268 (E. L. Eliel, Ed., Univ. of Notre Dame Press, Notre Dame, Ind., 1972).

As used herein and unless otherwise indicated, the term "stereomerically pure" means a composition that comprises one stereoisomer of a compound and is substantially free of other stereoisomers of that compound. For example, a stereomerically pure composition of a compound having one chiral center will be substantially free of the opposite enantiomer of the compound. A stereomerically pure composition of a compound having two chiral centers will be substantially free of other diastereomers of the compound. A typical stereomerically pure compound comprises greater than about 80% by weight of one stereoisomer of the compound and less than about 20% by weight of other stereoisomers of the compound, more preferably greater than about 90% by weight of one stereoisomer of the compound and less than about 10% by weight of the other stereoisomers of the compound, even more preferably greater than about 95% by weight of one stereoisomer of the compound and less than about 5% by weight of the other stereoisomers of the compound, and most preferably greater than about 97% by weight of one stereoisomer of the compound and less than about 3% by weight of the other stereoisomers of the compound. As used herein and unless otherwise indicated, the term "stereomerically enriched" means a composition that comprises greater than about 60% by weight of one stereoisomer of a compound, preferably greater than about 70% by weight, more preferably greater than about 80% by weight of one stereoisomer of a compound. As used herein and unless otherwise indicated, the term "enantiomerically pure" means a stereomerically pure composition of a compound having one chiral center. Similarly, the term "stereomerically enriched" means a stereomerically enriched composition of a compound having one chiral center.

It should be noted that if there is a discrepancy between a depicted structure and a name given that structure, the depicted structure is to be accorded more weight. In addition, if the stereochemistry of a structure or a portion of a structure is not indicated with, for example, bold or dashed lines, the structure or portion of the structure is to be interpreted as encompassing all stereoisomers of it.

5.2 Second Active Agents

Immunomodulatory compounds can be combined with other pharmacologically active compounds ("second active agents") in methods and compositions of the invention. It is

12

believed that certain combinations work synergistically in the treatment of particular types of cancer and certain diseases and conditions associated with, or characterized by, undesired angiogenesis. Immunomodulatory compounds can also work to alleviate adverse effects associated with certain second active agents, and some second active agents can be used to alleviate adverse effects associated with immunomodulatory compounds.

One or more second active ingredients or agents can be used in the methods and compositions of the invention together with an immunomodulatory compound. Second active agents can be large molecules (e.g., proteins) or small molecules (e.g., synthetic inorganic, organometallic, or organic molecules).

Examples of large molecule active agents include, but are not limited to, hematopoietic growth factors, cytokines, and monoclonal and polyclonal antibodies. Typical large molecule active agents are biological molecules, such as naturally occurring or artificially made proteins. Proteins that are particularly useful in this invention include proteins that stimulate the survival and/or proliferation of hematopoietic precursor cells and immunologically active poietic cells in vitro or in vivo. Others stimulate the division and differentiation of committed erythroid progenitors in cells in vitro or in vivo. Particular proteins include, but are not limited to: interleukins, such as IL-2 (including recombinant IL-II ("rIL2") and canarypox IL-2), IL-10, IL-12, and IL-18; interferons, such as interferon alfa-2a, interferon alfa-2b, interferon alfa-n1, interferon alfa-n3, interferon beta-I a, and interferon gamma-I b; GM-CSF and GM-CSF; and EPO.

Particular proteins that can be used in the methods and compositions of the invention include, but are not limited to: filgrastim, which is sold in the United States under the trade name Neupogen® (Amgen, Thousand Oaks, Calif.); sargramostim, which is sold in the United States under the trade name Leukine® (Immunex, Seattle, Wash.); and recombinant EPO, which is sold in the United States under the trade name Epogen® (Amgen, Thousand Oaks, Calif.).

Recombinant and mutated forms of GM-CSF can be prepared as described in U.S. Pat. Nos. 5,391,485; 5,393,870; and 5,229,496; all of which are incorporated herein by reference. Recombinant and mutated forms of G-CSF can be prepared as described in U.S. Pat. Nos. 4,810,643; 4,999,291; 5,528,823; and 5,580,755; all of which are incorporated herein by reference.

This invention encompasses the use of native, naturally occurring, and recombinant proteins. The invention further encompasses mutants and derivatives (e.g., modified forms) of naturally occurring proteins that exhibit, in vivo, at least some of the pharmacological activity of the proteins upon which they are based. Examples of mutants include, but are not limited to, proteins that have one or more amino acid residues that differ from the corresponding residues in the naturally occurring forms of the proteins. Also encompassed by the term "mutants" are proteins that lack carbohydrate moieties normally present in their naturally occurring forms (e.g., nonglycosylated forms). Examples of derivatives include, but are not limited to, pegylated derivatives and fusion proteins, such as proteins formed by fusing IgG1 or IgG3 to the protein or active portion of the protein of interest. See, e.g., Penichet, M. L. and Morrison, S. L., *J. Immunol. Methods* 248:91-101 (2001).

Antibodies that can be used in combination with compounds of the invention include monoclonal and polyclonal antibodies. Examples of antibodies include, but are not limited to, trastuzumab (Herceptin®), rituximab (Rituxan®), bevacizumab (Avastin™), pertuzumab (Omnitarg™), tositu-

US 7,968,569 B2

13

momab (Bexxar®), edrecolomab (Panorex®), and G250. Compounds of the invention can also be combined with, or used in combination with, anti-TNF- α antibodies.

Large molecule active agents may be administered in the form of anti-cancer vaccines. For example, vaccines that secrete, or cause the secretion of, cytokines such as IL-2, G-CSF, and GM-CSF can be used in the methods, pharmaceutical compositions, and kits of the invention. See, e.g., Emens, L. A., et al., *Curr. Opinion Mol. Ther.* 3(1):77-84 (2001).

In one embodiment of the invention, the large molecule active agent reduces, eliminates, or prevents an adverse effect associated with the administration of an immunomodulatory compound. Depending on the particular immunomodulatory compound and the disease or disorder begin treated, adverse effects can include, but are not limited to, drowsiness and somnolence, dizziness and orthostatic hypotension, neutropenia, infections that result from neutropenia, increased HIV-viral load, bradycardia, Stevens-Johnson Syndrome and toxic epidermal necrolysis, and seizures (e.g., grand mal convulsions). A specific adverse effect is neutropenia.

Second active agents that are small molecules can also be used to alleviate adverse effects associated with the administration of an immunomodulatory compound. However, like some large molecules, many are believed to be capable of providing a synergistic effect when administered with (e.g., before, after or simultaneously) an immunomodulatory compound. Examples of small molecule second active agents include, but are not limited to, anti-cancer agents, antibiotics, immunosuppressive agents, and steroids.

Examples of anti-cancer agents include, but are not limited to: acivicin; aclarubicin; acodazole hydrochloride; acronine; adozelesin; aldesleukin; altretamine; ambomycin; amet-antrone acetate; amsacrine; anastrozole; anthramycin; asparaginase; asperlin; azacitidine; azetepa; azotomycin; batimastat; benzodepa; bicalutamide; bisantrene hydrochloride; bisnafide dimesylate; bizelesin; bleomycin sulfate; brequinar sodium; bropirimine; busulfan; cactinomycin; calusterone; caracemide; carbetimer; carboplatin; carmustine; carubicin hydrochloride; carzelesin; cedefingol; celecoxib (COX-2 inhibitor); chlorambucil; cirolemycin; cisplatin; cladribine; crisnatol mesylate; cyclophosphamide; cytarabine; dacarbazine; dactinomycin; daunorubicin hydrochloride; decitabine; dexormaplatin; dezaguanine; dezaguanine mesylate; diaziqone; docetaxel; doxorubicin; doxorubicin hydrochloride; droloxifene; droloxifene citrate; dromostanolone propionate; duazomycin; edatrexate; eflomithine hydrochloride; elsamitucin; enloplatin; enpromate; epipropidine; epirubicin hydrochloride; erbulozole; esorubicin hydrochloride; estramustine; estramustine phosphate sodium; etanidazole; etoposide; etoposide phosphate; etoprine; fadrozole hydrochloride; fazarabine; fenretinide; floxuridine; fludarabine phosphate; fluorouracil; flurocitabine; fosquidone; fostriecin sodium; gemcitabine; gemcitabine hydrochloride; hydroxyurea; idarubicin hydrochloride; ifosfamide; ilmofofosine; iproplatin; irinotecan; irinotecan hydrochloride; lanreotide acetate; letrozole; leuprolide acetate; liarozole hydrochloride; lometrexol sodium; lomustine; losoxantrone hydrochloride; masoprocol; maytansine; mechlorethamine hydrochloride; megestrol acetate; melengestrol acetate; melphalan; menogaril; mercaptopurine; methotrexate; methotrexate sodium; metoprine; meturedepa; mitindomide; mitocarcin; mitocromin; mitogillin; mitomalcin; mitomycin; mitosper; mitotane; mitoxantrone hydrochloride; mycophenolic acid; nocodazole; nogalamycin; ormaplatin; oxisuran; paclitaxel; pegaspargase; peliomycin; pentamustine; peplomycin sulfate; perfosfamide; pipo-

14

broman; pipsulfan; piroxantrone hydrochloride; plicamycin; plomestane; porfimer sodium; porfiromycin; prednimustine; procarbazine hydrochloride; puromycin; puromycin hydrochloride; pyrazofurin; riboprine; safingol; safingol hydrochloride; semustine; simtrazene; sparfosate sodium; sparsomycin; spirogermanium hydrochloride; spiro-mustine; spiroplatin; streptonigrin; streptozocin; sulofenur; talisomycin; tecogalan sodium; taxotere; tegafur; telox-antrone hydrochloride; temoporfin; teniposide; teroxirone; testolactone; thiamiprine; thioguanine; thiotepea; tiaozofurin; tirapazamine; toremifene citrate; trestolone acetate; triciribine phosphate; trimetrexate; trimetrexate glucuronate; triptorelin; tubulozole hydrochloride; uracil mustard; uredepa; vapreotide; verteporfin; vinblastine sulfate; vincristine sulfate; vindesine; vindesine sulfate; vinepidine sulfate; vinglycinate sulfate; vinleurosine sulfate; vinorelbine tartrate; vinrosidine sulfate; vinzolidine sulfate; vorozole; zeniplatin; zinostatin; and zorubicin hydrochloride.

Other anti-cancer drugs include, but are not limited to: 20-epi-1,25 dihydroxyvitamin D3; 5-ethynyluracil; abiraterone; aclarubicin; acylfulvene; adecypenol; adozelesin; aldesleukin; ALL-TK antagonists; altretamine; ambamustine; amidox; amifostine; aminolevulinic acid; amrubicin; amsacrine; anagrelide; anastrozole; andrographolide; angiogenesis inhibitors; antagonist D; antagonist G; antarelix; antidorsalizing morphogenetic protein-1; antiandrogen, prostatic carcinoma; antiestrogen; antineoplastin; antisense oligonucleotides; aphidicolin glycinate; apoptosis gene modulators; apoptosis regulators; apurinic acid; ara-CDP-DL-PTBA; arginine deaminase; asulacrine; atamestane; atrimustine; axinastatin 1; axinastatin 2; axinastatin 3; azasetron; azatoxin; azatyrosine; baccatin III derivatives; balanol; batimastat; BCR/ABL antagonists; benzochlorins; benzoylstauroporine; beta lactam derivatives; beta-alethine; betaclamycin B; betulinic acid; bFGF inhibitor; bicalutamide; bisantrene; bisaziridinylspermine; bisnafide; bistratene A; bizelesin; breflate; bropirimine; budotitane; buthionine sulfoximine; calcipotriol; calphostin C; camptothecin derivatives; capecitabine; carboxamide-amino-triazole; carboxyamidotriazole; CaRest M3; CARN 700; cartilage derived inhibitor; carzelesin; casein kinase inhibitors (ICOS); castanospermine; cecropin B; cetrorelix; chlorlins; chloroquinoline sulfonamide; cicaprost; cis-porphyrin; cladribine; clomifene analogues; clotrimazole; collismycin A; collismycin B; combretastatin A4; combretastatin analogue; conagenin; crambescidin 816; crisnatol; cryptophycin 8; cryptophycin A derivatives; curacin A; cyclopentantraquinones; cycloplatan; cypemycin; cytarabine ocfosfate; cytolytic factor; cytostatin; dacliximab; decitabine; dehydridemnin B; deslorelin; dexamethasone; dexifosfamide; dexrazoxane; dextrapamil; diaziqone; didemnin B; didox; diethylnor-spermine; dihydro-5-azacytidine; dihydrotaxol, 9-; dioxamycin; diphenyl spiromustine; docetaxel; docosanol; dolasetron; doxifluridine; doxorubicin; droloxifene; dronabinol; duocarmycin SA; ebselen; ecomustine; edelfosine; edrecolomab; eflomithine; elemene; emitofur; epirubicin; epriesteride; estramustine analogue; estrogen agonists; estrogen antagonists; etanidazole; etoposide phosphate; exemestane; fadrozole; fazarabine; fenretinide; filgrastim; finasteride; flavopiridol; flezelastine; fluasterone; fludarabine; fluorodaunorubicin hydrochloride; forfenimex; formestane; fostriecin; fotemustine; gadolinium texaphyrin; gallium nitrate; galocitabine; ganirelix; gelatinase inhibitors; gemcitabine; glutathione inhibitors; hepsulfam; heregulin; hexamethylene bisacetamide; hypericin; ibandronic acid; idarubicin; idoxifene; idramantone; ilmofofosine; ilomastat; imatinib (e.g., Gleevec®), imiquimod; immunostimulant peptides;

US 7,968,569 B2

15

insulin-like growth factor-1 receptor inhibitor; interferon agonists; interferons; interleukins; iobenguane; iododoxorubicin; ipomeanol, 4-; iroplact; irsogladine; isobengazole; isohomohalicondrin B; itasetron; jasplakinolide; kahalalide F; lamellarin-N triacetate; lanreotide; leinamycin; lenograstim; lentinan sulfate; leptolstatin; letrozole; leukemia inhibiting factor; leukocyte alpha interferon; leuprolide+estrogen+progesterone; leuprorelin; levamisole; liarozole; linear polyamine analogue; lipophilic disaccharide peptide; lipophilic platinum compounds; lissoclinamide 7; lobaplatin; lombricine; lometrexol; lonidamine; losoxantrone; loxoribine; lurtotecan; lutetium texaphyrin; lysofylline; lytic peptides; maitansine; mannostatin A; marimastat; masoprocol; maspin; matrilysin inhibitors; matrix metalloproteinase inhibitors; menogaril; merbarone; meterelin; methioninase; metoclopramide; MLF inhibitor; mifepristone; miltefosine; mirimostim; mitoguanzone; mitolactol; mitomycin analogues; mitonafide; mitotoxin fibroblast growth factor-saporin; mitoxantrone; mofarotene; molgramostim; Erbitux, human chorionic gonadotrophin; monophosphoryl lipid A+mycobacterium cell wall sk; mopidamol; mustard anticancer agent; mycaperoxide B; mycobacterial cell wall extract; myriaporone; N-acetyldinaline; N-substituted benzamides; nafarelin; nagrestip; naloxone+pentazocine; napavin; naphterpin; nartograstim; nedaplatin; nemorubicin; neridronic acid; nilutamide; nisamycin; nitric oxide modulators; nitroxide antioxidant; nitrullyn; oblimersen (Genasense®); O⁶-benzylguanine; octreotide; okicenone; oligonucleotides; onapristone; ondansetron; ondansetron; oracin; oral cytokine inducer; ormaplatin; osaterone; oxaliplatin; oxaunomycin; paclitaxel; paclitaxel analogues; paclitaxel derivatives; palauamine; palmitoylrhizoxin; pamidronic acid; panaxyriol; panomifene; parabactin; pazelliptine; pegaspargase; peldesine; pentosan polysulfate sodium; pentostatin; pentrozole; perflubron; perfosfamide; perillyl alcohol; phenazinomycin; phenylacetate; phosphatase inhibitors; picibanil; pilocarpine hydrochloride; pirarubicin; piritrexim; placetin A; placetin B; plasminogen activator inhibitor; platinum complex; platinum compounds; platinum-triamine complex; porfimer sodium; porfirofomycin; prednisone; propyl bis-acridone; prostaglandin J2; proteasome inhibitors; protein A-based immune modulator; protein kinase C inhibitor; protein kinase C inhibitors, microalgal; protein tyrosine phosphatase inhibitors; purine nucleoside phosphorylase inhibitors; purpurins; pyrazoloacridine; pyridoxylated hemoglobin polyoxyethylene conjugate; raf antagonists; raltitrexed; ramosetron; ras famesyl protein transferase inhibitors; ras inhibitors; ras-GAP inhibitor; retelliptine demethylated; rhenium Re 186 etidronate; rhizoxin; ribozymes; RII retinamide; rohitukine; romurtide; roquinimex; rubiginone B1; ruboxyl; safinol; saintopin; SarCNU; sarcophytol A; sargramostim; Sdi 1 mimetics; semustine; senescence derived inhibitor 1; sense oligonucleotides; signal transduction inhibitors; sizofiran; sobuzoxane; sodium borocaptate; sodium phenylacetate; solverol; somatomedin binding protein; sonermin; sparfosic acid; spicamycin D; spiromustine; splenopentin; spongistatin 1; squalamine; stipiamide; stromelysin inhibitors; sulfinosine; superactive vasoactive intestinal peptide antagonist; suradista; suramin; swainsonine; tallimustine; tamoxifen methiodide; tauromustine; tazarotene; tecogalan sodium; tegafur; tellurapyrylium; telomerase inhibitors; temoporfin; teniposide; tetrachlorodecaoxide; tetrazomine; thaliblastine; thiocoraline; thrombopoietin; thrombopoietin mimetic; thymalfasin; thymopoietin receptor agonist; thymotrinan; thyroid stimulating hormone; tin ethyl etiopurpurin; tirapazamine; titanocene bichloride; topsentin; toremifene; translation inhibitors; tretinoiin; triacetyluridine; triciribine;

16

trimetrexate; triptorelin; tropisetron; turosteride; tyrosine kinase inhibitors; tyrphostins; UBC inhibitors; ubenimex; urogenital sinus-derived growth inhibitory factor; urokinase receptor antagonists; vapreotide; variolin B; velaresol; veramine; verdins; verteporfin; vinorelbine; vinxaltine; vitaxin; vorozole; zanoterone; zeniplatin; zilascorb; and zinostatin stimalamer.

Specific second active agents include, but are not limited to, oblimersen (Genasense®), remicade, docetaxel, celecoxib, melphalan, dexamethasone (Decadron®), steroids, gemcitabine, cisplatin, temozolomide, etoposide, cyclophosphamide, temodar, carboplatin, procarbazine, gliadel, tamoxifen, topotecan, methotrexate, Arisa®, taxol, taxotere, fluorouracil, leucovorin, irinotecan, xeloda, CPT-11, interferon alpha, pegylated interferon alpha (e.g., PEG INTRON-A), capecitabine, cisplatin, thiotepe, fludarabine, carboplatin, liposomal daunorubicin, cytarabine, doxetaxol, pacilitaxel, vinblastine, IL-2, GM-CSF, dacarbazine, vinorelbine, zoledronic acid, palmitronate, biacin, busulphan, prednisone, bisphosphonate, arsenic trioxide, vincristine, doxorubicin (Doxil®), paclitaxel, ganciclovir, adriamycin, estramustine sodium phosphate (Emcyt®), sulindac, and etoposide.

5.3 Methods of Treatments and Prevention

Methods of this invention encompass methods of treating, preventing and/or managing various types of cancer and diseases and disorders associated with, or characterized by, undesired angiogenesis. As used herein, unless otherwise specified, the term “treating” refers to the administration of a compound of the invention or other additional active agent after the onset of symptoms of the particular disease or disorder. As used herein, unless otherwise specified, the term “preventing” refers to the administration prior to the onset of symptoms, particularly to patients at risk of cancer, and other diseases and disorders associated with, or characterized by, undesired angiogenesis. The term “prevention” includes the inhibition of a symptom of the particular disease or disorder. Patients with familial history of cancer and diseases and disorders associated with, or characterized by, undesired angiogenesis are preferred candidates for preventive regimens. As used herein and unless otherwise indicated, the term “managing” encompasses preventing the recurrence of the particular disease or disorder in a patient who had suffered from it, and/or lengthening the time a patient who had suffered from the disease or disorder remains in remission.

As used herein, the term “cancer” includes, but is not limited to, solid tumors and blood born tumors. The term “cancer” refers to disease of skin tissues, organs, blood, and vessels, including, but not limited to, cancers of the bladder, bone or blood, brain, breast, cervix, chest, colon, endometrium, esophagus, eye, head, kidney, liver, lymph nodes, lung, mouth, neck, ovaries, pancreas, prostate, rectum, stomach, testis, throat, and uterus. Specific cancers include, but are not limited to, advanced malignancy, amyloidosis, neuroblastoma, meningioma, hemangiopericytoma, multiple brain metastase, glioblastoma multiforms, glioblastoma, brain stem glioma, poor prognosis malignant brain tumor, malignant glioma, recurrent malignant glioma, anaplastic astrocytoma, anaplastic oligodendroglioma, neuroendocrine tumor, rectal adenocarcinoma, Dukes C & D colorectal cancer, unresectable colorectal carcinoma, metastatic hepatocellular carcinoma, Kaposi’s sarcoma, karotype acute myeloblastic leukemia, Hodgkin’s lymphoma, non-Hodgkin’s lymphoma, cutaneous T-Cell lymphoma, cutaneous B-Cell lymphoma, diffuse large B-Cell lymphoma, low grade follicular lymphoma, malignant melanoma, malignant mesothelioma, malignant pleural effusion mesothelioma syndrome, peritoneal carcinoma, papillary serous carcinoma, gynec-

US 7,968,569 B2

17

logic sarcoma, soft tissue sarcoma, scleroderma, cutaneous vasculitis, Langerhans cell histiocytosis, leiomyosarcoma, fibrodysplasia ossificans progressive, hormone refractory prostate cancer, resected high-risk soft tissue sarcoma, unresectable hepatocellular carcinoma, Waldenström's macroglobulinemia, smoldering myeloma, indolent myeloma, fallopian tube cancer, androgen independent prostate cancer, androgen dependent stage IV non-metastatic prostate cancer, hormone-insensitive prostate cancer, chemotherapy-insensitive prostate cancer, papillary thyroid carcinoma, follicular thyroid carcinoma, medullary thyroid carcinoma, and leiomyoma. In a specific embodiment, the cancer is metastatic. In another embodiment, the cancer is refractory or resistance to chemotherapy or radiation; in particular, refractory to thalidomide.

As used herein to refer to diseases and conditions other than cancer, the terms "diseases or disorders associated with, or characterized by, undesired angiogenesis," "diseases or disorders associated with undesired angiogenesis," and "diseases or disorders characterized by undesired angiogenesis" refer to diseases, disorders and conditions that are caused, mediated or attended by undesired, unwanted or uncontrolled angiogenesis, including, but not limited to, inflammatory diseases, autoimmune diseases, genetic diseases, allergic diseases, bacterial diseases, ocular neovascular diseases, choroidal neovascular diseases, and retina neovascular diseases.

Examples of such diseases or disorders associated with undesired angiogenesis include, but are not limited to, diabetic retinopathy, retinopathy of prematurity, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, proliferative vitreoretinopathy, trachoma, myopia, optic pits, epidemic keratoconjunctivitis, atopic keratitis, superior limbic keratitis, pterygium keratitis sicca, sjogrens, acne rosacea, phlyctenulosis, syphilis, lipid degeneration, bacterial ulcer, fungal ulcer, Herpes simplex infection, Herpes zoster infection, protozoan infection, Kaposi sarcoma, Mooren ulcer, Terrien's marginal degeneration, mariginal keratolysis, rheumatoid arthritis, systemic lupus, polyarteritis, trauma, Wegeners sarcoidosis, Scleritis, Steven's Johnson disease, peripheral radial keratotomy, sickle cell anemia, sarcoid, pseudoxanthoma elasticum, Pagets disease, vein occlusion, artery occlusion, carotid obstructive disease, chronic uveitis, chronic vitritis, Lyme's disease, Eales disease, Bechets disease, retinitis, choroiditis, presumed ocular histoplasmosis, Bests disease, Stargarts disease, pars planitis, chronic retinal detachment, hyperviscosity syndromes, toxoplasmosis, rubeosis, sarcodiosis, sclerosis, soriatis, psoriasis, primary sclerosing cholangitis, proctitis, primary biliary srosis, idiopathic pulmonary fibrosis, and alcoholic hepatitis.

In specific embodiments of the invention, diseases or disorders associated with undesired angiogenesis do not include congestive heart failure, cardiomyopathy, pulmonary edema, endotoxin-mediated septic shock, acute viral myocarditis, cardiac allograft rejection, myocardial infarction, HIV, hepatitis, adult respiratory distress syndrome, bone-resorption disease, chronic obstructive pulmonary diseases, chronic pulmonary inflammatory disease, dermatitis, cystic fibrosis, septic shock, sepsis, endotoxic shock, hemodynamic shock, sepsis syndrome, post ischemic reperfusion injury, meningitis, psoriasis, fibrotic disease, cachexia, graft rejection, rheumatoid spondylitis, osteoporosis, Crohn's disease, ulcerative colitis, inflammatory-bowel disease, multiple sclerosis, systemic lupus erythrematosus, erythema nodosum leprosum in leprosy, radiation damage, asthma, hyperoxic alveolar injury, malaria, mycobacterial infection, and opportunistic infections resulting from HIV.

18

This invention encompasses methods of treating patients who have been previously treated for cancer or diseases or disorders associated with, or characterized by, undesired angiogenesis, but are non-responsive to standard therapies, as well as those who have not previously been treated. The invention also encompasses methods of treating patients regardless of patient's age, although some diseases or disorders are more common in certain age groups. The invention further encompasses methods of treating patients who have undergone surgery in an attempt to treat the disease or condition at issue, as well as those who have not. Because patients with cancer and diseases and disorders characterized by undesired angiogenesis have heterogenous clinical manifestations and varying clinical outcomes, the treatment given to a patient may vary, depending on his/her prognosis. The skilled clinician will be able to readily determine without undue experimentation specific secondary agents, types of surgery, and types of non-drug based standard therapy that can be effectively used to treat an individual patient with cancer and other diseases or disorders.

Methods encompassed by this invention comprise administering one or more immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, to a patient (e.g., a human) suffering, or likely to suffer, from cancer or a disease or disorder mediated by undesired angiogenesis.

In one embodiment of the invention, an immunomodulatory compound of the invention can be administered orally and in single or divided daily doses in an amount of from about 0.10 to about 150 mg/day. In a particular embodiment, 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™) may be administered in an amount of from about 0.1 to about 1 mg per day, or alternatively from about 0.1 to about 5 mg every other day. In a preferred embodiment, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) may be administered in an amount of from about 5 to 25 mg per day, or alternatively from about 10 to about 50 mg every other day.

In a specific embodiment, 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™) may be administered in an amount of about 1, 2, or 5 mg per day to patients with relapsed multiple myeloma. In a particular embodiment, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) may be administered initially in an amount of 5 mg/day and the dose can be escalated every week to 10, 20, 25, 30 and 50 mg/day. In a specific embodiment, Revimid™ can be administered in an amount of up to about 30 mg/day to patients with solid tumor. In a particular embodiment, Revimid™ can be administered in an amount of up to about 40 mg/day to patients with glioma.

5.3.1 Combination Therapy with a Second Active Agent

Specific methods of the invention comprise administering an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, in combination with one or more second active agents, and/or in combination with radiation therapy, blood transfusions, or surgery. Examples of immunomodulatory compounds of the invention are disclosed herein (see, e.g., section 5.1). Examples of second active agents are also disclosed herein (see, e.g., section 5.2).

Administration of the immunomodulatory compounds and the second active agents to a patient can occur simultaneously or sequentially by the same or different routes of administration. The suitability of a particular route of administration employed for a particular active agent will depend on the active agent itself (e.g., whether it can be administered orally

US 7,968,569 B2

19

without decomposing prior to entering the blood stream) and the disease being treated. A preferred route of administration for an immunomodulatory compound of the invention is orally. Preferred routes of administration for the second active agents or ingredients of the invention are known to those of ordinary skill in the art. See, e.g., *Physicians' Desk Reference*, 1755-1760 (56th ed., 2002).

In one embodiment of the invention, the second active agent is administered intravenously or subcutaneously and once or twice daily in an amount of from about 1 to about 1000 mg, from about 5 to about 500 mg, from about 10 to about 350 mg, or from about 50 to about 200 mg. The specific amount of the second active agent will depend on the specific agent used, the type of disease being treated or managed, the severity and stage of disease, and the amount(s) of immunomodulatory compounds of the invention and any optional additional active agents concurrently administered to the patient. In a particular embodiment, the second active agent is oblimersen (Genasense®), GM-CSF, G-CSF, EPO, taxotere, irinotecan, dacarbazine, transretinoic acid, topotecan, pentoxifylline, ciprofloxacin, dexamethasone, vincristine, doxorubicin, COX-2 inhibitor, IL2, IL8, IL18, IFN, Ara-C, vinorelbine, or a combination thereof.

In a particular embodiment, GM-CSF, G-CSF or EPO is administered subcutaneously during about five days in a four or six week cycle in an amount of from about 1 to about 750 mg/m²/day, preferably in an amount of from about 25 to about 500 mg/m²/day, more preferably in an amount of from about 50 to about 250 mg/m²/day, and most preferably in an amount of from about 50 to about 200 mg/m²/day. In a certain embodiment, GM-CSF may be administered in an amount of from about 60 to about 500 mcg/m² intravenously over 2 hours, or from about 5 to about 12 mcg/m²/day subcutaneously. In a specific embodiment, G-CSF may be administered subcutaneously in an amount of about 1 mcg/kg/day initially and can be adjusted depending on rise of total granulocyte counts. The maintenance dose of G-CSF may be administered in an amount of about 300 (in smaller patients) or 480 mcg subcutaneously. In a certain embodiment, EPO may be administered subcutaneously in an amount of 10,000 Unit 3 times per week.

In another embodiment, Revimid™ in an amount of about 25 mg/d and dacarbazine in an amount of about from 200 to 1,000 mg/m²/d are administered to patients with metastatic malignant melanoma. In a specific embodiment, Revimid™ is administered in an amount of from about 5 to about 25 mg/d to patients with metastatic malignant melanoma whose disease has progressed on treatment with dacarbazine, IL-2 or IFN. In a specific embodiment, Revimid™ is administered to patients with relapsed or refractory multiple myeloma in an amount of about 15 mg/d twice a day or about 30 mg/d four times a day in a combination with dexamethasone.

In another embodiment, an immunomodulatory compound is administered with melphalan and dexamethasone to patients with amyloidosis. In a specific embodiment, an immunomodulatory compound of the invention and steroids can be administered to patients with amyloidosis.

In another embodiment, an immunomodulatory compound is administered with gemcitabine and cisplatin to patients with locally advanced or metastatic transitional cell bladder cancer.

In another embodiment, an immunomodulatory compound is administered in combination with a second active ingredient as follows: temozolomide to pediatric patients with relapsed or progressive brain tumors or recurrent neuroblastoma; celecoxib, etoposide and cyclophosphamide for relapsed or progressive CNS cancer; temodar to patients with

20

recurrent or progressive meningioma, malignant meningioma, hemangiopericytoma, multiple brain metastases, relapsed brain tumors, or newly diagnosed glioblastoma multiformis; irinotecan to patients with recurrent glioblastoma; carboplatin to pediatric patients with brain stem glioma; procarbazine to pediatric patients with progressive malignant gliomas; cyclophosphamide to patients with poor prognosis malignant brain tumors, newly diagnosed or recurrent glioblastoma multiformis; Gliadel® for high grade recurrent malignant gliomas; temozolomide and tamoxifen for anaplastic astrocytoma; or topotecan for gliomas, glioblastoma, anaplastic astrocytoma or anaplastic oligodendroglioma.

In another embodiment, an immunomodulatory compound is administered with methotrexate and cyclophosphamide to patients with metastatic breast cancer.

In another embodiment, an immunomodulatory compound is administered with temozolomide to patients with neuroendocrine tumors.

In another embodiment, an immunomodulatory compound is administered with gemcitabine to patients with recurrent or metastatic head or neck cancer. In another embodiment, an immunomodulatory compound is administered with gemcitabine to patients with pancreatic cancer.

In another embodiment, an immunomodulatory compound is administered to patients with colon cancer in combination with Arisa®, taxol and/or taxotere.

In another embodiment, an immunomodulatory compound is administered with capecitabine to patients with refractory colorectal cancer or patients who fail first line therapy or have poor performance in colon or rectal adenocarcinoma.

In another embodiment, an immunomodulatory compound is administered in combination with fluorouracil, leucovorin, and irinotecan to patients with Dukes C & D colorectal cancer or to patients who have been previously treated for metastatic colorectal cancer.

In another embodiment, an immunomodulatory compound is administered to patients with refractory colorectal cancer in combination with capecitabine, xeloda, and/or CPT-11.

In another embodiment, an immunomodulatory compound of the invention is administered with capecitabine and irinotecan to patients with refractory colorectal cancer or to patients with unresectable or metastatic colorectal carcinoma.

In another embodiment, an immunomodulatory compound is administered alone or in combination with interferon alpha or capecitabine to patients with unresectable or metastatic hepatocellular carcinoma; or with cisplatin and thiotepa to patients with primary or metastatic liver cancer.

In another embodiment, an immunomodulatory compound is administered in combination with pegylated interferon alpha to patients with Kaposi's sarcoma.

In another embodiment, an immunomodulatory compound is administered in combination with fludarabine, carboplatin, and/or topotecan to patients with refractory or relapsed or high-risk acuted myelogenous leukemia.

In another embodiment, an immunomodulatory compound is administered in combination with liposomal daunorubicin, topotecan and/or cytarabine to patients with unfavorable karyotype acute myeloblastic leukemia.

In another embodiment, an immunomodulatory compound is administered in combination with gemcitabine and irinotecan to patients with non-small cell lung cancer. In one embodiment, an immunomodulatory compound is administered in combination with carboplatin and irinotecan to patients with non-small cell lung cancer. In one embodiment, an immunomodulatory compound is administered with dox-

US 7,968,569 B2

21

etaxol to patients with non-small cell lung cancer who have been previously treated with carbo/VP 16 and radiotherapy.

In another embodiment, an immunomodulatory compound is administered in combination with carboplatin and/or taxotere, or in combination with carboplatin, paclitaxel and/or thoracic radiotherapy to patients with non-small cell lung cancer. In a specific embodiment, an immunomodulatory compound is administered in combination with taxotere to patients with stage IIIB or IV non-small cell lung cancer.

In another embodiment, an immunomodulatory compound of the invention is administered in combination with oblimersen (Genasense®) to patients with small cell lung cancer.

In another embodiment, an immunomodulatory compound is administered alone or in combination with a second active ingredient such as vinblastine or fludarabine to patients with various types of lymphoma, including, but not limited to, Hodgkin's lymphoma, non-Hodgkin's lymphoma, cutaneous T-Cell lymphoma, cutaneous B-Cell lymphoma, diffuse large B-Cell lymphoma or relapsed or refractory low grade follicular lymphoma.

In another embodiment, an immunomodulatory compound is administered in combination with taxotere, IL-2, IFN, GM-CSF, and/or dacarbazine to patients with various types or stages of melanoma.

In another embodiment, an immunomodulatory compound is administered alone or in combination with vinorelbine to patients with malignant mesothelioma, or stage IIIB non-small cell lung cancer with pleural implants or malignant pleural effusion mesothelioma syndrome.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of multiple myeloma in combination with dexamethasone, zoledronic acid, palmitronate, GM-CSF, biacin, vinblastine, melphalan, busulphan, cyclophosphamide, IFN, palmidronate, prednisone, bisphosphonate, celecoxib, arsenic trioxide, PEG INTRON-A, vincristine, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with relapsed or refractory multiple myeloma in combination with doxorubicin (Doxil®), vincristine and/or dexamethasone (Decadron®).

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of ovarian cancer such as peritoneal carcinoma, papillary serous carcinoma, refractory ovarian cancer or recurrent ovarian cancer, in combination with taxol, carboplatin, doxorubicin, gemcitabine, cisplatin, xeloda, paclitaxel, dexamethasone, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of prostate cancer, in combination with xeloda, 5 FU/LV, gemcitabine, irinotecan plus gemcitabine, cyclophosphamide, vincristine, dexamethasone, GM-CSF, celecoxib, taxotere, ganciclovir, paclitaxel, adriamycin, docetaxel, estramustine, Emcyt, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of renal cell cancer, in combination with capecitabine, IFN, tamoxifen, IL-2, GM-CSF, Celebrex®, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of gynecologic, uterus or soft tissue sarcoma cancer in combination with IFN, a COX-2 inhibitor such as Celebrex®, and/or sulindac.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of

22

solid tumors in combination with celebrex, etoposide, cyclophosphamide, docetaxel, apicitabine, IFN, tamoxifen, IL-2, GM-CSF, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with scleroderma or cutaneous vasculitis in combination with celebrex, etoposide, cyclophosphamide, docetaxel, apicitabine, IFN, tamoxifen, IL-2, GM-CSF, or a combination thereof.

This invention also encompasses a method of increasing the dosage of an anti-cancer drug or agent that can be safely and effectively administered to a patient, which comprises administering to a patient (e.g., a human) an immunomodulatory compound of the invention, or a pharmaceutically acceptable derivative, salt, solvate, clathrate, hydrate, or pro-drug thereof. Patients that can benefit by this method are those likely to suffer from an adverse effect associated with anti-cancer drugs for treating a specific cancer of the skin, subcutaneous tissue, lymph nodes, brain, lung, liver, bone, intestine, colon, heart, pancreas, adrenal, kidney, prostate, breast, colorectal, or combinations thereof. The administration of an immunomodulatory compound of the invention alleviates or reduces adverse effects which are of such severity that it would otherwise limit the amount of anti-cancer drug.

In one embodiment, an immunomodulatory compound of the invention can be administered orally and daily in an amount of from about 0.1 to about 150 mg, and preferably from about 1 to about 50 mg, more preferably from about 2 to about 25 mg prior to, during, or after the occurrence of the adverse effect associated with the administration of an anti-cancer drug to a patient. In a particular embodiment, an immunomodulatory compound of the invention is administered in combination with specific agents such as heparin, aspirin, coumadin, or G-CSF to avoid adverse effects that are associated with anti-cancer drugs such as but not limited to neutropenia or thrombocytopenia.

In one embodiment, an immunomodulatory compound of the invention can be administered to patients with diseases and disorders associated with, or characterized by, undesired angiogenesis in combination with additional active ingredients including but not limited to anti-cancer drugs, anti-inflammatories, antihistamines, antibiotics, and steroids.

In another embodiment, this invention encompasses a method of treating, preventing and/or managing cancer, which comprises administering an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, in conjunction with (e.g. before, during, or after) conventional therapy including, but not limited to, surgery, immunotherapy, biological therapy, radiation therapy, or other non-drug based therapy presently used to treat, prevent or manage cancer. The combined use of the immunomodulatory compounds of the invention and conventional therapy may provide a unique treatment regimen that is unexpectedly effective in certain patients. Without being limited by theory, it is believed that immunomodulatory compounds of the invention may provide additive or synergistic effects when given concurrently with conventional therapy.

As discussed elsewhere herein, the invention encompasses a method of reducing, treating and/or preventing adverse or undesired effects associated with conventional therapy including, but not limited to, surgery, chemotherapy, radiation therapy, hormonal therapy, biological therapy and immunotherapy. One or more immunomodulatory compounds of the invention and other active ingredient can be administered to a patient prior to, during, or after the occurrence of the adverse effect associated with conventional therapy.

US 7,968,569 B2

23

In one embodiment, an immunomodulatory compound of the invention can be administered in an amount of from about 0.1 to about 150 mg, and preferably from about 1 to about 25 mg, more preferably from about 2 to about 10 mg orally and daily alone, or in combination with a second active agent disclosed herein (see, e.g., section 5.2), prior to, during, or after the use of conventional therapy.

In a specific embodiment of this method, an immunomodulatory compound of the invention and doxorubicin are administered to patients with non-small cell lung cancer who were previously treated with carboplatin and radiotherapy.

5.3.2 Use with Transplantation Therapy

Compounds of the invention can be used to reduce the risk of Graft Versus Host Disease (GVHD). Therefore, the invention encompasses a method of treating, preventing and/or managing cancer, which comprises administering the immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, in conjunction with transplantation therapy.

As those of ordinary skill in the art are aware, the treatment of cancer is often based on the stages and mechanism of the disease. For example, as inevitable leukemic transformation develops in certain stages of cancer, transplantation of peripheral blood stem cells, hematopoietic stem cell preparation or bone marrow may be necessary. The combined use of the immunomodulatory compound of the invention and transplantation therapy provides a unique and unexpected synergism. In particular, an immunomodulatory compound of the invention exhibits immunomodulatory activity that may provide additive or synergistic effects when given concurrently with transplantation therapy in patients with cancer.

An immunomodulatory compound of the invention can work in combination with transplantation therapy reducing complications associated with the invasive procedure of transplantation and risk of GVHD. This invention encompasses a method of treating, preventing and/or managing cancer which comprises administering to a patient (e.g., a human) an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, before, during, or after the transplantation of umbilical cord blood, placental blood, peripheral blood stem cell, hematopoietic stem cell preparation or bone marrow. Examples of stem cells suitable for use in the methods of the invention are disclosed in U.S. provisional patent application No. 60/372,348, filed Apr. 12, 2002 by R. Hariri et al., the entirety of which is incorporated herein by reference.

In one embodiment of this method, an immunomodulatory compound of the invention is administered to patients with multiple myeloma before, during, or after the transplantation of autologous peripheral blood progenitor cell.

In another embodiment, an immunomodulatory compound is administered to patients with relapsing multiple myeloma after the stem cell transplantation.

In another embodiment, an immunomodulatory compound and prednisone are administered as maintenance therapy to patients with multiple myeloma following the transplantation of autologous stem cell.

In another embodiment, an immunomodulatory compound and dexamethasone are administered as salvage therapy for low risk post transplantation to patients with multiple myeloma.

In another embodiment, an immunomodulatory compound and dexamethasone are administered as maintenance therapy to patients with multiple myeloma following the transplantation of autologous bone marrow.

24

In another embodiment, an immunomodulatory compound is administered following the administration of high dose of melphalan and the transplantation of autologous stem cell to patients with chemotherapy responsive multiple myeloma.

In another embodiment, an immunomodulatory compound and PEG INTRON-A are administered as maintenance therapy to patients with multiple myeloma following the transplantation of autologous CD34-selected peripheral stem cell.

In another embodiment, an immunomodulatory compound is administered with post transplant consolidation chemotherapy to patients with newly diagnosed multiple myeloma to evaluate anti-angiogenesis.

In another embodiment, an immunomodulatory compound and dexamethasone are administered as maintenance therapy after DCEP consolidation, following the treatment with high dose of melphalan and the transplantation of peripheral blood stem cell to 65 years of age or older patients with multiple myeloma.

5.3.3 Cycling Therapy

In certain embodiments, the prophylactic or therapeutic agents of the invention are cyclically administered to a patient. Cycling therapy involves the administration of an active agent for a period of time, followed by a rest for a period of time, and repeating this sequential administration. Cycling therapy can reduce the development of resistance to one or more of the therapies, avoid or reduce the side effects of one of the therapies, and/or improves the efficacy of the treatment.

Consequently, in one specific embodiment of the invention, an immunomodulatory compound of the invention is administered daily in a single or divided doses in a four to six week cycle with a rest period of about a week or two weeks. The invention further allows the frequency, number, and length of dosing cycles to be increased. Thus, another specific embodiment of the invention encompasses the administration of an immunomodulatory compound of the invention for more cycles than are typical when it is administered alone. In yet another specific embodiment of the invention, an immunomodulatory compound of the invention is administered for a greater number of cycles that would typically cause dose-limiting toxicity in a patient to whom a second active ingredient is not also being administered.

In one embodiment, an immunomodulatory compound of the invention is administered daily and continuously for three or four weeks at a dose of from about 0.1 to about 150 mg/d followed by a break of one or two weeks. Actimid™ is preferably administered daily and continuously at an initial dose of 0.1 to 5 mg/d with dose escalation (every week) by 1 to 10 mg/d to a maximum dose of 50 mg/d for as long as therapy is tolerated. In a particular embodiment, Revimid™ is administered in an amount of about 5, 10, or 25 mg/day, preferably in an amount of about 10 mg/day for three to four weeks, followed by one week or two weeks of rest in a four or six week cycle.

In one embodiment of the invention, an immunomodulatory compound of the invention and a second active ingredient are administered orally, with administration of an immunomodulatory compound of the invention occurring 30 to 60 minutes prior to a second active ingredient, during a cycle of four to six weeks. In another embodiment of the invention, the combination of an immunomodulatory compound of the invention and a second active ingredient is administered by intravenous infusion over about 90 minutes every cycle. In a specific embodiment, one cycle comprises the administration of from about 10 to about 25 mg/day of Revimid™ and from about 50 to about 200 mg/m²/day of a second active ingredient daily for three to four weeks and then one or two weeks of

US 7,968,569 B2

25

rest. In another specific embodiment, each cycle comprises the administration of from about 5 to about 10 mg/day of Actimid™ and from about 50 to about 200 mg/m²/day of a second active ingredient for 3 to 4 weeks followed by one or two weeks of rest. Typically, the number of cycles during which the combinatorial treatment is administered to a patient will be from about one to about 24 cycles, more typically from about two to about 16 cycles, and even more typically from about four to about three cycles.

5.4 Pharmaceutical Compositions and Dosage Forms

Pharmaceutical compositions can be used in the preparation of individual, single unit dosage forms. Pharmaceutical compositions and dosage forms of the invention comprise an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof. Pharmaceutical compositions and dosage forms of the invention can further comprise one or more excipients.

Pharmaceutical compositions and dosage forms of the invention can also comprise one or more additional active ingredients. Consequently, pharmaceutical compositions and dosage forms of the invention comprise the active ingredients disclosed herein (e.g., an immunomodulatory compound and a second active agent). Examples of optional second, or additional, active ingredients are disclosed herein (see, e.g., section 5.2).

Single unit dosage forms of the invention are suitable for oral, mucosal (e.g., nasal, sublingual, vaginal, buccal, or rectal), parenteral (e.g., subcutaneous, intravenous, bolus injection, intramuscular, or intraarterial), topical (e.g., eye drops or other ophthalmic preparations), transdermal or transcutaneous administration to a patient. Examples of dosage forms include, but are not limited to: tablets; caplets; capsules, such as soft elastic gelatin capsules; cachets; troches; lozenges; dispersions; suppositories; powders; aerosols (e.g., nasal sprays or inhalers); gels; liquid dosage forms suitable for oral or mucosal administration to a patient, including suspensions (e.g., aqueous or non-aqueous liquid suspensions, oil-in-water emulsions, or a water-in-oil liquid emulsions), solutions, and elixirs; liquid dosage forms suitable for parenteral administration to a patient; eye drops or other ophthalmic preparations suitable for topical administration; and sterile solids (e.g., crystalline or amorphous solids) that can be reconstituted to provide liquid dosage forms suitable for parenteral administration to a patient.

The composition, shape, and type of dosage forms of the invention will typically vary depending on their use. For example, a dosage form used in the acute treatment of a disease may contain larger amounts of one or more of the active ingredients it comprises than a dosage form used in the chronic treatment of the same disease. Similarly, a parenteral dosage form may contain smaller amounts of one or more of the active ingredients it comprises than an oral dosage form used to treat the same disease. These and other ways in which specific dosage forms encompassed by this invention will vary from one another will be readily apparent to those skilled in the art. See, e.g., *Remington's Pharmaceutical Sciences*, 18th ed., Mack Publishing, Easton Pa. (1990).

Typical pharmaceutical compositions and dosage forms comprise one or more excipients. Suitable excipients are well known to those skilled in the art of pharmacy, and non-limiting examples of suitable excipients are provided herein. Whether a particular excipient is suitable for incorporation into a pharmaceutical composition or dosage form depends on a variety of factors well known in the art including, but not limited to, the way in which the dosage form will be administered to a patient. For example, oral dosage forms such as

26

tablets may contain excipients not suited for use in parenteral dosage forms. The suitability of a particular excipient may also depend on the specific active ingredients in the dosage form. For example, the decomposition of some active ingredients may be accelerated by some excipients such as lactose, or when exposed to water. Active ingredients that comprise primary or secondary amines are particularly susceptible to such accelerated decomposition. Consequently, this invention encompasses pharmaceutical compositions and dosage forms that contain little, if any, lactose other mono- or disaccharides. As used herein, the term "lactose-free" means that the amount of lactose present, if any, is insufficient to substantially increase the degradation rate of an active ingredient.

Lactose-free compositions of the invention can comprise excipients that are well known in the art and are listed, for example, in the *U.S. Pharmacopeia* (USP) 25-NF20 (2002). In general, lactose-free compositions comprise active ingredients, a binder/filler, and a lubricant in pharmaceutically compatible and pharmaceutically acceptable amounts. Preferred lactose-free dosage forms comprise active ingredients, microcrystalline cellulose, pre-gelatinized starch, and magnesium stearate.

This invention further encompasses anhydrous pharmaceutical compositions and dosage forms comprising active ingredients, since water can facilitate the degradation of some compounds. For example, the addition of water (e.g., 5%) is widely accepted in the pharmaceutical arts as a means of simulating long-term storage in order to determine characteristics such as shelf-life or the stability of formulations over time. See, e.g., Jens T. Carstensen, *Drug Stability: Principles & Practice*, 2d. Ed., Marcel Dekker, NY, N.Y., 1995, pp. 379-80. In effect, water and heat accelerate the decomposition of some compounds. Thus, the effect of water on a formulation can be of great significance since moisture and/or humidity are commonly encountered during manufacture, handling, packaging, storage, shipment, and use of formulations.

Anhydrous pharmaceutical compositions and dosage forms of the invention can be prepared using anhydrous or low moisture containing ingredients and low moisture or low humidity conditions. Pharmaceutical compositions and dosage forms that comprise lactose and at least one active ingredient that comprises a primary or secondary amine are preferably anhydrous if substantial contact with moisture and/or humidity during manufacturing, packaging, and/or storage is expected.

An anhydrous pharmaceutical composition should be prepared and stored such that its anhydrous nature is maintained. Accordingly, anhydrous compositions are preferably packaged using materials known to prevent exposure to water such that they can be included in suitable formulary kits. Examples of suitable packaging include, but are not limited to, hermetically sealed foils, plastics, unit dose containers (e.g., vials), blister packs, and strip packs.

The invention further encompasses pharmaceutical compositions and dosage forms that comprise one or more compounds that reduce the rate by which an active ingredient will decompose. Such compounds, which are referred to herein as "stabilizers," include, but are not limited to, antioxidants such as ascorbic acid, pH buffers, or salt buffers.

Like the amounts and types of excipients, the amounts and specific types of active ingredients in a dosage form may differ depending on factors such as, but not limited to, the route by which it is to be administered to patients. However, typical dosage forms of the invention comprise an immunomodulatory compound of the invention or a pharmaceutically

acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof in an amount of from about 0.10 to about 150 mg. Typical dosage forms comprise an immunomodulatory compound of the invention or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof in an amount of about 0.1, 1, 2, 5, 7.5, 10, 12.5, 15, 17.5, 20, 25, 50, 100, 150 or 200 mg. In a particular embodiment, a preferred dosage form comprises 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™) in an amount of about 1, 2, 5, 10, 25 or 50 mg. In a specific embodiment, a preferred dosage form comprises 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) in an amount of about 5, 10, 25 or 50 mg. Typical dosage forms comprise the second active ingredient in an amount of 1 to about 1000 mg, from about 5 to about 500 mg, from about 10 to about 350 mg, or from about 50 to about 200 mg. Of course, the specific amount of the anti-cancer drug will depend on the specific agent used, the type of cancer being treated or managed, and the amount(s) of an immunomodulatory compound of the invention and any optional additional active agents concurrently administered to the patient.

5.4.1 Oral Dosage Forms

Pharmaceutical compositions of the invention that are suitable for oral administration can be presented as discrete dosage forms, such as, but are not limited to, tablets (e.g., chewable tablets), caplets, capsules, and liquids (e.g., flavored syrups). Such dosage forms contain predetermined amounts of active ingredients, and may be prepared by methods of pharmacy well known to those skilled in the art. See generally, *Remington's Pharmaceutical Sciences*, 18th ed., Mack Publishing, Easton Pa. (1990).

Typical oral dosage forms of the invention are prepared by combining the active ingredients in an intimate admixture with at least one excipient according to conventional pharmaceutical compounding techniques. Excipients can take a wide variety of forms depending on the form of preparation desired for administration. For example, excipients suitable for use in oral liquid or aerosol dosage forms include, but are not limited to, water, glycols, oils, alcohols, flavoring agents, preservatives, and coloring agents. Examples of excipients suitable for use in solid oral dosage forms (e.g., powders, tablets, capsules, and caplets) include, but are not limited to, starches, sugars, micro-crystalline cellulose, diluents, granulating agents, lubricants, binders, and disintegrating agents.

Because of their ease of administration, tablets and capsules represent the most advantageous oral dosage unit forms, in which case solid excipients are employed. If desired, tablets can be coated by standard aqueous or nonaqueous techniques. Such dosage forms can be prepared by any of the methods of pharmacy. In general, pharmaceutical compositions and dosage forms are prepared by uniformly and intimately admixing the active ingredients with liquid carriers, finely divided solid carriers, or both, and then shaping the product into the desired presentation if necessary.

For example, a tablet can be prepared by compression or molding. Compressed tablets can be prepared by compressing in a suitable machine the active ingredients in a free-flowing form such as powder or granules, optionally mixed with an excipient. Molded tablets can be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

Examples of excipients that can be used in oral dosage forms of the invention include, but are not limited to, binders, fillers, disintegrants, and lubricants. Binders suitable for use in pharmaceutical compositions and dosage forms include, but are not limited to, corn starch, potato starch, or other

starches, gelatin, natural and synthetic gums such as acacia, sodium alginate, alginic acid, other alginates, powdered tragacanth, guar gum, cellulose and its derivatives (e.g., ethyl cellulose, cellulose acetate, carboxymethyl cellulose calcium, sodium carboxymethyl cellulose), polyvinyl pyrrolidone, methyl cellulose, pre-gelatinized starch, hydroxypropyl methyl cellulose, (e.g., Nos. 2208, 2906, 2910), microcrystalline cellulose, and mixtures thereof.

Suitable forms of microcrystalline cellulose include, but are not limited to, the materials sold as AVICEL-PH-101, AVICEL-PH-103 AVICEL RC-581, AVICEL-PH-105 (available from FMC Corporation, American Viscose Division, Avicel Sales, Marcus Hook, PA), and mixtures thereof. A specific binder is a mixture of microcrystalline cellulose and sodium carboxymethyl cellulose sold as AVICEL RC-581. Suitable anhydrous or low moisture excipients or additives include AVICEL-PH-103™ and Starch 1500 LM.

Examples of fillers suitable for use in the pharmaceutical compositions and dosage forms disclosed herein include, but are not limited to, talc, calcium carbonate (e.g., granules or powder), microcrystalline cellulose, powdered cellulose, dextrates, kaolin, mannitol, silicic acid, sorbitol, starch, pre-gelatinized starch, and mixtures thereof. The binder or filler in pharmaceutical compositions of the invention is typically present in from about 50 to about 99 weight percent of the pharmaceutical composition or dosage form.

Disintegrants are used in the compositions of the invention to provide tablets that disintegrate when exposed to an aqueous environment. Tablets that contain too much disintegrant may disintegrate in storage, while those that contain too little may not disintegrate at a desired rate or under the desired conditions. Thus, a sufficient amount of disintegrant that is neither too much nor too little to detrimentally alter the release of the active ingredients should be used to form solid oral dosage forms of the invention. The amount of disintegrant used varies based upon the type of formulation, and is readily discernible to those of ordinary skill in the art. Typical pharmaceutical compositions comprise from about 0.5 to about 15 weight percent of disintegrant, preferably from about 1 to about 5 weight percent of disintegrant.

Disintegrants that can be used in pharmaceutical compositions and dosage forms of the invention include, but are not limited to, agar-agar, alginic acid, calcium carbonate, microcrystalline cellulose, croscarmellose sodium, crospovidone, polacrilin potassium, sodium starch glycolate, potato or tapioca starch, other starches, pre-gelatinized starch, other starches, clays, other alginates, other celluloses, gums, and mixtures thereof.

Lubricants that can be used in pharmaceutical compositions and dosage forms of the invention include, but are not limited to, calcium stearate, magnesium stearate, mineral oil, light mineral oil, glycerin, sorbitol, mannitol, polyethylene glycol, other glycols, stearic acid, sodium lauryl sulfate, talc, hydrogenated vegetable oil (e.g., peanut oil, cottonseed oil, sunflower oil, sesame oil, olive oil, corn oil, and soybean oil), zinc stearate, ethyl oleate, ethyl laureate, agar, and mixtures thereof. Additional lubricants include, for example, a syloid silica gel (AEROSIL200, manufactured by W.R. Grace Co. of Baltimore, Md.), a coagulated aerosol of synthetic silica (marketed by Degussa Co. of Plano, Tex.), CAB-O-SIL (a pyrogenic silicon dioxide product sold by Cabot Co. of Boston, Mass.), and mixtures thereof. If used at all, lubricants are typically used in an amount of less than about 1 weight percent of the pharmaceutical compositions or dosage forms into which they are incorporated.

A preferred solid oral dosage form of the invention comprises an immunomodulatory compound of the invention,

US 7,968,569 B2

29

anhydrous lactose, microcrystalline cellulose, polyvinylpyrrolidone, stearic acid, colloidal anhydrous silica, and gelatin.

5.4.2 Delayed Release Dosage Forms

Active ingredients of the invention can be administered by controlled release means or by delivery devices that are well known to those of ordinary skill in the art. Examples include, but are not limited to, those described in U.S. Pat. Nos. 3,845,770; 3,916,899; 3,536,809; 3,598,123; and 4,008,719, 5,674,533, 5,059,595, 5,591,767, 5,120,548, 5,073,543, 5,639,476, 5,354,556, and 5,733,566, each of which is incorporated herein by reference. Such dosage forms can be used to provide slow or controlled-release of one or more active ingredients using, for example, hydropropylmethyl cellulose, other polymer matrices, gels, permeable membranes, osmotic systems, multilayer coatings, microparticles, liposomes, microspheres, or a combination thereof to provide the desired release profile in varying proportions. Suitable controlled-release formulations known to those of ordinary skill in the art, including those described herein, can be readily selected for use with the active ingredients of the invention. The invention thus encompasses single unit dosage forms suitable for oral administration such as, but not limited to, tablets, capsules, gelcaps, and caplets that are adapted for controlled-release.

All controlled-release pharmaceutical products have a common goal of improving drug therapy over that achieved by their non-controlled counterparts. Ideally, the use of an optimally designed controlled-release preparation in medical treatment is characterized by a minimum of drug substance being employed to cure or control the condition in a minimum amount of time. Advantages of controlled-release formulations include extended activity of the drug, reduced dosage frequency, and increased patient compliance. In addition, controlled-release formulations can be used to affect the time of onset of action or other characteristics, such as blood levels of the drug, and can thus affect the occurrence of side (e.g., adverse) effects.

Most controlled-release formulations are designed to initially release an amount of drug (active ingredient) that promptly produces the desired therapeutic effect, and gradually and continually release of other amounts of drug to maintain this level of therapeutic or prophylactic effect over an extended period of time. In order to maintain this constant level of drug in the body, the drug must be released from the dosage form at a rate that will replace the amount of drug being metabolized and excreted from the body. Controlled-release of an active ingredient can be stimulated by various conditions including, but not limited to, pH, temperature, enzymes, water, or other physiological conditions or compounds.

5.4.3 Parenteral Dosage Forms

Parenteral dosage forms can be administered to patients by various routes including, but not limited to, subcutaneous, intravenous (including bolus injection), intramuscular, and intraarterial. Because their administration typically bypasses patients' natural defenses against contaminants, parenteral dosage forms are preferably sterile or capable of being sterilized prior to administration to a patient. Examples of parenteral dosage forms include, but are not limited to, solutions ready for injection, dry products ready to be dissolved or suspended in a pharmaceutically acceptable vehicle for injection, suspensions ready for injection, and emulsions.

Suitable vehicles that can be used to provide parenteral dosage forms of the invention are well known to those skilled in the art. Examples include, but are not limited to: Water for Injection USP; aqueous vehicles such as, but not limited to, Sodium Chloride Injection, Ringer's Injection, Dextrose

30

Injection, Dextrose and Sodium Chloride Injection, and Lactated Ringer's Injection; water-miscible vehicles such as, but not limited to, ethyl alcohol, polyethylene glycol, and polypropylene glycol; and non-aqueous vehicles such as, but not limited to, corn oil, cottonseed oil, peanut oil, sesame oil, ethyl oleate, isopropyl myristate, and benzyl benzoate.

Compounds that increase the solubility of one or more of the active ingredients disclosed herein can also be incorporated into the parenteral dosage forms of the invention. For example, cyclodextrin and its derivatives can be used to increase the solubility of an immunomodulatory compound of the invention and its derivatives. See, e.g., U.S. Pat. No. 5,134,127, which is incorporated herein by reference.

5.4.4 Topical and Mucosal Dosage Forms

Topical and mucosal dosage forms of the invention include, but are not limited to, sprays, aerosols, solutions, emulsions, suspensions, eye drops or other ophthalmic preparations, or other forms known to one of skill in the art. See, e.g., *Remington's Pharmaceutical Sciences*, 16th and 18th eds., Mack Publishing, Easton Pa. (1980 & 1990); and *Introduction to Pharmaceutical Dosage Forms*, 4th ed., Lea & Febiger, Philadelphia (1985). Dosage forms suitable for treating mucosal tissues within the oral cavity can be formulated as mouthwashes or as oral gels.

Suitable excipients (e.g., carriers and diluents) and other materials that can be used to provide topical and mucosal dosage forms encompassed by this invention are well known to those skilled in the pharmaceutical arts, and depend on the particular tissue to which a given pharmaceutical composition or dosage form will be applied. With that fact in mind, typical excipients include, but are not limited to, water, acetone, ethanol, ethylene glycol, propylene glycol, butane-1,3-diol, isopropyl myristate, isopropyl palmitate, mineral oil, and mixtures thereof to form solutions, emulsions or gels, which are non-toxic and pharmaceutically acceptable. Moisturizers or humectants can also be added to pharmaceutical compositions and dosage forms if desired. Examples of such additional ingredients are well known in the art. See, e.g., *Remington's Pharmaceutical Sciences*, 16th and 18th eds., Mack Publishing, Easton Pa. (1980 & 1990).

The pH of a pharmaceutical composition or dosage form may also be adjusted to improve delivery of one or more active ingredients. Similarly, the polarity of a solvent carrier, its ionic strength, or tonicity can be adjusted to improve delivery. Compounds such as stearates can also be added to pharmaceutical compositions or dosage forms to advantageously alter the hydrophilicity or lipophilicity of one or more active ingredients so as to improve delivery. In this regard, stearates can serve as a lipid vehicle for the formulation, as an emulsifying agent or surfactant, and as a delivery-enhancing or penetration-enhancing agent. Different salts, hydrates or solvates of the active ingredients can be used to further adjust the properties of the resulting composition.

5.4.5 Kits

Typically, active ingredients of the invention are preferably not administered to a patient at the same time or by the same route of administration. This invention therefore encompasses kits which, when used by the medical practitioner, can simplify the administration of appropriate amounts of active ingredients to a patient.

A typical kit of the invention comprises a dosage form of an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt salt, solvate, hydrate, stereoisomer, prodrug, or clathrate thereof. Kits encompassed by this invention can further comprise additional active ingredients such as oblimersen (Genasense®), melphalan, G-CSF, GM-CSF, EPO, topotecan, dacarbazine, irinotecan, taxotere, IFN,

US 7,968,569 B2

31

COX-2 inhibitor, pentoxifylline, ciprofloxacin, dexamethasone, IL2, IL8, IL18, Ara-C, vinorelbine, isotretinoin, 13 cis-retinoic acid, or a pharmacologically active mutant or derivative thereof, or a combination thereof. Examples of the additional active ingredients include, but are not limited to, those disclosed herein (see, e.g., section 5.2).

Kits of the invention can further comprise devices that are used to administer the active ingredients. Examples of such devices include, but are not limited to, syringes, drip bags, patches, and inhalers.

Kits of the invention can further comprise cells or blood for transplantation as well as pharmaceutically acceptable vehicles that can be used to administer one or more active ingredients. For example, if an active ingredient is provided in a solid form that must be reconstituted for parenteral administration, the kit can comprise a sealed container of a suitable vehicle in which the active ingredient can be dissolved to form a particulate-free sterile solution that is suitable for parenteral administration. Examples of pharmaceutically acceptable vehicles include, but are not limited to: Water for Injection USP; aqueous vehicles such as, but not limited to, Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, and Lactated Ringer's Injection; water-miscible vehicles such as, but not limited to, ethyl alcohol, polyethylene glycol, and polypropylene glycol; and non-aqueous vehicles such as, but not limited to, corn oil, cottonseed oil, peanut oil, sesame oil, ethyl oleate, isopropyl myristate, and benzyl benzoate.

6. EXAMPLES

Certain embodiments of the invention are illustrated by the following non-limiting examples.

6.1 Modulation of Cytokine Production

A series of non-clinical pharmacology and toxicology studies have been performed to support the clinical evaluation of an immunomodulatory compound of the invention in human subjects. These studies were performed in accordance with internationally recognized guidelines for study design and in compliance with the requirements of Good Laboratory Practice (GLP), unless otherwise noted.

Inhibition of TNF- α production following LPS-stimulation of human PBMC and human whole blood by 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (ActimidTM), 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione and thalidomide (RevimidTM) was investigated in vitro (Muller et al., *Bioorg. Med. Chem. Lett.* 9:1625-1630, 1999). The IC₅₀'s of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione for inhibiting production of TNF- α following LPS-stimulation of PBMC and human whole blood were ~24 nM (6.55 ng/mL) and ~25 nM (6.83 ng/mL), respectively. In vitro studies suggest a pharmacological activity profile for 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione that is similar to, but at least 200 times more potent than, thalidomide. In vitro studies have also demonstrated that concentrations of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione of 2.73 to 27.3 ng/mL (0.01 to 0.1 μ M) achieved 50% inhibition of the proliferation of MM.1S and Hs Sultan cells.

The IC₅₀'s of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione for inhibiting production of TNF- α following LPS-stimulation of PBMC and human whole blood were ~100 nM (25.9 ng/mL) and ~480 nM (103.6 ng/mL), respectively. Thalidomide, in contrast, had an IC₅₀ of ~194 μ M (50.2 μ g/mL) for inhibiting production of TNF- α following LPS-stimulation of PBMC. In vitro studies suggest a pharmacological activity profile for 3-(4-amino-1-oxo-1,3-

32

dihydro-isoindol-2-yl)-piperidine-2,6-dione that is similar to, but 50 to 2000 times more potent than, thalidomide. It has been shown that the compound is approximately 50-100 times more potent than thalidomide in stimulating the proliferation of T-cells following primary induction by T-cell receptor (TCR) activation. 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione is also approximately 50 to 100 times more potent than thalidomide in augmenting the production of IL-2 and IFN- γ following TCR activation of PBMC (IL-2) or T-cells (IFN- γ). In addition, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione exhibited dose-dependent inhibition of LPS-stimulated production of the pro-inflammatory cytokines TNF- α , IL-1 β , and IL-6 by PBMC while it increased production of the anti-inflammatory cytokine IL-10.

6.2 Inhibition of MM Cell Proliferation

The ability of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (RevimidTM) and thalidomide for comparison to effect the proliferation of MM cell lines has been investigated in an in vitro study. Uptake [³H]-thymidine by different MM cell lines (MM.1S, Hs Sultan, U266 and RPMI-8226) was measured as an indicator of cell proliferation. Cells were incubated in the presence of compounds for 48 hours; [³H]-thymidine was included for the last 8 hours of the incubation period. Addition of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione to MM. 1S and Hs Sultan cells resulted in 50% inhibition of cell proliferation at concentrations of 0.4 μ M and 1 μ M, respectively. In contrast, addition of thalidomide at concentrations up to 100 μ M resulted in only 15% and 20% inhibition of cell proliferation in MM.1S and Hs Sultan cells, respectively. These data are summarized in FIG. 1.

6.3 Toxicology Studies

The effects of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (RevimidTM) on cardiovascular and respiratory function are investigated in anesthetized dogs. Two groups of Beagle dogs (2/sex/group) are used. One group receives three doses of vehicle only and the other receives three ascending doses of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (2, 10, and 20 mg/kg). In all cases, doses of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione or vehicle are successively administered via infusion through the jugular vein separated by intervals of at least 30 minutes.

The cardiovascular and respiratory changes induced by 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione are minimal at all doses when compared to the vehicle control group. The only statistically significant difference between the vehicle and treatment groups is a small increase in arterial blood pressure (from 94 mmHg to 101 mmHg) following administration of the low dose of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. This effect lasts approximately 15 minutes and is not seen at higher doses. Deviations in femoral blood flow, respiratory parameters, and Qtc interval are common to both the control and treated groups and are not considered treatment-related.

6.4 Cycling Therapy in Patients

In a specific embodiment, an immunomodulatory compound of the invention are cyclically administered to patients with cancer. Cycling therapy involves the administration of a first agent for a period of time, followed by a rest for a period of time and repeating this sequential administration. Cycling therapy can reduce the development of resistance to one or more of the therapies, avoid or reduce the side effects of one of the therapies, and/or improves the efficacy of the treatment.

US 7,968,569 B2

33

In a specific embodiment, prophylactic or therapeutic agents are administered in a cycle of about 4 to 6 weeks, about once or twice every day. One cycle can comprise the administration of a therapeutic or prophylactic agent for three to four weeks and at least a week or two weeks of rest. The number of cycles administered is from about one to about 24 cycles, more typically from about two to about 16 cycles, and more typically from about four to about eight cycles.

For example, in a cycle of four weeks, on day 1, the administration of 25 mg/d of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione is started. On day 22, the administration of the compound is stopped for a week of rest. On day 29, the administration of 25 mg/d 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidin-2,6-dione is begun.

6.5 Clinical Studies in Patients

6.5.1 Treatment of Relapsed Multiple Myeloma

4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™) was administered to patients with relapsed/refractory multiple myeloma. The study was conducted in compliance with Good Clinical Practices. Patients were at least 18 years old, had been diagnosed with multiple myeloma (with paraprotein in serum and/or urine), and were considered refractory to treatment after at least two cycles of treatment, or have relapsed after two cycles of treatment.

Patients who have progressive disease, according to the Southwest Oncology Group (SWOG) criteria, on their prior regimen are considered treatment refractory. Relapse following remission is defined as >25% increase in M component from baseline levels; reappearance of the M paraprotein that had previously disappeared; or a definite increase in the size and number of lytic bone lesions recognized on radiographs. Patients may have had prior therapy with thalidomide, provided they were able to tolerate the treatment. A Zubrod performance status of 0 to 2 is required for all patients.

4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione is administered to patients at doses of 1, 2, 5, or 10 mg/day for up to four weeks; at each dose level, three patients are initially enrolled. Dosing occurs at approximately the same time each morning; all doses are administered in the fasted state (no eating for at least two hours prior to dosing and two hours after dosing). 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione doses are administered in an ascending fashion such that patients in the first cohort receive the lowest dose of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (1 mg/day) and escalation to the next higher dose level occurs only following the establishment of safety and tolerability at the current dose. If one out of three patients at any dose level experience dose limiting toxicity (DLT), three

34

additional patients are enrolled at that dose. If none of the three additional patients experience DLT, escalation to the next dose level occurs; dose escalations continue in a similar fashion until the MTD is established or the maximum daily dose (10 mg/day) is attained. However, if one of the three additional patients enrolled experiences DLT, the MTD has been reached. If two or more of the three additional patients enrolled experience DLT, the MTD is judged to have been exceeded and three additional patients are enrolled at the preceding dose level to confirm the MTD. Once the MTD has been identified, four additional patients are enrolled at that dose level so that a total of 10 patients is treated at the MTD.

Blood sampling for analysis of pharmacokinetic parameters is performed on Days 1 and 28 according to the following sampling schedule: pre-dose, 0.25, 0.5, 0.75, 1, 1.5, 2, 2.5, 3, 4, 6, 8, 10, 12, 18, and 24 hours post-dose. An additional blood sample is collected at each weekly visit for the determination of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione levels. Total urine collections are also made with urine pooled according to the following time intervals post-dose: 0 to 4, 4 to 8, 8 to 12, and 12 to 24 hours. Safety assessments are made by monitoring adverse events, vital signs, ECGs, clinical laboratory evaluations (blood chemistry, hematology, lymphocyte phenotyping, and urinalysis), and physical examination at specific times during the study.

Results of interim pharmacokinetic analyses obtained following single- and multiple-dose administration of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione to multiple myeloma patients are presented below in Tables 1 and 2. These data show that 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione was steadily absorbed at all dose levels in relapsed multiple myeloma patients. Maximum plasma concentrations occurred at a median T_{max} of between 2.5 and 2.8 hours post-dose at Day 1 and between 3 and 4 hours post-dose at Week 4. At all doses, plasma concentrations declined in a monophasic manner after reaching C_{max} . The start of the elimination phase occurred between 3 and 10 hours post-dose at Day 1 and Week 4, respectively.

These data also showed that after 4 weeks of dosing, 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione accumulated to a small extent (mean accumulation ratios ~1.02 to 1.52 and ~0.94 to 1.62 for C_{max} and $AUC_{(0-\infty)}$, respectively). There was almost a dose proportional increase in $AUC_{(0-\infty)}$ and C_{max} values with increasing dose. A five-fold higher dose of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione produced a 3.2- and 2.2-fold increase in C_{max} at Day 1 and Week 4, respectively. Similarly, a 5-fold increase in dose resulted in a 3.6- and 2.3-fold increase in $AUC_{(0-\infty)}$, at Day 1 and Week 4, respectively.

TABLE 1

| Pharmacokinetic parameters of Actimid™ in relapsed multiple myeloma patients | | | | |
|--|---------|----------------|----------------|-----------------|
| Parameter | | 1 mg | 2 mg | 5 mg |
| | | (N = 6) | (N = 2) | (N = 3) |
| Day 1 | | | | |
| C_{max} | ng/mL | 15.03 (4.04) | 24.4* (12.1) | 48.56 (14.03) |
| t_{max} | h | 3.3 (2.6) | 2.7* (0.3) | 2.3 (0.3) |
| $AUC_{(0-\infty)}$ | ng.h/mL | 152.90 (36.62) | 279.18 (51.10) | 593.10 (335.23) |
| $AUC_{(0-\tau)}$ | | 134.21 (27.14) | 249.57 (29.26) | 520.94 (267.32) |
| $t_{1/2}$ | h | 7.3 (3.4) | 6.3 (1.4) | 6.5 (2.2) |
| CL/F | mL/min | 114.75 (29.20) | 121.43 (22.22) | 182.31 (117.06) |
| Vz/f | L | 69.55 (44.97) | 65.31 (2.80) | 87.24 (22.61) |

t = 24 hours

N/A = not available

TABLE 2

| Pharmacokinetic parameters of Actimid™ following multiple oral doses (1, 2, and 5 mg/day) in relapsed multiple myeloma patients | | | | |
|---|---------|-----------------|-----------------|-----------------|
| Parameter | | 1 mg | 2 mg | 5 mg |
| | | (N = 5) | (N = 2) | (N = 3) |
| Week 4 | | | | |
| C_{max} | ng/mL | 23.20 (7.48) | 30.05* (15.64) | 58.07 (38.08) |
| t_{max} | h | 3.6 (1.5) | 2.8* (0.3) | 5.0 (2.6) |
| $AUC_{(0-\infty)}$ | ng.h/mL | N/A | N/A | N/A |
| $AUC_{(0-\tau)}$ | | 239.31 (122.59) | 269.36 (186.34) | 597.24 (354.23) |
| $t_{1/2}$ | h | 6.2* (0.6) | 7.7 (2.8) | 7.8 (4.0) |
| CL/F | mL/min | 87.85 (48.48) | 162.68 (112.54) | 207.50 (175.41) |
| Vz/f | L | 41.35* (8.84) | 95.04 (35.39) | 103.95 (27.25) |

τ = 24 hours

N/A = not available

*N = 3 patients

6.5.2 Treatment of Relapsed Multiple Myeloma

Two Phase 1 clinical studies of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) have been conducted to identify the maximum tolerated dose (MTD) in patients with refractory or relapsed multiple myeloma. These studies have also characterized the safety profile of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione when ascending doses of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione were given orally for up to 4 weeks. Patients started 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione treatment at 5 mg/day with subsequent escalation to 10, 25, and 50 mg/day. Patients were enrolled for 28 days at their assigned dose, with the option of extended treatment for those who did not exhibit disease progression or experience dose limiting toxicity (DLT). Patients were evaluated for adverse events at each visit and the severity of these events was graded according to the National Cancer Institute (NCI) Common Toxicity Criteria. Patients were discontinued if they experienced DLT (Grade 3 or greater non-hematological, or Grade 4 hematological toxicity).

In this study, 27 patients were enrolled. All patients had relapsed multiple myeloma and 18 (72%) were refractory to salvage therapy. Among these patients, 15 had undergone prior autologous stem cell transplantation and 16 patients had received prior thalidomide treatment. The median number of prior regimens was 3 (range 2 to 6).

Blood and urine samples were collected for analysis of pharmacokinetic parameters on Days 1 and 28. Blood samples were collected according to the following sampling schedule: pre-dose, 0.25, 0.5, 0.75, 1, 1.5, 2, 2.5, 3, 4, 6, 8, 10, 12, 18, and 24 hours post-dose. In addition, a blood sample was collected at each weekly clinic visit for 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione determination. Total urine was collected and pooled according to the following time intervals post-dose: 0 to 4, 4 to 8, 8 to 12, and 12 to 24 hours. Response to treatment was assessed by M-protein quantification (by immunoelectrophoresis) from serum and a 24-hour urine collection, with creatinine clearance and 24-hour protein calculations undertaken at screening, baseline, Weeks 2 and 4, and monthly thereafter (or upon early termination). Bone marrow aspirations and/or tissue biopsy are also performed at Months 3, 6 and 12 if a patient's paraprotein serum concentration or 24-hour urine protein excretion declined to the next lower level, based on best response criteria. Preliminary results for the 28-day treatment period are summarized below.

Preliminary pharmacokinetic analyses based on these two studies indicated that AUC and C_{max} values increase propor-

tionally with dose following single and multiple doses in multiple myeloma patients (as was seen in healthy volunteers). Further, there was no evidence of accumulation with multiple dosing as single dose $AUC_{(0-\infty)}$ was comparable to multiple dose $AUC_{(0-\tau)}$ following the same dose of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione.

Similar to healthy volunteer studies, double peaks were observed. Exposure in multiple myeloma patients appeared to be slightly higher based on C_{max} and AUC values as compared to healthy male volunteers while clearance in multiple myeloma patients was lower than it was in healthy volunteers, consistent with their poorer renal function (both as a consequence of their age and their disease). Finally, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione half-life in patients was shorter than in healthy volunteers (mean 8 hours, ranging up to 17 hours).

In this study, the first cohort of 3 patients was treated for 28 days at 5 mg/day without any dose limiting toxicity (DLT). The second cohort of 3 patients subsequently commenced therapy at 10 mg/day. Patients in the second 10 mg/day of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione cohort tolerated treatment well.

6.5.3 Treatment of Solid Tumors

Study with 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) was conducted in patients with varying types of solid tumors, including malignant melanoma (13), carcinoma of the pancreas (2), carcinoma-unknown primary (1), renal carcinoma (1), breast carcinoma (1) and NSCLC (2). Patients received 5 mg/day 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione for seven days and are subsequently escalated every seven days to 10 mg/day, 25 mg/day, and 50 mg/day for a total of 4 weeks of treatment. Patients who, experienced clinical benefit were permitted to continue on treatment as Named Patients.

The study initially enrolled 20 patients and was subsequently amended to enroll 16 additional patients (adrenal carcinoma, NSCLC, malignant mesothelioma, breast cancer, malignant melanoma (8), renal cell cancer (4)) at a higher dose. The 16 additional patients were given weekly escalating doses of 25 mg/day, 50 mg/day, 75 mg/day, 100 mg/day, 125 mg/day, and 150 mg/day over a 6-week period with continuing treatment for an additional six weeks.

The study of Phase 1 study was designed to determine a maximum tolerated dose (MTD) of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione in patients with refractory solid tumors and/or lymphoma, as well as to characterize the pharmacokinetic and side effect profiles of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione in this patient population. The study design dictates that

US 7,968,569 B2

37

at least 3 patients must be enrolled at a dose level and have completed 28 days of treatment prior to enrollment of patients at the next higher dose level. Patients in the first cohort began dosing at 5 mg/day of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. Patients will be escalated to 10, 20, 25, and 30 mg/day provided there is no toxicity.

In this study, the MTD is defined as the highest dose level in which fewer than two of six patients treated did not experience Grade 3 or greater non-hematological toxicity or Grade 4 or greater hematological toxicity. If, at any given dose level in either study, one out of three patients experiences toxicity, three additional patients must be treated at that particular dose. If, however, two out of six patients experience DLT, the MTD is judged to have been exceeded. No further dose escalations are to occur and additional patients are to be enrolled at the previous dose level. The dose of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione administered is escalated until the MTD is achieved or the maximum daily dose of is reached.

No DLTs were reported in the initial group of 20 patients enrolled in the study. Thirteen of the original 20 trial patients, along with 2 non-trial patients, continued on treatment as named patients at doses up to 150 mg/day.

6.5.4 Treatment of Gliomas

This study was performed to find toxicity in patients with recurrent, high-grade gliomas. The study is designed such that patients are given increasingly higher doses of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione until a maximum tolerated dose (MTD) is established. The study also seeks to obtain preliminary toxicity information and pharmacokinetic data on 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione, as well as to develop exploratory data concerning surrogate end points of angiogenic activity in vivo using functional neuro-imaging studies, and in vitro assays of serum angiogenic peptides.

Patients enrolled in the first cohort receive 2.5 mg/m²/day for a 4-week cycle. During each 4-week cycle of therapy, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione administered once daily for 3 weeks followed by a week of rest. Patients who complete a treatment cycle may receive another cycle of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione treatment if two criteria are met. First, the patient must have stable disease or have experienced a partial response or complete response, or the patient is benefiting from the therapy with 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione as evidenced by a decrease in tumor-related symptoms such as neurological deficits. Second, the patient must have recovered from toxicity related to 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione which occurred in the prior cycle by Day 42 or sooner (28-day cycle plus limit of 2 weeks to recover) as evidenced by a return to Grade \leq 1 toxicity level. Patients who experience DLT in the previous cycle should have their dose modified. DLT is defined as a non-hematological event Grade \geq 3 toxicity or hematological event of Grade 4 toxicity thought to be related to the study medication. Patients who experience DLT in the first cycle and have no response to therapy are removed from the study.

3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione doses are subsequently escalated to 5, 8, 11, 15, and 20 mg/m²/day to a maximum total daily dose of 40 mg. Patients continue to receive 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione on a 4-week cycle per dose level until one of the off-study criteria are met.

Three patients are enrolled in each cohort. If at least one DLT occurs, three additional patients are added to the cohort at that particular dose level. If two DLTs occur, the MTD,

38

defined as the dose at which fewer than one-third of patients at each dose level experiences DLT has been exceeded and four more patients are treated at the previous dose.

Patients who experience DLT during the first 4-week cycle are removed from the study, except if they have a response to therapy. For patients who have completed their first 4-week cycle of without DLT, but who subsequently experience Grade 3 or 4 hematological and/or nonhematological toxicity, treatment is suspended for a minimum of a week. If the toxicity resolves to $<$ Grade 2 within three weeks, the patient is treated at two dose levels lower than the dose that caused the toxicity (or a 50% reduction if the patient was treated at the first or second dose level). Patients in whom Grade 3 or 4 toxicity does not resolve to $<$ Grade 1 within three weeks, or those who have another Grade 3 toxicity at the reduced dose are removed from the study.

Pharmacokinetic sampling is performed prior the first dose of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Day 1) and 0.5, 1, 2, 4, 6, 8, 24, and 48 hours thereafter. Sampling is also conducted pre-dose on Days 7 and 21 and 0.5, 1, 2, 4, 6, 8, and 24 post-dose on Day 21 to evaluate steady-state 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione levels.

6.5.5 Treatment of Metastatic Melanoma

Patients with metastatic melanoma were started on 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revmid™) at 5 mg/day for seven days. The dose was then increased every seven days to 10 mg/day, 25 mg/day, and 50 mg/day, respectively, for a total of four weeks on therapy. Five of the 13 melanoma patients who were treated under this regimen either showed disease stabilization or a partial response in the first four weeks of treatment. Tumor response was seen in cutaneous and subcutaneous lesions (five patients), lymph nodes (two patients), and liver (one patient). The duration of response was approximately six months. The result suggests that the compound appears is a promising new anti-cancer agent and has both antiangiogenic and immunomodulatory properties.

6.5.6 Treatment of Relapsed or Refractory Multiple Myeloma

Patients with relapsed and refractory Dune-Salmon stage III multiple myeloma, who have either failed at least three previous regimens or presented with poor performance status, neutropenia or thrombocytopenia, are treated with up to four cycles of combination of melphalan (50 mg intravenously), an immunomodulatory compound of the invention (about 1 to 150 mg orally daily), and dexamethasone (40 mg/day orally on days 1 to 4) every four to six weeks. Maintenance treatment consisting of daily an immunomodulatory compound of the invention and monthly dexamethasone are continued until the disease progression. The therapy using an immunomodulatory compound of the invention in combination with melphalan and dexamethasone is highly active and generally tolerated in heavily pretreated multiple myeloma patients whose prognosis is otherwise poor.


The embodiments of the invention described above are intended to be merely exemplary, and those skilled in the art will recognize, or will be able to ascertain using no more than routine experimentation, numerous equivalents of specific compounds, materials, and procedures. All such equivalents are considered to be within the scope of the invention and are encompassed by the appended claims.

What is claimed is:

1. A method of treating multiple myeloma, which comprises cyclically administering to a patient having multiple myeloma about 5 to about 25 mg per day of a compound of the formula:

US 7,968,569 B2

39



or a pharmaceutically acceptable salt thereof, for 21 consecutive days followed by seven in combination with 40 mg per day dexamethasone.

2. The method of claim 1, wherein the multiple myeloma is smoldering myeloma, indolent myeloma, chemotherapy responsive multiple myeloma, refractory myeloma, relapsed myeloma, or relapsed and refractory Dune-Salmon stage III multiple myeloma.

3. The method of claim 1, wherein the compound is a pharmaceutically acceptable salt.

4. The method of claim 1, wherein the multiple myeloma is relapsed, refractory or resistant to previous therapy.

5. The method of claim 1, wherein the compound and dexamethasone are administered orally.

6. The method of claim 5, wherein the compound is administered in the form of a capsule or tablet.

7. The method of claim 1, wherein the compound is administered in an amount of from about 10 to about 25 mg per day.

8. The method of claim 1, wherein the compound is administered in an amount of about 5, 10, 20, or 25 mg per day.

9. The method of claim 7, wherein the compound is administered in an amount of about 25 mg per day.

40


10. The method of claim 1, wherein the compound is administered in an amount of 5 mg per day.

11. The method of claim 7, wherein the compound is administered in an amount of 10 mg per day.

5 12. The method of claim 1, wherein the compound is administered in a capsule of 5 mg, 10 mg, 15 mg or 25 mg.

13. A method of treating multiple myeloma, which comprises administering, on a 28 day cycle, to a patient having multiple myeloma:

10 (a) about 25 mg per day of a compound of the formula:



or a pharmaceutically acceptable salt thereof, for 21 consecutive days followed by seven consecutive days of rest from administration of said compound, and; (b) 40 mg per day of dexamethasone on days 1-4 every 28 days.

14. The method of claim 12, wherein the capsule comprises the compound, lactose anhydrous, microcrystalline cellulose, croscarmellose sodium and magnesium stearate.

15. The method of claim 1, wherein said dexamethasone is administered.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 7,968,569 B2
APPLICATION NO. : 10/438213
DATED : June 28, 2011
INVENTOR(S) : Jerome B. Zeldis

Page 1 of 1


It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

In claim 1, at column 39, line 11, after “followed by seven”, insert -- consecutive days of rest from administration of said compound during a 28 day cycle, --

In claim 1, at column 39, line 12, between “day” and “dexamethasone”, insert -- of --

In claim 15, at column 40, line 30, after “administered”, insert -- orally --

Signed and Sealed this
Ninth Day of August, 2011

A handwritten signature in black ink that reads "David J. Kappos". The signature is written in a cursive style with a large initial "D" and a stylized "K".

David J. Kappos
Director of the United States Patent and Trademark Office

EXHIBIT D



US008530498B1

(12) **United States Patent**
Zeldis

(10) **Patent No.:** **US 8,530,498 B1**
(45) **Date of Patent:** ***Sep. 10, 2013**

(54) **METHODS FOR TREATING MULTIPLE MYELOMA WITH 3-(4-AMINO-1-OXO-1,3-DIHYDROISOINDOL-2-YL)PIPERIDINE-2,6-DIONE**

(71) Applicant: **Celgene Corporation**, Summit, NJ (US)

(72) Inventor: **Jerome B. Zeldis**, Princeton, NJ (US)

(73) Assignee: **Celgene Corporation**, Summit, NJ (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **13/858,708**

(22) Filed: **Apr. 8, 2013**

Related U.S. Application Data

(63) Continuation of application No. 13/488,888, filed on Jun. 5, 2012, which is a continuation of application No. 12/640,702, filed on Dec. 17, 2009, now Pat. No. 8,198,306, which is a continuation of application No. 10/438,213, filed on May 15, 2003, now Pat. No. 7,968,569.

(60) Provisional application No. 60/380,842, filed on May 17, 2002, provisional application No. 60/424,600, filed on Nov. 6, 2002.

(51) **Int. Cl.**
A01N 43/40 (2006.01)
A61K 31/445 (2006.01)

(52) **U.S. Cl.**
USPC **514/320**; 514/323

(58) **Field of Classification Search**
USPC 514/323
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,536,809 A 10/1970 Applezweig
3,598,123 A 8/1971 Zaffaroni et al.
3,845,770 A 11/1974 Theeuwes et al.
3,916,899 A 11/1975 Theeuwes et al.
4,008,719 A 2/1977 Theeuwes et al.
4,810,643 A 3/1989 Souza
4,999,291 A 3/1991 Souza
5,059,595 A 10/1991 Le Grazie
5,073,543 A 12/1991 Marshall et al.
5,120,548 A 6/1992 McClelland et al.
5,134,127 A 7/1992 Stella et al.
5,229,496 A 7/1993 Deeley et al.
5,354,556 A 10/1994 Sparks et al.
5,385,901 A 1/1995 Kaplan et al.
5,391,485 A 2/1995 Deeley et al.
5,393,870 A 2/1995 Deeley et al.
5,528,823 A 6/1996 Rudy, Jr. et al.
5,580,755 A 12/1996 Souza
5,591,767 A 1/1997 Mohr et al.
5,593,990 A 1/1997 D'Amato
5,629,327 A 5/1997 D'Amato
5,635,517 A 6/1997 Muller et al.

5,639,476 A 6/1997 Oshlack et al.
5,674,533 A 10/1997 Santus et al.
5,698,579 A 12/1997 Muller
5,712,291 A 1/1998 D'Amato
5,731,325 A 3/1998 Andrulis, Jr. et al.
5,733,566 A 3/1998 Lewis
5,798,368 A 8/1998 Muller et al.
5,874,448 A 2/1999 Muller et al.
5,877,200 A 3/1999 Muller
5,929,117 A 7/1999 Muller et al.
5,955,476 A 9/1999 Muller et al.
6,020,358 A 2/2000 Muller et al.
6,071,948 A 6/2000 D'Amato
6,114,355 A 9/2000 D'Amato
6,140,346 A 10/2000 Andrulis, Jr. et al.
6,228,879 B1 5/2001 Green et al.
6,235,756 B1 5/2001 D'Amato
6,281,230 B1 8/2001 Muller et al.
6,316,471 B1 11/2001 Muller et al.
6,326,388 B1 12/2001 Man et al.
6,335,349 B1 1/2002 Muller et al.
6,380,239 B1 4/2002 Muller et al.
6,395,754 B1 5/2002 Muller et al.
6,403,613 B1 6/2002 Man et al.
6,420,414 B1 7/2002 D'Amato
6,458,810 B1 10/2002 Muller et al.
6,469,045 B1 10/2002 D'Amato

(Continued)

FOREIGN PATENT DOCUMENTS

WO WO 92/14455 9/1992
WO WO 94/20085 9/1994

(Continued)

OTHER PUBLICATIONS

Carstensen, 1995, *Drug Stability: Principles & Practice*, 2nd. ed., Marcel Dekker, New York, NY pp. 379-380.
Corral et al., 1999, "Immunomodulation by thalidomide and thalidomide analogues," *Ann. Rheum. Dis.* 58(Suppl 1):1107-1113.
Craig et al., 1967, "Potential anticancer agents. III. 2-phthalimidoaldehydes and derivatives," *Potential Anticancer Agents III* 10:1071-1073.
D'Amato et al., 2001, "Mechanism of action of thalidomide and 3-aminothalidomide in multiple myeloma," *Semin. Oncol.* 28:597-601.
D'Amato et al., 1994, "Thalidomide is an Inhibitor of Angiogenesis", *Proc. Natl. Acad. Sci.* 91:4082-4085.

(Continued)

Primary Examiner — Jeffrey S. Lundgren
Assistant Examiner — Chris Simmons
(74) *Attorney, Agent, or Firm* — Jones Day

(57) **ABSTRACT**

Methods of treating, preventing and/or managing cancer as well as and diseases and disorders associated with, or characterized by, undesired angiogenesis are disclosed. Specific methods encompass the administration of an immunomodulatory compound alone or in combination with a second active ingredient. The invention further relates to methods of reducing or avoiding adverse side effects associated with chemotherapy, radiation therapy, hormonal therapy, biological therapy or immunotherapy which comprise the administration of an immunomodulatory compound. Pharmaceutical compositions, single unit dosage forms, and kits suitable for use in methods of the invention are also disclosed.

13 Claims, 1 Drawing Sheet

US 8,530,498 B1

Page 2

(56)

References Cited

U.S. PATENT DOCUMENTS

| | | | |
|--------------|----|---------|------------------|
| 6,476,052 | B1 | 11/2002 | Muller et al. |
| 6,518,298 | B2 | 2/2003 | Green et al. |
| 6,555,554 | B2 | 4/2003 | Muller et al. |
| 6,673,828 | B1 | 1/2004 | Green et al. |
| 7,323,479 | B2 | 1/2008 | Zeldis |
| 7,393,862 | B2 | 7/2008 | Zeldis |
| 7,435,745 | B2 | 10/2008 | D'Amato |
| 7,468,363 | B2 | 12/2008 | Zeldis |
| 7,968,569 | B2 | 6/2011 | Zeldis |
| 8,188,118 | B2 | 5/2012 | Zeldis |
| 8,198,262 | B2 | 6/2012 | Zeldis |
| 2001/0018445 | A1 | 8/2001 | Huang et al. |
| 2001/0056114 | A1 | 12/2001 | D'Amato |
| 2002/0035090 | A1 | 3/2002 | Zeldis et al. |
| 2002/0045643 | A1 | 4/2002 | Muller et al. |
| 2002/0052398 | A1 | 5/2002 | D'Amato |
| 2002/0054899 | A1 | 5/2002 | Zeldis |
| 2002/0061923 | A1 | 5/2002 | D'Amato |
| 2002/0128228 | A1 | 9/2002 | Hwu |
| 2002/0161023 | A1 | 10/2002 | D'Amato |
| 2002/0173658 | A1 | 11/2002 | Muller et al. |
| 2002/0183360 | A1 | 12/2002 | Muller et al. |
| 2003/0013739 | A1 | 1/2003 | Masferrer et al. |
| 2003/0028028 | A1 | 2/2003 | Man et al. |
| 2003/0045552 | A1 | 3/2003 | Robarge et al. |
| 2003/0069428 | A1 | 4/2003 | Muller et al. |
| 2003/0096841 | A1 | 5/2003 | Robarge et al. |
| 2003/0139451 | A1 | 7/2003 | Shah et al. |
| 2003/0144325 | A1 | 7/2003 | Muller et al. |
| 2003/0181428 | A1 | 9/2003 | Green et al. |
| 2003/0187024 | A1 | 10/2003 | D'Amato |
| 2003/0191098 | A1 | 10/2003 | D'Amato |
| 2003/0235909 | A1 | 12/2003 | Hariri et al. |
| 2004/0029832 | A1 | 2/2004 | Zeldis |
| 2004/0077685 | A1 | 4/2004 | Figg et al. |
| 2004/0077686 | A1 | 4/2004 | Dannenberget al. |
| 2004/0087546 | A1 | 5/2004 | Zeldis |
| 2004/0091455 | A1 | 5/2004 | Zeldis |
| 2004/0122052 | A1 | 6/2004 | Muller et al. |
| 2004/0266809 | A1 | 12/2004 | Emanuel et al. |
| 2008/0145368 | A1 | 6/2008 | Zeldis |
| 2008/0292583 | A1 | 11/2008 | Zeldis |
| 2009/0010877 | A1 | 1/2009 | Zeldis |
| 2009/0123416 | A1 | 5/2009 | Zeldis |
| 2010/0093683 | A1 | 4/2010 | Zeldis |
| 2010/0196369 | A1 | 8/2010 | Zeldis |
| 2010/0260719 | A1 | 10/2010 | Zeldis |
| 2012/0035145 | A1 | 2/2012 | Zeldis |
| 2012/0135042 | A1 | 5/2012 | Zeldis |

FOREIGN PATENT DOCUMENTS

| | | |
|----|--------------|---------|
| WO | WO 98/03502 | 1/1998 |
| WO | WO 98/54170 | 12/1998 |
| WO | WO 01/70275 | 9/2001 |
| WO | WO 01/87307 | 11/2001 |
| WO | WO 02/15926 | 2/2002 |
| WO | WO 02/059106 | 8/2002 |
| WO | WO 02/064083 | 8/2002 |
| WO | WO 03/086373 | 10/2003 |

OTHER PUBLICATIONS

De et al., 1976, "Hansch analysis for some antineoplastic glutarimides," *J. Indian Chem. Soc. I.III*: 825-826.

De et al., 1976, "Possible antineoplastic agents: III. Synthesis of 6-alkyl-2-[4'-methoxyphthalimido] and 6-alkyl-3-[3'-4'-dimethoxyphenyl] glutarimides," *J. Indian Chem. Soc. I.III*: 1122-1125.

Dredge et al., 2002, "Novel thalidomide analogues display antiangiogenic activity independently of immunomodulatory effects," *Br. J. Cancer* 87(10):1166-1172.

Folkman et al., 1983, "Angiogenesis inhibition and tumor regression caused by heparin or a heparin fragment in the presence of cortisone," *Science* 221(4612):719-725.

Gershbein, 1991, "The thalidomide analog, EM 12, enhances 1,2-dimethylhydrazine-induction of rat colon adenocarcinomas," *Cancer Letters* 60: 129-133.

Grabstald et al., 1965, "Clinical experiences with thalidomide in patients with cancer," *Clinical Pharmacology and Therapeutics* 6:298-302.

Lentzsch et al., 2003, "Immunomodulatory analogs of thalidomide inhibit growth of Hs Sultan cells and angiogenesis in vivo," *Leukemia* 17(1):41-44.

Lentzsch et al., 2002, "S-3-amino-phthalimido-glutarimide inhibits angiogenesis and growth of B-cell neoplasias in mice", *Cancer Research* 62:2300-2305.

Miyachi et al., 1997, "Novel biological response modifiers: phthalimides with tumor necrosis factor-alpha production-regulating activity," *J. Med. Chem.* 40:2858-2865.

Muller et al., 1999, "Amino-substituted thalidomide analogs: potent inhibitors of TNF-alpha production," *Bioorg. Med. Chem. Lett.* 9(11):1625-1630.

Muller et al., 1998, "Thalidomide analogs and PDE4 inhibition," *Bioorg. Med. Chem. Lett.* 8(19):2669-2674.

Muller et al., 1996, "Structural modifications of thalidomide produce analogs with enhanced tumor necrosis factor inhibitory activity," *J. Med. Chem.* 39(17):3238-3240.

Olson et al., 1965, "Thalidomide (N-phthaloylglutaramide) in the treatment of advanced cancer," *Clinical Pharmacology and Therapeutics* 6(3):292-297.

Penichet et al., 2001, "Antibody-cytokine fusion proteins or the therapy of cancer," *J. Immunol. Methods* 248(1-2):91-101.

Physician's Desk Reference, 2002, 56th ed., pp. 1755-1760.

Raza et al., 2001, "Thalidomide produces transfusion independence in long-standing refractory anemias of patients with myelodysplastic syndromes," *Blood* 98(4):958-965.

Shah et al., 1999, "Synthesis and enantiomeric separation of 2-phthalimidino-glutaric acid analogues: potent inhibitors of tumor metastasis," *J. Med. Chem.* 42:3014-3017.

Shibata et al., 1995, "N-alkylphthalimides: structural requirement of thalidomide action on 12-O-tetradecanoylphorbol-13-acetate-induced tumor necrosis factor alpha production by human leukemia HL-60 cells," *Chem. Pharm. Bull.* 43(1):177-179.

Shimazawa et al., 1999, "Antiangiogenic activity of tumor necrosis factor-alpha production regulators derived from thalidomide," *Biol. Pharm. Bull.* 22(2):224-226.

Rubin et al., "Principles of Cancer Treatment-1", 12 ONCO IV 1, May 2003.

Wilen et al., 1977, *Tetrahedron* 33:2725.

Wilen, 1972, *Tables of Resolving Agents and Optical Resolutions*, E.L. Eliel, ed., Univ. of Notre Dame Press, Notre Dame, IN pp. 268.

Wolff ed., 1995, *Burger's Medicinal Chemistry and Drug Discovery*, 5th ed., pp. 172-178, 949-982.

N. Ake Jonsson, 1972, "Chemical Structure and Teratogenic Properties," *Acta Pharm.*, pp. 521-542.

Alexanian et al., 2004, "VTD (Velcade, thalidomide, dexamethasone) as primary therapy for newly-diagnosed multiple myeloma," *Am. Soc. Hematol. 46th Ann. Meeting Dec. 4-7, 2004*, San Diego, CA Abstract #210.

Anderson, 2000, "Thalidomide: Therapeutic potential in hematologic malignancies," *Seminars in Hematology* 37(I Supp 3): 1-4.

Attal et al., 2004, "Maintenance treatment with thalidomide after autologous transplantation for myeloma: First analysis of a prospective randomized study of the Intergroupe Francophone du Myelome (IFM 99 02)," *Am. Soc. Hematol. 46th Ann. Meeting Dec. 4-7, 2004*, San Diego, CA Abstract #535.

Bernardeschi et al., 2003, *J. Exp. Clin. Cancer Res.* 22(4):129-133.

Corral et al., 1999, "Differential cytokine modulation and T cell activation by two distinct classes of thalidomide analogues that are potent inhibitors of TNF-alpha," *J. Immunol.* 163(1):380-386.

Davies et al., 2001, "Thalidomide and immunomodulatory derivatives augment natural killer cell cytotoxicity in multiple myeloma," *Blood* 98(1):210-216.

Dimopoulos et al., 2004, "Primary treatment with pulsed melphalan, dexamethasone, thalidomide (MDT) for symptomatic patients with multiple myeloma ≥ 75 years of age," *Am. Soc. Hematol. 46th Ann. Meeting Dec. 4-7, 2004*, San Diego, CA Abstract #1482.

US 8,530,498 B1

Page 3

- Eisen et al., 2000, "Continuous low dose Thalidomide: a phase II study in advanced melanoma, renal cell, ovarian and breast cancer," *Br. J. Cancer* 82(4):812-817.
- Fakhouri et al., 2004, "Thalidomide in patients with multiple myeloma and renal failure," *Br. J. Haematol.* 125:90-102.
- Fenk et al., 2005, "Single-agent thalidomide for treatment of first relapse following high-dose chemotherapy in patients with multiple myeloma," *Leukemia* 19(1):156-159.
- Gupta et al., 2001, "Adherence of multiple myeloma cells to bone marrow stromal cells upregulates vascular endothelial growth factor secretion: therapeutic applications," *Leukemia* 15(12):1950-1961.
- Haslett et al., 2003, "Thalidomide and a thalidomide analogue drug costimulate virus-specific CD8+ T Cells in vitro," *J. Infect. Dis.* 187(6):946-955.
- Hideshima et al., 2000, "Thalidomide and its analogs overcome drug resistance of human multiple myeloma cells to conventional therapy," *Blood* 96(9):2943-2950.
- Offidani et al., 2003, "Thalidomide plus oral melphalan for advanced multiple myeloma: a phase II study," *Haematologica*, Dec. 2003;88(12):1432-1433.
- Palumbo et al., 2004, "A prospective randomized trial of oral melphalan prednisone, thalidomide (MPT) vs. oral melphalan, prednisone (MP): An interim analysis," *Am. Soc. Hematol. 46th Ann. Meeting* Dec. 4-7, 2004, San Diego, CA Abstract #207.
- Raje et al., 1999, "Thalidomide—a revival story," *N. Engl. J. Med.* 341(21):1606-1609.
- Rajkumar et al., 2004, "Thalidomide plus dexamethasone versus dexamethasone alone in newly diagnosed multiple myeloma (E1A00): Results of a phase III trial coordinated by the Eastern Cooperative Oncology Group," *Am. Soc. Hematol. 46th Ann. Meeting* Dec. 4-7, 2004, San Diego, CA Abstract #205.
- Rajkumar et al., 2000, "Prognostic value of bone marrow angiogenesis in multiple myeloma," *Clin. Cancer Res.* 6(8):3111-3116.
- Ribatti et al., 1999, "Bone marrow angiogenesis and mast cell density increase simultaneously with progression of human multiple myeloma," *Br. J. Cancer* 79(3-4):451-455.
- Singhal et al., 1999, "Antitumor activity of thalidomide in refractory multiple myeloma," *N. Engl. J. Med.* 341(21):1565-1571.
- Steins et al., 2002, "Efficacy and safety of thalidomide in patients with acute myeloid leukemia," *Blood* 99(3):834-839.
- Vacca et al., 1999, "Bone marrow neovascularization, plasma cell angiogenic potential, and matrix metalloproteinase-2 secretion parallel progression of human multiple myeloma," *Blood* 93(9):3064-3073.
- Wohrer et al., 2004, "Effective treatment of primary plasma cell leukemia with thalidomide and dexamethasone—a case report," *Hematol. J.* 5(4):361-363.
- Bach, 1963, "Thalidomide in Cancer Chemotherapy," *The Lancet*, No. 1271, p. 71.
- Bach, 1963, "Studies on the Possible Anti-Neoplastic Effect of Thalidomide," *Acta Pathologica Et Microbiologica Scandinavica* 59:491-499.
- Chaudhry, 1966, *Cancer Research*, "Effect of Prednisolone and Thalidomide on Induced Submandibular Gland Tumors in Hamster," 26(part 1):1884-86.
- DiPaolo, 1963, "Effect of Thalidomide on a Variety of Transplantable Tumors," *Cancer Chemotherapy Reports* No. 29, p. 99-102.
- DiPaolo, 1963, "In vitro Test Systems for Cancer Chemotherapy, II. Correlation of in vitro Inhibition of Dehydrogenase and Growth with in vivo Inhibition of Ehrlich Asoites Tumor," *Proceedings of the Society for Experimental Biology & Medicine*, 114:384-387.
- DiPaolo, 1964, "Thalidomide: Effects on Ehrlich Ascites Tumor Cells in vitro" *Science* 144:1583.
- Mauad, 1963, "Clinical Improvements Obtained in Advanced Cancer Patients with Treatment with Thalidomide Associated with Hormones," *Anais Paulistas de Medicina e Cirurgia* 86:13-40.
- Roe and Mitchley, 1963, "Thalidomide and Neoplasia" *Nature* 200:1016-1017.
- Liu et al., "Phase I study of CC-5013 (Revimid), a thalidomide derivative, in patients with refractory metastatic cancer," *American Society of Clinical Oncology*, Abstract #927, 2003.
- Zangari et al., "Results of phase I study of CC-5013 for the treatment of multiple myeloma (MM) patients who relapse after high dose chemotherapy (HDCT)," *American Society of Hematology*, Abstract #3226, 2001.
- Zeldis et al., "Update on the evolution of the 1MiD™," *International Society for Biological Therapy of Cancer, Oral Abstract*, 2003.
- Anderson, "Moving disease biology from the laboratory to the clinic," *Seminars in Oncology*, 2002 29:17-20.
- Barlogie et al., "Total Therapy II (TT11) for newly diagnosed multiple myeloma (MM): preliminary data on feasibility and efficacy in the first 231 enrolled patients; comparison with predecessor total therapy I (TT1) (N=231)," *Blood*, Abstract # 2857, Dec. 7-11, 2001, *American Society of Hematology*.
- Barlogie et al., "High-dose therapy immunomodulatory drugs in multiple myeloma," *Seminars in Oncology*, 2002, 29 (6):26-33.
- Barlogie et al., "Introduction: Thalidomide and the 1MiDs in multiple myeloma," *Seminars in Hematology*, 2003, 40 (4):1-2.
- Barlogie, "Thalidomide and CC-5013 in Multiple Myeloma: The University of Arkansas experience," *Seminars in Hematology*, 2003, 40 (4):33-38.
- Bartlett et al., "The evolution of thalidomide and its 1MiD derivatives as anticancer agents," *Nature Reviews Cancer*, 2004, 4 (4):1-9.
- Bartlett et al., "Phase I study to determine the safety, tolerability and immunostimulatory activity of thalidomide analogue CC-5013 in patients with metastatic malignant melanoma and other advanced cancers," *British Journal of Cancer*, 2004, 90:955-961.
- Battegay, "Angiogenesis: mechanistic insights, neovascular diseases, and therapeutic prospects," *J. Mol. Med.*, 1995, 73:333-346.
- Baz et al., "Doxil (D), vincristine (V), reduced frequency dexamethasone (d) and revlimid (R) (DVD-R) results in a high response rate in patients with refractory multiple myeloma (RMM)," *Blood*, Abstract #2559, *American Society of Hematology*, Dec. 10-13, 2005.
- Brennen et al., "Thalidomide and analogues: current proposed mechanisms and therapeutic usage," *Clinical Prostate Cancer*, 2004, 3 (1):54-61.
- Celgene Corporation, "Celgene advances immunomodulatory drug (1MiD™) clinical program," Press Release, Feb. 2000.
- Celgene Corporation, "Initial Phase I solid tumor data on Celgene's lead 1MiD™, Revimid™," Press Release, Jun. 2001.
- Celgene Corporation, "Celgene Corporation Receives orphan drug designation for Revimid™ for multiple myeloma," Press Release, Oct. 2001.
- Celgene Corporation, "Celgene Corporation announces third quarter results. Thalomid® (thalidomide) sales increase 24%. Prescriptions up 50%. Enhanced S.T.E.P.S.® launched. Pilot d-MPH data presented," Press Release, Oct. 2001.
- Celgene Corporation, "Celgene expands clinical development program for Revimid™. Five additional trials of Revimid initiated in hematological and solid tumor cancers," Press Release, Jun. 2002.
- Celgene Corporation, "Celgene Corporation announces third quarter results. Thalomid® (thalidomide) revenue increases 41% to \$30.5 million. Pivotal programs for Thalomid and Revimid™ finalized. Peer-reviewed publications of Thalomid and Revimid data. First JNK inhibitor advanced to Phase I clinical trial," Press Release, Oct. 2002.
- Celgene Corporation, "Blood reports Revimid™ has anti-tumor activity in patients with relapsed and refractory multiple myeloma," Press Release, Nov. 1, 2002.
- Celgene Corporation, "Celgene provides update on clinical pipeline. Celgene Announces first target indication for ACTIMID™, CC-8490. SelCID™ program to advance based on results from Phase I/II trial of CC-1088. First JNK inhibitor successfully completes phase I trial," Press Release, Jan. 2003.
- Celgene Corporation, "Celgene Corporation announces fourth quarter and full year results for 2002," Press Release, Jan. 2003.
- Celgene Corporation, "Celgene receives fast track status from FDA for Revimid™ in multiple myeloma," Press Release, Feb. 2003.
- Celgene Corporation, "Celgene receives fast track status from FDA for Revimid™ in myelodysplastic syndromes," Press Release, Apr. 2003.
- Celgene Corporation, "New Revimid™ clinical data shows potential as novel approach to treating myelodysplastic syndromes (MDS)," Press Release, May 2003.

US 8,530,498 B1

Page 4

- Celgene Corporation, "Celgene corporation reports strong operating performance in second quarter as total sales increase 100 percent and profits rise," Press Release, Jul. 2003.
- Celgene Corporation, "Celgene corporation reports record operating performance in third quarter as total revenue increases 117% and profits rise," Press Release, Oct. 2003.
- Celgene Corporation, "Celgene corporation advances ACTIMID™ (CC-4047) into phase II trial for prostate cancer," Press Release, Oct. 2003.
- Celgene Corporation, "Additional clinical data presented on Revimid™ in myelodysplastic syndromes at the American Society of Hematology 45th annual meeting," Press Release, Dec. 2003.
- Celgene Corporation, "Celgene corporation reviews 2003 achievements and announces 2004 financial outlook," Press Release, Jan. 2004.
- Celgene Corporation, "Revlimid™ receives orphan drug designation from the European commission for multiple myeloma," Press Release, Feb. 2004.
- Celgene Corporation, "Revlimid™ receives orphan drug designation from the European commission for myelodysplastic syndromes," Press Release, Mar. 2004.
- Celgene Corporation, "Celgene corporation reports record operating performance in first quarter with strong revenue growth and profits," Press Release, Apr. 2004.
- Celgene Corporation, "Celgene announces plans to stop phase III trials in melanoma due to lack of efficacy," Press Release, Apr. 2004.
- Dalgleish, et al., "New thalidomide analogues; anti-cancer, anti-angiogenic and immunostimulatory," *British Journal of Cancer*, 2001, 85 (1)25.
- Dalgleish et al., "Thalidomide analogues CC-5013 and CC-4047 induce T cell activation and IL-12 production in patients with both solid tumours and relapsed and refractory multiple myeloma," *British Journal of Cancer*, 2003, 88(Suppl 1), S25-S54.
- Davies et al., "Thalidomide (Thal) and immunomodulatory derivatives (IMiDs) augment natural killer (NK) cell cytotoxicity in multiple myeloma(MM)," Abstract # 3617, *American Society of Hematology*, Dec. 1-5, 2000.
- Davies et al., "Thalidomide (Thal) and immunomodulatory derivatives (IMiDs) augment natural killer (NK) cell cytotoxicity in multiple myeloma ~MM)," Abstract # P222, *VIIIth International Myeloma Workshop*, May 4-8, 2001.
- Dibbs et al., "Thalidomide and thalidomide analogs suppress TNF α secretion by myocytes," Abstract # 1284, *Circulation*, 1998.
- Dimopoulos et al., "Results of thalidomide and IMiDs in multiple myeloma," Abstract # P12.1.4, *International Multiple Myeloma Workshop*, May 23-27, 2003.
- Dimopoulos et al., "Treatment of plasma cell dyscrasias with thalidomide and its derivatives," *Journal of Clinical Oncology*, Dec. 1, 2003, 21 (23)4444-4454.
- Dimopoulos et al., "Study of lenalidomide plus dexamethasone versus dexamethasone alone in relapsed or refractory multiple myeloma (MM): Results of a phase 3 Study (MM-010)," Abstract # 6, *American Society of Hematology*, Dec. 10-13, 2005.
- Dredge et al., "A costimulatory thalidomide analog enhances the partial anti-tumor immunity of an autologous vaccination in a model of colorectal cancer," Abstract # 491, *American Association for Cancer Research*, Apr. 6-10, 2002.
- Dredge et al., "Adjuvants and the promotion of Th1-type cytokines in tumour immunotherapy," *Cancer Immunol. Immunother.*, 2002, 51:521-531.
- Dredge et al., "Immunological effects of thalidomide and its chemical and functional analogs," *Critical Reviews in Immunology*, 2002, 22 (5&6):425-437.
- Dredge et al., "Protective antitumor immunity induced by a costimulatory thalidomide analog in conjunction with whole tumor cell vaccination is mediated by increased Th1-type immunity," *The Journal of Immunology*, 2002, 168:4914-4919.
- Dredge et al., "Recent developments in antiangiogenic therapy," *Expert Opin. Biol. Ther.*, 2002, 2 (8):953-966.
- Dredge et al., "Angiogenesis inhibitors in cancer therapy," *Current Opinion in Investigational Drugs*, 2003, 4 (6):667-674.
- Dredge et al., "Thalidomide analogs as emerging anti-cancer drugs," *Anti-Cancer Drugs*, 2003, 14:331-335.
- Fickentscher et al., "Stereochemical properties and teratogenic activity of some tetrahydrophthalimides," *Molecular Pharmacology*, 1976, 13:133-141.
- Figg et al., "Inhibition of angiogenesis: treatment options for patients with metastatic prostate cancer," *Investigational New Drugs*, 2002, 20(2):183-194.
- Galustian et al., "Thalidomide-derived immunomodulatory drugs as therapeutic agents," *Expert Opin. Biol. Ther.*, 2004, 4 (12):1-8.
- Glaspy et al., "The potential role of thalidomide and thalidomide analogs in melanoma," *Clinical Advances in Hematology & Oncology*, 2004, 1-7.
- Gupta et al., "Adherence of multiple myeloma cells to bone marrow stromal cells upregulates vascular endothelial growth factor secretion: therapeutic applications," *Leukemia*, 2001, 15:1950-1961.
- Hayashi et al., "Mechanisms whereby immunomodulatory analogs of thalidomide augment autologous NK cell anti-myeloma immunity," Blood, Abstract # 3219, Dec. 6-10, 2002, *American Society of Hematology*.
- He, W., et al., 1993, Abstract of papers, 206th American Chemical Society, Chicago, IL; Med. Chem., paper 216.
- Helm et al., "Comparative teratological investigation of compounds of structurally and pharmacologically related to thalidomide," *Arzneimittel Forschung/Drug Research*, 1981, 31 (1)941-949.
- Hernandez-Illizaliturri et al., "Addition of immunomodulatory drugs CC5013 or CC4047 to rituximab enhances anti-tumor activity in a severe combined immunodeficiency (SCID) mouse lymphoma model," Abstract # 235, *American Society of Hematology*, Dec. 6-9, 2003.
- Hideshima et al., "Thalidomide and its analogs overcome drug resistance of human multiple myeloma cells to conventional therapy," Blood, 2000, 96:2943-2950, *American Society of Hematology*.
- Hideshima et al., "Thalidomide (Thal) and its analogs overcome drug resistance of human multiple myeloma (MM) cells to conventional therapy," Abstract 1313, *American Society of Hematology*, Dec. 1-5, 2000.
- Hunt et al., "Markers of endothelial and haemostatic activation in the use of CC-4047, a structural analogue of thalidamide, in relapsed myeloma," Blood, Abstract # 3216, Dec. 6-10, 2002, *American Society of Hematology*.
- Hussein et al., "Doxil (D), vincristine (V), reduced frequency dexamethasone (d) and Revlimid (DVd-R) a phase I/II trial in advanced relapsed/refractory multiple myeloma (Rmm) patients," Blood, Abstract #208, *American Society of Hematology*, Dec. 4-7, 2004.
- Hwu et al., "Thalidomide and its analogues in the treatment of metastatic melanoma," *Chemotherapy Foundation Symposium*, Abstract #44, 2002.
- Kyle, "Current therapy of multiple myeloma," *Internal Medicine*, 2002, 41 (3)175-180.
- Kyle et al., "Multiple myeloma," *New England Journal of Medicine*, 2004, 351:1860-1873.
- Leblanc et al., "Immunomodulatory drug costimulates T cells via the B7-CD28 pathway," Blood, 2004, 103:1787-1790, *American Society of Hematology*.
- Lentzsch et al., "In vivo activity of thalidomide and immunomodulatory drugs against multiple myeloma," *VIIIth International Myeloma Workshop*, Abstract #P225, May 4-8, 2001.
- Lentzsch et al., "Immunomodulatory derivative of thalidomide (IMiD CC-4047) determine the lineage commitment of hematopoietic progenitors by down regulation of GATA-1 and modulation of cytokine secretion," Abstract # 3073, *American Society of Hematology*, Dec. 6-9, 2003.
- Lentzsch et al., "Immunomodulatory derivative of thalidomide (IMiD CC-4047) down regulates CAAT/enhancer-binding protein β (C/EBP β) in multiple myeloma (MM)," Abstract # 3456, *American Society of Hematology*, Dec. 6-9, 2003.
- Luzzio et al., "Thalidomide analogues: derivatives of an orphan drug with diverse biological activity," *Expert Opin. Ther. Patents*, 2004, 14 (2):215-229.
- Man et al., " α -Fluoro-substituted thalidomide analogues," *Bioorganic & Medicinal Chemistry Letters* 13, 2003, 3415-3417.
- Marriott et al., "Immunotherapeutic and antitumour potential of thalidomide analogues," *Expert Opin. Biol. Ther.*, 2001, 1 (4):1-8.

US 8,530,498 B1

Page 5

- Marriott et al., "New thalidomide analogues; anti-cancer, anti-angiogenic and immunostimulatory," *British Journal of Cancer*, 85:25, Jul. 6, 2001.
- Marriott et al., "Thalidomide and its analogues have distinct and opposing effects on TNF- α and TNFR2 during co-stimulation of both CD4⁺ and CD8⁺ T cells," *Clin. Exp. Immunol.*, 2002, 130:75-84.
- Marriott et al., "A novel subclass of thalidomide analogue with anti-solid tumor activity in which caspase-dependent apoptosis is associated with altered expression of bcl-2 family proteins¹," *Cancer Research*, 2003, 63:593-599.
- Marriott et al., "Thalidomide derived immunomodulatory drugs (IMiDs) as potential therapeutic agents," *Current Drug Targets—Immune, Endocrine & Metabolic Disorders*, 2003, 3:181-186.
- Masellis et al., "Changes in gene expression in bone marrow mesenchymal progenitor cells as a consequence of IMiD therapy in multiple myeloma patients," *Blood*, Abstract # 1548, Dec. 7-11, 2001, *American Society of Hematology*.
- McCarty, "Thalidomide may impede cell migration in primates by down-regulating integrin β -chains: potential therapeutic utility in solid malignancies, proliferative retinopathy, inflammatory disorders, neointimal hyperplasia, and osteoporosis," *Medical Hypotheses*, 1997, 49:123-131.
- Mitsiades et al., "Apoptotic signaling induced by immunomodulatory thalidomide analogs (Imids) in human multiple myeloma cells: therapeutic implications," Abstract, # 3224, Dec. 7-11, 2001, *American Society of Hematology*.
- Mitsiades et al., "Apoptotic signaling induced by immunomodulatory thalidomide analogs in human multiple myeloma cells: therapeutic implications," *Blood*, 2002, 99:4525-4530, *American Society of Hematology*.
- Mitsiades et al., "CC-5013 Celgene," *Current Opinion in Investigational Drugs*, 2004, 5 (6):635-647.
- Moutouh et al., "Novel immunomodulatory drugs (IMiDs®): A potential, new therapy for β -hemoglobinopathies," Abstract # 3740, *American Society of Hematology*, Dec. 4-7, 2004.
- Patten et al., "The early use of the serum free light chain assay in patients with relapsed refractory myeloma receiving treatment with a thalidomide analogue (CC-4047)," Abstract # 1640, *American Society of Hematology*, Dec. 6-9, 2003.
- Payvandi et al., "Effects of a thalidomide analog on binding activity of transcription factors and cell cycle progression of multiple myeloma cell lines," *Blood*, Abstract #2487, Dec. 1-5, 2000, *American Society of Hematology*.
- Payvandi et al., "The thalidomide analogs IMiDs enhance expression of CD69 stimulatory receptor on natural killer cells," Abstract # 1793, *American Association for Cancer Research*, Mar. 24-28, 2001.
- Payvandi et al., "Thalidomide analogs IMiDs inhibit expression of cyclooxygenase-2 in multiple myeloma cell line and LPS stimulated PBMCs," *Blood*, Abstract # 2689, Dec. 7-11, 2001, *American Society of Hematology*.
- Payvandi et al., "Thalidomide and IMiDs inhibit microvessel formation from human arterial rings in the absence of human liver microsomes," *Blood*, Abstract # 5046, Dec. 6-10, 2002, *American Society of Hematology*.
- Payvandi et al., "CC-5013 inhibits the expression of adhesion molecules ICAM-I and CD44 and prevents metastasis of B16 F10 mouse melanoma cells in an animal model," *American Society of Clinical Oncology*, Abstract # 992, 2003.
- Payvandi et al., "Immunomodulatory drugs inhibit expression of cyclooxygenase-2 from TNF- α , IL-1 β , and LPS-stimulated human PBMC in a partially IL-10-dependent manner," *Cellular Immunology*, 2004, 81-88.
- Raje et al., "Combination of the mTOR inhibitor rapamycin and CC-5013 has synergistic activity in multiple myeloma," *Blood*, Dec. 15, 2004, 104 (13):4188-4193.
- Rajkumar et al., "Combination therapy with lenalidomide plus dexamethasone (Rev/Dex) for newly diagnosed myeloma," *Blood*, Dec. 15, 2005, 106 (13):4050-4053.
- Richardson et al., "A Phase I study of oral CC5013, an immunomodulatory thalidomide (Thal) derivative, in patients with relapsed and refractory multiple myeloma (MM)," *Blood*, Abstract #3225, Dec. 7-11, 2001, *American Society of Hematology*.
- Richardson et al., "Immunomodulatory drug CC-5013 overcomes drug resistance and is well tolerated in patients with relapsed multiple myeloma," *Blood*, 2002 100:3063-3067, *American Society of Hematology*.
- Richardson et al., "A multi-center, randomized, phase 2 study to evaluate the efficacy and safety of 2 CDC-5013 dose regimens when used alone or in combination with dexamethasone (Dex) for the treatment of relapsed or refractory multiple myeloma (MM)," *Blood*, Abstract # 825, *American Society of Hematology*, Dec. 6-9, 2003.
- Richardson et al., "Immunomodulatory analogs of thalidomide: an emerging new therapy in myeloma," *Journal of Clinical Oncology*, 2004, 22(16) 3212-3214.
- Richardson et al., "A multicenter, single-arm, open-label study to evaluate the efficacy and safety of single-agent lenalidomide in patients with relapsed and refractory multiple myeloma: preliminary results," *10th International Myeloma Workshop*, Apr. 10-14, 2005.
- Richardson et al., "Novel biological therapies for the treatment of multiple myeloma," *Best Practice & Research Clinical Haematology*, 2005, 18(4):619-634.
- Richardson et al., "A phase 1 trial of lenalidomide (REVLIMID®) with bortezomib (VELCADE®) in relapsed and refractory multiple myeloma," *Blood*, Abstract # 365, *American Society of Hematology*, Dec. 10-13, 2005.
- Rubin et al., "Principles of cancer treatment-1," 2003, 12 ONCO IV 1.
- Schafer et al., "Enhancement of cytokine production and AP-1 transcriptional activity in T cells by thalidomide-related immunomodulatory drugs," *Journal of Pharmacology and Experimental Therapeutics*, 2003, 305(3):1222-1232.
- Schey et al., "A phase 1 study of an immunomodulatory thalidomide analog, CC-4047, in relapsed or refractory multiple myeloma," *Journal of Clinical Oncology*, 2004, 22 (16):1-8.
- Schey et al., "A phase 1 study of an immunomodulatory thalidomide analogue (CC4047) in relapse/refractory multiple myeloma," *International Society for Experimental Hematology*, Abstract #248, 2002.
- Shaughnessy et al., "Global gene expression analysis shows loss of C-MYC and IL-6 receptor gene mRNA after exposure of myeloma to thalidomide and IMiD," Abstract # 2485, *The American Society of Hematology*, Dec. 1-5, 2000.
- Shire et al., "TNF- α inhibitors and rheumatoid arthritis," *Exp. Opin. Ther. Patents*, 1998, 8 (5):531-544.
- Sorbera et al., "CC-5013. Treatment of multiple myeloma. Treatment of Melanoma. Treatment of myelodysplastic syndrome. Angiogenesis inhibitor. TNF- α production inhibitor," *Drugs of the Future*, 2003, 28(5):425-431.
- Streety et al., "Thalidomide analogue CC-4047 is effective in the treatment of patients with relapsed and refractory multiple myeloma (MM) and induces T-cell activation and IL-12 production," Abstract #367, *International Multiple Myeloma Workshop*, May 23-27, 2003.
- Streety et al., "Changes in neutrophil phenotype following the administration of CC-4047 (Actimid) to patients with multiple myeloma," Abstract # 2543, *American Society of Hematology*, Dec. 6-9, 2003.
- Streety et al., "An update of the use and outcomes of the new immunomodulatory agent CC-4047 (Actimid) in patients with relapsed/refractory myeloma," Abstract #829, *American Society of Hematology*, Dec. 6-9, 2003.
- Teo et al., "A phase I, single-blind, placebo-controlled, ascending single oral dose, safety, tolerability and pharmacokinetic study of CDC-501, a novel immunomodulatory-oncologic agent, in healthy male subjects with a comparison of fed and fasted," *Clinical Pharmacology and Therapeutics*, 2002, 71 (2):93.
- Teo et al., "Chiral inversion of the second generation IMiD™ CC-4047 (ACTIMID™) in human plasma and phosphate-buffered saline," *Chirality*, 2003, 15:348-351.
- Thertulien et al., "Hybrid MEL/DT PACE autotransplant regimen for Multiple Myeloma (MM)—safety and efficacy data in pilot study of 15 patients," *Blood*, Abstract # 2869, *American Society of Hematology*, Dec. 7-11, 2001.
- Tohny et al., "A phase I study of oral CC-5013 (lenalidomide, Revlimid™), a thalidomide derivative, in patients with refractory metastatic cancer," *Clinical Prostate Cancer*, 2004, 2:241-243.

US 8,530,498 B1

Page 6

- Tricot et al., "Angiochemotherapy (ACT) for multiple myeloma (MM) with DT-PACE results in a high response rate, but in contrast to tandem transplants with melphalan does not affect durable disease control," *Blood*, Abstract # 3531, *American Society of Hematology*, Dec. 7-11, 2001.
- Tsenova et al., "Use of IMiD3, a thalidomide analog, as an adjunct to therapy for experimental tuberculous meningitis," *Antimicrobial Agents and Chemotherapy*, 2002, 46 (6):1887-1895.
- Weber, "Lenalidomide (CC-5013, Revlimid™) and other ImiDs," Abstract # PL.5.02, *International Multiple Myeloma Workshop*, Apr. 10-14, 2005.
- Weber et al., "A multicenter, randomized, parallel-group, double-blind, placebo-controlled study of lenalidomide plus dexamethasone versus dexamethasone alone in previously treated subjects with multiple myeloma," Abstract # PO.738, *International Multiple Myeloma Workshop*, Apr. 10-14, 2005.
- Ye et al., "Novel IMiD drugs enhance expansion and regulate differentiation of human cord blood CD34+ cells with cytokines," *Blood*, Abstract #4099, *American Society of Hematology*, Dec. 6-10, 2002.
- Zangari et al., "Risk factors for deep vein thrombosis (DVT) in a large group of myeloma patients (Pts) treated with thalidomide (Thal): The Arkansas Experience" *Blood*, Abstract # 681, *American Society of Hematology*, Dec. 7-11, 2001.
- Zangari et al., "Revimid 25 mg (Rev 25) x20 versus 50 mg (REV 50) x10 q 28 days with bridging of 5 mg x10 versus 10 mg x5 as post-transplant salvage therapy for multiple myeloma (MM)," *Blood*, Abstract # 1642, *American Society of Hematology*, Dec. 6-9, 2003.
- Zeldis et al., "Potential new therapeutics for Waldenstrom's macroglobulinemia," *Seminars in Oncology*, 2003, 30 (2):275-281.
- Zhang et al., "CC-5079, a novel microtubule and TNF- α inhibitor with anti-angiogenic and antimetastasis activity," Abstract # B012, *International Conference on Molecular Targets and Cancer Therapeutics*, Nov. 17-21, 2003.
- Anderson, "The Role of Immunomodulatory Drugs in Multiple Myeloma," *Seminars in Hematology*, vol. 40, No. 4, Suppl 4, 2003: pp. 23-32.
- Weber, "Thalidomide and Its Derivatives: New Promise for Multiple Myeloma," *Cancer Control*, vol. 10, No. 5, 375-383, 2003.
- Patt, Yehuda A.; Hassan, Manal M.; Lozano, Richard D.; Ellis, Lee M.; Peterson, J. Andrew; Waugh, Kimberly A.; *Durable Clinical Response of Refractory Hepatocellular Carcinoma to Orally Administered Thalidomide*. American Journal of Clinical Oncology, 2000.
- Richardson, Paul; Hideshima, Teru; Anderson, Kenneth; *Thalidomide: The Revival of a Drug with Therapeutic Promise in the Treatment of Cancer*; Principles & Practice of Oncology, vol. 15, No. 2, 2001.
- Thomas, Melodie; DOSS, Deborah, *Thalidomide Nursing Roundtable Update*, Monograph, Sep. 2002.
- Richardson, Paul; Hideshima, Teru; Anderson, Kenneth; *Thalidomide: Emerging Role in Cancer Medicine*; Annual Review of Medicine, 2002.
- Berenson, J.R.; Bergsagel, P. L.; Munshi, N.; *Initiation and Maintenance of Multiple Myeloma*; *Seminars in Hematology*, vol. 36, No. 1, Supp. 3, Jan. 1999, pp. 9-13.
- Gollob, J.A.; Schinipper, C.P.; Orsini, E.; Murphy, E.; Daley, J.F.; Lazo, S.B.; Frank, D.A.; *Characterization of a Novel Subset of CD8 T Cells That Expands in patients Receiving Interleukin-12*, 02, Am. Soc. For Clin. Investigation, Inc., vol. 102, No. 3, Aug. 1998, pp. 561-575.
- Cavanagh, L.L.; Barnetson, R.S.; Basten, A.; Halliday, G.M.; *Dendritic Epidermal T-Cell Involvement in Induction of CD8+ T-Cell-Mediated Immunity Against an Ultraviolet Radiation-Induced Skin Tumor* Int. J. Cancer: 70, 98-105, 1997.
- Thomas, D.A.; Aguayo, A.; Estey, E.; Albitar, M.; O'Brien, S.; Giles, F.J.; Beran, M.; Cortes, J.; Zeldis, J.; Keating, M.J.; Barlogie, B.; Kantarjian, H.M.; Thalidomide as anti-angiogenesis therapy (rx) in refractory or relapsed leukemia. Abstract #2269, American Society of Hematology, Dec. 3-7, 1999.
- Barlogie, B.; Desikan, R.; Munshi, N.; Siegel, D.; Mehta, J.; Singhal, S.; Anaissie, E.; Single Course D.T. Pace Anti-Angiochemotherapy Effects CR in Plasma Cell Leukemia and Fulminant Multiple Myeloma (MM). Abstract #4180. American Society of Hematology, Dec. 4-9, 1998.
- Hideshima, T.; Chauhan, D.; Shima, Y.; Noopur, R.; Davies, F.E.; Tai, Y.; Treon, S.P.; Lin, B.K.; Schlossman, R.L.; Richardson, P.G.; Gupta, D.; Muller, G.W.; Stirling, D.I.; Anderson, K.C.; Thalidomide (THAL) and its Analogs Overcome Drug Resistance of Human Multiple Myeloma (MM) Cells to Conventional Therapy. Abstract #1313. American Society of Hematology, Dec. 1-5, 2000.
- Payvandi, F.; Wu, L.; Gupta, D.; Hideshima, T.; Haley, M.; Muller, G.; Chen, R.; Anderson, K.C.; Stirling, D.; Effects of a Thalidomide Analog on Binding Activity of Transcription Factors and Cell Cycle Progression of Multiple Myeloma Cell Lines. Abstract #2487. American Society of Hematology, Dec. 1-5, 2000.
- Davies, F.E.; Raje, N.; Hideshima, T.; Lentzsch, S.; Young, G.; Tai, Y.; Lin, B.K.; Podar, K.; Chauhan, D.; Treon, S.P.; Gupta, D.; Mitsiades, C.; Mitsiades, N.; Hayashi, T.; Richardson, P.G.; Schlossman, R.L.; Muller, G.W.; Stirling, D.I.; Anderson, K.C.; Thalidomide (THAL) and Immunomodulatory Derivatives (IMiDs) Augment Natural Killer (NK) Cell Cytotoxicity in Multiple Myeloma (MM). Abstract #3617. American Society of Hematology, Dec. 1-5, 2000.
- Hideshima, T.; Chauhan, D.; Castro, A.; Hayashi, T.; Mitsiades, C.; Mitsiades, N.; Akiyama, M.; Richardson, P.G.; Schlossman, R.L.; Adams, J.; Anderson, K.C.; NF-KB as a Therapeutic Target in Multiple Myeloma (MM), Abstract #1581. American Society of Hematology, Dec. 7-11, 2001.
- Lentsch, S.; Rogers, M.; Leblanc, R.; Birsner, A.; Shah, J.; Anderson K.; D'Amato R.; 3-Amino-Phthalimido-Glutarimide (S-3APG) Inhibits Angiogenesis and Growth in Drug Resistant Multiple Myeloma (MM) in vivo. Abstract #1976, American Society of Hematology, Dec. 7-11, 2001.
- Park, Y.; Kim, S.A.; Kim, C.J.; Chung, J.H.; Mechanism of the Effect of Thalidomide on Human Multiple Myeloma Cells. Abstract #2685. American Society of Clinical Oncology, May 12-17, 2001.
- Payvandi, F.; Wu, E.; Haley M.; Gupta, D.; Zhang, L.; Schafer, P.; Muller, G.W.; Chen, R.; Anderson, K.C.; Stirling, D.; Thalidomide Analogs IMiDs Inhibit Expression of Cyclooxygenase-2 in Multiple Myeloma Cell Line and LPS Stimulated PBMCs. Abstract #2689. American Society of Hematology, Dec. 7-11, 2001.
- Mitsiades, N.; Mitsiades, C.; Poulaki V.; Akiyama, M.; Tai Y.; Lin, B.; Hayashi, T.; Catley, L.; Hideshima, T.; Chauhan, D.; Treon, S.P.; Anderson, K.C.; Apoptotic Signaling Induced by Immunomodulatory Thalidomide Analogs (Imids) in Human Multiple Myeloma Cells: Therapeutic Implications. Abstract #3224. American Society of Hematology, Dec. 7-11, 2001.
- Richardson, P.G.; Schlossman, R.L.; Hideshima, T.; Davies, F.; Leblanc, R.; Catley, L.; Doss, D.; Kelly, K.A.; Mckenney, M.; Mechlowicz, J.; Freeman, A.; Deocampo, R.; Rich. R.; Ryoo, J.; Chauhan, D.; Munshi, N.; Weller, E.; Zeldis, J.; Anderson, K.C.; A Phase I Study of Oral CC5013, an Immunomodulatory Thalidomide (Thal) Derivative, in Patients With Relapsed and Refractory Multiple Myeloma (MM), Abstract #3225. American Society of Hematology, Dec. 7-11, 2001.
- "Celgene drug promises activity in solid tumors," Marketletter, Jun. 18, 2001.
- Meregalli et al., "High-dose dexamethasone as first line therapy of multiple myeloma?," *Recenti Progressi in Medicina*, 1998, 89(1):18-20.
- Official Action in corresponding Canadian Application No. 2,476,983 dated Aug. 21, 2009.
- List, A., "New Approaches to the Treatment of Myelodysplasia," *The Oncologist*, 2002, 7(suppl. 1):39-49.
- Kurzrock, R., "Myelodysplastic syndrome overview," *Seminars in Hematology* (Abstract only), 2002, 39(3)(suppl. 2):18-25 Abstract only.
- Goerner, et al., "Morbidity and mortality of chronic GVHD after hematopoietic stem cell transplantation from HLA-identical siblings for patients with aplastic or refractory anemias," *Biology of Blood and Marrow Transplantation* (Abstract only), 2002, 8(1):47-56.
- Thomas, D., "Pilot studies of Thalidomide in Acute Myelogenous Leukemia, Myelodysplastic Syndromes, and Myeloproliferative Disorders," *Seminars in Hematology*, 2000, 37(1)(suppl. 3):26-34.
- Zorat, F. et al., "The clinical and biological effects of thalidomide in patients with myelodysplastic syndromes," *British Journal of Haematology*, 2001, 115:881-894.

US 8,530,498 B1

Page 7

- Official Action dated Feb. 10, 2009 in JP Application No. 2004-545192. (English translation provided.).
- Teramura, M., Men-ekiyokusei Ryouhou, *Current Therapy*, 2000, 18(5):140-144 (in Japanese).
- Kon-nichi no Chiryou Shishin, 1997 [Pocket Edition], Igaku Shoin, 1997, 513-514 (in Japanese).
- Okamoto, T., Kotsuzuikeisei Shoukougun to Men-eki Ijo, Bessatsu Nihon Rinshou, Syndrome Series for each area, No. 22, Blood Syndromes III, Nihon Rinshou, 213-216 (in Japanese), Oct. 1998.
- Merck Manual, 17th ed. Japanese version, 1999, 951-952.
- Notice of Allowance from U.S. Appl. No. 11/096,155 dated Jan. 12, 2010.
- Rajkumar et al., "Combination therapy with thalidomide plus dexamethasone for newly diagnosed multiple myeloma," *American Society of Hematology*, 43rd Annual Meeting, Dec. 7-11, 2001, Abstract #3525.
- Scheffler et al., "Safety and pharmacokinetics of CDC-501, a novel immunomodulatory-oncologic agent, after single then multiple, oral 100 mg twice daily doses," *American Society Clinical Pharmacology and Therapeutics*, Mar. 24-27, 2002, Abstract #WP111-63.
- Marriott et al., "Thalidomide analogue CDC-501 is safe and well tolerated by patients with end stage cancer and shows evidence of clinical responses and extensive immune activation," *Br. J. Cancer*, 2002, 86(Supp. 1):Abst 6.4.
- Kast, R.E., "Evidence of a mechanism by which etanercept increased TNF-alpha in multiple myeloma: New insights into the biology of TNF-alpha giving new treatment opportunities—the role of burproion," *Leukemia Research*, 2005, 29:1459-1463.
- Tsimberidou, A. et al., "Pilot study of recombinant human soluble tumor necrosis factor (TNF) receptor (p75) fusion protein (TNFR:Fc:Enbrel) in patients with refractory multiple myeloma: increase in plasma TNF α levels during treatment," *Leukemia Research*, 2003, 27:375-380.
- Dimopoulos, et al., "Long-term follow-up on overall survival from the MM-009 and MM-010 phase III trials of lenalidomide plus dexamethasone in patients with relapsed or refractory multiple myeloma," *Leukemia*, 2009, 1-6.
- Hideshima, T., et al., "A review of lenalidomide in combination with dexathasone for the treatment of multiple myeloma," *Therapeutics and Clinical Risk Management*, 2008, 4(1):129-136.
- Wang, M., et al., "Lenalidomide plus dexamethasone is more effective than dexamethasone alone in patients with relapsed or refractory multiple myeloma regardless of prior thalidomide exposure," *Blood*, 2008, 112(12):4445-4451.
- Gandhi, A., et al., "Dexamethasone Synergizes with Lenalidomide to Inhibit Multiple Myeloma Tumor Growth, But Reduces Lenalidomide-Induced Immunomodulation of T and NK Cell Function," *Current Cancer Drug Targets*, 2010, 10(1):1-13.
- Gay, F. et al., "Lenalidomide plus dexamethasone versus thalidomide plus dexamethasone in newly diagnosed multiple myeloma: a comparative analysis of 411 patients," *Blood*, 2010, 115(97):1343-150.
- Richardson, P. et al., "Thalidomide in multiple myeloma," *Biomed Pharmacother*, 2002, 56:115-28.
- Swartz, G. et al., "Pre-clinical evaluation of ENMD-0995: A thalidomide analog with activity against multiple myeloma and solid tumors," *Cell and Tumor Biology*, 2002, 43:181-182, Abstract# 910.
- Mazucco, R., "Angiogenesis and Anti-angiogenesis Therapeutics," *IDrugs*, 2002, 5(4): 320-322.
- Worker, C., "JP Morgan Hambrecht & Quist—20th Annual Healthcare Conference," *IDrugs*, 2002, 5(2):113-116.
- Treston, A. et al., "Pre-Clinical Evaluation of a Thalidomide Analog with Activity Against Multiple Myeloma and Solid Tumors—ENMD-0995 (S(-)-3-(3-amino-phthalimido)-glutarimide)," *Blood*, 2002, 100(11):816a, Abstract #3225.
- Mazucco, R. and Williams, L., "Immunotherapy, chemoprevention and angiogenesis," *IDrugs*, 2002, 5(5):408-411.
- Fernandes, P., "Anti-Cancer Drug Discovery and Development Summit," *IDrugs*, 2002, 5(8):757-764.
- Notification letter dated Aug. 30, 2010 from Natco Pharma Limited to Celgene Corporation re: Notification pursuant to § 505(j)(2)(B) of the Federal Food, Drug and Cosmetic Act.
- Complaint for Patent Infringement filed on Oct. 8, 2010 by Celgene Corporation in the U.S. District Court, District of New Jersey against Natco Pharma Limited.
- Answer to Complaint filed on Nov. 18, 2010 by Natco Pharma Limited in the U.S. District Court, District of New Jersey.
- Grosshans, E. and Illy, G., "Thalidomide Therapy for Inflammatory Dermatoses," *International Journal of Dermatology*, 1984, 23(9):598-602.
- Krenn, M. et al., "Improvements in Solubility and Stability of Thalidomide upon Complexation with Hydropropyl- β -Cyclodextrin," *Journal of Pharmaceutical Sciences*, 1992, 81(7):685-689.
- Schmahl, H. J. et al., "Pharmacokinetics of the Teratogenic and Nonteratogenic Thalidomide Analogs EM 12 and Supidimide in the Rat and Marmoset Monkey", in *Pharmacokinetics in Teratogenesis*, CRC Press, 1987, vol. 1, Ch. 12, pp. 181-192.
- Schumacher, H. et al., "The Teratogenic Activity of a Thalidomide Analogue, EM₁₂, in Rabbits, Rats, and Monkeys," *Teratology*, 1971, 5:233-240.
- Smith, R. et al., "Studies on the Relationship Between the Chemical Structure and Embryotoxic Activity of Thalidomide and Related Compounds," in *A Symposium on Embryopathic Activity of Drugs*, J. & A. Churchill Ltd., 1965, Session 6, pp. 194-209.
- Sheskin, J. and Sagher, F., "Trials with Thalidomide Derivatives in Leprosy Reactions," *Leprosy Review*, 1968, 39(4):203-205.
- Sheskin, J., "Study with Nine Thalidomide Derivatives in the Leprosy Reaction," *Pharmacology and Therapeutics*, 1978, 17:82-84.
- Raje, N. and Anderson, K., "Thalidomide and immunomodulatory drugs as cancer therapy," *Current Opinions in Oncology*, 2002, 14:635-640.
- Kumar, S. et al., "Thalidomide as an anti-cancer agent," *J. Cell. Mod. Med.*, 2002, 6(2):160-174.
- Singhal, S. and Mehta, J., "Thalidomide in Cancer," *BioDrugs*, 2001, 15(3):163-172.
- Notice of Opposition to EP 1 505 973 filed by Synthron B.V. on Nov. 30, 2010.
- Notice of Opposition to EP 1 505 973 filed by Straw:man Limited on Dec. 1, 2010.
- Samson, D. et al., "Infusion of Vincristine and Doxorubicin with Oral Dexamethasone as First-Line Therapy for Multiple Myeloma," *The Lancet*, 1989, 334(8668):882-885.
- Barlogie, B. et al., "Effective Treatment of Advanced Multiple Myeloma Refractory to Alkylating Agents," *N. Engl. J. Med.*, 1984, 310(21):1353-1356.
- Dimopoulos, M. et al., "Thalidomide and dexamethasone combination for refractory multiple myeloma," *Annals of Oncology*, 2001, 12:991-995.
- Zangari, M., et al., "Thrombogenic activity of doxorubicin in myeloma patients receiving thalidomide: implications for therapy," *Blood*, 2002, 100:1168-1171.
- List, A. et al., "High Erythropoietic Remitting Activity of the Immunomodulatory Thalidomide Analog, CC5013, in Patients with Myelodysplastic Syndrome (MDS)," Abstract #353, *Blood*, 2002, 100(11):96a.
- Mufti, G. et al., "Myelodysplastic Syndrome," *American Society of Hematology*, 2003, pp. 176-199.
- Extracts from drug databases: retrieved from <http://www.nextbio.com/b/search/ov/IMiD3%20cpd> on Nov. 26, 2010 and <http://pubchem.ncbi.nlm.nih.gov/summary/summary.cgi?cid=216326> on Nov. 26, 2010.
- Stockdale, 1998, Medicine, Rubenstein and Federman, eds., vol. 3, Ch. 12, Sections IV and X.
- Office Action mailed Jun. 18, 2008, U.S. Appl. No. 11/325,954.
- Gamberi et al., "Overall Safety and Treatment Duration in Lenalidomide (LEN)-, Thalidomide (THAL)-, and Bortezomib (BORT)-Treated Patients (Pts) within the European Post-Approval Safety Study (EU PASS) of Relapsed/Refractory Multiple Myeloma (RRMM)," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #4068.
- Korde et al., "Phase II Clinical and Correlative Study of Carfilzomib, Lenalidomide, and Dexamethasone (CRD) in Newly Diagnosed Mul-

US 8,530,498 B1

Page 8

multiple Myeloma (MM) Patients”, presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #732.

Kumar et al., “A Phase 1/2 Study of Weekly MLN9708, an Investigational Oral Proteasome Inhibitor, in Combination with Lenalidomide and Dexamethasone in Patients with Previously Untreated Multiple Myeloma (MM)”, presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #332.

Palumbo et al., “Pomalidomide Cyclophosphamide and Prednisone (PCP) Treatment for Relapsed/Refractory Multiple Myeloma”, presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #446.

Richardson et al., “A Phase 2 Study of Elotuzumab (Elo) in Combination with Lenalidomide and Low-Dose Dexamethasone (Ld) in Patients (pts) with Relapsed/Refractory Multiple Myeloma (R/R MM): Updated Results”, presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #202.

Sacchi et al., “A Phase I/II Study of Bendamustine, Low-Dose Dexamethasone, and Lenalidomide (BdL) for the Treatment of

Patients with Relapsed Multiple Myeloma”, presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #1851.



Sonneveld et al., “Escalated Dose Bortezomib Once Weekly Combined with Lenalidomide and Dexamethasone (eVRD) Followed by Lenalidomide Maintenance in First Relapse of Multiple Myeloma (MM), the HOVON 86 Phase 2 Trial”, presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #1853.

Suvannasankha et al., “A Phase I/II Trial Combining High-Dose Lenalidomide with Melphalan and Autologous Transplant for Multiple Myeloma: A Report of the Phase 1 Dose-Finding Study”, presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #3146.

Office Action in corresponding CN Application No. 201110256752.0 dated Feb. 8, 2013.

Stirling, D., “Thalidomide: A Novel Template for Anticancer Drugs,” *Seminars in Oncology*, Dec. 2001, 28(6):602-606.

Effects of Revimid™ and Thalidomide on MM Cell Proliferation



US 8,530,498 B1

1

**METHODS FOR TREATING MULTIPLE
MYELOMA WITH
3-(4-AMINO-1-OXO-1,3-DIHYDROISOINDOL-
2-YL)PIPERIDINE-2,6-DIONE**

This application is a continuation of U.S. patent application Ser. No. 13/488,888, filed Jun. 5, 2012, which is a continuation of U.S. patent application Ser. No. 12/640,702, filed Dec. 17, 2009, now U.S. Pat. No. 8,198,306, which is a continuation application of U.S. patent application Ser. No. 10/438,213, filed May 15, 2003, now U.S. Pat. No. 7,968,569, which claims the benefit of U.S. provisional application Nos. 60/380,842, filed May 17, 2002, and 60/424,600, filed Nov. 6, 2002, the entireties of which are incorporated herein by reference.

1. FIELD OF THE INVENTION

This invention relates to methods of treating, preventing and/or managing specific cancers, and other diseases including, but not limited to, those associated with, or characterized by, undesired angiogenesis, by the administration of one or more immunomodulatory compounds alone or in combination with other therapeutics. In particular, the invention encompasses the use of specific combinations, or “cocktails,” of drugs and other therapy, e.g., radiation to treat these specific cancers, including those refractory to conventional therapy. The invention also relates to pharmaceutical compositions and dosing regimens.

2. BACKGROUND OF THE INVENTION

2.1 Pathobiology of Cancer and Other Diseases

Cancer is characterized primarily by an increase in the number of abnormal cells derived from a given normal tissue, invasion of adjacent tissues by these abnormal cells, or lymphatic or blood-borne spread of malignant cells to regional lymph nodes and to distant sites (metastasis). Clinical data and molecular biologic studies indicate that cancer is a multistep process that begins with minor preneoplastic changes, which may under certain conditions progress to neoplasia. The neoplastic lesion may evolve clonally and develop an increasing capacity for invasion, growth, metastasis, and heterogeneity, especially under conditions in which the neoplastic cells escape the host's immune surveillance. Roitt, I., Brostoff, J and Kale, D., *Immunology*, 17.1-17.12 (3rd ed., Mosby, St. Louis, Mo., 1993).

There is an enormous variety of cancers which are described in detail in the medical literature. Examples includes cancer of the lung, colon, rectum, prostate, breast, brain, and intestine. The incidence of cancer continues to climb as the general population ages, as new cancers develop, and as susceptible populations (e.g., people infected with AIDS or excessively exposed to sunlight) grow. A tremendous demand therefore exists for new methods and compositions that can be used to treat patients with cancer.

Many types of cancers are associated with new blood vessel formation, a process known as angiogenesis. Several of the mechanisms involved in tumor-induced angiogenesis have been elucidated. The most direct of these mechanisms is the secretion by the tumor cells of cytokines with angiogenic properties. Examples of these cytokines include acidic and basic fibroblastic growth factor (a,b-FGF), angiogenin, vascular endothelial growth factor (VEGF), and TNF- α . Alternatively, tumor cells can release angiogenic peptides through the production of proteases and the subsequent breakdown of the extracellular matrix where some cytokines are stored

2

(e.g., b-FGF). Angiogenesis can also be induced indirectly through the recruitment of inflammatory cells (particularly macrophages) and their subsequent release of angiogenic cytokines (e.g., TNF- α , bFGF).

A variety of other diseases and disorders are also associated with, or characterized by, undesired angiogenesis. For example, enhanced or unregulated angiogenesis has been implicated in a number of diseases and medical conditions including, but not limited to, ocular neovascular diseases, choroidal neovascular diseases, retina neovascular diseases, rubeosis (neovascularization of the angle), viral diseases, genetic diseases, inflammatory diseases, allergic diseases, and autoimmune diseases. Examples of such diseases and conditions include, but are not limited to: diabetic retinopathy; retinopathy of prematurity; corneal graft rejection; neovascular glaucoma; retrolental fibroplasia; and proliferative vitreoretinopathy.

Accordingly, compounds that can control angiogenesis or inhibit the production of certain cytokines, including TNF- α , may be useful in the treatment and prevention of various diseases and conditions.

2.2 Methods of Treating Cancer

Current cancer therapy may involve surgery, chemotherapy, hormonal therapy and/or radiation treatment to eradicate neoplastic cells in a patient (see, for example, Stockdale, 1998, *Medicine*, vol. 3, Rubenstein and Federman, eds., Chapter 12, Section IV). Recently, cancer therapy could also involve biological therapy or immunotherapy. All of these approaches pose significant drawbacks for the patient. Surgery, for example, may be contraindicated due to the health of a patient or may be unacceptable to the patient. Additionally, surgery may not completely remove neoplastic tissue. Radiation therapy is only effective when the neoplastic tissue exhibits a higher sensitivity to radiation than normal tissue. Radiation therapy can also often elicit serious side effects. Hormonal therapy is rarely given as a single agent. Although hormonal therapy can be effective, it is often used to prevent or delay recurrence of cancer after other treatments have removed the majority of cancer cells. Biological therapies and immunotherapies are limited in number and may produce side effects such as rashes or swellings, flu-like symptoms, including fever, chills and fatigue, digestive tract problems or allergic reactions.

With respect to chemotherapy, there are a variety of chemotherapeutic agents available for treatment of cancer. A majority of cancer chemotherapeutics act by inhibiting DNA synthesis, either directly, or indirectly by inhibiting the biosynthesis of deoxyribonucleotide triphosphate precursors, to prevent DNA replication and concomitant cell division. Gilman et al., *Goodman and Gilman's: The Pharmacological Basis of Therapeutics*, Tenth Ed. (McGraw Hill, New York).

Despite availability of a variety of chemotherapeutic agents, chemotherapy has many drawbacks. Stockdale, *Medicine*, vol. 3, Rubenstein and Federman, eds., ch. 12, sect. 10, 1998. Almost all chemotherapeutic agents are toxic, and chemotherapy causes significant, and often dangerous side effects including severe nausea, bone marrow depression, and immunosuppression. Additionally, even with administration of combinations of chemotherapeutic agents, many tumor cells are resistant or develop resistance to the chemotherapeutic agents. In fact, those cells resistant to the particular chemotherapeutic agents used in the treatment protocol often prove to be resistant to other drugs, even if those agents act by different mechanism from those of the drugs used in the specific treatment. This phenomenon is referred to as pleio-

US 8,530,498 B1

3

tropic drug or multidrug resistance. Because of the drug resistance, many cancers prove refractory to standard chemotherapeutic treatment protocols.

Other diseases or conditions associated with, or characterized by, undesired angiogenesis are also difficult to treat. However, some compounds such as protamine, heparin and steroids have been proposed to be useful in the treatment of certain specific diseases. Taylor et al., *Nature* 297:307 (1982); Folkman et al., *Science* 221:719 (1983); and U.S. Pat. Nos. 5,001,116 and 4,994,443. Thalidomide and certain derivatives of it have also been proposed for the treatment of such diseases and conditions. U.S. Pat. Nos. 5,593,990, 5,629,327, 5,712,291, 6,071,948 and 6,114,355 to D'Amato.

Still, there is a significant need for safe and effective methods of treating, preventing and managing cancer and other diseases and conditions, particularly for diseases that are refractory to standard treatments, such as surgery, radiation therapy, chemotherapy and hormonal therapy, while reducing or avoiding the toxicities and/or side effects associated with the conventional therapies.

2.3 IMIDS™

A number of studies have been conducted with the aim of providing compounds that can safely and effectively be used to treat diseases associated with abnormal production of TNF- α . See, e.g., Marriott, J. B., et al., *Expert Opin. Biol. Ther.* 1(4):1-8 (2001); G. W. Muller, et al., *Journal of Medicinal Chemistry* 39(17): 3238-3240 (1996); and G. W. Muller, et al., *Bioorganic & Medicinal Chemistry Letters* 8: 2669-2674 (1998). Some studies have focused on a group of compounds selected for their capacity to potentially inhibit TNF- α production by LPS stimulated PBMC. L. G. Corral, et al., *Ann. Rheum. Dis.* 58: (Suppl 1) 1107-1113 (1999). These compounds, which are referred to as IMiDs™ (Celgene Corporation) or Immunomodulatory Drugs, show not only potent inhibition of TNF- α but also marked inhibition of LPS induced monocyte IL1 β and IL12 production. LPS induced IL6 is also inhibited by immunomodulatory compounds, albeit partially. These compounds are potent stimulators of LPS induced IL10. Id. Particular examples of IMiDs™ include, but are not limited to, the substituted 2-(2,6-dioxopiperidin-3-yl)phthalimides and substituted 2-(2,6-dioxopiperidin-3-yl)-1-oxoisindoles described in U.S. Pat. Nos. 6,281,230 and 6,316,471, both to G. W. Muller, et al.

3. SUMMARY OF THE INVENTION

This invention encompasses methods of treating and preventing certain types of cancer, including primary and metastatic cancer, as well as cancers that are refractory or resistant to conventional chemotherapy. The methods comprise administering to a patient in need of such treatment or prevention a therapeutically or prophylactically effective amount of an immunomodulatory compound, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof. The invention also encompasses methods of managing certain cancers (e.g., preventing or prolonging their recurrence, or lengthening the time of remission) which comprise administering to a patient in need of such management a prophylactically effective amount of an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

In particular methods of the invention, an immunomodulatory compound is administered in combination with a therapy conventionally used to treat, prevent or manage cancer. Examples of such conventional therapies include, but are

4

not limited to, surgery, chemotherapy, radiation therapy, hormonal therapy, biological therapy and immunotherapy.

This invention also encompasses methods of treating, managing or preventing diseases and disorders other than cancer that are associated with, or characterized by, undesired angiogenesis, which comprise administering to a patient in need of such treatment, management or prevention a therapeutically or prophylactically effective amount of an immunomodulatory compound, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

In other methods of the invention, an immunomodulatory compound is administered in combination with a therapy conventionally used to treat, prevent or manage diseases or disorders associated with, or characterized by, undesired angiogenesis. Examples of such conventional therapies include, but are not limited to, surgery, chemotherapy, radiation therapy, hormonal therapy, biological therapy and immunotherapy.

This invention encompasses pharmaceutical compositions, single unit dosage forms, dosing regimens and kits which comprise an immunomodulatory compound, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, and a second, or additional, active agent. Second active agents include specific combinations, or "cocktails," of drugs.

4. BRIEF DESCRIPTION OF FIGURE

FIG. 1 shows a comparison of the effects of 3-(4-amino-1-oxo-1,3-dihydro-isindol-2-yl)-piperidine-2,6-dione (Revimid™) and thalidomide in inhibiting the proliferation of multiple myeloma (MM) cell lines in an in vitro study. The uptake of [³H]-thymidine by different MM cell lines (MM.1S, Hs Sultan, U266 and RPMI-8226) was measured as an indicator of the cell proliferation.

5. DETAILED DESCRIPTION OF THE INVENTION

A first embodiment of the invention encompasses methods of treating, managing, or preventing cancer which comprises administering to a patient in need of such treatment or prevention a therapeutically or prophylactically effective amount of an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

In particular methods encompassed by this embodiment, the immunomodulatory compound is administered in combination with another drug ("second active agent") or method of treating, managing, or preventing cancer. Second active agents include small molecules and large molecules (e.g., proteins and antibodies), examples of which are provided herein, as well as stem cells. Methods, or therapies, that can be used in combination with the administration of the immunomodulatory compound include, but are not limited to, surgery, blood transfusions, immunotherapy, biological therapy, radiation therapy, and other non-drug based therapies presently used to treat, prevent or manage cancer.

Another embodiment of the invention encompasses methods of treating, managing or preventing diseases and disorders other than cancer that are characterized by undesired angiogenesis. These methods comprise the administration of a therapeutically or prophylactically effective amount of an immunomodulatory compound, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

US 8,530,498 B1

5

Examples of diseases and disorders associated with, or characterized by, undesired angiogenesis include, but are not limited to, inflammatory diseases, autoimmune diseases, viral diseases, genetic diseases, allergic diseases, bacterial diseases, ocular neovascular diseases, choroidal neovascular diseases, retina neovascular diseases, and rubeosis (neovascularization of the angle).

In particular methods encompassed by this embodiment, the immunomodulatory compound is administered in combination with a second active agent or method of treating, managing, or preventing the disease or condition. Second active agents include small molecules and large molecules (e.g., proteins and antibodies), examples of which are provided herein, as well as stem cells. Methods, or therapies, that can be used in combination with the administration of the immunomodulatory compound include, but are not limited to, surgery, blood transfusions, immunotherapy, biological therapy, radiation therapy, and other non-drug based therapies presently used to treat, prevent or manage disease and conditions associated with, or characterized by, undesired angiogenesis.

The invention also encompasses pharmaceutical compositions (e.g., single unit dosage forms) that can be used in methods disclosed herein. Particular pharmaceutical compositions comprise an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, and a second active agent.

5.1 Immunomodulatory Compounds

Compounds used in the invention include immunomodulatory compounds that are racemic, stereomerically enriched or stereomerically pure, and pharmaceutically acceptable salts, solvates, hydrates, stereoisomers, clathrates, and prodrugs thereof. Preferred compounds used in the invention are small organic molecules having a molecular weight less than about 1,000 g/mol, and are not proteins, peptides, oligonucleotides, oligosaccharides or other macromolecules.

As used herein and unless otherwise indicated, the terms "immunomodulatory compounds" and "IMiDs™" (Celgene Corporation) encompasses small organic molecules that markedly inhibit TNF- α , LPS induced monocyte IL1 β and IL12, and partially inhibit IL6 production. Specific immunomodulatory compounds are discussed below.

TNF- α is an inflammatory cytokine produced by macrophages and monocytes during acute inflammation. TNF- α is responsible for a diverse range of signaling events within cells. TNF- α may play a pathological role in cancer. Without being limited by theory, one of the biological effects exerted by the immunomodulatory compounds of the invention is the reduction of synthesis of TNF- α . Immunomodulatory compounds of the invention enhance the degradation of TNF- α mRNA.


Further, without being limited by theory, immunomodulatory compounds used in the invention may also be potent co-stimulators of T cells and increase cell proliferation dramatically in a dose dependent manner. Immunomodulatory compounds of the invention may also have a greater co-stimulatory effect on the CD8+ T cell subset than on the CD4+ T cell subset. In addition, the compounds preferably have anti-inflammatory properties, and efficiently co-stimulate T cells.

Specific examples of immunomodulatory compounds of the invention, include, but are not limited to, cyano and carboxy derivatives of substituted styrenes such as those disclosed in U.S. Pat. No. 5,929,117; 1-oxo-2-(2,6-dioxo-3-fluoropiperidin-3-yl)isoindolines and 1,3-dioxo-2-(2,6-dioxo-3-fluoropiperidine-3-yl)isoindolines such as those

6

described in U.S. Pat. No. 5,874,448; the tetra substituted 2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolines described in U.S. Pat. No. 5,798,368; 1-oxo and 1,3-dioxo-2-(2,6-dioxopiperidin-3-yl) isoindolines (e.g., 4-methyl derivatives of thalidomide and EM-12), including, but not limited to, those disclosed in U.S. Pat. No. 5,635,517; and a class of non-polypeptide cyclic amides disclosed in U.S. Pat. Nos. 5,698,579 and 5,877,200; analogs and derivatives of thalidomide, including hydrolysis products, metabolites, derivatives and precursors of thalidomide, such as those described in U.S. Pat. Nos. 5,593,990, 5,629,327, and 6,071,948 to D'Amato; aminothalidomide, as well as analogs, hydrolysis products, metabolites, derivatives and precursors of aminothalidomide, and substituted 2-(2,6-dioxopiperidin-3-yl)phthalimides and substituted 2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindoles such as those described in U.S. Pat. Nos. 6,281,230 and 6,316,471; isoindole-imide compounds such as those described in U.S. patent application Ser. No. 09/972,487 filed on Oct. 5, 2001, U.S. patent application Ser. No. 10/032,286 filed on Dec. 21, 2001, and International Application No. PCT/US01/50401 (International Publication No. WO 02/059106). The entireties of each of the patents and patent applications identified herein are incorporated herein by reference. Immunomodulatory compounds of the invention do not include thalidomide.

Other specific immunomodulatory compounds of the invention include, but are not limited to, 1-oxo- and 1,3-dioxo-2-(2,6-dioxopiperidin-3-yl)isoindolines substituted with amino in the benzo ring as described in U.S. Pat. No. 5,635,517 which is incorporated herein by reference. These compounds have the structure I:





in which one of X and Y is C=O, the other of X and Y is C=O or CH₂, and R² is hydrogen or lower alkyl, in particular methyl. Specific immunomodulatory compounds include, but are not limited to:

- 1-oxo-2-(2,6-dioxopiperidin-3-yl)-4-aminoisoindoline;
- 1-oxo-2-(2,6-dioxopiperidin-3-yl)-5-aminoisoindoline;
- 1-oxo-2-(2,6-dioxopiperidin-3-yl)-6-aminoisoindoline;
- 1-oxo-2-(2,6-dioxopiperidin-3-yl)-7-aminoisoindoline;
- 1,3-dioxo-2-(2,6-dioxopiperidin-3-yl)-4-aminoisoindoline;
- and
- 1,3-dioxo-2-(2,6-dioxopiperidin-3-yl)-5-aminoisoindoline.

Other specific immunomodulatory compounds of the invention belong to a class of substituted 2-(2,6-dioxopiperidin-3-yl)phthalimides and substituted 2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindoles, such as those described in U.S. Pat. Nos. 6,281,230; 6,316,471; 6,335,349; and 6,476,052, and International Patent Application No. PCT/US97/13375 (International Publication No. WO 98/03502), each of which is incorporated herein by reference. Compounds representative of this class are of the formulas:


US 8,530,498 B1

7



wherein R^1 is hydrogen or methyl. In a separate embodiment, the invention encompasses the use of enantiomerically pure forms (e.g. optically pure (R) or (S) enantiomers) of these compounds.

Still other specific immunomodulatory compounds of the invention belong to a class of isoindole-imides disclosed in U.S. patent application Ser. Nos. 10/032,286 and 09/972,487, and International Application No. PCT/US01/50401 (International Publication No. WO 02/059106), each of which are incorporated herein by reference. Representative compounds are of formula II:



and pharmaceutically acceptable salts, hydrates, solvates, clathrates, enantiomers, diastereomers, racemates, and mixtures of stereoisomers thereof, wherein:

one of X and Y is C=O and the other is CH_2 or C=O;

R^1 is H, (C_1-C_8) alkyl, (C_3-C_7) cycloalkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, benzyl, aryl, (C_0-C_4) alkyl- (C_1-C_6) heterocycloalkyl, (C_0-C_4) alkyl- (C_2-C_5) heteroaryl, $C(O)R^3$, $C(S)R^3$, $C(O)OR^4$, (C_1-C_8) alkyl- $N(R^6)_2$, (C_1-C_8) alkyl- OR^5 , (C_1-C_8) alkyl- $C(O)OR^5$, $C(O)NHR^3$, $C(S)NHR^3$, $C(O)NR^3R^3$, $C(S)NR^3R^3$ or (C_1-C_8) alkyl- $O(CO)R^5$;

8

R^2 is H, F, benzyl, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, or (C_2-C_8) alkynyl;

R^3 and $R^{3'}$ are independently (C_1-C_8) alkyl, (C_3-C_7) cycloalkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, benzyl, aryl, (C_0-C_4) alkyl- (C_1-C_6) heterocycloalkyl, (C_0-C_4) alkyl- (C_2-C_5) heteroaryl, (C_0-C_8) alkyl- $N(R^6)_2$, (C_1-C_8) alkyl- OR^5 , (C_1-C_8) alkyl- $C(O)OR^5$, (C_1-C_8) alkyl- $O(CO)R^5$, or $C(O)OR^5$;

R^4 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, (C_1-C_4) alkyl- OR^5 , benzyl, aryl, (C_0-C_4) alkyl- (C_1-C_6) heterocycloalkyl, or (C_0-C_4) alkyl- (C_2-C_5) heteroaryl;

R^5 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, benzyl, aryl, or (C_2-C_5) heteroaryl;

each occurrence of R^6 is independently H, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, benzyl, aryl, (C_2-C_5) heteroaryl, or (C_0-C_8) alkyl- $C(O)OR^5$ or the R^6 groups can join to form a heterocycloalkyl group;

n is 0 or 1; and

* represents a chiral-carbon center.

In specific compounds of formula II, when n is 0 then R^1 is (C_3-C_7) cycloalkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, benzyl, aryl, (C_0-C_4) alkyl- (C_1-C_6) heterocycloalkyl, (C_0-C_4) alkyl- (C_2-C_5) heteroaryl, $C(O)R^3$, $C(O)OR^4$, (C_1-C_8) alkyl- $N(R^6)_2$, (C_1-C_8) alkyl- OR^5 , (C_1-C_8) alkyl- $C(O)OR^5$, $C(S)NHR^3$, or (C_1-C_8) alkyl- $O(CO)R^5$;


R^2 is H or (C_1-C_8) alkyl; and

R^3 is (C_1-C_8) alkyl, (C_3-C_7) cycloalkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, benzyl, aryl, (C_0-C_4) alkyl- (C_1-C_6) heterocycloalkyl, (C_0-C_4) alkyl- (C_2-C_5) heteroaryl, (C_5-C_8) alkyl- $N(R^6)_2$; (C_0-C_8) alkyl- $NH-C(O)OR^5$; (C_1-C_8) alkyl- OR^5 , (C_1-C_8) alkyl- $C(O)OR^5$, (C_1-C_8) alkyl- $O(CO)R^5$, or $C(O)OR^5$; and the other variables have the same definitions.

In other specific compounds of formula II, R^2 is H or (C_1-C_4) alkyl.


In other specific compounds of formula II, R^1 is (C_1-C_8) alkyl or benzyl.

In other specific compounds of formula II, R^1 is H, (C_1-C_8) alkyl, benzyl, CH_2OCH_3 , $CH_2CH_2OCH_3$, or



In another embodiment of the compounds of formula II, R^1 is

II



wherein Q is O or S, and each occurrence of R^7 is independently H, (C_1-C_8) alkyl, benzyl, CH_2OCH_3 , or $CH_2CH_2OCH_3$.

In other specific compounds of formula II, R^1 is $C(O)R^3$.

In other specific compounds of formula II, R^3 is (C_0-C_4) alkyl- (C_2-C_5) heteroaryl, (C_1-C_8) alkyl, aryl, or (C_0-C_4) alkyl- OR^5 .

In other specific compounds of formula II, heteroaryl is pyridyl, furyl, or thienyl.


US 8,530,498 B1

9

In other specific compounds of formula II, R¹ is C(O)OR⁴.

In other specific compounds of formula II, the H of C(O)NHC(O) can be replaced with (C₁-C₄)alkyl, aryl, or benzyl.

Still other specific immunomodulatory compounds of the invention belong to a class of isoindole-imides disclosed in U.S. patent application Ser. No. 09/781,179, International Publication No. WO 98/54170, and U.S. Pat. No. 6,395,754, each of which are incorporated herein by reference. Representative compounds are of formula III:



and pharmaceutically acceptable salts, hydrates, solvates, clathrates, enantiomers, diastereomers, racemates, and mixtures of stereoisomers thereof, wherein:

one of X and Y is C=O and the other is CH₂ or C=O;

R is H or CH₂OCOR⁵;

(i) each of R¹, R², R³, or R⁴, independently of the others, is halo, alkyl of 1 to 4 carbon atoms, or alkoxy of 1 to 4 carbon atoms or (ii) one of R¹, R², R³, or R⁴ is nitro or —NHR⁵ and the remaining of R¹, R², R³, or R⁴ are hydrogen;

R⁵ is hydrogen or alkyl of 1 to 8 carbons

R⁶ hydrogen, alkyl of 1 to 8 carbon atoms, benzo, chloro, or fluoro;


R⁷ is R⁷—CHR¹⁰—N(R⁸R⁹);

R⁷ is m-phenylene or p-phenylene or —(C_nH_{2n})— in which n has a value of 0 to 4;

each of R⁸ and R⁹ taken independently of the other is hydrogen or alkyl of 1 to 8 carbon atoms, or R⁸ and R⁹ taken together are tetramethylene, pentamethylene, hexamethylene, or —CH₂CH₂[X]₁CH₂CH₂— in which [X]₁ is —O—, —S—, or —NH—;


R¹⁰ is hydrogen, alkyl of to 8 carbon atoms, or phenyl; and * represents a chiral-carbon center.

The most preferred immunomodulatory compounds of the invention are 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione and 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. The compounds can be obtained via standard, synthetic methods (see e.g., U.S. Pat. No. 5,635,517, incorporated herein by reference). The compounds are available from Celgene Corporation, Warren, N.J. 4-(Amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (ACTIMID™) has the following chemical structure:



The compound 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (REVIMID™) has the following chemical structure:

10



Compounds of the invention can either be commercially purchased or prepared according to the methods described in the patents or patent publications disclosed herein. Further, optically pure compounds can be asymmetrically synthesized or resolved using known resolving agents or chiral columns as well as other standard synthetic organic chemistry techniques.

As used herein and unless otherwise indicated, the term “pharmaceutically acceptable salt” encompasses non-toxic acid and base addition salts of the compound to which the term refers. Acceptable non-toxic acid addition salts include those derived from organic and inorganic acids or bases known in the art, which include, for example, hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid, methanesulphonic acid, acetic acid, tartaric acid, lactic acid, succinic acid, citric acid, malic acid, maleic acid, sorbic acid, aconitic acid, salicylic acid, phthalic acid, embolic acid, enanthic acid, and the like.

Compounds that are acidic in nature are capable of forming salts with various pharmaceutically acceptable bases. The bases that can be used to prepare pharmaceutically acceptable base addition salts of such acidic compounds are those that form non-toxic base addition salts, i.e., salts containing pharmacologically acceptable cations such as, but not limited to, alkali metal or alkaline earth metal salts and the calcium, magnesium, sodium or potassium salts in particular. Suitable organic bases include, but are not limited to, N,N-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumaine (N-methylglucamine), lysine, and procaine.

As used herein and unless otherwise indicated, the term “prodrug” means a derivative of a compound that can hydrolyze, oxidize, or otherwise react under biological conditions (in vitro or in vivo) to provide the compound. Examples of prodrugs include, but are not limited to, derivatives of immunomodulatory compounds of the invention that comprise biohydrolyzable moieties such as biohydrolyzable amides, biohydrolyzable esters, biohydrolyzable carbamates, biohydrolyzable carbonates, biohydrolyzable ureides, and biohydrolyzable phosphate analogues. Other examples of prodrugs include derivatives of immunomodulatory compounds of the invention that comprise —NO, —NO₂, —ONO, or —ONO₂ moieties. Prodrugs can typically be prepared using well-known methods, such as those described in 1 *Burger's Medicinal Chemistry and Drug Discovery*, 172-178, 949-982 (Manfred E. Wolff ed., 5th ed. 1995), and *Design of Prodrugs* (H. Bundgaard ed., Elsevier, New York 1985).

As used herein and unless otherwise indicated, the terms “biohydrolyzable amide,” “biohydrolyzable ester,” “biohydrolyzable carbamate,” “biohydrolyzable carbonate,” “biohydrolyzable ureide,” “biohydrolyzable phosphate” mean an amide, ester, carbamate, carbonate, ureide, or phosphate, respectively, of a compound that either: 1) does not interfere with the biological activity of the compound but can confer upon that compound advantageous properties in vivo, such as uptake, duration of action, or onset of action; or 2) is biologi-

US 8,530,498 B1

11

cally inactive but is converted in vivo to the biologically active compound. Examples of biohydrolyzable esters include, but are not limited to, lower alkyl esters, lower acyloxyalkyl esters (such as acetoxyethyl, acetoxyethyl, aminocarbonyloxymethyl, pivaloyloxymethyl, and pivaloyloxyethyl esters), lactonyl esters (such as phthalidyl and thiophthalidyl esters), lower alkoxyacyloxyalkyl esters (such as methoxycarbonyl-oxymethyl, ethoxycarbonyloxyethyl and isopropoxycarbonyloxyethyl esters), alkoxyalkyl esters, choline esters, and acylamino alkyl esters (such as acetamidomethyl esters). Examples of biohydrolyzable amides include, but are not limited to, lower alkyl amides, α -amino acid amides, alkoxyacyl amides, and alkylaminoalkylcarbonyl amides. Examples of biohydrolyzable carbamates include, but are not limited to, lower alkylamines, substituted ethylenediamines, amino acids, hydroxyalkylamines, heterocyclic and heteroaromatic amines, and polyether amines.

Various immunomodulatory compounds of the invention contain one or more chiral centers, and can exist as racemic mixtures of enantiomers or mixtures of diastereomers. This invention encompasses the use of stereomerically pure forms of such compounds, as well as the use of mixtures of those forms. For example, mixtures comprising equal or unequal amounts of the enantiomers of a particular immunomodulatory compounds of the invention may be used in methods and compositions of the invention. These isomers may be asymmetrically synthesized or resolved using standard techniques such as chiral columns or chiral resolving agents. See, e.g., Jacques, J., et al., *Enantiomers, Racemates and Resolutions* (Wiley-Interscience, New York, 1981); Wilen, S. H., et al., *Tetrahedron* 33:2725 (1977); Eliel, E. L., *Stereochemistry of Carbon Compounds* (McGraw-Hill, NY, 1962); and Wilen, S. H., *Tables of Resolving Agents and Optical Resolutions* p. 268 (E. L. Eliel, Ed., Univ. of Notre Dame Press, Notre Dame, Ind., 1972).

As used herein and unless otherwise indicated, the term "stereomerically pure" means a composition that comprises one stereoisomer of a compound and is substantially free of other stereoisomers of that compound. For example, a stereomerically pure composition of a compound having one chiral center will be substantially free of the opposite enantiomer of the compound. A stereomerically pure composition of a compound having two chiral centers will be substantially free of other diastereomers of the compound. A typical stereomerically pure compound comprises greater than about 80% by weight of one stereoisomer of the compound and less than about 20% by weight of other stereoisomers of the compound, more preferably greater than about 90% by weight of one stereoisomer of the compound and less than about 10% by weight of the other stereoisomers of the compound, even more preferably greater than about 95% by weight of one stereoisomer of the compound and less than about 5% by weight of the other stereoisomers of the compound, and most preferably greater than about 97% by weight of one stereoisomer of the compound and less than about 3% by weight of the other stereoisomers of the compound. As used herein and unless otherwise indicated, the term "stereomerically enriched" means a composition that comprises greater than about 60% by weight of one stereoisomer of a compound, preferably greater than about 70% by weight, more preferably greater than about 80% by weight of one stereoisomer of a compound. As used herein and unless otherwise indicated, the term "enantiomerically pure" means a stereomerically pure composition of a compound having one chiral center. Similarly, the term "stereomerically enriched" means a stereomerically enriched composition of a compound having one chiral center.

12

It should be noted that if there is a discrepancy between a depicted structure and a name given that structure, the depicted structure is to be accorded more weight. In addition, if the stereochemistry of a structure or a portion of a structure is not indicated with, for example, bold or dashed lines, the structure or portion of the structure is to be interpreted as encompassing all stereoisomers of it.

5.2 Second Active Agents

Immunomodulatory compounds can be combined with other pharmacologically active compounds ("second active agents") in methods and compositions of the invention. It is believed that certain combinations work synergistically in the treatment of particular types of cancer and certain diseases and conditions associated with, or characterized by, undesired angiogenesis. Immunomodulatory compounds can also work to alleviate adverse effects associated with certain second active agents, and some second active agents can be used to alleviate adverse effects associated with immunomodulatory compounds.

One or more second active ingredients or agents can be used in the methods and compositions of the invention together with an immunomodulatory compound. Second active agents can be large molecules (e.g., proteins) or small molecules (e.g., synthetic inorganic, organometallic, or organic molecules).

Examples of large molecule active agents include, but are not limited to, hematopoietic growth factors, cytokines, and monoclonal and polyclonal antibodies. Typical large molecule active agents are biological molecules, such as naturally occurring or artificially made proteins. Proteins that are particularly useful in this invention include proteins that stimulate the survival and/or proliferation of hematopoietic precursor cells and immunologically active poietic cells in vitro or in vivo. Others stimulate the division and differentiation of committed erythroid progenitors in cells in vitro or in vivo. Particular proteins include, but are not limited to: interleukins, such as IL-2 (including recombinant IL-II ("rIL.2") and canarypox TL-2), IL-10, IL-12, and IL-18; interferons, such as interferon alfa-2a, interferon alfa-2b, interferon alfa-n1, interferon alfa-n3, interferon beta-1 a, and interferon gamma-I b; GM-CSF and GM-CSF; and EPO.

Particular proteins that can be used in the methods and compositions of the invention include, but are not limited to: filgrastim, which is sold in the United States under the trade name Neupogen® (Amgen, Thousand Oaks, Calif.); sargramostim, which is sold in the United States under the trade name Leukine® (Immunex, Seattle, Wash.); and recombinant EPO, which is sold in the United States under the trade name Epogen® (Amgen, Thousand Oaks, Calif.).

Recombinant and mutated forms of GM-CSF can be prepared as described in U.S. Pat. Nos. 5,391,485; 5,393,870; and 5,229,496; all of which are incorporated herein by reference. Recombinant and mutated forms of G-CSF can be prepared as described in U.S. Pat. Nos. 4,810,643; 4,999,291; 5,528,823; and 5,580,755; all of which are incorporated herein by reference.

This invention encompasses the use of native, naturally occurring, and recombinant proteins. The invention further encompasses mutants and derivatives (e.g., modified forms) of naturally occurring proteins that exhibit, in vivo, at least some of the pharmacological activity of the proteins upon which they are based. Examples of mutants include, but are not limited to, proteins that have one or more amino acid residues that differ from the corresponding residues in the naturally occurring forms of the proteins. Also encompassed by the term "mutants" are proteins that lack carbohydrate moieties normally present in their naturally occurring forms

US 8,530,498 B1

13

(e.g., nonglycosylated forms). Examples of derivatives include, but are not limited to, pegylated derivatives and fusion proteins, such as proteins formed by fusing IgG1 or IgG3 to the protein or active portion of the protein of interest. See, e.g., Penichet, M. L. and Morrison, S. L., *J. J. Immunol. Methods* 248:91-101 (2001).

Antibodies that can be used in combination with compounds of the invention include monoclonal and polyclonal antibodies. Examples of antibodies include, but are not limited to, trastuzumab (Herceptin®), rituximab (Rituxan®), bevacizumab (Avastin™), pertuzumab (Omnitarg™), tositumomab (Bexxar®), edrecolomab (Panorex®), and G250. Compounds of the invention can also be combined with, or used in combination with, anti-TNF- α antibodies.

Large molecule active agents may be administered in the form of anti-cancer vaccines. For example, vaccines that secrete, or cause the secretion of, cytokines such as IL-2, G-CSF, and GM-CSF can be used in the methods, pharmaceutical compositions, and kits of the invention. See, e.g., Emens, L. A., et al., *Curr. Opinion Mol. Ther.* 3(1):77-84 (2001).

In one embodiment of the invention, the large molecule active agent reduces, eliminates, or prevents an adverse effect associated with the administration of an immunomodulatory compound. Depending on the particular immunomodulatory compound and the disease or disorder begin treated, adverse effects can include, but are not limited to, drowsiness and somnolence, dizziness and orthostatic hypotension, neutropenia, infections that result from neutropenia, increased HIV-viral load, bradycardia, Stevens-Johnson Syndrome and toxic epidermal necrolysis, and seizures (e.g., grand mal convulsions). A specific adverse effect is neutropenia.

Second active agents that are small molecules can also be used to alleviate adverse effects associated with the administration of an immunomodulatory compound. However, like some large molecules, many are believed to be capable of providing a synergistic effect when administered with (e.g., before, after or simultaneously) an immunomodulatory compound. Examples of small molecule second active agents include, but are not limited to, anti-cancer agents, antibiotics, immunosuppressive agents, and steroids.

Examples of anti-cancer agents include, but are not limited to: acivicin; aclarubicin; acodazole hydrochloride; acronine; adozelesin; aldesleukin; altretamine; ambomycin; ametantrone acetate; amsacrine; anastrozole; anthramycin; asparaginase; asperlin; azacitidine; azetepa; azotomycin; batimastat; benzodepa; bicalutamide; bisantrene hydrochloride; bisnafide dimesylate; bizelesin; bleomycin sulfate; brequinar sodium; bropirimine; busulfan; cactinomycin; calusterone; caracemide; carbetimer; carboplatin; carmustine; carubicin hydrochloride; carzelesin; cedefingol; celecoxib (COX-2 inhibitor); chlorambucil; cirolemycin; cisplatin; cladribine; crisnatol mesylate; cyclophosphamide; cytarabine; dacarbazine; dactinomycin; daunorubicin hydrochloride; decitabine; dexormaplatin; dezaguanine; dezaguanine mesylate; diaziquone; docetaxel; doxorubicin; doxorubicin hydrochloride; droloxifene; droloxifene citrate; dromostanolone propionate; duazomycin; edatrexate; eflornithine hydrochloride; elsamitucin; enloplatin; enpromate; epiropridine; epirubicin hydrochloride; erbulozole; esorubicin hydrochloride; estramustine; estramustine phosphate sodium; etanidazole; etoposide; etoposide phosphate; etoprine; fadrozole hydrochloride; fazarabine; fenretinide; floxuridine; fludarabine phosphate; fluorouracil; fluorocitabine; fosquidone; fostriecin sodium; gemcitabine; gemcitabine hydrochloride; hydroxyurea; idarubicin hydrochloride; ifosfamide; ilmofofosine; iproplatin; irinotecan; irinotecan

14

hydrochloride; lanreotide acetate; letrozole; leuprolide acetate; liarozole hydrochloride; lometrexol sodium; lomustine; losoxantrone hydrochloride; masoprocol; maytansine; mechlorethamine hydrochloride; megestrol acetate; melengestrol acetate; melphalan; menogaril; mercaptopurine; methotrexate; methotrexate sodium; metoprine; meturedepa; mitindomide; mitocarcin; mitocromin; mitogillin; mitomalcin; mitomycin; mitosper; mitotane; mitoxantrone hydrochloride; mycophenolic acid; nocodazole; nogalamycin; ormaplatin; oxisuran; paclitaxel; pegaspargase; peliomycin; pentamustine; peplomycin sulfate; perfosfamide; pipobroman; pipsulfan; piroxantrone hydrochloride; plicamycin; plomestane; porfimer sodium; porfiromycin; prednimustine; procarbazine hydrochloride; puromycin; puromycin hydrochloride; pyrazofurin; riboprine; safingol; safingol hydrochloride; semustine; simtrazene; sparfosate sodium; sparsomycin; spirogermanium hydrochloride; spiro-mustine; spiroplatin; streptonigrin; streptozocin; sulofenur; talisomycin; tecogalan sodium; taxotere; tegafur; teloxantrone hydrochloride; temoporfin; teniposide; teroxirone; testolactone; thiamiprine; thioguanine; thiotepa; tiazofurin; tirapazamine; toremifene citrate; tretolone acetate; triciribine phosphate; trimetrexate; trimetrexate glucuronate; triptorelin; tubulozole hydrochloride; uracil mustard; uredepa; vaporeotide; verteporfin; vinblastine sulfate; vincristine sulfate; vindesine; vindesine sulfate; vinepidine sulfate; vinglycinate sulfate; vinleurosine sulfate; vinorelbine tartrate; vinrosidine sulfate; vinzolidine sulfate; vorozole; zeniplatin; zinostatin; and zorubicin hydrochloride.

Other anti-cancer drugs include, but are not limited to: 20-epi-1,25 dihydroxyvitamin D3; 5-ethynyluracil; abiraterone; aclarubicin; acylfulvene; adecypenol; adozelesin; aldesleukin; ALL-TK antagonists; altretamine; ambamustine; amidox; amifostine; aminolevulinic acid; amrubicin; amsacrine; anagrelide; anastrozole; andrographolide; angiogenesis inhibitors; antagonist D; antagonist G; antarelix; antidorsalizing morphogenetic protein-1; antiandrogen, prostatic carcinoma; antiestrogen; antineoplaston; antisense oligonucleotides; aphidicolin glycinate; apoptosis gene modulators; apoptosis regulators; apurinic acid; ara-CDP-DL-PTBA; arginine deaminase; asulacrine; atamcstane; atrimustine; axinastatin 1; axinastatin 2; axinastatin 3; azasetron; azatoxin; azatyrosine; baccatin ITT derivatives; balanol; batimastat; BCR/ABL antagonists; benzochlorins; benzoylstauroporine; beta lactam derivatives; beta-alethine; betaclamycin B; betulinic acid; bFGF inhibitor; bicalutamide; bisantrene; bisaziridinylspermine; bisnafide; bistratene A; bizelesin; breflate; bropirimine; budotitane; buthionine sulfoximine; calcipotriol; calphostin C; camptothecin derivatives; capecitabine; carboxamide-amino-triazole; carboxamidotriazole; CaRest M3; CARN 700; cartilage derived inhibitor; carzelesin; casein kinase inhibitors (ICOS); castanospermine; cecropin B; cetrorelix; chlorins; chloroquinoline sulfonamide; cicaprost; cis-porphyrin; cladribine; clomifene analogues; clotrimazole; collismycin A; collismycin B; combretastatin A4; combretastatin analogue; conagein; crambescidin 816; crisnatol; cryptophycin 8; cryptophycin A derivatives; curacin A; cyclopentantraquinones; cycloplatin; cypemycin; cytarabine ocfosfate; cytolytic factor; cytostatin; dacliximab; decitabine; dehydrididemnin B; deslorelin; dexamethasone; dexifosfamide; dexrazoxane; dexverapamil; diaziquone; didemnin B; didox; diethylnor-spermine; dihydro-5-azacytidine; dihydrotaxol, 9-; dioxamycin; diphenyl spiromustine; docetaxel; docosanol; dolasetron; doxifluridine; doxorubicin; droloxifene; dronabinol; duocarmycin SA; ebsclen; ecomustine; edelfosine; edrecolomab; eflornithine; elemene; emitefur; epirubicin; epristeride;

US 8,530,498 B1

15

estramustine analogue; estrogen agonists; estrogen antagonists; etanidazole; etoposide phosphate; exemestane; fadrozole; fazarabine; fenretinide; filgrastim; finasteride; flavopiridol; fluzelastine; fluasterone; fludarabine; fluorodaunorubicin hydrochloride; forfenimex; formestane; fostriecin; fotemustine; gadolinium texaphyrin; gallium nitrate; galocitabine; ganirelix; gelatinase inhibitors; gemcitabine; glutathione inhibitors; hepsulfam; heregulin; hexamethylene bisacetamide; hypericin; ibandronic acid; idarubicin; idoxifene; idramantone; ilmofosine; ilomastat; imatinib (e.g., Gleevec®), imiquimod; immunostimulant peptides; insulin-like growth factor-1 receptor inhibitor; interferon agonists; interferons; interleukins; iobenguane; iododoxorubicin; ipomeanol, 4-; iroplact; irsogladine; isobengazole; isohomohalicondrin B; itasetron; jaspilakinolide; kahalalide F; lamellarin-N triacetate; lanreotide; leinamycin; lenograstim; lentinan sulfate; leptolstatin; letrozole; leukemia inhibiting factor; leukocyte alpha interferon; leuprolide+estrogen+progesterone; leuprorelin; levamisole; liarozole; linear polyamine analogue; lipophilic disaccharide peptide; lipophilic platinum compounds; lissoclinamide 7; lobaplatin; lombricine; lometrexol; lonidamine; losoxantrone; loxoribine; lurtotecan; lutetium texaphyrin; lysofylline; lytic peptides; maitansine; mannostatin A; marimastat; masoprocol; maspin; matrilysin inhibitors; matrix metalloproteinase inhibitors; menogaril; merbarone; meterelin; methioninase; metoclopramide; MIF inhibitor; mifepristone; miltefosine; mirimostim; mitoguanzone; mitolactol; mitomycin analogues; mitonafide; mitotoxin fibroblast growth factor-saporin; mitoxantrone; mofarotene; molgramostim; Erbitux, human chorionic gonadotrophin; monophosphoryl lipid A+myobacterium cell wall sk; mopidamol; mustard anticancer agent; mycaperoxide B; mycobacterial cell wall extract; myriaporone; N-acetyldinaline; N-substituted benzamides; nafarelin; nagrestip; naloxone+pentazocine; napavin; naphterpin; nartograstim; nedaplatin; nemorubicin; neridronic acid; nilutamide; nisamycin; nitric oxide modulators; nitroxide antioxidant; nitrullyn; oblimersen (Genasense®); O⁶-benzylguanine; octreotide; okicenone; oligonucleotides; onapristone; ondansetron; ondansetron; oracin; oral cytokine inducer; ormaplatin; osaterone; oxaliplatin; oxaunomycin; paclitaxel; paclitaxel analogues; paclitaxel derivatives; palauamine; palmitoylrhizoxin; pamidronic acid; panaxytriol; panomifene; parabactin; pazelliptine; pegaspargase; peldesine; pentosan polysulfate sodium; pentostatin; pentozole; perflubron; perfosfamide; perillyl alcohol; phenazinomycin; phenylacetate; phosphatase inhibitors; picibanil; pilocarpine hydrochloride; pirarubicin; piritrexim; placetin A; placetin B; plasminogen activator inhibitor; platinum complex; platinum compounds; platinum-triamine complex; porfimer sodium; porfiriomycin; prednisone; propyl bis-acridone; prostaglandin J2; proteasome inhibitors; protein A-based immune modulator; protein kinase C inhibitor; protein kinase C inhibitors, microalgal; protein tyrosine phosphatase inhibitors; purine nucleoside phosphorylase inhibitors; purpurins; pyrazoloacridine; pyridoxylated hemoglobin polyoxyethylene conjugate; raf antagonists; raltitrexed; ramosestron; ras farnesyl protein transferase inhibitors; ras inhibitors; ras-GAP inhibitor; retelliptine demethylated; rhenium Re 186 etidronate; rhizoxin; ribozymes; RII retinamide; rohitukine; romurtide; roquinimex; rubiginone B1; ruboxyl; safingol; saintopin; SarCNU; sarcophytol A; sargramostim; Sdi 1 mimetics; semustine; senescence derived inhibitor 1; sense oligonucleotides; signal transduction inhibitors; sizofuran; sobuzoxane; sodium borocaptate; sodium phenylacetate; solverol; somatomedin binding protein; sonermin; sparfosic acid; spicamycin D; spiromustine; splenopentin;

16

spongistatin 1; squalamine; stipiamide; stromelysin inhibitors; sulfinosine; superactive vasoactive intestinal peptide antagonist; suradista; suramin; swainsonine; tallimustine; tamoxifen methiodide; taumustine; tazartotene; tecogalan sodium; tegafur; tellurapyrylium; telomerase inhibitors; temoporfin; teniposide; tetrachlorodecaoxide; tetrazomine; thaliblastine; thiocoraline; thrombopoietin; thrombopoietin mimetic; thymalfasin; thymopoietin receptor agonist; thymotrigan; thyroid stimulating hormone; tin ethyl etiopurpurin; tirapazamine; titanocene bichloride; topsentin; toremifene; translation inhibitors; tretinoin; triacetyluridine; triciribine; trimetrexate; triptorelin; tropisetron; turosteride; tyrosine kinase inhibitors; tyrphostins; UBC inhibitors; ubenimex; urogenital sinus-derived growth inhibitory factor; urokinase receptor antagonists; vapreotide; variolin B; velaresol; veramine; verteporfin; vinorelbine; vinoxaltine; vitaxin; vorozole; zanoterone; zenioplatin; zilascorb; and zinostatin stimalamer.

Specific second active agents include, but are not limited to, oblimersen (Genasense®), remicade, docetaxel, celecoxib, melphalan, dexamethasone (Decadron®), steroids, gemcitabine, cisplatin, temozolomide, etoposide, cyclophosphamide, temodar, carboplatin, procarbazine, gliadel, tamoxifen, topotecan, methotrexate, Arisa®, taxol, taxotere, fluorouracil, leucovorin, irinotecan, xeloda, CPT-11, interferon alpha, pegylated interferon alpha (e.g., PEG INTRON-A), capecitabine, cisplatin, thiotepa, fludarabine, carboplatin, liposomal daunorubicin, cytarabine, doxetaxol, paclitaxel, vinblastine, IL-2, GM-CSF, dacarbazine, vinorelbine, zoledronic acid, palmitronate, biaxin, busulphan, prednisone, bisphosphonate, arsenic trioxide, vincristine, doxorubicin (Doxil®), paclitaxel, ganciclovir, adriamycin, estramustine sodium phosphate (Emcyt®), sulindac, and etoposide.

5.3 Methods of Treatments and Prevention

Methods of this invention encompass methods of treating, preventing and/or managing various types of cancer and diseases and disorders associated with, or characterized by, undesired angiogenesis. As used herein, unless otherwise specified, the term “treating” refers to the administration of a compound of the invention or other additional active agent after the onset of symptoms of the particular disease or disorder. As used herein, unless otherwise specified, the term “preventing” refers to the administration prior to the onset of symptoms, particularly to patients at risk of cancer, and other diseases and disorders associated with, or characterized by, undesired angiogenesis. The term “prevention” includes the inhibition of a symptom of the particular disease or disorder. Patients with familial history of cancer and diseases and disorders associated with, or characterized by, undesired angiogenesis are preferred candidates for preventive regimens. As used herein and unless otherwise indicated, the term “managing” encompasses preventing the recurrence of the particular disease or disorder in a patient who had suffered from it, and/or lengthening the time a patient who had suffered from the disease or disorder remains in remission.

As used herein, the term “cancer” includes, but is not limited to, solid tumors and blood born tumors. The term “cancer” refers to disease of skin tissues, organs, blood, and vessels, including, but not limited to, cancers of the bladder, bone or blood, brain, breast, cervix, chest, colon, endometrium, esophagus, eye, head, kidney, liver, lymph nodes, lung, mouth, neck, ovaries, pancreas, prostate, rectum, stomach, testis, throat, and uterus. Specific cancers include, but are not limited to, advanced malignancy, amyloidosis, neuroblastoma, meningioma, hemangiopericytoma, multiple brain metastase, glioblastoma multiforms, glioblastoma, brain stem glioma, poor prognosis malignant brain tumor,

US 8,530,498 B1

17

malignant glioma, recurrent malignant glioma, anaplastic astrocytoma, anaplastic oligodendroglioma, neuroendocrine tumor, rectal adenocarcinoma, Dukes C & D colorectal cancer, unresectable colorectal carcinoma, metastatic hepatocellular carcinoma, Kaposi's sarcoma, karotype acute myeloblastic leukemia, Hodgkin's lymphoma, non-Hodgkin's lymphoma, cutaneous T-Cell lymphoma, cutaneous B-Cell lymphoma, diffuse large B-Cell lymphoma, low grade follicular lymphoma, malignant melanoma, malignant mesothelioma, malignant pleural effusion mesothelioma syndrome, peritoneal carcinoma, papillary serous carcinoma, gynecologic sarcoma, soft tissue sarcoma, scleroderma, cutaneous vasculitis, Langerhans cell histiocytosis, leiomyosarcoma, fibrodysplasia ossificans progressive, hormone refractory prostate cancer, resected high-risk soft tissue sarcoma, unresectable hepatocellular carcinoma, Waldenstrom's macroglobulinemia, smoldering myeloma, indolent myeloma, fallopian tube cancer, androgen independent prostate cancer, androgen dependent stage IV non-metastatic prostate cancer, hormone-insensitive prostate cancer, chemotherapy-insensitive prostate cancer, papillary thyroid carcinoma, follicular thyroid carcinoma, medullary thyroid carcinoma, and leiomyoma. In a specific embodiment, the cancer is metastatic. In another embodiment, the cancer is refractory or resistance to chemotherapy or radiation; in particular, refractory to thalidomide.

As used herein to refer to diseases and conditions other than cancer, the terms "diseases or disorders associated with, or characterized by, undesired angiogenesis," "diseases or disorders associated with undesired angiogenesis," and "diseases or disorders characterized by undesired angiogenesis" refer to diseases, disorders and conditions that are caused, mediated or attended by undesired, unwanted or uncontrolled angiogenesis, including, but not limited to, inflammatory diseases, autoimmune diseases, genetic diseases, allergic diseases, bacterial diseases, ocular neovascular diseases, choroidal neovascular diseases, and retina neovascular diseases.

Examples of such diseases or disorders associated with undesired angiogenesis include, but are not limited to, diabetic retinopathy, retinopathy of prematurity, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, proliferative vitreoretinopathy, trachoma, myopia, optic pits, epidemic keratoconjunctivitis, atopic keratitis, superior limbic keratitis, pterygium keratitis sicca, sjogrens, acne rosacea, phlyctenulosis, syphilis, lipid degeneration, bacterial ulcer, fungal ulcer, Herpes simplex infection, Herpes zoster infection, protozoan infection, Kaposi sarcoma, Mooren ulcer, Terrien's marginal degeneration, marginal keratolysis, rheumatoid arthritis, systemic lupus, polyarteritis, trauma, Wegeners sarcoidosis, Scleritis, Steven's Johnson disease, peripheral radial keratotomy, sickle cell anemia, sarcoid, pseudoxanthoma elasticum, Pagets disease, vein occlusion, artery occlusion, carotid obstructive disease, chronic uveitis, chronic vitritis, Lyme's disease, Eales disease, Behcet's disease, retinitis, choroiditis, presumed ocular histoplasmosis, Bests disease, Stargarts disease, pars planitis, chronic retinal detachment, hyperviscosity syndromes, toxoplasmosis, rubeosis, sarcoidosis, sclerosis, soriatis, psoriasis, primary sclerosing cholangitis, proctitis, primary biliary srosis, idiopathic pulmonary fibrosis, and alcoholic hepatitis.

In specific embodiments of the invention, diseases or disorders associated with undesired angiogenesis do not include congestive heart failure, cardiomyopathy, pulmonary edema, endotoxin-mediated septic shock, acute viral myocarditis, cardiac allograft rejection, myocardial infarction, HIV, hepatitis, adult respiratory distress syndrome, bone-resorption disease, chronic obstructive pulmonary diseases, chronic pul-

18

monary inflammatory disease, dermatitis, cystic fibrosis, septic shock, sepsis, endotoxic shock, hemodynamic shock, sepsis syndrome, post ischemic reperfusion injury, meningitis, psoriasis, fibrotic disease, cachexia, graft rejection, rheumatoid spondylitis, osteoporosis, Crohn's disease, ulcerative colitis, inflammatory-bowel disease, multiple sclerosis, systemic lupus erythrematosus, erythema nodosum leprosum in leprosy, radiation damage, asthma, hyperoxic alveolar injury, malaria, mycobacterial infection, and opportunistic infections resulting from HIV.

This invention encompasses methods of treating patients who have been previously treated for cancer or diseases or disorders associated with, or characterized by, undesired angiogenesis, but are non-responsive to standard therapies, as well as those who have not previously been treated. The invention also encompasses methods of treating patients regardless of patient's age, although some diseases or disorders are more common in certain age groups. The invention further encompasses methods of treating patients who have undergone surgery in an attempt to treat the disease or condition at issue, as well as those who have not. Because patients with cancer and diseases and disorders characterized by undesired angiogenesis have heterogenous clinical manifestations and varying clinical outcomes, the treatment given to a patient may vary, depending on his/her prognosis. The skilled clinician will be able to readily determine without undue experimentation specific secondary agents, types of surgery, and types of non-drug based standard therapy that can be effectively used to treat an individual patient with cancer and other diseases or disorders.

Methods encompassed by this invention comprise administering one or more immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, to a patient (e.g., a human) suffering, or likely to suffer, from cancer or a disease or disorder mediated by undesired angiogenesis.

In one embodiment of the invention, an immunomodulatory compound of the invention can be administered orally and in single or divided daily doses in an amount of from about 0.10 to about 150 mg/day. In a particular embodiment, 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™) may be administered in an amount of from about 0.1 to about 1 mg per day, or alternatively from about 0.1 to about 5 mg every other day. In a preferred embodiment, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) may be administered in an amount of from about 5 to 25 mg per day, or alternatively from about 10 to about 50 mg every other day.

In a specific embodiment, 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™) may be administered in an amount of about 1, 2, or 5 mg per day to patients with relapsed multiple myeloma. In a particular embodiment, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) may be administered initially in an amount of 5 mg/day and the dose can be escalated every week to 10, 20, 25, 30 and 50 mg/day. In a specific embodiment, Revimid™ can be administered in an amount of up to about 30 mg/day to patients with solid tumor. In a particular embodiment, Revimid™ can be administered in an amount of up to about 40 mg/day to patients with glioma.

5.3.1 Combination Therapy with a Second Active Agent

Specific methods of the invention comprise administering an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, in combination with one or more second active agents, and/or in combination with radia-

US 8,530,498 B1

19

tion therapy, blood transfusions, or surgery. Examples of immunomodulatory compounds of the invention are disclosed herein (see, e.g., section 5.1). Examples of second active agents are also disclosed herein (see, e.g., section 5.2).

Administration of the immunomodulatory compounds and the second active agents to a patient can occur simultaneously or sequentially by the same or different routes of administration. The suitability of a particular route of administration employed for a particular active agent will depend on the active agent itself (e.g., whether it can be administered orally without decomposing prior to entering the blood stream) and the disease being treated. A preferred route of administration for an immunomodulatory compound of the invention is orally. Preferred routes of administration for the second active agents or ingredients of the invention are known to those of ordinary skill in the art. See, e.g., *Physicians' Desk Reference*, 1755-1760 (56th ed., 2002).

In one embodiment of the invention, the second active agent is administered intravenously or subcutaneously and once or twice daily in an amount of from about 1 to about 1000 mg, from about 5 to about 500 mg, from about 10 to about 350 mg, or from about 50 to about 200 mg. The specific amount of the second active agent will depend on the specific agent used, the type of disease being treated or managed, the severity and stage of disease, and the amount(s) of immunomodulatory compounds of the invention and any optional additional active agents concurrently administered to the patient. In a particular embodiment, the second active agent is oblimersen (Genasense®), GM-CSF, G-CSF, EPO, taxotere, irinotecan, dacarbazine, transretinoic acid, topotecan, pentoxifylline, ciprofloxacin, dexamethasone, vincristine, doxorubicin, COX-2 inhibitor, IL2, IL8, IL18, IFN, Ara-C, vinorelbine, or a combination thereof.

In a particular embodiment, GM-CSF, G-CSF or EPO is administered subcutaneously during about five days in a four or six week cycle in an amount of from about 1 to about 750 mg/m²/day, preferably in an amount of from about 25 to about 500 mg/m²/day, more preferably in an amount of from about 50 to about 250 mg/m²/day, and most preferably in an amount of from about 50 to about 200 mg/m²/day. In a certain embodiment, GM-CSF may be administered in an amount of from about 60 to about 500 mcg/m² intravenously over 2 hours, or from about 5 to about 12 mcg/m²/day subcutaneously. In a specific embodiment, G-CSF may be administered subcutaneously in an amount of about 1 mcg/kg/day initially and can be adjusted depending on rise of total granulocyte counts. The maintenance dose of G-CSF may be administered in an amount of about 300 (in smaller patients) or 480 mcg subcutaneously. In a certain embodiment, EPO may be administered subcutaneously in an amount of 10,000 Unit 3 times per week.

In another embodiment, Revimid™ in an amount of about 25 mg/d and dacarbazine in an amount of about from 200 to 1,000 mg/m²/d are administered to patients with metastatic malignant melanoma. In a specific embodiment, Revimid™ is administered in an amount of from about 5 to about 25 mg/d to patients with metastatic malignant melanoma whose disease has progressed on treatment with dacarbazine, IL-2 or IFN. In a specific embodiment, Revimid™ is administered to patients with relapsed or refractory multiple myeloma in an amount of about 15 mg/d twice a day or about 30 mg/d four times a day in a combination with dexamethasone.

In another embodiment, an immunomodulatory compound is administered with melphalan and dexamethasone to patients with amyloidosis. In a specific embodiment, an immunomodulatory compound of the invention and steroids can be administered to patients with amyloidosis.

20

In another embodiment, an immunomodulatory compound is administered with gemcitabine and cisplatin to patients with locally advanced or metastatic transitional cell bladder cancer.

In another embodiment, an immunomodulatory compound is administered in combination with a second active ingredient as follows: temozolomide to pediatric patients with relapsed or progressive brain tumors or recurrent neuroblastoma; celecoxib, etoposide and cyclophosphamide for relapsed or progressive CNS cancer; temodar to patients with recurrent or progressive meningioma, malignant meningioma, hemangiopericytoma, multiple brain metastases, relapsed brain tumors, or newly diagnosed glioblastoma multiformis; irinotecan to patients with recurrent glioblastoma; carboplatin to pediatric patients with brain stem glioma; procarbazine to pediatric patients with progressive malignant gliomas; cyclophosphamide to patients with poor prognosis malignant brain tumors, newly diagnosed or recurrent glioblastoma multiformis; Gliadel® for high grade recurrent malignant gliomas; temozolomide and tamoxifen for anaplastic astrocytoma; or topotecan for gliomas, glioblastoma, anaplastic astrocytoma or anaplastic oligodendroglioma.

In another embodiment, an immunomodulatory compound is administered with methotrexate and cyclophosphamide to patients with metastatic breast cancer.

In another embodiment, an immunomodulatory compound is administered with temozolomide to patients with neuroendocrine tumors.

In another embodiment, an immunomodulatory compound is administered with gemcitabine to patients with recurrent or metastatic head or neck cancer. In another embodiment, an immunomodulatory compound is administered with gemcitabine to patients with pancreatic cancer.

In another embodiment, an immunomodulatory compound is administered to patients with colon cancer in combination with Arisa®, taxol and/or taxotere.

In another embodiment, an immunomodulatory compound is administered with capecitabine to patients with refractory colorectal cancer or patients who fail first line therapy or have poor performance in colon or rectal adenocarcinoma.

In another embodiment, an immunomodulatory compound is administered in combination with fluorouracil, leucovorin, and irinotecan to patients with Dukes C & D colorectal cancer or to patients who have been previously treated for metastatic colorectal cancer.

In another embodiment, an immunomodulatory compound is administered to patients with refractory colorectal cancer in combination with capecitabine, xeloda, and/or CPT-11.

In another embodiment, an immunomodulatory compound of the invention is administered with capecitabine and irinotecan to patients with refractory colorectal cancer or to patients with unresectable or metastatic colorectal carcinoma.

In another embodiment, an immunomodulatory compound is administered alone or in combination with interferon alpha or capecitabine to patients with unresectable or metastatic hepatocellular carcinoma; or with cisplatin and thiotepa to patients with primary or metastatic liver cancer.

In another embodiment, an immunomodulatory compound is administered in combination with pegylated interferon alpha to patients with Kaposi's sarcoma.

In another embodiment, an immunomodulatory compound is administered in combination with fludarabine, carboplatin, and/or topotecan to patients with refractory or relapsed or high-risk acuted myelogenous leukemia.

In another embodiment, an immunomodulatory compound is administered in combination with liposomal daunorubicin,

US 8,530,498 B1

21

topotecan and/or cytarabine to patients with unfavorable karyotype acute myeloblastic leukemia.

In another embodiment, an immunomodulatory compound is administered in combination with gemcitabine and irinotecan to patients with non-small cell lung cancer. In one embodiment, an immunomodulatory compound is administered in combination with carboplatin and irinotecan to patients with non-small cell lung cancer. In one embodiment, an immunomodulatory compound is administered with doxetaxol to patients with non-small cell lung cancer who have been previously treated with carbo/VP 16 and radiotherapy.

In another embodiment, an immunomodulatory compound is administered in combination with carboplatin and/or taxotere, or in combination with carboplatin, paclitaxel and/or thoracic radiotherapy to patients with non-small cell lung cancer. In a specific embodiment, an immunomodulatory compound is administered in combination with taxotere to patients with stage IIIB or IV non-small cell lung cancer.

In another embodiment, an immunomodulatory compound of the invention is administered in combination with oblimersen (Genasense®) to patients with small cell lung cancer.

In another embodiment, an immunomodulatory compound is administered alone or in combination with a second active ingredient such as vinblastine or fludarabine to patients with various types of lymphoma, including, but not limited to, Hodgkin's lymphoma, non-Hodgkin's lymphoma, cutaneous T-Cell lymphoma, cutaneous B-Cell lymphoma, diffuse large B-Cell lymphoma or relapsed or refractory low grade follicular lymphoma.

In another embodiment, an immunomodulatory compound is administered in combination with taxotere, IL-2, IFN, GM-CSF, and/or dacarbazine to patients with various types or stages of melanoma.

In another embodiment, an immunomodulatory compound is administered alone or in combination with vinorelbine to patients with malignant mesothelioma, or stage IIIB non-small cell lung cancer with pleural implants or malignant pleural effusion mesothelioma syndrome.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of multiple myeloma in combination with dexamethasone, zoledronic acid, palmitronate, GM-CSF, biacin, vinblastine, melphalan, busulphan, cyclophosphamide, IFN, palmidronate, prednisone, bisphosphonate, celecoxib, arsenic trioxide, PEG INTRON-A, vincristine, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with relapsed or refractory multiple myeloma in combination with doxorubicin (Doxil®), vincristine and/or dexamethasone (Decadron®).

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of ovarian cancer such as peritoneal carcinoma, papillary serous carcinoma, refractory ovarian cancer or recurrent ovarian cancer, in combination with taxol, carboplatin, doxorubicin, gemcitabine, cisplatin, xeloda, paclitaxel, dexamethasone, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of prostate cancer, in combination with xeloda, 5 FU/LV, gemcitabine, irinotecan plus gemcitabine, cyclophosphamide, vincristine, dexamethasone, GM-CSF, celecoxib, taxotere, ganciclovir, paclitaxel, adriamycin, docetaxel, estramustine, Emcyt, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of

22

renal cell cancer, in combination with capecitabine, IFN, tamoxifen, IL-2, GM-CSF, Celebrex®, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of gynecologic, uterus or soft tissue sarcoma cancer in combination with IFN, a COX-2 inhibitor such as Celebrex®, and/or sulindac.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of solid tumors in combination with celebrex, etoposide, cyclophosphamide, docetaxel, apecitabine, IFN, tamoxifen, IL-2, GM-CSF, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with scleroderma or cutaneous vasculitis in combination with celebrex, etoposide, cyclophosphamide, docetaxel, apecitabine, IFN, tamoxifen, IL-2, GM-CSF, or a combination thereof.

This invention also encompasses a method of increasing the dosage of an anti-cancer drug or agent that can be safely and effectively administered to a patient, which comprises administering to a patient (e.g., a human) an immunomodulatory compound of the invention, or a pharmaceutically acceptable derivative, salt, solvate, clathrate, hydrate, or prodrug thereof. Patients that can benefit by this method are those likely to suffer from an adverse effect associated with anti-cancer drugs for treating a specific cancer of the skin, subcutaneous tissue, lymph nodes, brain, lung, liver, bone, intestine, colon, heart, pancreas, adrenal, kidney, prostate, breast, colorectal, or combinations thereof. The administration of an immunomodulatory compound of the invention alleviates or reduces adverse effects which are of such severity that it would otherwise limit the amount of anti-cancer drug.

In one embodiment, an immunomodulatory compound of the invention can be administered orally and daily in an amount of from about 0.1 to about 150 mg, and preferably from about 1 to about 50 mg, more preferably from about 2 to about 25 mg prior to, during, or after the occurrence of the adverse effect associated with the administration of an anti-cancer drug to a patient. In a particular embodiment, an immunomodulatory compound of the invention is administered in combination with specific agents such as heparin, aspirin, coumadin, or G-CSF to avoid adverse effects that are associated with anti-cancer drugs such as but not limited to neutropenia or thrombocytopenia.

In one embodiment, an immunomodulatory compound of the invention can be administered to patients with diseases and disorders associated with, or characterized by, undesired angiogenesis in combination with additional active ingredients including but not limited to anti-cancer drugs, anti-inflammatories, antihistamines, antibiotics, and steroids.

In another embodiment, this invention encompasses a method of treating, preventing and/or managing cancer, which comprises administering an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, in conjunction with (e.g. before, during, or after) conventional therapy including, but not limited to, surgery, immunotherapy, biological therapy, radiation therapy, or other non-drug based therapy presently used to treat, prevent or manage cancer. The combined use of the immunomodulatory compounds of the invention and conventional therapy may provide a unique treatment regimen that is unexpectedly effective in certain patients. Without being limited by theory, it is believed that immunomodulatory compounds of the invention may provide additive or synergistic effects when given concurrently with conventional therapy.

US 8,530,498 B1

23

As discussed elsewhere herein, the invention encompasses a method of reducing, treating and/or preventing adverse or undesired effects associated with conventional therapy including, but not limited to, surgery, chemotherapy, radiation therapy, hormonal therapy, biological therapy and immunotherapy. One or more immunomodulatory compounds of the invention and other active ingredient can be administered to a patient prior to, during, or after the occurrence of the adverse effect associated with conventional therapy.

In one embodiment, an immunomodulatory compound of the invention can be administered in an amount of from about 0.1 to about 150 mg, and preferably from about 1 to about 25 mg, more preferably from about 2 to about 10 mg orally and daily alone, or in combination with a second active agent disclosed herein (see, e.g., section 5.2), prior to, during, or after the use of conventional therapy.

In a specific embodiment of this method, an immunomodulatory compound of the invention and doxorubicin are administered to patients with non-small cell lung cancer who were previously treated with carbo/VP 16 and radiotherapy.

5.3.2 Use with Transplantation Therapy

Compounds of the invention can be used to reduce the risk of Graft Versus Host Disease (GVHD). Therefore, the invention encompasses a method of treating, preventing and/or managing cancer, which comprises administering the immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, in conjunction with transplantation therapy.

As those of ordinary skill in the art are aware, the treatment of cancer is often based on the stages and mechanism of the disease. For example, as inevitable leukemic transformation develops in certain stages of cancer, transplantation of peripheral blood stem cells, hematopoietic stem cell preparation or bone marrow may be necessary. The combined use of the immunomodulatory compound of the invention and transplantation therapy provides a unique and unexpected synergism. In particular, an immunomodulatory compound of the invention exhibits immunomodulatory activity that may provide additive or synergistic effects when given concurrently with transplantation therapy in patients with cancer.

An immunomodulatory compound of the invention can work in combination with transplantation therapy reducing complications associated with the invasive procedure of transplantation and risk of GVHD. This invention encompasses a method of treating, preventing and/or managing cancer which comprises administering to a patient (e.g., a human) an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, before, during, or after the transplantation of umbilical cord blood, placental blood, peripheral blood stem cell, hematopoietic stem cell preparation or bone marrow. Examples of stem cells suitable for use in the methods of the invention are disclosed in U.S. provisional patent application No. 60/372,348, filed Apr. 12, 2002 by R. Hariri et al., the entirety of which is incorporated herein by reference.

In one embodiment of this method, an immunomodulatory compound of the invention is administered to patients with multiple myeloma before, during, or after the transplantation of autologous peripheral blood progenitor cell.

In another embodiment, an immunomodulatory compound is administered to patients with relapsing multiple myeloma after the stem cell transplantation.

24

In another embodiment, an immunomodulatory compound and prednisone are administered as maintenance therapy to patients with multiple myeloma following the transplantation of autologous stem cell.

In another embodiment, an immunomodulatory compound and dexamethasone are administered as salvage therapy for low risk post transplantation to patients with multiple myeloma.

In another embodiment, an immunomodulatory compound and dexamethasone are administered as maintenance therapy to patients with multiple myeloma following the transplantation of autologous bone marrow.

In another embodiment, an immunomodulatory compound is administered following the administration of high dose of melphalan and the transplantation of autologous stem cell to patients with chemotherapy responsive multiple myeloma.

In another embodiment, an immunomodulatory compound and PEG INTRO-A are administered as maintenance therapy to patients with multiple myeloma following the transplantation of autologous CD34-selected peripheral stem cell.

In another embodiment, an immunomodulatory compound is administered with post transplant consolidation chemotherapy to patients with newly diagnosed multiple myeloma to evaluate anti-angiogenesis.

In another embodiment, an immunomodulatory compound and dexamethasone are administered as maintenance therapy after DCEP consolidation, following the treatment with high dose of melphalan and the transplantation of peripheral blood stem cell to 65 years of age or older patients with multiple myeloma.

5.3.3 Cycling Therapy

In certain embodiments, the prophylactic or therapeutic agents of the invention are cyclically administered to a patient. Cycling therapy involves the administration of an active agent for a period of time, followed by a rest for a period of time, and repeating this sequential administration. Cycling therapy can reduce the development of resistance to one or more of the therapies, avoid or reduce the side effects of one of the therapies, and/or improves the efficacy of the treatment.

Consequently, in one specific embodiment of the invention, an immunomodulatory compound of the invention is administered daily in a single or divided doses in a four to six week cycle with a rest period of about a week or two weeks. The invention further allows the frequency, number, and length of dosing cycles to be increased. Thus, another specific embodiment of the invention encompasses the administration of an immunomodulatory compound of the invention for more cycles than are typical when it is administered alone. In yet another specific embodiment of the invention, an immunomodulatory compound of the invention is administered for a greater number of cycles that would typically cause dose-limiting toxicity in a patient to whom a second active ingredient is not also being administered.

In one embodiment, an immunomodulatory compound of the invention is administered daily and continuously for three or four weeks at a dose of from about 0.1 to about 150 mg/d followed by a break of one or two weeks. Actimid™ is preferably administered daily and continuously at an initial dose of 0.1 to 5 mg/d with dose escalation (every week) by 1 to 10 mg/d to a maximum dose of 50 mg/d for as long as therapy is tolerated. In a particular embodiment, Revimid™ is administered in an amount of about 5, 10, or 25 mg/day, preferably in an amount of about 10 mg/day for three to four weeks, followed by one week or two weeks of rest in a four or six week cycle.

In one embodiment of the invention, an immunomodulatory compound of the invention and a second active ingredient are administered orally, with administration of an immunomodulatory compound of the invention occurring 30 to 60 minutes prior to a second active ingredient, during a cycle of four to six weeks. In another embodiment of the invention, the combination of an immunomodulatory compound of the invention and a second active ingredient is administered by intravenous infusion over about 90 minutes every cycle. In a specific embodiment, one cycle comprises the administration of from about 10 to about 25 mg/day of Revimid™ and from about 50 to about 200 mg/m²/day of a second active ingredient daily for three to four weeks and then one or two weeks of rest. In another specific embodiment, each cycle comprises the administration of from about 5 to about 10 mg/day of Actimid™ and from about 50 to about 200 mg/m²/day of a second active ingredient for 3 to 4 weeks followed by one or two weeks of rest. Typically, the number of cycles during which the combinatorial treatment is administered to a patient will be from about one to about 24 cycles, more typically from about two to about 16 cycles, and even more typically from about four to about three cycles.

5.4 Pharmaceutical Compositions and Dosage Forms

Pharmaceutical compositions can be used in the preparation of individual, single unit dosage forms. Pharmaceutical compositions and dosage forms of the invention comprise an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof. Pharmaceutical compositions and dosage forms of the invention can further comprise one or more excipients.

Pharmaceutical compositions and dosage forms of the invention can also comprise one or more additional active ingredients. Consequently, pharmaceutical compositions and dosage forms of the invention comprise the active ingredients disclosed herein (e.g., an immunomodulatory compound and a second active agent). Examples of optional second, or additional, active ingredients are disclosed herein (see, e.g., section 5.2).

Single unit dosage forms of the invention are suitable for oral, mucosal (e.g., nasal, sublingual, vaginal, buccal, or rectal), parenteral (e.g., subcutaneous, intravenous, bolus injection, intramuscular, or intraarterial), topical (e.g., eye drops or other ophthalmic preparations), transdermal or transcutaneous administration to a patient. Examples of dosage forms include, but are not limited to: tablets; caplets; capsules, such as soft elastic gelatin capsules; cachets; troches; lozenges; dispersions; suppositories; powders; aerosols (e.g., nasal sprays or inhalers); gels; liquid dosage forms suitable for oral or mucosal administration to a patient, including suspensions (e.g., aqueous or non-aqueous liquid suspensions, oil-in-water emulsions, or a water-in-oil liquid emulsions), solutions, and elixirs; liquid dosage forms suitable for parenteral administration to a patient; eye drops or other ophthalmic preparations suitable for topical administration; and sterile solids (e.g., crystalline or amorphous solids) that can be reconstituted to provide liquid dosage forms suitable for parenteral administration to a patient.

The composition, shape, and type of dosage forms of the invention will typically vary depending on their use. For example, a dosage form used in the acute treatment of a disease may contain larger amounts of one or more of the active ingredients it comprises than a dosage form used in the chronic treatment of the same disease. Similarly, a parenteral dosage form may contain smaller amounts of one or more of the active ingredients it comprises than an oral dosage form used to treat the same disease. These and other ways in which

specific dosage forms encompassed by this invention will vary from one another will be readily apparent to those skilled in the art. See, e.g., *Remington's Pharmaceutical Sciences*, 18th ed., Mack Publishing, Easton Pa. (1990).

Typical pharmaceutical compositions and dosage forms comprise one or more excipients. Suitable excipients are well known to those skilled in the art of pharmacy, and non-limiting examples of suitable excipients are provided herein. Whether a particular excipient is suitable for incorporation into a pharmaceutical composition or dosage form depends on a variety of factors well known in the art including, but not limited to, the way in which the dosage form will be administered to a patient. For example, oral dosage forms such as tablets may contain excipients not suited for use in parenteral dosage forms. The suitability of a particular excipient may also depend on the specific active ingredients in the dosage form. For example, the decomposition of some active ingredients may be accelerated by some excipients such as lactose, or when exposed to water. Active ingredients that comprise primary or secondary amines are particularly susceptible to such accelerated decomposition. Consequently, this invention encompasses pharmaceutical compositions and dosage forms that contain little, if any, lactose other mono- or disaccharides. As used herein, the term "lactose-free" means that the amount of lactose present, if any, is insufficient to substantially increase the degradation rate of an active ingredient.

Lactose-free compositions of the invention can comprise excipients that are well known in the art and are listed, for example, in the *U.S. Pharmacopeia* (USP) 25-NF20 (2002). In general, lactose-free compositions comprise active ingredients, a binder/filler, and a lubricant in pharmaceutically compatible and pharmaceutically acceptable amounts. Preferred lactose-free dosage forms comprise active ingredients, microcrystalline cellulose, pre-gelatinized starch, and magnesium stearate.

This invention further encompasses anhydrous pharmaceutical compositions and dosage forms comprising active ingredients, since water can facilitate the degradation of some compounds. For example, the addition of water (e.g., 5%) is widely accepted in the pharmaceutical arts as a means of simulating long-term storage in order to determine characteristics such as shelf-life or the stability of formulations over time. See, e.g., Jens T. Carstensen, *Drug Stability: Principles & Practice*, 2d. Ed., Marcel Dekker, NY, N.Y., 1995, pp. 379-80. In effect, water and heat accelerate the decomposition of some compounds. Thus, the effect of water on a formulation can be of great significance since moisture and/or humidity are commonly encountered during manufacture, handling, packaging, storage, shipment, and use of formulations.

Anhydrous pharmaceutical compositions and dosage forms of the invention can be prepared using anhydrous or low moisture containing ingredients and low moisture or low humidity conditions. Pharmaceutical compositions and dosage forms that comprise lactose and at least one active ingredient that comprises a primary or secondary amine are preferably anhydrous if substantial contact with moisture and/or humidity during manufacturing, packaging, and/or storage is expected.

An anhydrous pharmaceutical composition should be prepared and stored such that its anhydrous nature is maintained. Accordingly, anhydrous compositions are preferably packaged using materials known to prevent exposure to water such that they can be included in suitable formulary kits. Examples

US 8,530,498 B1

27

of suitable packaging include, but are not limited to, hermetically sealed foils, plastics, unit dose containers (e.g., vials), blister packs, and strip packs.

The invention further encompasses pharmaceutical compositions and dosage forms that comprise one or more compounds that reduce the rate by which an active ingredient will decompose. Such compounds, which are referred to herein as “stabilizers,” include, but are not limited to, antioxidants such as ascorbic acid, pH buffers, or salt buffers.

Like the amounts and types of excipients, the amounts and specific types of active ingredients in a dosage form may differ depending on factors such as, but not limited to, the route by which it is to be administered to patients. However, typical dosage forms of the invention comprise an immunomodulatory compound of the invention or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof in an amount of from about 0.10 to about 150 mg. Typical dosage forms comprise an immunomodulatory compound of the invention or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof in an amount of about 0.1, 1, 2, 5, 7.5, 10, 12.5, 15, 17.5, 20, 25, 50, 100, 150 or 200 mg. In a particular embodiment, a preferred dosage form comprises 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™) in an amount of about 1, 2, 5, 10, 25 or 50 mg. In a specific embodiment, a preferred dosage form comprises 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) in an amount of about 5, 10, 25 or 50 mg. Typical dosage forms comprise the second active ingredient in an amount of 1 to about 1000 mg, from about 5 to about 500 mg, from about 10 to about 350 mg, or from about 50 to about 200 mg. Of course, the specific amount of the anti-cancer drug will depend on the specific agent used, the type of cancer being treated or managed, and the amount(s) of an immunomodulatory compound of the invention and any optional additional active agents concurrently administered to the patient.

5.4.1 Oral Dosage Forms

Pharmaceutical compositions of the invention that are suitable for oral administration can be presented as discrete dosage forms, such as, but are not limited to, tablets (e.g., chewable tablets), caplets, capsules, and liquids (e.g., flavored syrups). Such dosage forms contain predetermined amounts of active ingredients, and may be prepared by methods of pharmacy well known to those skilled in the art. See generally, *Remington's Pharmaceutical Sciences*, 18th ed., Mack Publishing, Easton Pa. (1990).

Typical oral dosage forms of the invention are prepared by combining the active ingredients in an intimate admixture with at least one excipient according to conventional pharmaceutical compounding techniques. Excipients can take a wide variety of forms depending on the form of preparation desired for administration. For example, excipients suitable for use in oral liquid or aerosol dosage forms include, but are not limited to, water, glycols, oils, alcohols, flavoring agents, preservatives, and coloring agents. Examples of excipients suitable for use in solid oral dosage forms (e.g., powders, tablets, capsules, and caplets) include, but are not limited to, starches, sugars, micro-crystalline cellulose, diluents, granulating agents, lubricants, binders, and disintegrating agents.

Because of their ease of administration, tablets and capsules represent the most advantageous oral dosage unit forms, in which case solid excipients are employed. If desired, tablets can be coated by standard aqueous or nonaqueous techniques. Such dosage forms can be prepared by any of the methods of pharmacy. In general, pharmaceutical compositions and dosage forms are prepared by uniformly and inti-

28

mately admixing the active ingredients with liquid carriers, finely divided solid carriers, or both, and then shaping the product into the desired presentation if necessary.

For example, a tablet can be prepared by compression or molding. Compressed tablets can be prepared by compressing in a suitable machine the active ingredients in a free-flowing form such as powder or granules, optionally mixed with an excipient. Molded tablets can be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

Examples of excipients that can be used in oral dosage forms of the invention include, but are not limited to, binders, fillers, disintegrants, and lubricants. Binders suitable for use in pharmaceutical compositions and dosage forms include, but are not limited to, corn starch, potato starch, or other starches, gelatin, natural and synthetic gums such as acacia, sodium alginate, alginic acid, other alginates, powdered tragacanth, guar gum, cellulose and its derivatives (e.g., ethyl cellulose, cellulose acetate, carboxymethyl cellulose calcium, sodium carboxymethyl cellulose), polyvinyl pyrrolidone, methyl cellulose, pre-gelatinized starch, hydroxypropyl methyl cellulose, (e.g., Nos. 2208, 2906, 2910), microcrystalline cellulose, and mixtures thereof.

Suitable forms of microcrystalline cellulose include, but are not limited to, the materials sold as AVICEL-PH-101, AVICEL-PH-103 AVICEL RC-581, AVICEL-PH-105 (available from FMC Corporation, American Viscose Division, Avicel Sales, Marcus Hook, PA), and mixtures thereof. A specific binder is a mixture of microcrystalline cellulose and sodium carboxymethyl cellulose sold as AVICEL RC-581. Suitable anhydrous or low moisture excipients or additives include AVICEL-PH-103™ and Starch 1500 LM.

Examples of fillers suitable for use in the pharmaceutical compositions and dosage forms disclosed herein include, but are not limited to, talc, calcium carbonate (e.g., granules or powder), microcrystalline cellulose, powdered cellulose, dextrates, kaolin, mannitol, silicic acid, sorbitol, starch, pre-gelatinized starch, and mixtures thereof. The binder or filler in pharmaceutical compositions of the invention is typically present in from about 50 to about 99 weight percent of the pharmaceutical composition or dosage form.

Disintegrants are used in the compositions of the invention to provide tablets that disintegrate when exposed to an aqueous environment. Tablets that contain too much disintegrant may disintegrate in storage, while those that contain too little may not disintegrate at a desired rate or under the desired conditions. Thus, a sufficient amount of disintegrant that is neither too much nor too little to detrimentally alter the release of the active ingredients should be used to form solid oral dosage forms of the invention. The amount of disintegrant used varies based upon the type of formulation, and is readily discernible to those of ordinary skill in the art. Typical pharmaceutical compositions comprise from about 0.5 to about 15 weight percent of disintegrant, preferably from about 1 to about 5 weight percent of disintegrant.

Disintegrants that can be used in pharmaceutical compositions and dosage forms of the invention include, but are not limited to, agar-agar, alginic acid, calcium carbonate, microcrystalline cellulose, croscarmellose sodium, crospovidone, polacrillin potassium, sodium starch glycolate, potato or tapioca starch, other starches, pre-gelatinized starch, other starches, clays, other alginates, other celluloses, gums, and mixtures thereof.

Lubricants that can be used in pharmaceutical compositions and dosage forms of the invention include, but are not limited to, calcium stearate, magnesium stearate, mineral oil, light mineral oil, glycerin, sorbitol, mannitol, polyethylene

US 8,530,498 B1

29

glycol, other glycols, stearic acid, sodium lauryl sulfate, talc, hydrogenated vegetable oil (e.g., peanut oil, cottonseed oil, sunflower oil, sesame oil, olive oil, corn oil, and soybean oil), zinc stearate, ethyl oleate, ethyl laureate, agar, and mixtures thereof. Additional lubricants include, for example, a syloid silica gel (AEROSIL200, manufactured by W.R. Grace Co. of Baltimore, Md.), a coagulated aerosol of synthetic silica (marketed by Degussa Co. of Plano, Tex.), CAB-O-SIL (a pyrogenic silicon dioxide product sold by Cabot Co. of Boston, Mass.), and mixtures thereof. If used at all, lubricants are typically used in an amount of less than about 1 weight percent of the pharmaceutical compositions or dosage forms into which they are incorporated.

A preferred solid oral dosage form of the invention comprises an immunomodulatory compound of the invention, anhydrous lactose, microcrystalline cellulose, polyvinylpyrrolidone, stearic acid, colloidal anhydrous silica, and gelatin.

5.4.2 Delayed Release Dosage Forms

Active ingredients of the invention can be administered by controlled release means or by delivery devices that are well known to those of ordinary skill in the art. Examples include, but are not limited to, those described in U.S. Pat. Nos. 3,845,770; 3,916,899; 3,536,809; 3,598,123; and 4,008,719, 5,674,533, 5,059,595, 5,591,767, 5,120,548, 5,073,543, 5,639,476, 5,354,556, and 5,733,566, each of which is incorporated herein by reference. Such dosage forms can be used to provide slow or controlled-release of one or more active ingredients using, for example, hydropropylmethyl cellulose, other polymer matrices, gels, permeable membranes, osmotic systems, multilayer coatings, microparticles, liposomes, microspheres, or a combination thereof to provide the desired release profile in varying proportions. Suitable controlled-release formulations known to those of ordinary skill in the art, including those described herein, can be readily selected for use with the active ingredients of the invention. The invention thus encompasses single unit dosage forms suitable for oral administration such as, but not limited to, tablets, capsules, gelcaps, and caplets that are adapted for controlled-release.

All controlled-release pharmaceutical products have a common goal of improving drug therapy over that achieved by their non-controlled counterparts. Ideally, the use of an optimally designed controlled-release preparation in medical treatment is characterized by a minimum of drug substance being employed to cure or control the condition in a minimum amount of time. Advantages of controlled-release formulations include extended activity of the drug, reduced dosage frequency, and increased patient compliance. In addition, controlled-release formulations can be used to affect the time of onset of action or other characteristics, such as blood levels of the drug, and can thus affect the occurrence of side (e.g., adverse) effects.

Most controlled-release formulations are designed to initially release an amount of drug (active ingredient) that promptly produces the desired therapeutic effect, and gradually and continually release of other amounts of drug to maintain this level of therapeutic or prophylactic effect over an extended period of time. In order to maintain this constant level of drug in the body, the drug must be released from the dosage form at a rate that will replace the amount of drug being metabolized and excreted from the body. Controlled-release of an active ingredient can be stimulated by various conditions including, but not limited to, pH, temperature, enzymes, water, or other physiological conditions or compounds.

30

5.4.3 Parenteral Dosage Forms

Parenteral dosage forms can be administered to patients by various routes including, but not limited to, subcutaneous, intravenous (including bolus injection), intramuscular, and intraarterial. Because their administration typically bypasses patients' natural defenses against contaminants, parenteral dosage forms are preferably sterile or capable of being sterilized prior to administration to a patient. Examples of parenteral dosage forms include, but are not limited to, solutions ready for injection, dry products ready to be dissolved or suspended in a pharmaceutically acceptable vehicle for injection, suspensions ready for injection, and emulsions.

Suitable vehicles that can be used to provide parenteral dosage forms of the invention are well known to those skilled in the art. Examples include, but are not limited to: Water for Injection USP; aqueous vehicles such as, but not limited to, Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, and Lactated Ringer's Injection; water-miscible vehicles such as, but not limited to, ethyl alcohol, polyethylene glycol, and polypropylene glycol; and non-aqueous vehicles such as, but not limited to, corn oil, cottonseed oil, peanut oil, sesame oil, ethyl oleate, isopropyl myristate, and benzyl benzoate.

Compounds that increase the solubility of one or more of the active ingredients disclosed herein can also be incorporated into the parenteral dosage forms of the invention. For example, cyclodextrin and its derivatives can be used to increase the solubility of an immunomodulatory compound of the invention and its derivatives. See, e.g., U.S. Pat. No. 5,134,127, which is incorporated herein by reference.

5.4.4 Topical and Mucosal Dosage Forms

Topical and mucosal dosage forms of the invention include, but are not limited to, sprays, aerosols, solutions, emulsions, suspensions, eye drops or other ophthalmic preparations, or other forms known to one of skill in the art. See, e.g., *Remington's Pharmaceutical Sciences*, 16th and 18th eds., Mack Publishing, Easton Pa. (1980 & 1990); and *Introduction to Pharmaceutical Dosage Forms*, 4th ed., Lea & Febiger, Philadelphia (1985). Dosage forms suitable for treating mucosal tissues within the oral cavity can be formulated as mouthwashes or as oral gels.

Suitable excipients (e.g., carriers and diluents) and other materials that can be used to provide topical and mucosal dosage forms encompassed by this invention are well known to those skilled in the pharmaceutical arts, and depend on the particular tissue to which a given pharmaceutical composition or dosage form will be applied. With that fact in mind, typical excipients include, but are not limited to, water, acetone, ethanol, ethylene glycol, propylene glycol, butane-1,3-diol, isopropyl myristate, isopropyl palmitate, mineral oil, and mixtures thereof to form solutions, emulsions or gels, which are non-toxic and pharmaceutically acceptable. Moisturizers or humectants can also be added to pharmaceutical compositions and dosage forms if desired. Examples of such additional ingredients are well known in the art. See, e.g., *Remington's Pharmaceutical Sciences*, 16th and 18th eds., Mack Publishing, Easton Pa. (1980 & 1990).

The pH of a pharmaceutical composition or dosage form may also be adjusted to improve delivery of one or more active ingredients. Similarly, the polarity of a solvent carrier, its ionic strength, or tonicity can be adjusted to improve delivery. Compounds such as stearates can also be added to pharmaceutical compositions or dosage forms to advantageously alter the hydrophilicity or lipophilicity of one or more active ingredients so as to improve delivery. In this regard, stearates can serve as a lipid vehicle for the formulation, as an emulsifying agent or surfactant, and as a delivery-

US 8,530,498 B1

31

enhancing or penetration-enhancing agent. Different salts, hydrates or solvates of the active ingredients can be used to further adjust the properties of the resulting composition.

5.4.5 Kits

Typically, active ingredients of the invention are preferably not administered to a patient at the same time or by the same route of administration. This invention therefore encompasses kits which, when used by the medical practitioner, can simplify the administration of appropriate amounts of active ingredients to a patient.

A typical kit of the invention comprises a dosage form of an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, pro-drug, or clathrate thereof. Kits encompassed by this invention can further comprise additional active ingredients such as oblimersen (Genasense®), melphalan, G-CSF, GM-CSF, EPO, topotecan, dacarbazine, irinotecan, taxotere, IFN, COX-2 inhibitor, pentoxifylline, ciprofloxacin, dexamethasone, IL2, IL8, IL18, Ara-C, vinorelbine, isotretinoin, 13 cis-retinoic acid, or a pharmacologically active mutant or derivative thereof, or a combination thereof. Examples of the additional active ingredients include, but are not limited to, those disclosed herein (see, e.g., section 5.2).

Kits of the invention can further comprise devices that are used to administer the active ingredients. Examples of such devices include, but are not limited to, syringes, drip bags, patches, and inhalers.

Kits of the invention can further comprise cells or blood for transplantation as well as pharmaceutically acceptable vehicles that can be used to administer one or more active ingredients. For example, if an active ingredient is provided in a solid form that must be reconstituted for parenteral administration, the kit can comprise a sealed container of a suitable vehicle in which the active ingredient can be dissolved to form a particulate-free sterile solution that is suitable for parenteral administration. Examples of pharmaceutically acceptable vehicles include, but are not limited to: Water for Injection USP; aqueous vehicles such as, but not limited to, Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, and Lactated Ringer's Injection; water-miscible vehicles such as, but not limited to, ethyl alcohol, polyethylene glycol, and polypropylene glycol; and non-aqueous vehicles such as, but not limited to, corn oil, cottonseed oil, peanut oil, sesame oil, ethyl oleate, isopropyl myristate, and benzyl benzoate.

6. EXAMPLES

Certain embodiments of the invention are illustrated by the following non-limiting examples.

6.1 Modulation of Cytokine Production

A series of non-clinical pharmacology and toxicology studies have been performed to support the clinical evaluation of an immunomodulatory compound of the invention in human subjects. These studies were performed in accordance with internationally recognized guidelines for study design and in compliance with the requirements of Good Laboratory Practice (GLP), unless otherwise noted.

Inhibition of TNF- α production following LPS-stimulation of human PBMC and human whole blood by 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™), 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione and thalidomide (Revimid™) was investigated in vitro (Muller et al., *Bioorg. Med. Chem. Lett.* 9:1625-1630, 1999). The IC₅₀'s of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione for inhibiting production of TNF- α following LPS-stimulation of PBMC and

32

human whole blood were ~24 nM (6.55 ng/mL) and ~25 nM (6.83 ng/mL), respectively. In vitro studies suggest a pharmacological activity profile for 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione that is similar to, but at least 200 times more potent than, thalidomide. In vitro studies have also demonstrated that concentrations of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione of 2.73 to 27.3 ng/mL (0.01 to 0.1 μ M) achieved 50% inhibition of the proliferation of MM.1S and Hs Sultan cells.

The IC₅₀'s of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione for inhibiting production of TNF- α following LPS-stimulation of PBMC and human whole blood were ~100 nM (25.9 ng/mL) and ~480 nM (103.6 ng/mL), respectively. Thalidomide, in contrast, had an IC₅₀ of ~194 μ M (50.2 μ g/mL) for inhibiting production of TNF- α following LPS-stimulation of PBMC. In vitro studies suggest a pharmacological activity profile for 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione that is similar to, but 50 to 2000 times more potent than, thalidomide. It has been shown that the compound is approximately 50-100 times more potent than thalidomide in stimulating the proliferation of T-cells following primary induction by T-cell receptor (TCR) activation. 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione is also approximately 50 to 100 times more potent than thalidomide in augmenting the production of IL-2 and IFN- γ following TCR activation of PBMC (IL-2) or T-cells (IFN- γ). In addition, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione exhibited dose-dependent inhibition of LPS-stimulated production of the pro-inflammatory cytokines TNF- α , IL-1 β , and IL-6 by PBMC while it increased production of the anti-inflammatory cytokine IL-10.

6.2 Inhibition of MM Cell Proliferation

The ability of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) and thalidomide for comparison to effect the proliferation of MM cell lines has been investigated in an in vitro study. Uptake [³H]-thymidine by different MM cell lines (MM.1S, Hs Sultan, U266 and RPMI-8226) was measured as an indicator of cell proliferation. Cells were incubated in the presence of compounds for 48 hours; [³H]-thymidine was included for the last 8 hours of the incubation period. Addition of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione to MM.1S and Hs Sultan cells resulted in 50% inhibition of cell proliferation at concentrations of 0.4 μ M and 1 μ M, respectively. In contrast, addition of thalidomide at concentrations up to 100 μ M resulted in only 15% and 20% inhibition of cell proliferation in MM.1S and Hs Sultan cells, respectively. These data are summarized in FIG. 1.

6.3 Toxicology Studies

The effects of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) on cardiovascular and respiratory function are investigated in anesthetized dogs. Two groups of Beagle dogs (2/sex/group) are used. One group receives three doses of vehicle only and the other receives three ascending doses of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (2, 10, and 20 mg/kg). In all cases, doses of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione or vehicle are successively administered via infusion through the jugular vein separated by intervals of at least 30 minutes.

The cardiovascular and respiratory changes induced by 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione are minimal at all doses when compared to the vehicle control group. The only statistically significant difference between the vehicle and treatment groups is a small increase in arterial blood pressure (from 94 mmHg to 101

mmHg) following administration of the low dose of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. This effect lasts approximately 15 minutes and is not seen at higher doses. Deviations in femoral blood flow, respiratory parameters, and Qtc interval are common to both the control and treated groups and are not considered treatment-related.

6.4 Cycling Therapy in Patients

In a specific embodiment, an immunomodulatory compound of the invention are cyclically administered to patients with cancer. Cycling therapy involves the administration of a first agent for a period of time, followed by a rest for a period of time and repeating this sequential administration. Cycling therapy can reduce the development of resistance to one or more of the therapies, avoid or reduce the side effects of one of the therapies, and/or improves the efficacy of the treatment.

In a specific embodiment, prophylactic or therapeutic agents are administered in a cycle of about 4 to 6 weeks, about once or twice every day. One cycle can comprise the administration of a therapeutic or prophylactic agent for three to four weeks and at least a week or two weeks of rest. The number of cycles administered is from about one to about 24 cycles, more typically from about two to about 16 cycles, and more typically from about four to about eight cycles.

For example, in a cycle of four weeks, on day 1, the administration of 25 mg/d of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione is started. On day 22, the administration of the compound is stopped for a week of rest. On day 29, the administration of 25 mg/d 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione is begun.

6.5 Clinical Studies in Patients

6.5.1 Treatment of Relapsed Multiple Myeloma

4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™) was administered to patients with relapsed/refractory multiple myeloma. The study was conducted in compliance with Good Clinical Practices. Patients were at least 18 years old, had been diagnosed with multiple myeloma (with paraprotein in serum and/or urine), and were considered refractory to treatment after at least two cycles of treatment, or have relapsed after two cycles of treatment.

Patients who have progressive disease, according to the Southwest Oncology Group (SWOG) criteria, on their prior regimen are considered treatment refractory. Relapse following remission is defined as >25% increase in M component from baseline levels; reappearance of the M paraprotein that had previously disappeared; or a definite increase in the size and number of lytic bone lesions recognized on radiographs. Patients may have had prior therapy with thalidomide, provided they were able to tolerate the treatment. A Zubrod performance status of 0 to 2 is required for all patients.

4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione is administered to patients at doses of 1, 2, 5, or 10 mg/day for up to four weeks; at each dose level, three patients are initially enrolled. Dosing occurs at approximately the same time each morning; all doses are administered in the fasted state (no eating for at least two hours prior to dosing and two hours after dosing). 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione doses are administered in an ascending fashion such that patients in the first cohort receive the lowest dose of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (1 mg/day) and escalation to the next higher dose level occurs only following the establishment of safety and tolerability at the current dose. If one out of three patients at any dose level experience dose limiting toxicity (DLT), three additional patients are enrolled at that dose. If none of the three additional patients experience DLT, escalation to the next dose level occurs; dose escalations continue in a similar

fashion until the MTD is established or the maximum daily dose (10 mg/day) is attained. However, if one of the three additional patients enrolled experiences DLT, the MTD has been reached. If two or more of the three additional patients enrolled experience DLT, the MTD is judged to have been exceeded and three additional patients are enrolled at the preceding dose level to confirm the MTD. Once the MTD has been identified, four additional patients are enrolled at that dose level so that a total of 10 patients is treated at the MTD.

Blood sampling for analysis of pharmacokinetic parameters is performed on Days 1 and 28 according to the following sampling schedule: pre-dose, 0.25, 0.5, 0.75, 1, 1.5, 2, 2.5, 3, 4, 6, 8, 10, 12, 18, and 24 hours post-dose. An additional blood sample is collected at each weekly visit for the determination of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione levels. Total urine collections are also made with urine pooled according to the following time intervals post-dose: 0 to 4, 4 to 8, 8 to 12, and 12 to 24 hours. Safety assessments are made by monitoring adverse events, vital signs, ECGs, clinical laboratory evaluations (blood chemistry, hematology, lymphocyte phenotyping, and urinalysis), and physical examination at specific times during the study.

Results of interim pharmacokinetic analyses obtained following single- and multiple-dose administration of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione to multiple myeloma patients are presented below in Tables 1 and 2. These data show that 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione was steadily absorbed at all dose levels in relapsed multiple myeloma patients. Maximum plasma concentrations occurred at a median T_{max} of between 2.5 and 2.8 hours post-dose at Day 1 and between 3 and 4 hours post-dose at Week 4. At all doses, plasma concentrations declined in a monophasic manner after reaching C_{max} . The start of the elimination phase occurred between 3 and 10 hours post-dose at Day 1 and Week 4, respectively.

These data also showed that after 4 weeks of dosing, 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione accumulated to a small extent (mean accumulation ratios ~1.02 to 1.52 and ~0.94 to 1.62 for C_{max} and $AUC_{(0-\tau)}$, respectively). There was almost a dose proportional increase in $AUC_{(0-\tau)}$ and C_{max} values with increasing dose. A five-fold higher dose of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione produced a 3.2- and 2.2-fold increase in C_{max} at Day 1 and Week 4, respectively. Similarly, a 5-fold increase in dose resulted in a 3.6- and 2.3-fold increase in $AUC_{(0-\tau)}$, at Day 1 and Week 4, respectively.

TABLE 1

| Pharmacokinetic parameters of Actimid™ in relapsed multiple myeloma patients | | | | |
|--|-----------|----------------|----------------|-----------------|
| Parameter | | 1 mg | 2 mg | 5 mg |
| | | (N = 6) | (N = 2) | (N = 3) |
| Day 1 | | | | |
| C_{max} | ng/mL | 15.03 (4.04) | 24.4* (12.1) | 48.56 (14.03) |
| t_{max} | h | 3.3 (2.6) | 2.7* (0.3) | 2.3 (0.3) |
| $AUC_{(0-\infty)}$ | ng · h/mL | 152.90 (36.62) | 279.18 (51.10) | 593.10 (335.23) |
| $AUC_{(0-\tau)}$ | | 134.21 (27.14) | 249.57 (29.26) | 520.94 (267.32) |
| $t_{1/2}$ | h | 7.3 (3.4) | 6.3 (1.4) | 6.5 (2.2) |
| CL/F | mL/min | 114.75 (29.20) | 121.43 (22.22) | 182.31 (117.06) |
| Vz/f | L | 69.55 (44.97) | 65.31 (2.80) | 87.24 (22.61) |

t = 24 hours

N/A = not available

US 8,530,498 B1

35

TABLE 2

| Pharmacokinetic parameters of Actimid™ following multiple oral doses (1, 2, and 5 mg/day) in relapsed multiple myeloma patients | | | | |
|---|-----------|-----------------|-----------------|-----------------|
| Parameter | | 1 mg | 2 mg | 5 mg |
| | | (N = 5) | (N = 2) | (N = 3) |
| Week 4 | | | | |
| C_{max} | ng/mL | 23.20 (7.48) | 30.05* (15.64) | 58.07 (38.08) |
| t_{max} | h | 3.6 (1.5) | 2.8* (0.3) | 5.0 (2.6) |
| $AUC_{(0-\infty)}$ | ng · h/mL | N/A | N/A | N/A |
| $AUC_{(0-\tau)}$ | | 239.31 (122.59) | 269.36 (186.34) | 597.24 (354.23) |
| $t_{1/2}$ | h | 6.2* (0.6) | 7.7 (2.8) | 7.8 (4.0) |
| CL/F | mL/min | 87.85 (48.48) | 162.68 (112.54) | 207.50 (175.41) |
| Vz/f | L | 41.35* (8.84) | 95.04 (35.39) | 103.95 (27.25) |

 τ = 24 hours

N/A = not available

*N = 3 patients

6.5.2 Treatment of Relapsed Multiple Myeloma

Two Phase 1 clinical studies of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) have been conducted to identify the maximum tolerated dose (MTD) in patients with refractory or relapsed multiple myeloma. These studies have also characterized the safety profile of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione when ascending doses of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione were given orally for up to 4 weeks. Patients started 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione treatment at 5 mg/day with subsequent escalation to 10, 25, and 50 mg/day. Patients were enrolled for 28 days at their assigned dose, with the option of extended treatment for those who did not exhibit disease progression or experience dose limiting toxicity (DLT). Patients were evaluated for adverse events at each visit and the severity of these events was graded according to the National Cancer Institute (NCI) Common Toxicity Criteria. Patients were discontinued if they experienced DLT (Grade 3 or greater non-hematological, or Grade 4 hematological toxicity).

In this study, 27 patients were enrolled. All patients had relapsed multiple myeloma and 18 (72%) were refractory to salvage therapy. Among these patients, 15 had undergone prior autologous stem cell transplantation and 16 patients had received prior thalidomide treatment. The median number of prior regimens was 3 (range 2 to 6).

Blood and urine samples were collected for analysis of pharmacokinetic parameters on Days 1 and 28. Blood samples were collected according to the following sampling schedule: pre-dose, 0.25, 0.5, 0.75, 1, 1.5, 2, 2.5, 3, 4, 6, 8, 10, 12, 18, and 24 hours post-dose. In addition, a blood sample was collected at each weekly clinic visit for 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione determination. Total urine was collected and pooled according to the following time intervals post-dose: 0 to 4, 4 to 8, 8 to 12, and 12 to 24 hours. Response to treatment was assessed by M-protein quantification (by immunoelectrophoresis) from serum and a 24-hour urine collection, with creatinine clearance and 24-hour protein calculations undertaken at screening, baseline, Weeks 2 and 4, and monthly thereafter (or upon early termination). Bone marrow aspirations and/or tissue biopsy are also performed at Months 3, 6 and 12 if a patient's paraprotein serum concentration or 24-hour urine protein excretion declined to the next lower level, based on best response criteria. Preliminary results for the 28-day treatment period are summarized below.

Preliminary pharmacokinetic analyses based on these two studies indicated that AUC and C_{max} values increase proportionally with dose following single and multiple doses in

36

multiple myeloma patients (as was seen in healthy volunteers). Further, there was no evidence of accumulation with multiple dosing as single dose $AUC_{(0-\infty)}$ was comparable to multiple dose $AUC_{(0-\tau)}$ following the same dose of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione.

Similar to healthy volunteer studies, double peaks were observed. Exposure in multiple myeloma patients appeared to be slightly higher based on C_{max} and AUC values as compared to healthy male volunteers while clearance in multiple myeloma patients was lower than it was in healthy volunteers, consistent with their poorer renal function (both as a consequence of their age and their disease). Finally, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione half-life in patients was shorter than in healthy volunteers (mean 8 hours, ranging up to 17 hours).

In this study, the first cohort of 3 patients was treated for 28 days at 5 mg/day without any dose limiting toxicity (DLT). The second cohort of 3 patients subsequently commenced therapy at 10 mg/day. Patients in the second 10 mg/day of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione cohort tolerated treatment well.

6.5.3 Treatment of Solid Tumors

Study with 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) was conducted in patients with varying types of solid tumors, including malignant melanoma (13), carcinoma of the pancreas (2), carcinoid-unknown primary (1), renal carcinoma (1), breast carcinoma (1) and NSCLC (2). Patients received 5 mg/day 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione for seven days and are subsequently escalated every seven days to 10 mg/day, 25 mg/day, and 50 mg/day for a total of 4 weeks of treatment. Patients who, experienced clinical benefit were permitted to continue on treatment as Named Patients.

The study initially enrolled 20 patients and was subsequently amended to enroll 16 additional patients (adrenal carcinoma, NSCLC, malignant mesothelioma, breast cancer, malignant melanoma (8), renal cell cancer (4)) at a higher dose. The 16 additional patients were given weekly escalating doses of 25 mg/day, 50 mg/day, 75 mg/day, 100 mg/day, 125 mg/day, and 150 mg/day over a 6-week period with continuing treatment for an additional six weeks.

The study of Phase 1 study was designed to determine a maximum tolerated dose (MTD) of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione in patients with refractory solid tumors and/or lymphoma, as well as to characterize the pharmacokinetic and side effect profiles of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione in this patient population. The study design dictates that at least 3 patients must be enrolled at a dose level and have completed 28 days of treatment prior to enrollment of patients at the next higher dose level. Patients in the first cohort began dosing at 5 mg/day of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. Patients will be escalated to 10, 20, 25, and 30 mg/day provided there is no toxicity.

In this study, the MTD is defined as the highest dose level in which fewer than two of six patients treated did not experience Grade 3 or greater non-hematological toxicity or Grade 4 or greater hematological toxicity. If, at any given dose level in either study, one out of three patients experiences toxicity, three additional patients must be treated at that particular dose. If, however, two out of six patients experience DLT, the MTD is judged to have been exceeded. No further dose escalations are to occur and additional patients are to be enrolled at the previous dose level. The dose of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione administered is escalated until the MTD is achieved or the maximum daily dose of is reached.

US 8,530,498 B1

37

No DLTs were reported in the initial group of 20 patients enrolled in the study. Thirteen of the original 20 trial patients, along with 2 non-trial patients, continued on treatment as named patients at doses up to 150 mg/day.

6.5.4 Treatment of Gliomas

This study was performed to find toxicity in patients with recurrent, high-grade gliomas. The study is designed such that patients are given increasingly higher doses of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione until a maximum tolerated dose (MTD) is established. The study also seeks to obtain preliminary toxicity information and pharmacokinetic data on 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione, as well as to develop exploratory data concerning surrogate end points of angiogenic activity in vivo using functional neuro-imaging studies, and in vitro assays of scrum angiogenic peptides.

Patients enrolled in the first cohort receive 2.5 mg/m²/day for a 4-week cycle. During each 4-week cycle of therapy, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione administered once daily for 3 weeks followed by a week of rest. Patients who complete a treatment cycle may receive another cycle of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione treatment if two criteria are met. First, the patient must have stable disease or have experienced a partial response or complete response, or the patient is benefiting from the therapy with 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione as evidenced by a decrease in tumor-related symptoms such as neurological deficits. Second, the patient must have recovered from toxicity related to 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione which occurred in the prior cycle by Day 42 or sooner (28-day cycle plus limit of 2 weeks to recover) as evidenced by a return to Grade \leq 1 toxicity level. Patients who experience DLT in the previous cycle should have their dose modified. DLT is defined as a non-hematological event Grade \geq 3 toxicity or hematological event of Grade 4 toxicity thought to be related to the study medication. Patients who experience DLT in the first cycle and have no response to therapy are removed from the study.

3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione doses are subsequently escalated to 5, 8, 11, 15, and 20 mg/m²/day to a maximum total daily dose of 40 mg. Patients continue to receive 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione on a 4-week cycle per dose level until one of the off-study criteria are met.

Three patients are enrolled in each cohort. If at least one DLT occurs, three additional patients are added to the cohort at that particular dose level. If two DLTs occur, the MTD, defined as the dose at which fewer than one-third of patients at each dose level experiences DLT has been exceeded and four more patients are treated at the previous dose.

Patients who experience DLT during the first 4-week cycle are removed from the study, except if they have a response to therapy. For patients who have completed their first 4-week cycle of without DLT, but who subsequently experience Grade 3 or 4 hematological and/or nonhematological toxicity, treatment is suspended for a minimum of a week. If the toxicity resolves to <Grade 2 within three weeks, the patient is treated at two dose levels lower than the dose that caused the toxicity (or a 50% reduction if the patient was treated at the first or second dose level). Patients in whom Grade 3 or 4 toxicity does not resolve to <Grade 1 within three weeks, or those who have another Grade 3 toxicity at the reduced dose are removed from the study.

Pharmacokinetic sampling is performed prior the first dose of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Day 1) and 0.5, 1, 2, 4, 6, 8, 24, and 48 hours

38

thereafter. Sampling is also conducted pre-dose on Days 7 and 21 and 0.5, 1, 2, 4, 6, 8, and 24 post-dose on Day 21 to evaluate steady-state 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione levels.

6.5.5 Treatment of Metastatic Melanoma

Patients with metastatic melanoma were started on 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revmid™) at 5 mg/day for seven days. The dose was then increased every seven days to 10 mg/day, 25 mg/day, and 50 mg/day, respectively, for a total of four weeks on therapy. Five of the 13 melanoma patients who were treated under this regimen either showed disease stabilization or a partial response in the first four weeks of treatment. Tumor response was seen in cutaneous and subcutaneous lesions (five patients), lymph nodes (two patients), and liver (one patient). The duration of response was approximately six months. The result suggests that the compound appears is a promising new anti-cancer agent and has both antiangiogenic and immunomodulatory properties.

6.5.6 Treatment of Relapsed or Refractory Multiple Myeloma


Patients with relapsed and refractory Dune-Salmon stage III multiple myeloma, who have either failed at least three previous regimens or presented with poor performance status, neutropenia or thrombocytopenia, are treated with up to four cycles of combination of melphalan (50 mg intravenously), an immunomodulatory compound of the invention (about 1 to 150 mg orally daily), and dexamethasone (40 mg/day orally on days 1 to 4) every four to six weeks. Maintenance treatment consisting of daily an immunomodulatory compound of the invention and monthly dexamethasone are continued until the disease progression. The therapy using an immunomodulatory compound of the invention in combination with melphalan and dexamethasone is highly active and generally tolerated in heavily pretreated multiple myeloma patients whose prognosis is otherwise poor.

The embodiments of the invention described above are intended to be merely exemplary, and those skilled in the art will recognize, or will be able to ascertain using no more than routine experimentation, numerous equivalents of specific compounds, materials, and procedures. All such equivalents are considered to be within the scope of the invention and are encompassed by the appended claims.

What is claimed is:

1. A method of treating multiple myeloma, which comprises administering, on a 28 day cycle, to a patient having multiple myeloma:

(a) about 25 mg per day of a compound of the formula:



or a pharmaceutically acceptable salt or solvate thereof, for 21 consecutive days followed by seven consecutive days of rest from administration of the compound, and;

(b) 40 mg per day of dexamethasone on days 1 through 4 in the 28 day cycle, wherein the patient has received previous therapy for multiple myeloma.

2. The method of claim 1, wherein the multiple myeloma is relapsed multiple myeloma.

US 8,530,498 B1


39

3. The method of claim 1, wherein the multiple myeloma is refractory multiple myeloma.

4. The method of claim 1, wherein the multiple myeloma is relapsed and refractory multiple myeloma.

5. The method of claim 1, wherein the patient has demonstrated disease progression on the previous therapy.

6. The method of claim 1, wherein the compound is



40

and is not a pharmaceutically acceptable salt or solvate thereof.

7. The method of claim 1, wherein the compound is a pharmaceutically acceptable salt.

8. The method of claim 1, wherein the compound is a pharmaceutically acceptable solvate.

9. The method of claim 1, wherein the compound is administered orally.

10. The method of claim 1, wherein the compound is administered in the form of a capsule.

11. The method of claim 1, wherein the compound is administered in the form of a tablet.

12. The method of claim 10, wherein the capsule comprises the compound, lactose anhydrous, microcrystalline cellulose, croscarmellose sodium and magnesium stearate.

13. The method of claim 1, wherein the dexamethasone is administered orally.

* * * * *

EXHIBIT E



US008648095B2

(12) **United States Patent**
Zeldis

(10) **Patent No.:** **US 8,648,095 B2**
(45) **Date of Patent:** ***Feb. 11, 2014**

(54) **METHODS FOR TREATING MULTIPLE MYELOMA USING 3-(4-AMINO-1-OXO-1,3-DIHYDROISOINDOL-2-YL)-PIPERIDINE-2,6-DIONE IN COMBINATION WITH PROTEASOME INHIBITOR**

(75) Inventor: **Jerome B. Zeldis**, Princeton, NJ (US)

(73) Assignee: **Celgene Corporation**, Summit, NJ (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **13/488,888**

(22) Filed: **Jun. 5, 2012**

(65) **Prior Publication Data**

US 2012/0244214 A1 Sep. 27, 2012

Related U.S. Application Data

(63) Continuation of application No. 12/640,702, filed on Dec. 17, 2009, now Pat. No. 8,198,306, which is a continuation of application No. 10/438,213, filed on May 15, 2003, now Pat. No. 7,968,569.

(60) Provisional application No. 60/380,842, filed on May 17, 2002, provisional application No. 60/424,600, filed on Nov. 6, 2002.

(51) **Int. Cl.**
A01N 43/40 (2006.01)

(52) **U.S. Cl.**
USPC **514/321**; 514/323

(58) **Field of Classification Search**
USPC 514/323
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,536,809 A 10/1970 Applezweig
3,598,123 A 8/1971 Zaffaroni et al.
3,845,770 A 11/1974 Theeuwes et al.
3,916,899 A 11/1975 Theeuwes et al.
4,008,719 A 2/1977 Theeuwes et al.
4,810,643 A 3/1989 Souza
4,999,291 A 3/1991 Souza
5,059,595 A 10/1991 Le Grazie
5,073,543 A 12/1991 Marshall et al.
5,120,548 A 6/1992 McClelland et al.
5,134,127 A 7/1992 Stella et al.
5,229,496 A 7/1993 Deeley et al.
5,354,556 A 10/1994 Sparks et al.
5,385,901 A 1/1995 Kaplan et al.
5,387,421 A * 2/1995 Amidon et al. 424/472
5,391,485 A 2/1995 Deeley et al.
5,393,870 A 2/1995 Deeley et al.
5,528,823 A 6/1996 Rudy, Jr. et al.
5,580,755 A 12/1996 Souza
5,591,767 A 1/1997 Mohr et al.
5,593,990 A 1/1997 D'Amato
5,629,327 A 5/1997 D'Amato

5,635,517 A * 6/1997 Muller et al. 514/323
5,639,476 A 6/1997 Oshlack et al.
5,674,533 A 10/1997 Santus et al.
5,698,579 A 12/1997 Muller
5,712,291 A 1/1998 D'Amato
5,731,325 A 3/1998 Andrulis, Jr. et al.
5,733,566 A 3/1998 Lewis
5,798,368 A 8/1998 Muller et al.
5,874,448 A 2/1999 Muller et al.
5,877,200 A 3/1999 Muller
5,929,117 A 7/1999 Muller et al.
5,955,476 A 9/1999 Muller et al.
6,020,358 A 2/2000 Muller et al.
6,071,948 A 6/2000 D'Amato
6,114,355 A 9/2000 D'Amato
6,140,346 A 10/2000 Andrulis, Jr. et al.
6,228,879 B1 5/2001 Green et al.
6,235,756 B1 5/2001 D'Amato
6,281,230 B1 8/2001 Muller et al.
6,316,471 B1 11/2001 Muller et al.
6,326,388 B1 12/2001 Man et al.
6,335,349 B1 1/2002 Muller et al.
6,380,239 B1 4/2002 Muller et al.
6,395,754 B1 5/2002 Muller et al.

(Continued)

FOREIGN PATENT DOCUMENTS

WO WO 92/14455 9/1992
WO WO 94/20085 9/1994

(Continued)

OTHER PUBLICATIONS

Elliot et al. ("The Proteasome: A New Target for Novel Drug Therapies". (2001) American Journal of Clinical Pathology, 116, 637-646).*

Bjorkstrand et al. (Leukemia&lymphoma (1994), 15, (3-4), 265-72).*

Lacy et al. (Journal of Clinical Oncology, 27(30): 5008-5014).*

(Continued)

Primary Examiner — Jeffrey S. Lundgren

Assistant Examiner — Chris Simmons

(74) *Attorney, Agent, or Firm* — Jones Day

(57) **ABSTRACT**

Methods of treating, preventing and/or managing cancer as well as and diseases and disorders associated with, or characterized by, undesired angiogenesis are disclosed. Specific methods encompass the administration of an immunomodulatory compound alone or in combination with a second active ingredient. The invention further relates to methods of reducing or avoiding adverse side effects associated with chemotherapy, radiation therapy, hormonal therapy, biological therapy or immunotherapy which comprise the administration of an immunomodulatory compound. Pharmaceutical compositions, single unit dosage forms, and kits suitable for use in methods of the invention are also disclosed.

US 8,648,095 B2

Page 2

(56)

References Cited

U.S. PATENT DOCUMENTS

| | | | | |
|--------------|-----|---------|-------------------|---------|
| 6,403,613 | B1 | 6/2002 | Man et al. | |
| 6,420,414 | B1 | 7/2002 | D'Amato | |
| 6,458,810 | B1 | 10/2002 | Muller et al. | |
| 6,469,045 | B1 | 10/2002 | D'Amato | |
| 6,476,052 | B1 | 11/2002 | Muller et al. | |
| 6,518,298 | B2 | 2/2003 | Green et al. | |
| 6,555,554 | B2 | 4/2003 | Muller et al. | |
| 6,673,828 | B1 | 1/2004 | Green et al. | |
| 7,323,479 | B2 | 1/2008 | Zeldis | |
| 7,393,862 | B2 | 7/2008 | Zeldis | |
| 7,435,745 | B2 | 10/2008 | D'Amato | |
| 7,468,363 | B2 | 12/2008 | Zeldis | |
| 7,968,569 | B2* | 6/2011 | Zeldis | 514/323 |
| 8,188,118 | B2* | 5/2012 | Zeldis | 514/321 |
| 2001/0018445 | A1 | 8/2001 | Huang et al. | |
| 2001/0056114 | A1 | 12/2001 | D'Amato | |
| 2002/0035090 | A1 | 3/2002 | Zeldis et al. | |
| 2002/0045643 | A1 | 4/2002 | Muller et al. | |
| 2002/0052398 | A1 | 5/2002 | D'Amato | |
| 2002/0054899 | A1 | 5/2002 | Zeldis | |
| 2002/0061923 | A1 | 5/2002 | D'Amato | |
| 2002/0128228 | A1 | 9/2002 | Hwu | |
| 2002/0161023 | A1 | 10/2002 | D'Amato | |
| 2002/0173658 | A1 | 11/2002 | Muller et al. | |
| 2002/0183360 | A1 | 12/2002 | Muller et al. | |
| 2003/0013739 | A1 | 1/2003 | Masferrer et al. | |
| 2003/0028028 | A1 | 2/2003 | Man et al. | |
| 2003/0045552 | A1 | 3/2003 | Robarge et al. | |
| 2003/0069428 | A1 | 4/2003 | Muller et al. | |
| 2003/0096841 | A1 | 5/2003 | Robarge et al. | |
| 2003/0139451 | A1 | 7/2003 | Shah et al. | |
| 2003/0144325 | A1 | 7/2003 | Muller et al. | |
| 2003/0181428 | A1 | 9/2003 | Green et al. | |
| 2003/0187024 | A1 | 10/2003 | D'Amato | |
| 2003/0191098 | A1 | 10/2003 | D'Amato | |
| 2003/0235909 | A1 | 12/2003 | Hariri et al. | |
| 2004/0029832 | A1 | 2/2004 | Zeldis | |
| 2004/0077685 | A1 | 4/2004 | Figg et al. | |
| 2004/0077686 | A1 | 4/2004 | Dannenberg et al. | |
| 2004/0087546 | A1 | 5/2004 | Zeldis | |
| 2004/0091455 | A1 | 5/2004 | Zeldis | |
| 2004/0122052 | A1 | 6/2004 | Muller et al. | |
| 2004/0266809 | A1 | 12/2004 | Emanuel et al. | |
| 2008/0145368 | A1 | 6/2008 | Zeldis | |
| 2008/0292583 | A1 | 11/2008 | Zeldis | |
| 2008/0317708 | A1 | 12/2008 | Zeldis | |
| 2009/0010877 | A1 | 1/2009 | Zeldis | |
| 2009/0123416 | A1 | 5/2009 | Zeldis | |
| 2010/0093683 | A1 | 4/2010 | Zeldis | |
| 2010/0196369 | A1 | 8/2010 | Zeldis | |
| 2010/0260719 | A1 | 10/2010 | Zeldis | |
| 2012/0035145 | A1 | 2/2012 | Zeldis | |
| 2012/0135042 | A1 | 5/2012 | Zeldis | |

FOREIGN PATENT DOCUMENTS

| | | |
|----|--------------|---------|
| WO | WO 98/03502 | 1/1998 |
| WO | WO 98/54170 | 12/1998 |
| WO | WO 01/70275 | 9/2001 |
| WO | WO 01/87307 | 11/2001 |
| WO | WO 02/15926 | 2/2002 |
| WO | WO 02/059106 | 8/2002 |
| WO | WO 02/064083 | 8/2002 |
| WO | WO 03/086373 | 10/2003 |

OTHER PUBLICATIONS

Canal et al. (Pathology Oncology Research, (1998), 4(3):171-178).*

Elliot et al. (American Journal of Clinical Pathology, (2001), 116, 637-646).*

Hideshima et al. (Seminars in Oncology, (2001), 28(6):607-612).*

Mitsiades et al. (Expert Opin. Investig. Drugs (2001); 10(8):1521-1530).*

Schlossman et al. (Current Opinion in Oncology, (Mar. 1999), 11(2):102-108).*

U.S. Appl. No. 60/499,723, Markian.

U.S. Appl. No. 60/372,348, Hariri et al.

U.S. Appl. No. 10/732,867, D'Amato et al.

U.S. Appl. No. 09/545,654, D'Amato.

U.S. Appl. No. 09/287,377, D'Amato.

PCT/US03/11578, PCT.

Carstensen, 1995, *Drug Stability: Principles & Practice*, 2nd. ed., Marcel Dekker, New York, NY pp. 379-380.

Corral et al., 1999, "Immunomodulation by thalidomide and thalidomide analogues," *Ann. Rheum. Dis.* 58(Suppl 1):1107-1113.

Craig et al., 1967, "Potential anticancer agents. III. 2-phthalimidoaldehydes and derivatives," *Potential Anticancer Agents III* 10:1071-1073.

D'Amato et al., 2001, "Mechanism of action of thalidomide and 3-aminothalidomide in multiple myeloma," *Semin. Oncol.* 28:597-601.

D'Amato et al., 1994, "Thalidomide is an Inhibitor of Angiogenesis", *Proc. Natl. Acad. Sci.* 91:4082-4085.

De et al., 1976, "Hansch analysis for some antineoplastic alutarimides," *J. Indian Chem. Soc. I.III:* 825-826.

De et al., 1976, "Possible antineoplastic agents: III. Synthesis of 6-alkyl-2-[4'-methoxyphthalimido] and 6-alkyl-3-[3'-4'-dimethoxyphenyl] glutarimides." *J. Indian Chem. Soc. I.III:*1122-1125.

Dredge et al., 2002, "Novel thalidomide analogues display anti-angiogenic activity independently of immunomodulatory effects," *Br. J. Cancer* 87 (10):1166-1172.

Folkman et al., 1983, "Angiogenesis inhibition and tumor regression caused by heparin or a heparin fragment in the presence of cortisone," *Science* 221(4612):719-725.

Gershbein, 1991, "The thalidomide analog, EM 12, enhances 1,2-dimethylhydrazine-induction of rat colon adenocarcinomas," *Cancer Letters* 60: 129-133.

Grabstald et al., 1965, "Clinical experiences with thalidomide in patients with cancer," *Clinical Pharmacology and Therapeutics* 6:298-302.

Lentzsch et al., 2003, "Immunomodulatory analogs of thalidomide inhibit growth of Hs Sultan cells and angiogenesis in vivo," *Leukemia* 17(1):41-44.

Lentzsch et al., 2002, "S-3-amino-phthalimido-glutarimide inhibits angiogenesis and growth of B-cell neoplasias in mice", *Cancer Research* 62:2300-2305.

Miyachi et al., 1997, "Novel biological response modifiers: phthalimides with tumor necrosis factor-alpha production-regulating activity," *J. Med. Chem.* 40:2858-2865.

Muller et al., 1999, "Amino-substituted thalidomide analogs: potent inhibitors of TNF-alpha production," *Bioorg. Med. Chem. Lett.* 9(11):1625-1630.

Muller et al., 1998, "Thalidomide analogs and PDE4 inhibition," *Bioorg. Med. Chem. Lett.* 8(19):2669-2674.

Muller et al., 1996, "Structural modifications of thalidomide produce analogs with enhanced tumor necrosis factor inhibitory activity," *J. Med. Chem.* 39(17):3238-3240.

Olson et al., 1965, "Thalidomide (N-phthaloylglutamide) in the treatment of advanced cancer," *Clinical Pharmacology and Therapeutics* 6(3):292-297.

Penichet et al., 2001, "Antibody-cytokine fusion proteins for the therapy of cancer," *J. Immunol. Methods* 248(1-2):91-101.

Physician's Desk Reference, 2002, 56th ed., pp. 1755-1760.

Raza et al., 2001, "Thalidomide produces transfusion independence in long-standing refractory anemias of patients with myelodysplastic syndromes," *Blood* 98(4):958-965.

Shah et al., 1999, "Synthesis and enantiomeric separation of 2-phthalimidino-glutaric acid analogues: potent inhibitors of tumor metastasis," *J. Med. Chem.* 42:3014-3017.

Shibata et al., 1995, "N-alkylphthalimides: structural requirement of thalidomide action on 12-O-tetradecanoylphorbol-13-acetate-induced tumor necrosis factor production by human leukemia HL-60 cells," *Chem. Pharm. Bull.* 43(1):177-179.

Shimazawa et al., 1999, "Antiangiogenic activity of tumor necrosis factor-alpha production regulators derived from thalidomide," *Biol. Pharm. Bull.* 22(2):224-226.

(56) **References Cited**

OTHER PUBLICATIONS

- Rubin et al., "Principles of Cancer Treatment-1", 12 ONCO IV May 1, 2003.
- Wilens et al., 1977, *Tetrahedron* 33:2725.
- Wilens, 1972, *Tables of Resolving Agents and Optical Resolutions*, E.L. Eliel, ed., Univ. of Notre Dame Press, Notre Dame, IN pp. 268.
- Wolff ed., 1995, *Burger's Medicinal Chemistry and Drug Discovery*, 5th ed., pp. 172-178, 949-982.
- N. Ake Jonsson. 1972, "Chemical Structure and Teratogenic Properties," *Acta Pharm.*, pp. 521-542.
- Alexanian et al., 2004, "VTD (Velcade, thalidomide, dexamethasone) as primary therapy for newly-diagnosed multiple myeloma," *Am. Soc. Hematol. 46th Ann. Meeting Dec. 4-7, 2004, San Diego, CA Abstract #210.*
- Anderson. 2000, "Thalidomide: Therapeutic potential in hematologic malignancies," *Seminars in Hematology* 37(1 Supp 3): 1-4.
- Attal et al., 2004, "Maintenance treatment with thalidomide after autologous transplantation for myeloma: First analysis of a prospective randomized study of the Intergroupe Francophone du Myelome (IFM 9902)," *Am. Soc. Hematol. 46th Ann. Meeting Dec. 4-7, 2004, San Diego, CA Abstract #535.*
- Bernardeschi et al., 2003, *J. Exp. Clin. Cancer Res.* 22(4):129-133.
- Corral et al., 1999, "Differential cytokine modulation and T cell activation by two distinct classes of thalidomide analogues that are potent inhibitors of TNF-alpha," *J. Immunol.* 163(1):380-386.
- Davies et al., 2001, "Thalidomide and immunomodulatory derivatives augment natural killer cell cytotoxicity in multiple myeloma," *Blood* 98(1):210-216.
- Dimopoulos et al., 2004, "Primary treatment with pulsed melphalan, dexamethasone, thalidomide (MDT) for symptomatic patients with multiple myeloma ≥ 75 years of age," *Am. Soc. Hematol. 46th Ann. Meeting Dec. 4-7, 2004, San Diego, CA Abstract #1482.*
- Eisen et al., 2000, "Continuous low dose Thalidomide: a phase II study in advanced melanoma, renal cell, ovarian and breast cancer," *Br. J. Cancer* 82(4):812-817.
- Fakhouri et al., 2004, "Thalidomide in patients with multiple myeloma and renal failure," *Br. J. Haematol.* 125:90-102.
- Fenk et al., 2005, "Single-agent thalidomide for treatment of first relapse following high-dose chemotherapy in patients with multiple myeloma," *Leukemia* 19(1):156-159.
- Gupta et al., 2001, "Adherence of multiple myeloma cells to bone marrow stromal cells upregulates vascular endothelial growth factor secretion: therapeutic applications," *Leukemia* 15(12):1950-1961.
- Haslett et al., 2003, "Thalidomide and a thalidomide analogue drug costimulate virus-specific CD8+ T cells in vitro." *J. Infect. Dis.* 187(6):946-955.
- Hideshima et al., 2000, "Thalidomide and its analogs overcome drug resistance of human multiple myeloma cells to conventional therapy," *Blood* 96(9):2943-2950.
- Offitani et al., 2003. Thalidomide plus oral melphalan for advanced multiple myeloma: a phase II study. *Haematologica.* Dec. 2003;88(12):1432-1433.
- Palumbo et al., 2004, "A prospective randomized trial of oral melphalan prednisone, thalidomide (MPT) vs. oral melphalan, prednisone (MP): An interim analysis," *Am. Soc. Hematol. 46th Ann. Meeting Dec. 4-7, 2004, San Diego, CA Abstract #207.*
- Raje et al., 1999, "Thalidomide—a revival story," *N. Engl. J. Med.* 341(21):1606-1609.
- Rajkumar et al., 2004, "Thalidomide plus dexamethasone versus dexamethasone alone in newly diagnosed multiple myeloma (EIA00): Results of a phase III trial coordinated by the Eastern Cooperative Oncology Group," *Am. Soc. Hematol. 46th Ann. Meeting Dec. 4-7, 2004, San Diego, CA Abstract #205.*
- Rajkumar et al., 2000, "Prognostic value of bone marrow angiogenesis in multiple myeloma," *Clin. Cancer Res.* 6(8):3111-3116.
- Ribatti et al., 1999, "Bone marrow angiogenesis and mast cell density increase simultaneously with progression of human multiple myeloma," *Br. J. Cancer* 79(3-4):451-455.
- Singhal et al., 1999, Antitumor activity of thalidomide in refractory multiple myeloma, *N. Engl. J. Med.* 341(21):1565-1571.
- Steins et al., 2002, "Efficacy and safety of thalidomide in patients with acute myeloid leukemia," *Blood* 99(3):834-839.
- Vacca et al., 1999, "Bone marrow neovascularization, plasma cell angiogenic potential, and matrix metalloproteinase-2 secretion parallel progression of human multiple myeloma," *Blood* 93(9):3064-3073.
- Wohrer et al., 2004, "Effective treatment of primary plasma cell leukemia with thalidomide and dexamethasone—a case report," *Hematol. J.* 5(4):361-363.
- Bach, 1963. "Thalidomide in Cancer Chemotherapy," *The Lancet*, No. 1271, p. 71.
- Bach, 1963, "Studies on the Possible Anti-Neoplastic Effect of Thalidomide," *Acta Pathologica Et Microbiologica Scandinavica* 59:491-499.
- Chaundhry, 1966, *Cancer Research*, "Effect of Prednisolone and Thalidomide on Induced Submandibular Gland Tumors in Hamster," 26(part 1):1884-86.
- DiPaolo, 1963, "Effect of Thalidomide on a Variety of Transplantable Tumors," *Cancer Chemotherapy Reports* No. 29, p. 99-102.
- DiPaolo, 1963, "In vitro Test Systems for Cancer Chemotherapy, II. Correlation of in vitro Inhibition of Dehydrogenase and Growth with in vivo Inhibition of Ehrlich Ascites Tumor," *Proceedings of the Society for Experimental Biology & Medicine*, 114:384-387.
- DiPaolo, 1964, "Thalidomide: Effects on Ehrlich Ascites Tumor Cells in vitro" *Science* 144:1583.
- Mauad, 1963, "Clinical Improvements Obtained in Advanced Cancer Patients with Treatment with Thalidomide Associated with Hormones," *Anais Paulistus de Medicina e Cirurgia* 86:13-40.
- Roe and Mitchley, 1963, "Thalidomide and Neoplasia," *Nature* 200:1016-1017.
- Liu et al., "Phase I study of CC-5013 (Revimid), a thalidomide derivative, in patients with refractory metastatic cancer," *American Society of Clinical Oncology*, Abstract #927, 2003.
- Zangari et al., "Results of phase I study of CC-5013 for the treatment of multiple myeloma (MM) patients who relapse after high dose chemotherapy (IIDCT)," *American Society of Hematology*, Abstract #3226, 2001.
- Zeldis et al., "Update on the evolution of the IMiD™," *International Society for Biological Therapy of Cancer*, Oral Abstract, 2003.
- Anderson "Moving disease biology from the laboratory to the clinic," *Seminars in Oncology*, 2002 29:17-20.
- Barlogie et al., "Total Therapy II (TTII) for newly diagnosed multiple myeloma (MM): preliminary data on feasibility and efficacy in the first 231 enrolled patients; comparison with predecessor total therapy I (TTI) (N-231)," *Blood*, Abstract #2857, Dec. 7-11, 2001, *American Society of Hematology*.
- Barlogie et al., "High-dose therapy immunomodulatory drugs in multiple myeloma," *Seminars in Oncology*, 2002, 29 (6):26-33.
- Barlogie et al., "Introduction: Thalidomide and the IMiDs in multiple myeloma," *Seminars in Hematology*, 2003, 40 (4):1-2.
- Barlogie, "Thalidomide and CC-5013 in Multiple Myeloma: The University of Arkansas experience," *Seminars in Hematology*, 2003, 40 (4):33-38.
- Bartlett et al., "The evolution of thalidomide and its IMiD derivatives as anticancer agents," *Nature Reviews Cancer*, 2004, 4 (4): 1-9.
- Bartlett et al., "Phase I study to determine the safety, tolerability and immunostimulatory activity of thalidomide analogue CC-5013 in patients with metastatic malignant melanoma and other advanced cancers," *British Journal of Cancer*, 2004, 90:955-961.
- Battegay, "Angiogenesis: mechanistic insights, neovascular diseases, and therapeutic prospects," *J. Mol. Med.*, 1995, 73:333-346.
- Baz et al., "Doxil (D), vincristine (V), reduced frequency dexamethasone (d) and revlimid (R) (DVD-R) results in a high response rate in patients with refractory multiple myeloma (RMM)," *Blood*, Abstract #2559, *American Society of Hematology*, Dec. 10-13, 2005.
- Brennen et al., "Thalidomide and analogues: current proposed mechanisms and therapeutic usage," *Clinical Prostate Cancer*, 2004, 3 (1):54-61.
- Celgene Corporation, "Celgene advances immunomodulatory drug (IMiD™) clinical program," Press Release, Feb. 2000.

(56)

References Cited

OTHER PUBLICATIONS

Celgene Corporation, "Initial Phase I solid tumor data on Celgene's lead IMiD™, Revimid™," Press Release, Jun. 2001.

Celgene Corporation, "Celgene Corporation receives orphan drug designation for Revimid™ for multiple myeloma," Press Release, Oct. 2001.

Celgene Corporation, "Celgene Corporation announces third quarter results. Thalomid® (thalidomide) sales increase 24%. Prescriptions up 50%. Enhanced S.T.E.P.S.® launched. Pilot d-MPH data presented," Press Release, Oct. 2001.

Celgene Corporation, "Celgene expands clinical development program for Revimid™. Five additional trials of Revimid initiated in hematological and solid tumor cancers," Press Release, Jun. 2002.

Celgene Corporation, "Celgene Corporation announces third quarter results. THALOMID® (thalidomide) revenue increases 41% to \$30.5 million. Pivotal programs for THALOMID and REVIMID™ finalized. Peer-reviewed publications of THALOMID and REVIMID data. First JNK inhibitor advanced to Phase I clinical trial," Press Release, Oct. 2002.

Celgene Corporation, "Blood reports Revimid™ has anti-tumor activity in patients with relapsed and refractory multiple myeloma," Press Release, Nov. 1, 2002.

Celgene Corporation, "Celgene provides update on clinical pipeline. Celgene Announces first target indication for ACTIMID™, CC-8490. SelCID™ program to advance based on results from Phase I/II trial of CC-1088. First JNK inhibitor successfully completes phase I trial," Press Release, Jan. 2003.

Celgene Corporation, "Celgene Corporation announces fourth quarter and full year results for 2002," Press Release, Jan. 2003.

Celgene Corporation, "Celgene receives fast track status from FDA for Revimid™ in multiple myeloma," Press Release, Feb. 2003.

Celgene Corporation, "Celgene receives fast track status from FDA for Revimid™ in myelodysplastic syndromes," Press Release, Apr. 2003.

Celgene Corporation, "New Revimid™ clinical data shows potential as novel approach to treating myelodysplastic syndromes (MDS)," Press Release, May 2003.

Celgene Corporation, "Celgene corporation reports strong operating performance in second quarter as total sales increase 100 percent and profits rise," Press Release, Jul. 2003.

Celgene Corporation, "Celgene corporation reports record operating performance in third quarter as total revenue increases 117% and profits rise," Press Release, Oct. 2003.

Celgene Corporation, "Celgene corporation advances ACTIMID™ (CC-4047) into phase II trial for prostate cancer," Press Release, Oct. 2003.

Celgene Corporation, "Additional clinical data presented on Revimid™ in myelodysplastic syndromes at the American Society of Hematology 45th annual meeting," Press Release, Dec. 2003.

Celgene Corporation, "Celgene corporation reviews 2003 achievements and announces 2004 financial outlook," Press Release, Jan. 2004.

Celgene Corporation, "Revimid™ receives orphan drug designation from the European commission for multiple myeloma," Press Release, Feb. 2004.

Celgene Corporation, "Revimid™ receives orphan drug designation from the European commission for myelodysplastic syndromes," Press Release, Mar. 2004.

Celgene Corporation, "Celgene corporation reports record operating performance in first quarter with strong revenue growth and profits," Press Release, Apr. 2004.

Celgene Corporation, "Celgene announces plans to stop phase III trials in melanoma due to lack of efficacy," Press Release, Apr. 2004.

Dalgleish, et al., "New thalidomide analogues; anti-cancer, anti-angiogenic and immunostimulatory," *British Journal of Cancer*, 2001, 85 (1)25.

Dalgleish et al., "Thalidomide analogues CC-5013 and CC-4047 induce T cell activation and IL-12 production in patients with both solid tumours and relapsed and refractory multiple myeloma," *British Journal of Cancer*, 2003, 88(Suppl 1), S25-S54.

Davies et al., "Thalidomide (Thal) and immunomodulatory derivatives (IMiDs) augment natural killer (NK) cell cytotoxicity in multiple myeloma (MM)," Abstract #3617, *American Society of Hematology*, Dec. 1-5, 2000.

Da Vies et al., "Thalidomide (Thal) and immunomodulatory derivatives (IMiDs) augment natural killer (NK) cell cytotoxicity in multiple myeloma (MM)," Abstract #P222, *VIIIth International Myeloma Workshop*, May 4-8, 2001.

Dibbs et al., "Thalidomide and thalidomide analogs suppress TNF α secretion by myocytes," Abstract #1284, *Circulation*, 1998.

Dimopoulos et al., "Results of thalidomide and IMiDs in multiple myeloma," Abstract #P12.1.4, *International Multiple Myeloma Workshop*, May 23-27, 2003.

Dimopoulos et al., "Treatment of plasma cell dyscrasias with thalidomide and its derivatives," *Journal of Clinical Oncology*, Dec. 1, 2003, 21 (23)4444-4454.

Dimopoulos et al., "Study of lenalidomide plus dexamethasone versus dexamethasone alone in relapsed or refractory multiple myeloma (MM): Results of a phase 3 Study (MM-010)," Abstract #6, *American Society of Hematology*, Dec. 10-13, 2005.

Dredge et al., "A costimulatory thalidomide analog enhances the partial anti-tumor immunity of an autologous vaccination in a model of colorectal cancer," Abstract #491, *American Association for Cancer Research*, Apr. 6-10, 2002.

Dredge et al., "Adjuvants and the promotion of Th1-type cytokines in tumour immunotherapy," *Cancer Immunol. Immunother.*, 2002, 51:521-531.

Dredge et al., "Immunological effects of thalidomide and its chemical and functional analogs," *Critical Reviews in Immunology*, 2002, 22 (5&6):425-437.

Dredge et al., "Protective antitumor immunity induced by a costimulatory thalidomide analog in conjunction with whole tumor cell vaccination is mediated by increased Th1-type immunity¹," *The Journal of Immunology*, 2002, 168:4914-4919.

Dredge et al., "Recent developments in antiangiogenic therapy," *Expert Opin. Biol. Ther.*, 2002, 2 (8):953-966.

Dredge et al., "Angiogenesis inhibitors in cancer therapy," *Current Opinion in Investigational Drugs*, 2003, 4 (6):667-674.

Dredge et al., "Thalidomide analogs as emerging anti-cancer drugs," *Anti-Cancer Drugs*, 2003, 14:331-335.

Fickentscher et al., "Stereochemical properties and teratogenic activity of some tetrahydrothalimides," *Molecular Pharmacology*, 1976, 13:133-141.

Figg et al., "Inhibition of angiogenesis: treatment options for patients with metastatic prostate cancer," *Investigational New Drugs*, 2002, 20(2):183-194.

Galustian et al., "Thalidomide-derived immunomodulatory drugs as therapeutic agents," *Expert Opin. Biol. Ther.*, 2004, 4 (12):1-8.

Glaspay et al., "The potential role of thalidomide and thalidomide analogs in melanoma," *Clinical Advances in Hematology & Oncology*, 2004, 1-7.

Gupta et al., "Adherence of multiple myeloma cells to bone marrow stromal cells upregulates vascular endothelial growth factor secretion; therapeutic applications," *Leukemia*, 2001, 15:1950-1961.

Hayashi et al., "Mechanisms whereby immunomodulatory analogs of thalidomide augment autologous NK cell anti-myeloma immunity," *Blood*, Abstract #3219, Dec. 6-10, 2002, *American Society of Hematology*.

He, W., et al., 1993, Abstract of papers, 206th American Chemical Society, Chicago, IL: Med. Chem., paper 216.

Helm et al., "Comparative teratological investigation of compounds of structurally and pharmacologically related to thalidomide," *Arzneimittel Forschung Drug Research*, 1981, 31 (1)941-949.

Hernandez-Illizaliturri et al., "Addition of immunomodulatory drugs CC5013 or CC4047 to rituximab enhances anti-tumor activity in a severe combined immunodeficiency (SCID) mouse lymphoma model," Abstract #235, *American Society of Hematology*, Dec. 6-9, 2003.

Hideshima et al., "Thalidomide and its analogs overcome drug resistance of human multiple myeloma cells to conventional therapy," *Blood*, 2000, 96:2943-2950, *American Society of Hematology*.

(56)

References Cited

OTHER PUBLICATIONS

- Hideshima et al., "Thalidomide (Thal) and its analogs overcome drug resistance of human multiple myeloma (MM) cells to conventional therapy," Abstract 1313, *American Society of Hematology*, Dec. 1-5, 2000.
- Hunt et al., "Markers of endothelial and haemostatic activation in the use of CC-4047, a structural analogue of thalidamide, in relapsed myeloma," *Blood*, Abstract #3216, Dec. 6-10, 2002, American Society of Hematology.
- Hussein et al., "Doxil (D), vincristine (V), reduced frequency dexamethasone (d) and Revlimid (DvD-R) a phase I/II trial in advanced relapsed/refractory multiple myeloma (Rmm) patients," *Blood*, Abstract #208, American Society of Hematology, Dec. 4-7, 2004.
- Hwu et al., "Thalidomide and its analogues in the treatment of metastatic melanoma," *Chemotherapy Foundation Symposium*, Abstract #44, 2002.
- Kyle, "Current therapy of multiple myeloma," *Internal Medicine*, 2002, 41 (3):175-180.
- Kyle et al., "Multiple myeloma," *New England Journal of Medicine*, 2004, 351:1860-1873.
- Leblanc et al., "Immunomodulatory drug costimulates T cells via the B7-CD28 pathway," *Blood*, 2004, 103:1787-1790, *American Society of Hematology*.
- Lentzsch et al., "In vivo activity of thalidomide and immunomodulatory drugs against multiple myeloma," *VIIIth International Myeloma Workshop*, Abstract #P225, May 4-8, 2001.
- Lentzsch et al., "Immunomodulatory derivative of thalidomide (IMiD CC-4047) determine the lineage commitment of hematopoietic progenitors by down regulation of GATA-I and modulation of cytokine secretion," Abstract #3073, *American Society of Hematology*, Dec. 6-9, 2003.
- Lentzsch et al., "Immunomodulatory derivative of thalidomide (IMiD CC-4047) down regulates CAAT/enhancer-binding protein β (C/EBP β) in multiple myeloma (MM)," Abstract #3456, *American Society of Hematology*, Dec. 6-9, 2003.
- Luzzio et al., "Thalidomide analogues: derivatives of an orphan drug with diverse biological activity," *Expert Opin. Ther. Patents*, 2004, 14 (2):215-229.
- Man et al., " α -Fluoro-substituted thalidomide analogues," *Bioorganic & Medicinal Chemistry Letters* 13, 2003, 3415-3417.
- Marriott et al., "Immunotherapeutic and antitumour potential of thalidomide analogues," *Expert Opin. Biol. Ther.*, 2001. 1(4):1-8.
- Marriott et al., "New thalidomide analogues; anti-cancer, anti-angiogenic and immunostimulatory," *British Journal of Cancer*, 85:25, Jul. 6, 2001.
- Marriott et al., "Thalidomide and its analogues have distinct and opposing effects on TNF- α and TNFR2 during co-stimulation of both CD4⁺ and CD8⁺ T cells," *Clin. Exp. Immunol.*, 2002, 130:75-84.
- Marriott et al., "A novel subclass of thalidomide analogue with anti-solid tumor activity in which caspase-dependent apoptosis is associated with altered expression of bcl-2 family proteins¹," *Cancer Research*, 2003, 63:593-599.
- Marriott et al., "Thalidomide derived immunomodulatory drugs (IMiDs) as potential therapeutic agents," *Current Drug Targets—Immune, Endocrine & Metabolic Disorders*, 2003 3:181-186.
- Masellis et al., "Changes in gene expression in bone marrow mesenchymal progenitor cells as a consequence of IMiD therapy in multiple myeloma patients," *Blood*, Abstract #1548, Dec. 7-11, 2001, American Society of Hematology.
- McCarty, "Thalidomide may impede cell migration in primates by down-regulating integrin β -chains: potential therapeutic utility in solid malignancies, proliferative retinopathy, inflammatory disorders, neointimal hyperplasia, and osteoporosis," *Medical Hypotheses*, 1997, 49:123-131.
- Mitsiades et al., "Apoptotic signaling induced by immunomodulatory thalidomide analogs (ImiDs) in human multiple myeloma cells: therapeutic implications," Abstract #3224, Dec. 7-11, 2001, *American Society of Hematology*.
- Mitsiades et al., "Apoptotic signaling induced by immunomodulatory thalidomide analogs in human multiple myeloma cells: therapeutic implications," *Blood*, 2002, 99:4525-4530, American Society of Hematology.
- Mitsiades et al., "CC-5013 Celgene," *Current Opinion in Investigational Drugs*, 2004, 5 (6):635-647.
- Moutouh et al., "Novel immunomodulatory drugs (IMiDs®): A potential, new therapy for β -hemoglobinopathies," Abstract #3740, *American Society of Hematology*, Dec. 4-7, 2004.
- Patten et al., "The early use of the serum free light chain assay in patients with relapsed refractory myeloma receiving treatment with a thalidomide analogue (CC-4047)," Abstract #1640, *American Society of Hematology*, Dec. 6-9, 2003.
- Payvandi et al., "Effects of a thalidomide analog on binding activity of transcription factors and cell cycle progression of multiple myeloma cell lines," *Blood*, Abstract #2487, Dec. 1-5, 2000, American Society of Hematology.
- Payvandi et al., "The thalidomide analogs IMiDs enhance expression of CD69 stimulatory receptor on natural killer cells," Abstract #1793, *American Association for Cancer Research*, Mar. 24-28, 2001.
- Payvandi et al., "Thalidomide analogs IMiDs inhibit expression of cyclooxygenase-2 in multiple myeloma cell line and LPS stimulated PBMCs," *Blood*, Abstract #2689, Dec. 7-11, 2001, American Society of Hematology.
- Payvandi et al., "Thalidomide and IMiDs inhibit microvessel formation from human arterial rings in the absence of human liver microsomes," *Blood*, Abstract #5046, Dec. 6-10, 2002, American Society of Hematology.
- Payvandi et al., "CC-5013 inhibits the expression of adhesion molecules ICAM-1 and CD44 and prevents metastasis of B16 F10 mouse melanoma cells in an animal model," *American Society of Clinical Oncology*, Abstract #992, 2003.
- Payvandi et al., "Immunomodulatory drugs inhibit expression of cyclooxygenase-2 from TNF- α , IL-1 β , and LPS-stimulated human PBMC in a partially IL-10-dependent manner," *Cellular Immunology*, 2004, 81-88.
- Raje et al., "Combination of the mTOR inhibitor rapamycin and CC-5013 has synergistic activity in multiple myeloma," *Blood*, Dec. 15, 2004, 104 (13):4188-4193.
- Rajkumar et al., "Combination therapy with lenalidomide plus dexamethasone (Rev/Dex) for newly diagnosed myeloma," *Blood*, Dec. 15, 2005, 106 (13):4050-4053.
- Richardson et al., "A Phase I study of oral CC5013, an immunomodulatory thalidomide (Thal) derivative, in patients with relapsed and refractory multiple myeloma (MM)," *Blood*, Abstract #3225, Dec. 7-11, 2001, American Society of Hematology.
- Richardson et al., "Immunomodulatory drug CC5013, overcomes drug resistance and is well tolerated in patients with relapsed multiple myeloma," *Blood*, 2002 100:3063-3067, American Society of Hematology.
- Richardson et al., "A multi-center, randomized, phase 2 study to evaluate the efficacy and safety of 2 CDC-5013 dose regimens when used alone or in combination with dexamethasone (Dex) for the treatment of relapsed or refractory multiple myeloma (MM)," *Blood*, Abstract #825, American Society of Hematology, Dec. 6-9, 2003.
- Richardson et al., "Immunomodulatory analogs of thalidomide: an emerging new therapy in myeloma," *Journal of Clinical Oncology*, 2004, 22(16) 3212-3214.
- Richardson et al., "A multicenter, single-arm, open-label study to evaluate the efficacy and safety of single-agent lenalidomide in patients with relapsed and refractory multiple myeloma: preliminary results," *10th International Myeloma Workshop*, Apr. 10-14, 2005.
- Richardson et al., "Novel biological therapies for the treatment of multiple myeloma," *Best Practice & Research Clinical Hematology*, 2005, 18 (4):619-634.
- Richardson et al., "A phase I trial of lenalidomide (REVLIMID®) with bortezomib (VELCADE®) in relapsed and refractory multiple myeloma," *Blood*, Abstract #365, American Society of Hematology, Dec. 10-13, 2005.
- Rubin et al., "Principles of cancer treatment-1," 2003, 12 ONCO IV I.

(56)

References Cited

OTHER PUBLICATIONS

- Schafer et al., "Enhancement of cytokine production and AP-1 transcriptional activity in T cells by thalidomide-related immunomodulatory drugs," *Journal of Pharmacology and Experimental Therapeutics*, 2003, 305(3):1222-1232.
- Schey et al., "A phase I study of an immunomodulatory thalidomide analog, CC-4047, in relapsed or refractory multiple myeloma," *Journal of Clinical Oncology*, 2004, 22 (16): 1-8.
- Schey et al., "A phase I study of an immunomodulatory thalidomide analogue (CC4047) in relapse/refractory multiple myeloma," *International Society for Experimental Hematology*, Abstract #248, 2002.
- Shaughnessy et al., "Global gene expression analysis shows loss of C-MYC and IL-6 receptor gene mRNA after exposure of myeloma to thalidomide and IMiD," Abstract #2485, *The American Society of Hematology*, Dec. 1-5, 2000.
- Shire et al., "TNF- α inhibitors and rheumatoid arthritis," *Exp. Opin. Ther. Patents*, 1998, 8 (5):531-544.
- Sorbera et al., "CC-5013. Treatment of multiple myeloma. Treatment of Melanoma. Treatment of myelodysplastic syndrome. Angiogenesis inhibitor. TNF- α production inhibitor," *Drugs of the Future*, 2003, 28(5):425-431.
- Streetly et al., "Thalidomide analogue CC-4047 is effective in the treatment of patients with relapsed and refractory multiple myeloma (MM) and induces T-cell activation and IL-12 production," Abstract #367, *International Multiple Myeloma Workshop*, May 23-27, 2003.
- Streetly et al., "Changes in neutrophil phenotype following the administration of CC-4047 (Actimid) to patients with multiple myeloma," Abstract #2543, *American Society of Hematology*, Dec. 6-9, 2003.
- Streetly et al., "An update of the use and outcomes of the new immunomodulatory agent CC-4047 (Actimid) in patients with relapsed/refractory myeloma," Abstract #829, *American Society of Hematology*, Dec. 6-9, 2003.
- Teo et al., "A phase I, single-blind, placebo-controlled, ascending single oral dose, safety, tolerability and pharmacokinetic study of CDC-501, a novel immunomodulatory-oncologic agent, in healthy male subjects with a comparison of fed and fasted," *Clinical Pharmacology and Therapeutics*, 2002, 71 (2):93.
- Teo et al., "Chiral inversion of the second generation IMiD™ CC-4047 (ACTIMID™) in human plasma and phosphate-buffered saline," *Chirality*, 2003, 15:348-351.
- Theertulien et al., "Hybrid MEL/DT PACE autotransplant regimen for Multiple Myeloma (MM)—safety and efficacy data in pilot study of 15 patients," *Blood*, Abstract #2869, American Society of Hematology, Dec. 7-11, 2001.
- Tohny et al., "A phase I study of oral CC-5013 (lenalidomide, Revlimid™), a thalidomide derivative, in patients with refractory metastatic cancer," *Clinical Prostate Cancer*, 2004, 2:241-243.
- Tricot et al., "Angiochemotherapy (ACT) for multiple myeloma (MM) with DT-PACE results in a high response rate, but in contrast to tandem transplants with melphalan does not affect durable disease control," *Blood*, Abstract #3531, American Society of Hematology, Dec. 7-11, 2001.
- Tsenova et al., "Use of IMiD3, a thalidomide analog, as an adjunct to therapy for experimental tuberculous meningitis," *Antimicrobial Agents and Chemotherapy*, 2002, 46 (6):1887-1895.
- Weber, "Lenalidomide (CC-5013, Revlimid™) and other ImiDs," Abstract #PL5.02, *International Multiple Myeloma Workshop*, Apr. 10-14, 2005.
- Weber et al., "A multicenter, randomized, parallel-group, double-blind, placebo-controlled study of lenalidomide plus dexamethasone versus dexamethasone alone in previously treated subjects with multiple myeloma," Abstract #PO.738, *International Multiple Myeloma Workshop*, Apr. 10-14, 2005.
- Ye et al., "Novel IMiD drugs enhance expansion and regulate differentiation of human cord blood CD34+ cells with cytokines," *Blood*, Abstract #4099, American Society of Hematology, Dec. 6-10, 2002.
- Zangari et al., "Risk factors for deep vein thrombosis (DVT) in a large group of myeloma patients (Pts) treated with thalidomide (Thal): The Arkansas Experience," *Blood*, Abstract #681, American Society of Hematology, Dec. 7-11, 2001.
- Zangari et al., "Revimid 25 mg (REV 25) x 20 versus 50 mg (REV 50) x 10 q 28 days with bridging of 5 mg x 10 versus 10 mg x 5 as post-transplant salvage therapy for multiple myeloma (MM)," *Blood*, Abstract #1642, American Society of Hematology, Dec. 6-9, 2003.
- Zeldis et al., "Potential new therapeutics for Waldenstrom's macroglobulinemia," *Seminars in Oncology*, 2003, 30 (2):275-281.
- Zhang et al., "CC-5079, a novel microtubule and TNF- α inhibitor with anti-angiogenic and antimetastasis activity," Abstract #B012, *International Conference on Molecular Targets and Cancer Therapeutics*, Nov. 17-21, 2003.
- Anderson, "The Role of Immunomodulatory Drugs in Multiple Myeloma," *Seminars in Hematology*, vol. 40, No. 4, Suppl 4, 2003: pp. 23-32.
- Weber, "Thalidomide and Its Derivatives: New Promise for Multiple Myeloma," *Cancer Control*, vol. 10, No. 5, 375-383, 2003.
- Patt, Yehuda A.; Hassan, Manal M.; Lozano, Richard D.; Ellis, Lee M.; Peterson, J. Andrew; Waugh, Kimberly A.; *Durable Clinical Response of Refractory Hepatocellular Carcinoma to Orally Administered Thalidomide*. American Journal of Clinical Oncology, 2000.
- Richardson, Paul; Hideshima, Teru; Anderson, Kenneth; *Thalidomide: The Revival of a Drug with Therapeutic Promise in the Treatment of Cancer*; Principles & Practice of Oncology, vol. 15, No. 2, 2001.
- Thomas, Melodie; Doss, Deborah, *Thalidomide Nursing Roundtable Update*, Monograph, Sep. 2002.
- Richardson, Paul; Hideshima, Teru; Anderson, Kenneth; *Thalidomide: Emerging Role in Cancer Medicine*; Annual Review of Medicine, 2002.
- Berenson, J.R.; Bergsagel, P. L.; Munshi, N.; *Initiation and Maintenance of Multiple Myeloma*; Seminars in Hematology, vol. 36, No. 1, Supp. 3, Jan. 1999, pp. 9-13.
- Gollob, J.A.; Schinpper, C.P.; Orsini, E.; Murphy, E.; Daley, J.F.; Lazo, S.B.; Frank, D.A.; *Characterization of a Novel Subset of CD8 T Cells That Expands in patients Receiving Interleukin-12*, 02, Am. Soc. For Clin. Investigation, Inc., vol. 102, No. 3, Aug. 1998, pp. 561-575.
- Cavanagh, L.L.; Barnetson, R.S.; Basten, A.; Halliday, G.M.; *Dendritic Epidermal T-Cell Involvement in Induction of CD8+ T-Cell-Mediated Immunity Against an Ultraviolet Radiation-Induced Skin Tumor* Int. J. Cancer: 70, 98-105, 1997.
- Thomas, D.A., Aguayo, A., Estey, E., Albitar, M., O'Brien, S., Giles, F.J., Beran, M., Cortes, J., Zeldis, J., Keating, M.J., Barlogie, B., Kantarjian, H.M., *Thalidomide as anti-angiogenesis therapy (rx) in refractory or relapsed leukemia*. Abstract #2269, American Society of Hematology, Dec. 3-7, 1999.
- Barlogie, B., Desikan, R., Munshi, N., Siegel, D., Mehta, J., Singhal, S., Anaissie, E., *Single Course D.T. Pace Anti-Angiochemotherapy Effects CR in Plasma Cell Leukemia and Fulminant Multiple Myeloma (MM)*. Abstract #4180. American Society of Hematology, Dec. 4-9, 1998.
- Hideshima, T., Chauhan, D., Shima, Y., Noopur, R., Davies, F.E., Tai, Y., Treon, S.P., Lin, B.K., Schlossman, R.L., Richardson, P.G., Gupta, D., Muller, G.W., Stirling, D.I., Anderson, K.C., *Thalidomide (THAL) and its Analogs Overcome Drug Resistance of Human Multiple Myeloma (MM) Cells to Conventional Therapy*. Abstract #1313. American Society of Hematology, Dec. 1-5, 2000.
- Payvandi, F., Wu, L., Gupta, D., Hideshima, T., Haley, M., Muller, G., Chen, R., Anderson, K.C., Stirling, D., *Effects of a Thalidomide Analog on Binding Activity of Transcription Factors and Cell Cycle Progression of Multiple Myeloma Cell Lines*. Abstract #2487. American Society of Hematology, Dec. 1-5, 2000.
- Davies, F.E., Raju, N., Hideshima, T., Lentzsch, S., Young, G., Tai, Y., Lin, B.K., Podar, K., Chauhan, D., Treon, S.P., Gupta, D., Mitsiades, C., Mitsiades, N., Hayashi, T., Richardson, P.G., Schlossman, R.L., Muller, G.W., Stirling, D. I., Anderson, K.C., *Thalidomide (THAL) and Immunomodulatory Derivatives (IMiDs) Augment Natural Killer (NK) Cell Cytotoxicity in Multiple Myeloma (MM)*, Abstract #3617, American Society of Hematology, Dec. 1-5, 2000.

(56)

References Cited

OTHER PUBLICATIONS

- Hideshima, T., Chauhan, D., Castro, A., Hayashi, T., Mitsiades, C., Mitsiades, N., Akiyama, M., Richardson, P.G., Schlossman, R.L., Adams, J., Anderson, K.C., NF-KB as a Therapeutic Target in Multiple Myeloma (MM). Abstract #1581. American Society of Hematology, Dec. 7-11, 2001.
- Lentsch, S., Rogers, M., LeBlanc, R., Birsner, A., Shah, J., Anderson K., D'Amato R., 3-Amino-Phthalimido-Glutarimide (S-3APG) Inhibits Angiogenesis and Growth in Drug Resistant Multiple Myeloma (MM) in vivo. Abstract #1976, American Society of Hematology, Dec. 7-11, 2001.
- Park, Y., Kim, S.A., Kim, C.J., Chung, J.H., Mechanism of the Effect of Thalidomide on Human Multiple Myeloma Cells. Abstract #2685. American Society of Clinical Oncology, May 12-17, 2001.
- Payvandi, F., Wu, L., Haley M., Gupta, D., Zhang, L., Schafer, P., Muller, G. W., Chen, R., Anderson, K.C., Stirling, D., Thalidomide Analogs IMiDS Inhibit Expression of Cyclooxygenase-2 in Multiple Myeloma Cell Line and LPS Stimulated PBMCs. Abstract #2689. American Society of Hematology, Dec. 7-11, 2001.
- Mitsiades, N., Mitsiades, C., Poulaki, V., Akiyama, M., Tai, Y., Lin, B., Hayashi, T., Catley, L., Hideshima, T., Chauhan, D., Treon, S.P., Anderson, K.C., Apoptotic Signaling Induced by Immunomodulatory Thalidomide Analogs (Imids) in Human Multiple Myeloma Cells; Therapeutic Implications. Abstract #3224. American Society of Hematology, Dec. 7-11, 2001.
- Richardson, P.G., Schlossman, R.L. Hideshima, T., Davies, F., LeBlanc, R., Catley, L., Doss, D., Kelly, K.A., McKenney, M., Mechlowicz, J., Freeman, A., Deocampo, R., Rich, R., Ryoo, J., Chauhan, D., Munshi, N., Weller, E., ZELD1S, J., Anderson, K.C., A Phase I Study of Oral CC5013, an Immunomodulatory Thalidomide (Thal) Derivative, in Patients With Relapsed and Refractory Multiple Myeloma (MM). Abstract #3225. American Society of Hematology, Dec. 7-11, 2001.
- "Celgene drug promises activity in solid tumors," Marketletter, Jun. 18, 2001.
- Meregalli et al., "High-dose dexamethasone as first line therapy of multiple myeloma?," *Recenti Progressi in Medicina*, 1998, 89(1):18-20.
- Official Action in corresponding Canadian Application No. 2,476,983 dated Aug. 21, 2009.
- List, A., "New Approaches to the Treatment of Myelodysplasia," *The Oncologist*, 2002, 7(suppl. 1):39-49.
- Kurzrock, R., "Myelodysplastic syndrome overview," *Seminars in Hematology* (Abstract only), 2002, 39(3)(suppl. 2):18-25 Abstract only.
- Goerner, et al., "Morbidity and mortality of chronic GVHD after hematopoietic stem cell transplantation from HLA-identical siblings for patients with aplastic or refractory anemias," *Biology of Blood and Marrow Transplantation* (Abstract only), 2002, 8(1):47-56.
- Thomas, D., "Pilot studies of Thalidomide in Acute Myelogenous Leukemia, Myelodysplastic Syndromes, and Myeloproliferative Disorders," *Seminars in Hematology*, 2000, 37(1)(suppl. 3):26-34.
- Zorat, F. et al., "The clinical and biological effects of thalidomide in patients with myelodysplastic syndromes," *British Journal of Haematology*, 2001, 115:881-894.
- Official Action dated Feb. 10, 2009 in JP Application No. 2004-545192. (English translation provided.).
- Teramura, M., Men-ekiyokusei Ryouhou, *Current Therapy*, 2000, 18(5):140-144 (in Japanese).
- Kon-nichi no Chiryou Shishin, 1997 [Pocket Edition], Igaku Shoin, 1997, 513-514 (in Japanese).
- Okamoto, T., Kotsuzuikeisei Shoukougun to Men-eki Ijo, Bessatsu Nihon Rinsho, Syndrome Series for each area, No. 22, Blood Syndromes III, Nihon Rinsho, 213-216 (in Japanese), Oct. 1998.
- Merck Manual, 17th ed. Japanese version, 1999, 951-952.
- Notice of Allowance from U.S. Appl. No. 11/096,155 dated Jan. 12, 2010.
- Rajkumar et al., "Combination therapy with thalidomide plus dexamethasone for newly diagnosed multiple myeloma," *American Society of Hematology*, 43rd Annual Meeting, Dec. 7-11, 2001, Abstract #3525.
- Scheffler et al., "Safety and pharmacokinetics of CDC-501, a novel immunomodulatory-oncologic agent, after single then multiple, oral 100 mg twice daily doses," *American Society for Clinical Pharmacology and Therapeutics*, Mar. 24-27, 2002, Abstract #WP111-63.
- Marriott et al., "Thalidomide analogue CDC-501 is safe and well tolerated by patients with end stage cancer and shows evidence of clinical responses and extensive immune activation," *Br. J. Cancer*, 2002, 86(Suppl. 1):Abst 6.4.
- Kant, R.E., "Evidence of a mechanism by which etanercept increased TNF-alpha in multiple myeloma: New insights into the biology of TNF-alpha giving new treatment opportunities—the role of burproion" *Leukemia Research*, 2005, 29:1459-1463.
- Tsimberidou, A. et al., "Pilot study of recombinant human soluble tumor necrosis factor (TNF) receptor (p75) fusion protein (TNFR:Fc;Enbrel) in patients with refractory multiple myeloma: increase in plasma TNF α levels during treatment," *Leukemia Research*, 2003, 27:375-380.
- Dimopoulos, et al., "Long-term follow-up on overall survival from the MM-009 and MM-010 phase III trials of lenalidomide plus dexamethasone in patients with relapsed or refractory multiple myeloma," *Leukemia*, 2009, 1-6.
- Hideshima, T., et al., "A review of lenalidomide in combination with dexamethasone for the treatment of multiple myeloma," *Therapeutics and Clinical Risk Management*, 2008, 4(1):129-136.
- Wang, M., et al., "Lenalidomide plus dexamethasone is more effective than dexamethasone alone in patients with relapsed or refractory multiple myeloma regardless of prior thalidomide exposure," *Blood*, 2008, 112(12):4445-4451.
- Gandhi, A., et al., "Dexamethasone Synergizes with Lenalidomide to Inhibit Multiple Myeloma Tumor Growth, But Reduces Lenalidomide-Induced Immunomodulation of T and NK Cell Function," *Current Cancer Drug Targets*, 2010, 10(1):1-13.
- Gay, E. et al., "Lenalidomide plus dexamethasone versus thalidomide plus dexamethasone in newly diagnosed multiple myeloma: a comparative analysis of 411 patients," *Blood*, 2010, 115(97):1343-150.
- Richardson, P. et al., "Thalidomide in multiple myeloma," *Biomed Pharmacother*, 2002, 56:115-28.
- Swartz, G. et al., "Pre-clinical evaluation of ENMD-0995: A thalidomide analog with activity against multiple myeloma and solid tumors," *Cell and Tumor Biology*, 2002, 43:181-182, Abstract#910.
- Mazucco, R., "Angiogenesis and Anti-angiogenesis Therapeutics," *IDrugs*, 2002, 5(4): 320-322.
- Worker, C., "JP Morgan Hambreeht & Quist—20th Annual Healthcare Conference," *IDrugs*, 2002, 5(2):113-116.
- Treston, A. et al., "Pre-Clinical Evaluation of a Thalidomide Analog with Activity Against Multiple Myeloma and Solid Tumors—ENMD-0995 (S-(-)-3-(3-amino-phthalimido)-glutarimide)," *Blood*, 2002, 100(11):816a, Abstract #3225.
- Mazucco, R. and Williams, L., "Immunotherapy, chemoprevention and angiogenesis," *IDrugs*, 2002, 5(5):408-411.
- Fernandes, P., "Anti-Cancer Drug Discovery and Development Summit," *IDrugs*, 2002, 5(8):757-764.
- Notification letter dated Aug. 30, 2010 from Natco Pharma Limited to Celgene Corporation re: Notification pursuant to § 505(j)(2)(B) of the Federal Food, Drug and Cosmetic Act.
- Complaint for Patent Infringement filed on Oct. 8, 2010 by Celgene Corporation in the U.S. District Court, District of New Jersey against Natco Pharma Limited.
- Answer to Complaint filed on Nov. 18, 2010 by Natco Pharma Limited in the U.S. District Court, District of New Jersey.
- Grosshans, E. and Illy, G., "Thalidomide Therapy for Inflammatory Dermatoses," *International Journal of Dermatology*, 1984, 23(9):598-602.
- Krenn, M. et al., "Improvements in Solubility and Stability of Thalidomide upon Complexation with Hydropropyl- β -Cyclodextrin," *Journal of Pharmaceutical Sciences*, 1992, 81(7):685-689.
- Schmahl, H. J. et al., "Pharmacokinetics of the Teratogenic and Nonteratogenic Thalidomide Analogs EM 12 and Supidimide in the

(56)



References Cited

OTHER PUBLICATIONS

- Rat and Marmoset Monkey”, in *Pharmacokinetics in Teratogenesis*, CRC Press, 1987, vol. I, Ch. 12, pp. 181-192.
- Schumacher, H. et al., “The Teratogenic Activity of Thalidomide Analogue, EM₁₂, in Rabbits, Rats, and Monkeys,” *Teratology*, 1971, 5:233-240.
- Smith, R. et al., “Studies on the Relationship Between the Chemical Structure and Embryotoxic Activity of Thalidomide and Related Compounds,” in *A Symposium on Embryopathic Activity of Drugs*, J. & A. Churchill Ltd., 1965, Session 6, pp. 194-209.
- Sheskin, J. and Sagher, F., “Trials with Thalidomide Derivatives in Leprosy Reactions,” *Leprosy Review*, 1968, 39(4):203-205.
- Sheskin, J., “Study with Nine Thalidomide Derivatives in the Lepra Reaction,” *Pharmacology and Therapeutics*, 1978, 17:82-84.
- Raje, N. and Anderson, K., “Thalidomide and immunomodulatory drugs as cancer therapy,” *Current Opinions in Oncology*, 2002, 14:635-640.
- Kumar, S. et al., “Thalidomide as an anti-cancer agent,” *J Cell. Mod. Med.*, 2002, 6(2):160-174.
- Singhal, S. and Mehta, J., “Thalidomide in Cancer,” *BioDrugs*, 2001, 15(3):163-172.
- Notice of Opposition to EP 1 505 973 filed by Synthron B.V. on Nov. 30, 2010.
- Notice of Opposition to EP 1 505 973 filed by Strawman Limited on Dec. 1, 2010.
- Samson, D. et al., “Infusion of Vincristine and Doxorubicin with Oral Dexamethasone as First-Line Therapy for Multiple Myeloma.” *The Lancet*, 1989, 334(8668):882-885.
- Barlogie, B. et al., “Effective Treatment of Advanced Multiple Myeloma Refractory to Alkylating Agents,” *N. Engl. J. Med.*, 1984, 310(21):1353-1356.
- Dimopoulos, M. et al., “Thalidomide and dexamethasone combination for refractory multiple myeloma,” *Annals of Oncology*, 2001, 12:991-995.
- Zangari, M., et al., “Thrombogenic activity of doxorubicin in myeloma patients receiving thalidomide: implications for therapy,” *Blood*, 2002, 100:1168-1171.
- List, A. et al., “High Erythropoietic Remitting Activity of the Immunomodulatory Thalidomide Analog, CC5013, in Patients with Myelodysplastic Syndrome (MDS);” Abstract #353, *Blood*, 2002, 100(11):96a.
- Mufti, G. et al., “Myelodysplastic Syndrome,” *American Society of Hematology*, 2003, pp. 176-199.
- Extracts from drug databases: retrieved from <http://www.nextbio.com/b/search/ov/IMI3%20cpd> on Nov. 26, 2010 and <http://pubchem.ncbi.nlm.nih.gov/summary.cgi?cid=216326> on Nov. 26, 2010.
- Stockdale, 1998, *Medicine*, Rubenstein and Federman, eds., vol. 3, Ch. 12, Sections IV and X.
- “List of Approved Oncology Drugs with Approved Indications,” <http://www.accessdata.fda.gov/scripts/cder/onctools/druglist.cfm>.
- Office Action mailed Jun. 15, 2008, U.S. Appl. No. 11/325,954.
- Gamberi et al., “Overall Safety and Treatment Duration in Lenalidomide (LEN)-, Thalidomide (THAL)-, and Bortezomib (BORT)-Treated Patients (Pts) within the European Post-Approval Safety Study (EU PASS) of Relapsed/Refractory Multiple Myeloma (RRM)”, presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #4068.
- Korde et al., “Phase II Clinical and Correlative Study of Carfilzomib, Lenalidomide, and Dexamethasone (CRd) in Newly Diagnosed Multiple Myeloma (MM) Patients”, presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #732.
- Kumar et al., “A Phase 1/2 Study of Weekly MLN9708, an Investigational Oral Proteasome Inhibitor, in Combination with Lenalidomide and Dexamethasone in Patients with Previously Untreated Multiple Myeloma (MM)”, presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #332.
- Palumbo et al., “Pomalidomide Cyclophosphamide and Prednisone (PCP) Treatment for Relapsed/Refractory Multiple Myeloma”, presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #446.
- Richardson et al., “A Phase 2 Study of Elotuzumab (Elo) in Combination with Lenalidomide and Low-Dose Dexamethasone (Ld) in Patients (pts) with Relapsed/Refractory Multiple Myeloma (R/R MM): Updated Results”, presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #202.
- Sacchi et al., “A Phase I/II Study of Bendamustine, Low-Dose Dexamethasone, and Lenalidomide (BdL) for the Treatment of Patients with Relapsed Multiple Myeloma”, presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #1851.
- Sonneveld et al., “Escalated Dose Bortezomib Once Weekly Combined with Lenalidomide and Dexamethasone (eVRD) Followed by Lenalidomide Maintenance in First Relapse of Multiple Myeloma (MM), the HOVON 86 Phase 2 Trial”, presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #1853.
- Suvannasankha et al., “A Phase I/II Trial Combining High-Dose Lenalidomide with Melphalan and Autologous Transplant for Multiple Myeloma: A Report of the Phase I Dose-Finding Study”, presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #3146.
- Office Action in corresponding CN Application No. 201110256752.0 dated Feb. 8, 2013.
- Stirling, D., “Thalidomide: A Novel Template for Anticancer Drugs,” *Seminars in Oncology*, Dec. 2001, 28(6):602-606.
- Adams, J. et al., “Proteasome Inhibitors: A Novel Class of Potent and Effective Antitumor Agents,” *Cancer Res*, 1999, 59:2615-2622.
- Hideshima, T. et al., “The Proteasome Inhibitor PS-341 Inhibits Growth, Induces Apoptosis, and Overcomes Drug Resistance in Human Multiple Myeloma Cells,” *Cancer Research*, Apr. 2001, 61:3071-3076.
- Elofsson, M. et al., “Towards subunit-specific proteasome inhibitors: synthesis and evaluation of peptide α' , β' -epoxyketones,” *Chemistry & Biology*, Nov. 1999, 6:811-822.
- Celgene Press Release, “Celgene Will Discontinue Phase III ORIGIN® Trial in Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukemia,” published on Celgene Newsroom, <http://newsroom.celgene.com> on Jul. 18, 2013 at 7:30 am EDT.
- Mateos, M.-V., Ph.D. et al., “Lenalidomide plus Dexamethasone for High-Risk Smoldering Multiple Myeloma,” *New England Journal of Medicine*, Aug. 2013, 369(5):438-447.
- English translation of Japanese IP High Court decision in Application No. JP 2004-505051, dated Apr. 11, 2013.

* cited by examiner

Effects of Revimid™ and Thalidomide on MM Cell Proliferation



US 8,648,095 B2

1

**METHODS FOR TREATING MULTIPLE
MYELOMA USING 3-(4-AMINO-1-OXO-1,3-
DIHYDROISOINDOL-2-YL)-PIPERIDINE-2,6-
DIONE IN COMBINATION WITH
PROTEASOME INHIBITOR**

This application is a continuation of U.S. patent application Ser. No. 12/640,702, filed Dec. 17, 2009, now U.S. Pat. No. 8,198,306, which is a continuation application of U.S. patent application Ser. No. 10/438,213, filed May 15, 2003, now U.S. Pat. No. 7,968,569, issued Jun. 28, 2011, which claims the benefit of U.S. provisional application Nos. 60/380,842, filed May 17, 2002, and 60/424,600, filed Nov. 6, 2002, the entireties of which are incorporated herein by reference.

1. FIELD OF THE INVENTION

This invention relates to methods of treating, preventing and/or managing specific cancers, and other diseases including, but not limited to, those associated with, or characterized by, undesired angiogenesis, by the administration of one or more immunomodulatory compounds alone or in combination with other therapeutics. In particular, the invention encompasses the use of specific combinations, or "cocktails," of drugs and other therapy, e.g., radiation to treat these specific cancers, including those refractory to conventional therapy. The invention also relates to pharmaceutical compositions and dosing regimens.

2. BACKGROUND OF THE INVENTION

2.1 Pathobiology of Cancer and Other Diseases

Cancer is characterized primarily by an increase in the number of abnormal cells derived from a given normal tissue, invasion of adjacent tissues by these abnormal cells, or lymphatic or blood-borne spread of malignant cells to regional lymph nodes and to distant sites (metastasis). Clinical data and molecular biologic studies indicate that cancer is a multistep process that begins with minor preneoplastic changes, which may under certain conditions progress to neoplasia. The neoplastic lesion may evolve clonally and develop an increasing capacity for invasion, growth, metastasis, and heterogeneity, especially under conditions in which the neoplastic cells escape the host's immune surveillance. Roitt, I., Brostoff, J and Kale, D., *Immunology*, 17.1-17.12 (3rd ed., Mosby, St. Louis, Mo., 1993).

There is an enormous variety of cancers which are described in detail in the medical literature. Examples includes cancer of the lung, colon, rectum, prostate, breast, brain, and intestine. The incidence of cancer continues to climb as the general population ages, as new cancers develop, and as susceptible populations (e.g., people infected with AIDS or excessively exposed to sunlight) grow. A tremendous demand therefore exists for new methods and compositions that can be used to treat patients with cancer.

Many types of cancers are associated with new blood vessel formation, a process known as angiogenesis. Several of the mechanisms involved in tumor-induced angiogenesis have been elucidated. The most direct of these mechanisms is the secretion by the tumor cells of cytokines with angiogenic properties. Examples of these cytokines include acidic and basic fibroblastic growth factor (a,b-FGF), angiogenin, vascular endothelial growth factor (VEGF), and TNF- α . Alternatively, tumor cells can release angiogenic peptides through the production of proteases and the subsequent breakdown of the extracellular matrix where some cytokines are stored

2

(e.g., b-FGF). Angiogenesis can also be induced indirectly through the recruitment of inflammatory cells (particularly macrophages) and their subsequent release of angiogenic cytokines (e.g., TNF- α , bFGF).

A variety of other diseases and disorders are also associated with, or characterized by, undesired angiogenesis. For example, enhanced or unregulated angiogenesis has been implicated in a number of diseases and medical conditions including, but not limited to, ocular neovascular diseases, choroidal neovascular diseases, retina neovascular diseases, rubeosis (neovascularization of the angle), viral diseases, genetic diseases, inflammatory diseases, allergic diseases, and autoimmune diseases. Examples of such diseases and conditions include, but are not limited to: diabetic retinopathy; retinopathy of prematurity; corneal graft rejection; neovascular glaucoma; retrolental fibroplasia; and proliferative vitreoretinopathy.

Accordingly, compounds that can control angiogenesis or inhibit the production of certain cytokines, including TNF- α , may be useful in the treatment and prevention of various diseases and conditions.

2.2 Methods of Treating Cancer

Current cancer therapy may involve surgery, chemotherapy, hormonal therapy and/or radiation treatment to eradicate neoplastic cells in a patient (see, for example, Stockdale, 1998, *Medicine*, vol. 3, Rubenstein and Federman, eds., Chapter 12, Section IV). Recently, cancer therapy could also involve biological therapy or immunotherapy. All of these approaches pose significant drawbacks for the patient. Surgery, for example, may be contraindicated due to the health of a patient or may be unacceptable to the patient. Additionally, surgery may not completely remove neoplastic tissue. Radiation therapy is only effective when the neoplastic tissue exhibits a higher sensitivity to radiation than normal tissue. Radiation therapy can also often elicit serious side effects. Hormonal therapy is rarely given as a single agent. Although hormonal therapy can be effective, it is often used to prevent or delay recurrence of cancer after other treatments have removed the majority of cancer cells. Biological therapies and immunotherapies are limited in number and may produce side effects such as rashes or swellings, flu-like symptoms, including fever, chills and fatigue, digestive tract problems or allergic reactions.

With respect to chemotherapy, there are a variety of chemotherapeutic agents available for treatment of cancer. A majority of cancer chemotherapeutics act by inhibiting DNA synthesis, either directly, or indirectly by inhibiting the biosynthesis of deoxyribonucleotide triphosphate precursors, to prevent DNA replication and concomitant cell division. Gilman et al., *Goodman and Gilman's: The Pharmacological Basis of Therapeutics*, Tenth Ed. (McGraw Hill, New York).

Despite availability of a variety of chemotherapeutic agents, chemotherapy has many drawbacks. Stockdale, *Medicine*, vol. 3, Rubenstein and Federman, eds., ch. 12, sect. 10, 1998. Almost all chemotherapeutic agents are toxic, and chemotherapy causes significant, and often dangerous side effects including severe nausea, bone marrow depression, and immunosuppression. Additionally, even with administration of combinations of chemotherapeutic agents, many tumor cells are resistant or develop resistance to the chemotherapeutic agents. In fact, those cells resistant to the particular chemotherapeutic agents used in the treatment protocol often prove to be resistant to other drugs, even if those agents act by different mechanism from those of the drugs used in the specific treatment. This phenomenon is referred to as pleio-

US 8,648,095 B2

3

tropic drug or multidrug resistance. Because of the drug resistance, many cancers prove refractory to standard chemotherapeutic treatment protocols.

Other diseases or conditions associated with, or characterized by, undesired angiogenesis are also difficult to treat. However, some compounds such as protamine, heparin and steroids have been proposed to be useful in the treatment of certain specific diseases. Taylor et al., *Nature* 297:307 (1982); Folkman et al., *Science* 221:719 (1983); and U.S. Pat. Nos. 5,001,116 and 4,994,443. Thalidomide and certain derivatives of it have also been proposed for the treatment of such diseases and conditions. U.S. Pat. Nos. 5,593,990, 5,629,327, 5,712,291, 6,071,948 and 6,114,355 to D'Amato.

Still, there is a significant need for safe and effective methods of treating, preventing and managing cancer and other diseases and conditions, particularly for diseases that are refractory to standard treatments, such as surgery, radiation therapy, chemotherapy and hormonal therapy, while reducing or avoiding the toxicities and/or side effects associated with the conventional therapies.

2.3 IMiDs™

A number of studies have been conducted with the aim of providing compounds that can safely and effectively be used to treat diseases associated with abnormal production of TNF- α . See, e.g., Marriott, J. B., et al., *Expert Opin. Biol. Ther.* 1(4):1-8 (2001); G. W. Muller, et al., *Journal of Medicinal Chemistry* 39(17): 3238-3240 (1996); and G. W. Muller, et al., *Bioorganic & Medicinal Chemistry Letters* 8: 2669-2674 (1998). Some studies have focused on a group of compounds selected for their capacity to potentially inhibit TNF- α production by LPS stimulated PBMC. L. G. Corral, et al., *Ann. Rheum. Dis.* 58:(Suppl 1) 1107-1113 (1999). These compounds, which are referred to as IMiDs™ (Celgene Corporation) or Immunomodulatory Drugs, show not only potent inhibition of TNF- α but also marked inhibition of LPS induced monocyte IL1 β and IL12 production. LPS induced IL6 is also inhibited by immunomodulatory compounds, albeit partially. These compounds are potent stimulators of LPS induced IL10. Id. Particular examples of IMiDs™ include, but are not limited to, the substituted 2-(2,6-dioxopiperidin-3-yl)phthalimides and substituted 2-(2,6-dioxopiperidin-3-yl)-1-oxoisindoles described in U.S. Pat. Nos. 6,281,230 and 6,316,471, both to G. W. Muller, et al.

3. SUMMARY OF THE INVENTION

This invention encompasses methods of treating and preventing certain types of cancer, including primary and metastatic cancer, as well as cancers that are refractory or resistant to conventional chemotherapy. The methods comprise administering to a patient in need of such treatment or prevention a therapeutically or prophylactically effective amount of an immunomodulatory compound, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof. The invention also encompasses methods of managing certain cancers (e.g., preventing or prolonging their recurrence, or lengthening the time of remission) which comprise administering to a patient in need of such management a prophylactically effective amount of an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

In particular methods of the invention, an immunomodulatory compound is administered in combination with a therapy conventionally used to treat, prevent or manage cancer. Examples of such conventional therapies include, but are

4

not limited to, surgery, chemotherapy, radiation therapy, hormonal therapy, biological therapy and immunotherapy.

This invention also encompasses methods of treating, managing or preventing diseases and disorders other than cancer that are associated with, or characterized by, undesired angiogenesis, which comprise administering to a patient in need of such treatment, management or prevention a therapeutically or prophylactically effective amount of an immunomodulatory compound, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

In other methods of the invention, an immunomodulatory compound is administered in combination with a therapy conventionally used to treat, prevent or manage diseases or disorders associated with, or characterized by, undesired angiogenesis. Examples of such conventional therapies include, but are not limited to, surgery, chemotherapy, radiation therapy, hormonal therapy, biological therapy and immunotherapy.

This invention encompasses pharmaceutical compositions, single unit dosage forms, dosing regimens and kits which comprise an immunomodulatory compound, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, and a second, or additional, active agent. Second active agents include specific combinations, or "cocktails," of drugs.

4. BRIEF DESCRIPTION OF FIGURE

FIG. 1 shows a comparison of the effects of 3-(4-amino-1-oxo-1,3-dihydro-isindol-2-yl)-piperidine-2,6-dione (Revimid™) and thalidomide in inhibiting the proliferation of multiple myeloma (MM) cell lines in an in vitro study. The uptake of [³H]-thymidine by different MM cell lines (MM.1S, Hs Sultan, U266 and RPMI-8226) was measured as an indicator of the cell proliferation.

5. DETAILED DESCRIPTION OF THE INVENTION

A first embodiment of the invention encompasses methods of treating, managing, or preventing cancer which comprises administering to a patient in need of such treatment or prevention a therapeutically or prophylactically effective amount of an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

In particular methods encompassed by this embodiment, the immunomodulatory compound is administered in combination with another drug ("second active agent") or method of treating, managing, or preventing cancer. Second active agents include small molecules and large molecules (e.g., proteins and antibodies), examples of which are provided herein, as well as stem cells. Methods, or therapies, that can be used in combination with the administration of the immunomodulatory compound include, but are not limited to, surgery, blood transfusions, immunotherapy, biological therapy, radiation therapy, and other non-drug based therapies presently used to treat, prevent or manage cancer.

Another embodiment of the invention encompasses methods of treating, managing or preventing diseases and disorders other than cancer that are characterized by undesired angiogenesis. These methods comprise the administration of a therapeutically or prophylactically effective amount of an immunomodulatory compound, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

US 8,648,095 B2

5

Examples of diseases and disorders associated with, or characterized by, undesired angiogenesis include, but are not limited to, inflammatory diseases, autoimmune diseases, viral diseases, genetic diseases, allergic diseases, bacterial diseases, ocular neovascular diseases, choroidal neovascular diseases, retina neovascular diseases, and rubeosis (neovascularization of the angle).

In particular methods encompassed by this embodiment, the immunomodulatory compound is administered in combination with a second active agent or method of treating, managing, or preventing the disease or condition. Second active agents include small molecules and large molecules (e.g., proteins and antibodies), examples of which are provided herein, as well as stem cells. Methods, or therapies, that can be used in combination with the administration of the immunomodulatory compound include, but are not limited to, surgery, blood transfusions, immunotherapy, biological therapy, radiation therapy, and other non-drug based therapies presently used to treat, prevent or manage disease and conditions associated with, or characterized by, undesired angiogenesis.

The invention also encompasses pharmaceutical compositions (e.g., single unit dosage forms) that can be used in methods disclosed herein. Particular pharmaceutical compositions comprise an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, and a second active agent.

5.1 Immunomodulatory Compounds

Compounds used in the invention include immunomodulatory compounds that are racemic, stereomerically enriched or stereomerically pure, and pharmaceutically acceptable salts, solvates, hydrates, stereoisomers, clathrates, and prodrugs thereof. Preferred compounds used in the invention are small organic molecules having a molecular weight less than about 1,000 g/mol, and are not proteins, peptides, oligonucleotides, oligosaccharides or other macromolecules.

As used herein and unless otherwise indicated, the terms "immunomodulatory compounds" and "IMiDs™" (Celgene Corporation) encompasses small organic molecules that markedly inhibit TNF- α , LPS induced monocyte IL1 β and IL12, and partially inhibit IL6 production. Specific immunomodulatory compounds are discussed below.

TNF- α is an inflammatory cytokine produced by macrophages and monocytes during acute inflammation. TNF- α is responsible for a diverse range of signaling events within cells. TNF- α may play a pathological role in cancer. Without being limited by theory, one of the biological effects exerted by the immunomodulatory compounds of the invention is the reduction of synthesis of TNF- α . Immunomodulatory compounds of the invention enhance the degradation of TNF- α mRNA.


Further, without being limited by theory, immunomodulatory compounds used in the invention may also be potent co-stimulators of T cells and increase cell proliferation dramatically in a dose dependent manner. Immunomodulatory compounds of the invention may also have a greater co-stimulatory effect on the CD8+ T cell subset than on the CD4+ T cell subset. In addition, the compounds preferably have anti-inflammatory properties, and efficiently co-stimulate T cells.

Specific examples of immunomodulatory compounds of the invention, include, but are not limited to, cyano and carboxy derivatives of substituted styrenes such as those dis-

6

closed in U.S. Pat. No. 5,929,117; 1-oxo-2-(2,6-dioxo-3-fluoropiperidin-3-yl) isoindolines and 1,3-dioxo-2-(2,6-dioxo-3-fluoropiperidine-3-yl)isoindolines such as those described in U.S. Pat. No. 5,874,448; the tetra substituted 2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolines described in U.S. Pat. No. 5,798,368; 1-oxo and 1,3-dioxo-2-(2,6-dioxopiperidin-3-yl)isoindolines (e.g., 4-methyl derivatives of thalidomide and EM-12), including, but not limited to, those disclosed in U.S. Pat. No. 5,635,517; and a class of non-polypeptide cyclic amides disclosed in U.S. Pat. Nos. 5,698,579 and 5,877,200; analogs and derivatives of thalidomide, including hydrolysis products, metabolites, derivatives and precursors of thalidomide, such as those described in U.S. Pat. Nos. 5,593,990, 5,629,327, and 6,071,948 to D'Amato; aminothalidomide, as well as analogs, hydrolysis products, metabolites, derivatives and precursors of aminothalidomide, and substituted 2-(2,6-dioxopiperidin-3-yl)phthalimides and substituted 2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindoles such as those described in U.S. Pat. Nos. 6,281,230 and 6,316,471; isoindole-imide compounds such as those described in U.S. patent application Ser. No. 09/972,487 filed on Oct. 5, 2001, U.S. patent application Ser. No. 10/032,286 filed on Dec. 21, 2001, and International Application No. PCT/US01/50401 (International Publication No. WO 02/059106). The entireties of each of the patents and patent applications identified herein are incorporated herein by reference. Immunomodulatory compounds of the invention do not include thalidomide.

Other specific immunomodulatory compounds of the invention include, but are not limited to, 1-oxo- and 1,3-dioxo-2-(2,6-dioxopiperidin-3-yl)isoindolines substituted with amino in the benzo ring as described in U.S. Pat. No. 5,635,517 which is incorporated herein by reference. These compounds have the structure I:




in which one of X and Y is C=O, the other of X and Y is C=O or CH₂, and R² is hydrogen or lower alkyl, in particular methyl. Specific immunomodulatory compounds include, but are not limited to:

1-oxo-2-(2,6-dioxopiperidin-3-yl)-4-aminoisoindoline;
 1-oxo-2-(2,6-dioxopiperidin-3-yl)-5-aminoisoindoline;
 1-oxo-2-(2,6-dioxopiperidin-3-yl)-6-aminoisoindoline;
 1-oxo-2-(2,6-dioxopiperidin-3-yl)-7-aminoisoindoline;
 1,3-dioxo-2-(2,6-dioxopiperidin-3-yl)-4-aminoisoindoline;
 and
 1,3-dioxo-2-(2,6-dioxopiperidin-3-yl)-5-aminoisoindoline.

Other specific immunomodulatory compounds of the invention belong to a class of substituted 2-(2,6-dioxopiperidin-3-yl)phthalimides and substituted 2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindoles, such as those described in U.S. Pat. Nos. 6,281,230; 6,316,471; 6,335,349; and 6,476,052, and International Patent Application No. PCT/US97/13375 (International Publication No. WO 98/03502), each of which is incorporated herein by reference. Compounds representative of this class are of the formulas:


US 8,648,095 B2

7



wherein R¹ is hydrogen or methyl. In a separate embodiment, the invention encompasses the use of enantiomerically pure forms (e.g. optically pure (R) or (S) enantiomers) of these compounds.

Still other specific immunomodulatory compounds of the invention belong to a class of isoindole-imides disclosed in U.S. patent application Ser. Nos. 10/032,286 and 09/972,487, and International Application No. PCT/US01/50401 (International Publication No. WO 02/059106), each of which are incorporated herein by reference. Representative compounds are of formula II:



and pharmaceutically acceptable salts, hydrates, solvates, clathrates, enantiomers, diastereomers, racemates, and mixtures of stereoisomers thereof, wherein:

one of X and Y is C=O and the other is CH₂ or C≡O;

R¹ is H, (C₁-C₈)alkyl, (C₃-C₇)cycloalkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, benzyl, aryl, (C₀-C₄)alkyl-(C₁-C₆)heterocycloalkyl, (C₀-C₄)alkyl-(C₂-C₅)heteroaryl, C(O)R³, C(S)R³, C(O)OR⁴, (C₁-C₈)alkyl-N(R⁶)₂, (C₁-C₈)alkyl-OR⁵, (C₁-C₈)alkyl-C(O)OR⁵, C(O)NHR³, C(S)NHR³, C(O)NR³R^{3'}, C(S)NR³R^{3'} or (C₁-C₈)alkyl-O(CO)R⁵;

R² is H, F, benzyl, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, or (C₂-C₈)alkynyl;

R³ and R^{3'} are independently (C₁-C₈)alkyl, (C₃-C₇)cycloalkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, benzyl, aryl, (C₀-

8

C₄)alkyl-(C₁-C₆)heterocycloalkyl, (C₀-C₄)alkyl-(C₂-C₅)heteroaryl, (C₀-C₈)alkyl-N(R⁶)₂, (C₁-C₈)alkyl-OR⁵, (C₁-C₈)alkyl-C(O)OR⁵, (C₁-C₈)alkyl-O(CO)R⁵, or C(O)OR⁵;

R⁴ is (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, (C₁-C₄)alkyl-OR⁵, benzyl, aryl, (C₀-C₄)alkyl-(C₁-C₆)heterocycloalkyl, or (C₀-C₄)alkyl-(C₂-C₅)heteroaryl;

R⁵ is (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, benzyl, aryl, or (C₂-C₅)heteroaryl;

each occurrence of R⁶ is independently H, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, benzyl, aryl, (C₂-C₅)heteroaryl, or (C₀-C₈)alkyl-C(O)O—R⁵ or the R⁶ groups can join to form a heterocycloalkyl group;

n is 0 or 1; and

* represents a chiral-carbon center.

In specific compounds of formula II, when n is 0 then R¹ is (C₃-C₇)cycloalkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, benzyl, aryl, (C₀-C₄)alkyl-(C₁-C₆)heterocycloalkyl, (C₀-C₄)alkyl-(C₂-C₅)heteroaryl, C(O)R³, C(O)OR⁴, (C₁-C₈)alkyl-N(R⁶)₂, (C₁-C₈)alkyl-OR⁵, (C₁-C₈)alkyl-C(O)OR⁵, C(S)NHR³, or (C₁-C₈)alkyl-O(CO)R⁵;


R² is H or (C₁-C₈)alkyl; and

R³ is (C₁-C₈)alkyl, (C₃-C₇)cycloalkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, benzyl, aryl, (C₀-C₄)alkyl-(C₁-C₆)heterocycloalkyl, (C₀-C₄)alkyl-(C₂-C₅)heteroaryl, (C₅-C₈)alkyl-N(R⁶)₂; (C₀-C₈)alkyl-NH—C(O)O—R⁵; (C₁-C₈)alkyl-OR⁵, (C₁-C₈)alkyl-C(O)OR⁵, (C₁-C₈)alkyl-O(CO)R⁵, or C(O)OR⁵; and the other variables have the same definitions.


In other specific compounds of formula II, R² is H or (C₁-C₄)alkyl.

In other specific compounds of formula II, R¹ is (C₁-C₈)alkyl or benzyl.

In other specific compounds of formula II, R¹ is H, (C₁-C₈)alkyl, benzyl, CH₂OCH₃, CH₂CH₂OCH₃, or



In another embodiment of the compounds of formula II, R¹ is



wherein Q is O or S, and each occurrence of R⁷ is independently H, (C₁-C₈)alkyl, benzyl, CH₂OCH₃, or CH₂CH₂OCH₃.

In other specific compounds of formula II, R¹ is C(O)R³.

In other specific compounds of formula II, R³ is (C₀-C₄)alkyl-(C₂-C₅)heteroaryl, (C₁-C₈)alkyl, aryl, or (C₀-C₄)alkyl-OR⁵.

In other specific compounds of formula II, heteroaryl is pyridyl, furyl, or thienyl.


In other specific compounds of formula II, R¹ is C(O)OR⁴.

In other specific compounds of formula II, the H of C(O)NHC(O) can be replaced with (C₁-C₄)alkyl, aryl, or benzyl.

US 8,648,095 B2

9

Still other specific immunomodulatory compounds of the invention belong to a class of isoindole-imides disclosed in U.S. patent application Ser. No. 09/781,179, International Publication No. WO 98/54170, and U.S. Pat. No. 6,395,754, each of which are incorporated herein by reference. Representative compounds are of formula III:



III

and pharmaceutically acceptable salts, hydrates, solvates, clathrates, enantiomers, diastereomers, racemates, and mixtures of stereoisomers thereof, wherein:

one of X and Y is C=O and the other is CH₂ or C=O;

R is H or CH₂OCOR';

(i) each of R¹, R², R³, or R⁴, independently of the others, is halo, alkyl of 1 to 4 carbon atoms, or alkoxy of 1 to 4 carbon atoms or (ii) one of R¹, R², R³, or R⁴ is nitro or —NHR⁵ and the remaining of R¹, R², R³, or R⁴ are hydrogen;

R⁵ is hydrogen or alkyl of 1 to 8 carbons

R⁶ hydrogen, alkyl of 1 to 8 carbon atoms, benzo, chloro, or fluoro;

R⁷ is R⁷—CHR¹⁰—N(R⁸R⁹);


R⁷ is m-phenylene or p-phenylene or —(C_nH_{2n})— in which n has a value of 0 to 4;

each of R⁸ and R⁹ taken independently of the other is hydrogen or alkyl of 1 to 8 carbon atoms, or R⁸ and R⁹ taken together are tetramethylene, pentamethylene, hexamethylene, or —CH₂CH₂[X]X₁CH₂CH₂— in which [X]X₁ is —O—, —S—, or —NH—;

R¹⁰ is hydrogen, alkyl of 1 to 8 carbon atoms, or phenyl; and


* represents a chiral-carbon center.

The most preferred immunomodulatory compounds of the invention are 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione and 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. The compounds can be obtained via standard, synthetic methods (see e.g., U.S. Pat. No. 5,635,517, incorporated herein by reference). The compounds are available from Celgene Corporation, Warren, N.J. 4-(Amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (ACTIMID™) has the following chemical structure:



The compound 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (REVIMID™) has the following chemical structure:

10



5

10

15

Compounds of the invention can either be commercially purchased or prepared according to the methods described in the patents or patent publications disclosed herein. Further, optically pure compounds can be asymmetrically synthesized or resolved using known resolving agents or chiral columns as well as other standard synthetic organic chemistry techniques.

As used herein and unless otherwise indicated, the term “pharmaceutically acceptable salt” encompasses non-toxic acid and base addition salts of the compound to which the term refers. Acceptable non-toxic acid addition salts include those derived from organic and inorganic acids or bases known in the art, which include, for example, hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid, methanesulphonic acid, acetic acid, tartaric acid, lactic acid, succinic acid, citric acid, malic acid, maleic acid, sorbic acid, aconitic acid, salicylic acid, phthalic acid, embolic acid, enanthic acid, and the like.

25

30

Compounds that are acidic in nature are capable of forming salts with various pharmaceutically acceptable bases. The bases that can be used to prepare pharmaceutically acceptable base addition salts of such acidic compounds are those that form non-toxic base addition salts, i.e., salts containing pharmacologically acceptable cations such as, but not limited to, alkali metal or alkaline earth metal salts and the calcium, magnesium, sodium or potassium salts in particular. Suitable organic bases include, but are not limited to, N,N-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine), lysine, and procaine.

40

45

50

55

As used herein and unless otherwise indicated, the term “prodrug” means a derivative of a compound that can hydrolyze, oxidize, or otherwise react under biological conditions (in vitro or in vivo) to provide the compound. Examples of prodrugs include, but are not limited to, derivatives of immunomodulatory compounds of the invention that comprise biohydrolyzable moieties such as biohydrolyzable amides, biohydrolyzable esters, biohydrolyzable carbamates, biohydrolyzable carbonates, biohydrolyzable ureides, and biohydrolyzable phosphate analogues. Other examples of prodrugs include derivatives of immunomodulatory compounds of the invention that comprise —NO, —NO₂, —ONO, or —ONO₂ moieties. Prodrugs can typically be prepared using well-known methods, such as those described in 1 *Burger's Medicinal Chemistry and Drug Discovery*, 172-178, 949-982 (Manfred E. Wolff ed., 5th ed. 1995), and *Design of Prodrugs* (H. Bundgaard ed., Elsevier, New York 1985).

60

65

As used herein and unless otherwise indicated, the terms “biohydrolyzable amide,” “biohydrolyzable ester,” “biohydrolyzable carbamate,” “biohydrolyzable carbonate,” “biohydrolyzable ureide,” “biohydrolyzable phosphate” mean an amide, ester, carbamate, carbonate, ureide, or phosphate, respectively, of a compound that either: 1) does not interfere with the biological activity of the compound but can confer upon that compound advantageous properties in vivo, such as uptake, duration of action, or onset of action; or 2) is biologi-

US 8,648,095 B2

11

cally inactive but is converted in vivo to the biologically active compound. Examples of biohydrolyzable esters include, but are not limited to, lower alkyl esters, lower acyloxyalkyl esters (such as acetoxyethyl, acetoxyethyl, aminocarbonyloxymethyl, pivaloyloxymethyl, and pivaloyloxyethyl esters), lactonyl esters (such as phthalidyl and thiophthalidyl esters), lower alkoxyacyloxyalkyl esters (such as methoxycarbonyloxymethyl, ethoxycarbonyloxyethyl and isopropoxycarbonyloxyethyl esters), alkoxyalkyl esters, choline esters, and acylamino alkyl esters (such as acetamidomethyl esters). Examples of biohydrolyzable amides include, but are not limited to, lower alkyl amides, α -amino acid amides, alkoxyacyl amides, and alkylaminoalkylcarbonyl amides. Examples of biohydrolyzable carbamates include, but are not limited to, lower alkylamines, substituted ethylenediamines, amino acids, hydroxyalkylamines, heterocyclic and heteroaromatic amines, and polyether amines.

Various immunomodulatory compounds of the invention contain one or more chiral centers, and can exist as racemic mixtures of enantiomers or mixtures of diastereomers. This invention encompasses the use of stereomerically pure forms of such compounds, as well as the use of mixtures of those forms. For example, mixtures comprising equal or unequal amounts of the enantiomers of a particular immunomodulatory compounds of the invention may be used in methods and compositions of the invention. These isomers may be asymmetrically synthesized or resolved using standard techniques such as chiral columns or chiral resolving agents. See, e.g., Jacques, J., et al., *Enantiomers, Racemates and Resolutions* (Wiley-Interscience, New York, 1981); Wilen, S. H., et al., *Tetrahedron* 33:2725 (1977); Eliel, E. L., *Stereochemistry of Carbon Compounds* (McGraw-Hill, NY, 1962); and Wilen, S. H., *Tables of Resolving Agents and Optical Resolutions* p. 268 (E. L. Eliel, Ed., Univ. of Notre Dame Press, Notre Dame, Ind., 1972).

As used herein and unless otherwise indicated, the term "stereomerically pure" means a composition that comprises one stereoisomer of a compound and is substantially free of other stereoisomers of that compound. For example, a stereomerically pure composition of a compound having one chiral center will be substantially free of the opposite enantiomer of the compound. A stereomerically pure composition of a compound having two chiral centers will be substantially free of other diastereomers of the compound. A typical stereomerically pure compound comprises greater than about 80% by weight of one stereoisomer of the compound and less than about 20% by weight of other stereoisomers of the compound, more preferably greater than about 90% by weight of one stereoisomer of the compound and less than about 10% by weight of the other stereoisomers of the compound, even more preferably greater than about 95% by weight of one stereoisomer of the compound and less than about 5% by weight of the other stereoisomers of the compound, and most preferably greater than about 97% by weight of one stereoisomer of the compound and less than about 3% by weight of the other stereoisomers of the compound. As used herein and unless otherwise indicated, the term "stereomerically enriched" means a composition that comprises greater than about 60% by weight of one stereoisomer of a compound, preferably greater than about 70% by weight, more preferably greater than about 80% by weight of one stereoisomer of a compound. As used herein and unless otherwise indicated, the term "enantiomerically pure" means a stereomerically pure composition of a compound having one chiral center. Similarly, the term "stereomerically enriched" means a stereomerically enriched composition of a compound having one chiral center.

12

It should be noted that if there is a discrepancy between a depicted structure and a name given that structure, the depicted structure is to be accorded more weight. In addition, if the stereochemistry of a structure or a portion of a structure is not indicated with, for example, bold or dashed lines, the structure or portion of the structure is to be interpreted as encompassing all stereoisomers of it.

5.2 Second Active Agents

Immunomodulatory compounds can be combined with other pharmacologically active compounds ("second active agents") in methods and compositions of the invention. It is believed that certain combinations work synergistically in the treatment of particular types of cancer and certain diseases and conditions associated with, or characterized by, undesired angiogenesis. Immunomodulatory compounds can also work to alleviate adverse effects associated with certain second active agents, and some second active agents can be used to alleviate adverse effects associated with immunomodulatory compounds.

One or more second active ingredients or agents can be used in the methods and compositions of the invention together with an immunomodulatory compound. Second active agents can be large molecules (e.g., proteins) or small molecules (e.g., synthetic inorganic, organometallic, or organic molecules).

Examples of large molecule active agents include, but are not limited to, hematopoietic growth factors, cytokines, and monoclonal and polyclonal antibodies. Typical large molecule active agents are biological molecules, such as naturally occurring or artificially made proteins. Proteins that are particularly useful in this invention include proteins that stimulate the survival and/or proliferation of hematopoietic precursor cells and immunologically active poietic cells in vitro or in vivo. Others stimulate the division and differentiation of committed erythroid progenitors in cells in vitro or in vivo. Particular proteins include, but are not limited to: interleukins, such as IL-2 (including recombinant IL-II ("rIL2") and canarypox IL-2), IL-10, IL-12, and IL-18; interferons, such as interferon alfa-2a, interferon alfa-2b, interferon alfa-n1, interferon alfa-n3, interferon beta-1 a, and interferon gamma-I b; GM-CSF and GM-CSF; and EPO.

Particular proteins that can be used in the methods and compositions of the invention include, but are not limited to: filgrastim, which is sold in the United States under the trade name Neupogen® (Amgen, Thousand Oaks, Calif.); sargramostim, which is sold in the United States under the trade name Leukine® (Immunex, Seattle, Wash.); and recombinant EPO, which is sold in the United States under the trade name Epogen® (Amgen, Thousand Oaks, Calif.).

Recombinant and mutated forms of GM-CSF can be prepared as described in U.S. Pat. Nos. 5,391,485; 5,393,870; and 5,229,496; all of which are incorporated herein by reference. Recombinant and mutated forms of G-CSF can be prepared as described in U.S. Pat. Nos. 4,810,643; 4,999,291; 5,528,823; and 5,580,755; all of which are incorporated herein by reference.

This invention encompasses the use of native, naturally occurring, and recombinant proteins. The invention further encompasses mutants and derivatives (e.g., modified forms) of naturally occurring proteins that exhibit, in vivo, at least some of the pharmacological activity of the proteins upon which they are based. Examples of mutants include, but are not limited to, proteins that have one or more amino acid residues that differ from the corresponding residues in the naturally occurring forms of the proteins. Also encompassed by the term "mutants" are proteins that lack carbohydrate moieties normally present in their naturally occurring forms

US 8,648,095 B2

13

(e.g., nonglycosylated forms). Examples of derivatives include, but are not limited to, pegylated derivatives and fusion proteins, such as proteins formed by fusing IgG1 or IgG3 to the protein or active portion of the protein of interest. See, e.g., Penichet, M. L. and Morrison, S. L., *J. Immunol. Methods* 248:91-101 (2001).

Antibodies that can be used in combination with compounds of the invention include monoclonal and polyclonal antibodies. Examples of antibodies include, but are not limited to, trastuzumab (Herceptin®), rituximab (Rituxan®), bevacizumab (Avastin™), pertuzumab (Omnitarg™), tositumomab (Bexxar®), edrecolomab (Panorex®), and G250. Compounds of the invention can also be combined with, or used in combination with, anti-TNF- α antibodies.

Large molecule active agents may be administered in the form of anti-cancer vaccines. For example, vaccines that secrete, or cause the secretion of, cytokines such as IL-2, G-CSF, and GM-CSF can be used in the methods, pharmaceutical compositions, and kits of the invention. See, e.g., Emens, L. A., et al., *Curr. Opin. Mol. Ther.* 3(1):77-84 (2001).

In one embodiment of the invention, the large molecule active agent reduces, eliminates, or prevents an adverse effect associated with the administration of an immunomodulatory compound. Depending on the particular immunomodulatory compound and the disease or disorder being treated, adverse effects can include, but are not limited to, drowsiness and somnolence, dizziness and orthostatic hypotension, neutropenia, infections that result from neutropenia, increased HIV-viral load, bradycardia, Stevens-Johnson Syndrome and toxic epidermal necrolysis, and seizures (e.g., grand mal convulsions). A specific adverse effect is neutropenia.

Second active agents that are small molecules can also be used to alleviate adverse effects associated with the administration of an immunomodulatory compound. However, like some large molecules, many are believed to be capable of providing a synergistic effect when administered with (e.g., before, after or simultaneously) an immunomodulatory compound. Examples of small molecule second active agents include, but are not limited to, anti-cancer agents, antibiotics, immunosuppressive agents, and steroids.

Examples of anti-cancer agents include, but are not limited to: acivicin; aclarubicin; acodazole hydrochloride; acronine; adozelesin; aldesleukin; altretamine; ambomycin; ametantrone acetate; amsacrine; anastrozole; anthramycin; asparaginase; asperlin; azacitidine; azetepa; azotomycin; batimastat; benzodepa; bicalutamide; bisantrene hydrochloride; bisnafide dimesylate; bizelesin; bleomycin sulfate; brequinar sodium; bropirimine; busulfan; cactinomycin; calusterone; caracemide; carbetimer; carboplatin; carmustine; carubicin hydrochloride; carzelesin; cedefingol; celecoxib (COX-2 inhibitor); chlorambucil; cirolemycin; cisplatin; cladribine; crisnatol mesylate; cyclophosphamide; cytarabine; dacarbazine; dactinomycin; daunorubicin hydrochloride; decitabine; dexormaplatin; dezaguanine; dezaguanine mesylate; diaziquone; docetaxel; doxorubicin; doxorubicin hydrochloride; droloxifene; droloxifene citrate; dromostanolone propionate; duazomycin; edatrexate; eflornithine hydrochloride; elsamitucin; enloplatin; enpromate; epiropropidine; epirubicin hydrochloride; erbulozole; esorubicin hydrochloride; estramustine; estramustine phosphate sodium; etanidazole; etoposide; etoposide phosphate; etoprine; fadrozole hydrochloride; fazarabine; fenretinide; floxuridine; fludarabine phosphate; fluorouracil; fluorocitabine; fosquidone; fostriecin sodium; gemcitabine; gemcitabine hydrochloride; hydroxyurea; idarubicin hydrochloride; ifosfamide; ilmofofosine; iproplatin; irinotecan; irinotecan

14

hydrochloride; lanreotide acetate; letrozole; leuprolide acetate; liarozole hydrochloride; lometrexol sodium; lomustine; losoxantrone hydrochloride; masoprocil; maytansine; mechlorethamine hydrochloride; megestrol acetate; melengestrol acetate; melphalan; menogaril; mercaptopurine; methotrexate; methotrexate sodium; metoprine; meturedepa; mitindomide; mitocarcin; mitocromin; mitogillin; mitomalcin; mitomycin; mitosper; mitotane; mitoxantrone hydrochloride; mycophenolic acid; nocodazole; nogalamycin; ormaplatin; oxisuran; paclitaxel; pegaspargase; peliomycin; pentamustine; peplomycin sulfate; perfosfamide; pipobroman; pipsulfan; piroxantrone hydrochloride; plicamycin; plomestane; porfimer sodium; porfiromycin; prednimustine; procarbazine hydrochloride; puromycin; puromycin hydrochloride; pyrazofurin; riboprine; safingol; safingol hydrochloride; semustine; simtrazene; sparfosate sodium; sparsomycin; spirogermanium hydrochloride; spiro-mustine; spiroplatin; streptonigrin; streptozocin; sulofenur; talisomycin; tecogalan sodium; taxotere; tegafur; teloxantrone hydrochloride; temoporfin; teniposide; teroxirone; testolactone; thiamiprine; thioguanine; thiotepa; tiazofurin; tirapazamine; toremifene citrate; tretolone acetate; triciribine phosphate; trimetrexate; trimetrexate glucuronate; triptorelin; tubulozole hydrochloride; uracil mustard; uredepa; vaporeotide; verteporfin; vinblastine sulfate; vincristine sulfate; vindesine; vindesine sulfate; vinepidine sulfate; vinglycinate sulfate; vinleurosine sulfate; vinorelbine tartrate; vinrosidine sulfate; vinzolidine sulfate; vorozole; zeniplatin; zinostatin; and zorubicin hydrochloride.

Other anti-cancer drugs include, but are not limited to: 20-epi-1,25 dihydroxyvitamin D3; 5-ethynyluracil; abiraterone; aclarubicin; acylfulvene; adecypenol; adozelesin; aldesleukin; ALL-TK antagonists; altretamine; ambamustine; amidox; amifostine; aminolevulinic acid; amrubicin; amsacrine; anagrelide; anastrozole; andrographolide; angiogenesis inhibitors; antagonist D; antagonist G; antarelix; antidorsalizing morphogenetic protein-1; antiandrogen, prostatic carcinoma; antiestrogen; antineoplaston; antisense oligonucleotides; aphidicolin glycinate; apoptosis gene modulators; apoptosis regulators; apurinic acid; ara-CDP-DL-PTBA; arginine deaminase; asulacrine; atamestane; atrimustine; axinastatin 1; axinastatin 2; axinastatin 3; azasetron; azatoxin; azatyrosine; baccatin III derivatives; balanol; batimastat; BCR/ABL antagonists; benzochlorins; benzoylstauroporine; beta lactam derivatives; beta-alethine; betaclamycin B; betulinic acid; bFGF inhibitor; bicalutamide; bisantrene; bisaziridinylspermine; bisnafide; bistratene A; bizelesin; breflate; bropirimine; budotitane; buthionine sulfoximine; calcipotriol; calphostin C; camptothecin derivatives; capecitabine; carboxamide-amino-triazole; carboxamidotriazole; CaRest M3; CARN 700; cartilage derived inhibitor; carzelesin; casein kinase inhibitors (ICOS); castanospermine; cecropin B; cetrorelix; chlorins; chloroquinoline sulfonamide; cicaprost; cis-porphyrin; cladribine; clomifene analogues; clotrimazole; collismycin A; collismycin B; combretastatin A4; combretastatin analogue; conagein; crambescidin 816; crisnatol; cryptophycin 8; cryptophycin A derivatives; curacin A; cyclopentantraquinones; cycloplatin; cypemycin; cytarabine ocfosfate; cytolytic factor; cytosatin; dacliximab; decitabine; dehydridemnin B; deslorelin; dexamethasone; dexifosfamide; dexrazoxane; dexverapamil; diaziquone; didemnin B; didox; diethylnor-spermine; dihydro-5-azacytidine; dihydrotaxol, 9-; dioxamycin; diphenyl spiromustine; docetaxel; docosanol; dolasetron; doxifluridine; doxorubicin; droloxifene; dronabinol; duocarmycin SA; ebselen; ecomustine; edelfosine; edrecolomab; eflornithine; elemene; emitefur; epirubicin; epristeride;

US 8,648,095 B2

15

estramustine analogue; estrogen agonists; estrogen antagonists; etanidazole; etoposide phosphate; exemestane; fadrozole; fazarabine; fenretinide; filgrastim; finasteride; flavopiridol; flzelastine; fluasterone; fludarabine; fluorodaunorubicin hydrochloride; forfenimex; formestane; fostriecin; fotemustine; gadolinium texaphyrin; gallium nitrate; galocitabine; ganirelix; gelatinase inhibitors; gemcitabine; glutathione inhibitors; hepsulfam; heregulin; hexamethylene bisacetamide; hypericin; ibandronic acid; idarubicin; idoxifene; idramantone; ilmofosine; ilomastat; imatinib (e.g., Gleevec®), imiquimod; immunostimulant peptides; insulin-like growth factor-1 receptor inhibitor; interferon agonists; interferons; interleukins; iobenguane; iododoxorubicin; ipomeanol, 4-; iroplact; irsogladine; isobengazole; isohomohalicondrin B; itasetron; jasplakinolide; kahalalide F; lamellarin-N triacetate; lanreotide; leinamycin; lenograstim; lentinan sulfate; leptolstatin; letrozole; leukemia inhibiting factor; leukocyte alpha interferon; leuprolide+estrogen+progesterone; leuprorelin; levamisole; liarozole; linear polyamine analogue; lipophilic disaccharide peptide; lipophilic platinum compounds; lissoclinamide 7; lobaplatin; lombricine; lometrexol; lonidamine; losoxantrone; loxoribine; lurtotecan; lutetium texaphyrin; lysofylline; lytic peptides; maitansine; mannostatin A; marimastat; masoprocol; maspin; matrilysin inhibitors; matrix metalloproteinase inhibitors; menogaril; merbarone; meterelin; methioninase; metoclopramide; MIF inhibitor; mifepristone; miltefosine; mirimostim; mitoguanzone; mitolactol; mitomycin analogues; mitonafide; mitotoxin fibroblast growth factor-saporin; mitoxantrone; mofarotene; molgramostim; Erbitux, human chorionic gonadotrophin; monophosphoryl lipid A+myobacterium cell wall sk; mopidamol; mustard anticancer agent; mycaperoxide B; mycobacterial cell wall extract; myriaporone; N-acetyldinaline; N-substituted benzamides; nafarelin; nagrestip; naloxone+pentazocine; napavin; naphterpin; nartograstim; nedaplatin; nemorubicin; neridronic acid; nilutamide; nisamycin; nitric oxide modulators; nitroxide antioxidant; nitrullyn; oblimersen (Genasense®); O⁶-benzylguanine; octreotide; okicenone; oligonucleotides; onapristone; ondansetron; ondansetron; oracin; oral cytokine inducer; ormaplatin; osaterone; oxaliplatin; oxaunomycin; paclitaxel; paclitaxel analogues; paclitaxel derivatives; palauamine; palmitoylrhizoxin; pamidronic acid; panaxytriol; panomifene; parabactin; pazelliptine; pegaspargase; peldesine; pentosan polysulfate sodium; pentostatin; pentozole; perflubron; perfosfamide; perillyl alcohol; phenazinomycin; phenylacetate; phosphatase inhibitors; picibanil; pilocarpine hydrochloride; pirarubicin; piritrexim; placetin A; placetin B; plasminogen activator inhibitor; platinum complex; platinum compounds; platinum-triamine complex; porfimer sodium; porfirofomycin; prednisone; propyl bis-acridone; prostaglandin J2; proteasome inhibitors; protein A-based immune modulator; protein kinase C inhibitor; protein kinase C inhibitors, microalgal; protein tyrosine phosphatase inhibitors; purine nucleoside phosphorylase inhibitors; purpurins; pyrazoloacridine; pyridoxylated hemoglobin polyoxyethylene conjugate; raf antagonists; raltitrexed; ramosestron; ras farnesyl protein transferase inhibitors; ras inhibitors; ras-GAP inhibitor; retelliptine demethylated; rhenium Re 186 etidronate; rhizoxin; ribozymes; RII retinamide; rohitukine; romurtide; roquinimex; rubiginone B1; ruboxyl; safingol; saintopin; SarCNU; sarcophytol A; sargramostim; Sdi 1 mimetics; semustine; senescence derived inhibitor 1; sense oligonucleotides; signal transduction inhibitors; sizofuran; sobuzoxane; sodium borocaptate; sodium phenylacetate; solverol; somatomedin binding protein; sonermin; sparfosic acid; spicamycin D; spiromustine; splenopentin;

16

spongistatin 1; squalamine; stipiamide; stromelysin inhibitors; sulfinosine; superactive vasoactive intestinal peptide antagonist; suradista; suramin; swainsonine; tallimustine; tamoxifen methiodide; taumustine; tazartotene; tecogalan sodium; tegafur; tellurapyrylium; telomerase inhibitors; temoporfin; teniposide; tetrachlorodecaoxide; tetrazomine; thaliblastine; thiocoraline; thrombopoietin; thrombopoietin mimetic; thymalfasin; thymopoietin receptor agonist; thymotrivan; thyroid stimulating hormone; tin ethyl etiopurpurin; tirapazamine; titanocene bichloride; topsentin; toremifene; translation inhibitors; tretinoin; triacetyluridine; triciribine; trimetrexate; triptorelin; tropisetron; turosteride; tyrosine kinase inhibitors; tyrphostins; UBC inhibitors; ubenimex; urogenital sinus-derived growth inhibitory factor; urokinase receptor antagonists; vapreotide; variolin B; velaresol; veramine; verteporfin; vinorelbine; vinoxaltine; vitaxin; vorozole; zanoterone; zenioplatin; zilascorb; and zinostatin stimalamer.

Specific second active agents include, but are not limited to, oblimersen (Genasense®), remicade, docetaxel, celecoxib, melphalan, dexamethasone (Decadron®), steroids, gemcitabine, cisplatin, temozolomide, etoposide, cyclophosphamide, temodar, carboplatin, procarbazine, gliadel, tamoxifen, topotecan, methotrexate, Arisa®, taxol, taxotere, fluorouracil, leucovorin, irinotecan, xeloda, CPT-11, interferon alpha, pegylated interferon alpha (e.g., PEG INTRON-A), capecitabine, cisplatin, thiotepa, fludarabine, carboplatin, liposomal daunorubicin, cytarabine, doxetaxol, paclitaxel, vinblastine, IL-2, GM-CSF, dacarbazine, vinorelbine, zoledronic acid, palmitronate, biaxin, busulphan, prednisone, bisphosphonate, arsenic trioxide, vincristine, doxorubicin (Doxil®), paclitaxel, ganciclovir, adriamycin, estramustine sodium phosphate (Emcyt®), sulindac, and etoposide.

5.3 Methods of Treatments and Prevention

Methods of this invention encompass methods of treating, preventing and/or managing various types of cancer and diseases and disorders associated with, or characterized by, undesired angiogenesis. As used herein, unless otherwise specified, the term “treating” refers to the administration of a compound of the invention or other additional active agent after the onset of symptoms of the particular disease or disorder. As used herein, unless otherwise specified, the term “preventing” refers to the administration prior to the onset of symptoms, particularly to patients at risk of cancer, and other diseases and disorders associated with, or characterized by, undesired angiogenesis. The term “prevention” includes the inhibition of a symptom of the particular disease or disorder. Patients with familial history of cancer and diseases and disorders associated with, or characterized by, undesired angiogenesis are preferred candidates for preventive regimens. As used herein and unless otherwise indicated, the term “managing” encompasses preventing the recurrence of the particular disease or disorder in a patient who had suffered from it, and/or lengthening the time a patient who had suffered from the disease or disorder remains in remission.

As used herein, the term “cancer” includes, but is not limited to, solid tumors and blood born tumors. The term “cancer” refers to disease of skin tissues, organs, blood, and vessels, including, but not limited to, cancers of the bladder, bone or blood, brain, breast, cervix, chest, colon, endometrium, esophagus, eye, head, kidney, liver, lymph nodes, lung, mouth, neck, ovaries, pancreas, prostate, rectum, stomach, testis, throat, and uterus. Specific cancers include, but are not limited to, advanced malignancy, amyloidosis, neuroblastoma, meningioma, hemangiopericytoma, multiple brain metastase, glioblastoma multiforms, glioblastoma, brain stem glioma, poor prognosis malignant brain tumor,

US 8,648,095 B2

17

malignant glioma, recurrent malignant glioma, anaplastic astrocytoma, anaplastic oligodendroglioma, neuroendocrine tumor, rectal adenocarcinoma, Dukes C & D colorectal cancer, unresectable colorectal carcinoma, metastatic hepatocellular carcinoma, Kaposi's sarcoma, karotype acute myeloblastic leukemia, Hodgkin's lymphoma, non-Hodgkin's lymphoma, cutaneous T-Cell lymphoma, cutaneous B-Cell lymphoma, diffuse large B-Cell lymphoma, low grade follicular lymphoma, malignant melanoma, malignant mesothelioma, malignant pleural effusion mesothelioma syndrome, peritoneal carcinoma, papillary serous carcinoma, gynecologic sarcoma, soft tissue sarcoma, scleroderma, cutaneous vasculitis, Langerhans cell histiocytosis, leiomyosarcoma, fibrodysplasia ossificans progressive, hormone refractory prostate cancer, resected high-risk soft tissue sarcoma, unresectable hepatocellular carcinoma, Waldenstrom's macroglobulinemia, smoldering myeloma, indolent myeloma, fallopian tube cancer, androgen independent prostate cancer, androgen dependent stage IV non-metastatic prostate cancer, hormone-insensitive prostate cancer, chemotherapy-insensitive prostate cancer, papillary thyroid carcinoma, follicular thyroid carcinoma, medullary thyroid carcinoma, and leiomyoma. In a specific embodiment, the cancer is metastatic. In another embodiment, the cancer is refractory or resistance to chemotherapy or radiation; in particular, refractory to thalidomide.

As used herein to refer to diseases and conditions other than cancer, the terms "diseases or disorders associated with, or characterized by, undesired angiogenesis," "diseases or disorders associated with undesired angiogenesis," and "diseases or disorders characterized by undesired angiogenesis" refer to diseases, disorders and conditions that are caused, mediated or attended by undesired, unwanted or uncontrolled angiogenesis, including, but not limited to, inflammatory diseases, autoimmune diseases, genetic diseases, allergic diseases, bacterial diseases, ocular neovascular diseases, choroidal neovascular diseases, and retina neovascular diseases.

Examples of such diseases or disorders associated with undesired angiogenesis include, but are not limited to, diabetic retinopathy, retinopathy of prematurity, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, proliferative vitreoretinopathy, trachoma, myopia, optic pits, epidemic keratoconjunctivitis, atopic keratitis, superior limbic keratitis, pterygium keratitis sicca, sjogrens, acne rosacea, phlyctenulosis, syphilis, lipid degeneration, bacterial ulcer, fungal ulcer, Herpes simplex infection, Herpes zoster infection, protozoan infection, Kaposi sarcoma, Mooren ulcer, Terrien's marginal degeneration, marginal keratolysis, rheumatoid arthritis, systemic lupus, polyarteritis, trauma, Wegeners sarcoidosis, Scleritis, Steven's Johnson disease, peripheral radial keratotomy, sickle cell anemia, sarcoid, pseudoxanthoma elasticum, Pagets disease, vein occlusion, artery occlusion, carotid obstructive disease, chronic uveitis, chronic vitritis, Lyme's disease, Eales disease, Behcet's disease, retinitis, choroiditis, presumed ocular histoplasmosis, Bests disease, Stargarts disease, pars planitis, chronic retinal detachment, hyperviscosity syndromes, toxoplasmosis, rubeosis, sarcoidosis, sclerosis, soriatis, psoriasis, primary sclerosing cholangitis, proctitis, primary biliary srosis, idiopathic pulmonary fibrosis, and alcoholic hepatitis.

In specific embodiments of the invention, diseases or disorders associated with undesired angiogenesis do not include congestive heart failure, cardiomyopathy, pulmonary edema, endotoxin-mediated septic shock, acute viral myocarditis, cardiac allograft rejection, myocardial infarction, HIV, hepatitis, adult respiratory distress syndrome, bone-resorption disease, chronic obstructive pulmonary diseases, chronic pul-

18

monary inflammatory disease, dermatitis, cystic fibrosis, septic shock, sepsis, endotoxic shock, hemodynamic shock, sepsis syndrome, post ischemic reperfusion injury, meningitis, psoriasis, fibrotic disease, cachexia, graft rejection, rheumatoid spondylitis, osteoporosis, Crohn's disease, ulcerative colitis, inflammatory-bowel disease, multiple sclerosis, systemic lupus erythrematosus, erythema nodosum leprosum in leprosy, radiation damage, asthma, hyperoxic alveolar injury, malaria, mycobacterial infection, and opportunistic infections resulting from HIV.

This invention encompasses methods of treating patients who have been previously treated for cancer or diseases or disorders associated with, or characterized by, undesired angiogenesis, but are non-responsive to standard therapies, as well as those who have not previously been treated. The invention also encompasses methods of treating patients regardless of patient's age, although some diseases or disorders are more common in certain age groups. The invention further encompasses methods of treating patients who have undergone surgery in an attempt to treat the disease or condition at issue, as well as those who have not. Because patients with cancer and diseases and disorders characterized by undesired angiogenesis have heterogenous clinical manifestations and varying clinical outcomes, the treatment given to a patient may vary, depending on his/her prognosis. The skilled clinician will be able to readily determine without undue experimentation specific secondary agents, types of surgery, and types of non-drug based standard therapy that can be effectively used to treat an individual patient with cancer and other diseases or disorders.

Methods encompassed by this invention comprise administering one or more immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, to a patient (e.g., a human) suffering, or likely to suffer, from cancer or a disease or disorder mediated by undesired angiogenesis.

In one embodiment of the invention, an immunomodulatory compound of the invention can be administered orally and in single or divided daily doses in an amount of from about 0.10 to about 150 mg/day. In a particular embodiment, 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™) may be administered in an amount of from about 0.1 to about 1 mg per day, or alternatively from about 0.1 to about 5 mg every other day. In a preferred embodiment, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) may be administered in an amount of from about 5 to 25 mg per day, or alternatively from about 10 to about 50 mg every other day.

In a specific embodiment, 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™) may be administered in an amount of about 1, 2, or 5 mg per day to patients with relapsed multiple myeloma. In a particular embodiment, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) may be administered initially in an amount of 5 mg/day and the dose can be escalated every week to 10, 20, 25, 30 and 50 mg/day. In a specific embodiment, Revimid™ can be administered in an amount of up to about 30 mg/day to patients with solid tumor. In a particular embodiment, Revimid™ can be administered in an amount of up to about 40 mg/day to patients with glioma.

5.3.1 Combination Therapy with a Second Active Agent

Specific methods of the invention comprise administering an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, in combination with one or more second active agents, and/or in combination with radia-

US 8,648,095 B2

19

tion therapy, blood transfusions, or surgery. Examples of immunomodulatory compounds of the invention are disclosed herein (see, e.g., section 5.1). Examples of second active agents are also disclosed herein (see, e.g., section 5.2).

Administration of the immunomodulatory compounds and the second active agents to a patient can occur simultaneously or sequentially by the same or different routes of administration. The suitability of a particular route of administration employed for a particular active agent will depend on the active agent itself (e.g., whether it can be administered orally without decomposing prior to entering the blood stream) and the disease being treated. A preferred route of administration for an immunomodulatory compound of the invention is orally. Preferred routes of administration for the second active agents or ingredients of the invention are known to those of ordinary skill in the art. See, e.g., *Physicians' Desk Reference*, 1755-1760 (56th ed., 2002).

In one embodiment of the invention, the second active agent is administered intravenously or subcutaneously and once or twice daily in an amount of from about 1 to about 1000 mg, from about 5 to about 500 mg, from about 10 to about 350 mg, or from about 50 to about 200 mg. The specific amount of the second active agent will depend on the specific agent used, the type of disease being treated or managed, the severity and stage of disease, and the amount(s) of immunomodulatory compounds of the invention and any optional additional active agents concurrently administered to the patient. In a particular embodiment, the second active agent is oblimersen (Genasense®), GM-CSF, G-CSF, EPO, taxotere, irinotecan, dacarbazine, transretinoic acid, topotecan, pentoxifylline, ciprofloxacin, dexamethasone, vincristine, doxorubicin, COX-2 inhibitor, IL2, IL8, IL18, IFN, Ara-C, vinorelbine, or a combination thereof.

In a particular embodiment, GM-CSF, G-CSF or EPO is administered subcutaneously during about five days in a four or six week cycle in an amount of from about 1 to about 750 mg/m²/day, preferably in an amount of from about 25 to about 500 mg/m²/day, more preferably in an amount of from about 50 to about 250 mg/m²/day, and most preferably in an amount of from about 50 to about 200 mg/m²/day. In a certain embodiment, GM-C SF may be administered in an amount of from about 60 to about 500 mcg/m² intravenously over 2 hours, or from about 5 to about 12 mcg/m²/day subcutaneously. In a specific embodiment, G-CSF may be administered subcutaneously in an amount of about 1 mcg/kg/day initially and can be adjusted depending on rise of total granulocyte counts. The maintenance dose of G-CSF may be administered in an amount of about 300 (in smaller patients) or 480 mcg subcutaneously. In a certain embodiment, EPO may be administered subcutaneously in an amount of 10,000 Unit 3 times per week.

In another embodiment, Revimid™ in an amount of about 25 mg/d and dacarbazine in an amount of about from 200 to 1,000 mg/m²/d are administered to patients with metastatic malignant melanoma. In a specific embodiment, Revimid™ is administered in an amount of from about 5 to about 25 mg/d to patients with metastatic malignant melanoma whose disease has progressed on treatment with dacarbazine, IL-2 or IFN. In a specific embodiment, Revimid™ is administered to patients with relapsed or refractory multiple myeloma in an amount of about 15 mg/d twice a day or about 30 mg/d four times a day in a combination with dexamethasone.

In another embodiment, an immunomodulatory compound is administered with melphalan and dexamethasone to patients with amyloidosis. In a specific embodiment, an immunomodulatory compound of the invention and steroids can be administered to patients with amyloidosis.

20

In another embodiment, an immunomodulatory compound is administered with gemcitabine and cisplatin to patients with locally advanced or metastatic transitional cell bladder cancer.

In another embodiment, an immunomodulatory compound is administered in combination with a second active ingredient as follows: temozolomide to pediatric patients with relapsed or progressive brain tumors or recurrent neuroblastoma; celecoxib, etoposide and cyclophosphamide for relapsed or progressive CNS cancer; temodar to patients with recurrent or progressive meningioma, malignant meningioma, hemangiopericytoma, multiple brain metastases, relapsed brain tumors, or newly diagnosed glioblastoma multiformis; irinotecan to patients with recurrent glioblastoma; carboplatin to pediatric patients with brain stem glioma; procarbazine to pediatric patients with progressive malignant gliomas; cyclophosphamide to patients with poor prognosis malignant brain tumors, newly diagnosed or recurrent glioblastoma multiformis; Gliadel® for high grade recurrent malignant gliomas; temozolomide and tamoxifen for anaplastic astrocytoma; or topotecan for gliomas, glioblastoma, anaplastic astrocytoma or anaplastic oligodendroglioma.

In another embodiment, an immunomodulatory compound is administered with methotrexate and cyclophosphamide to patients with metastatic breast cancer.

In another embodiment, an immunomodulatory compound is administered with temozolomide to patients with neuroendocrine tumors.

In another embodiment, an immunomodulatory compound is administered with gemcitabine to patients with recurrent or metastatic head or neck cancer. In another embodiment, an immunomodulatory compound is administered with gemcitabine to patients with pancreatic cancer.

In another embodiment, an immunomodulatory compound is administered to patients with colon cancer in combination with Arisa®, taxol and/or taxotere.

In another embodiment, an immunomodulatory compound is administered with capecitabine to patients with refractory colorectal cancer or patients who fail first line therapy or have poor performance in colon or rectal adenocarcinoma.

In another embodiment, an immunomodulatory compound is administered in combination with fluorouracil, leucovorin, and irinotecan to patients with Dukes C & D colorectal cancer or to patients who have been previously treated for metastatic colorectal cancer.

In another embodiment, an immunomodulatory compound is administered to patients with refractory colorectal cancer in combination with capecitabine, xeloda, and/or CPT-11.

In another embodiment, an immunomodulatory compound of the invention is administered with capecitabine and irinotecan to patients with refractory colorectal cancer or to patients with unresectable or metastatic colorectal carcinoma.

In another embodiment, an immunomodulatory compound is administered alone or in combination with interferon alpha or capecitabine to patients with unresectable or metastatic hepatocellular carcinoma; or with cisplatin and thiotepa to patients with primary or metastatic liver cancer.

In another embodiment, an immunomodulatory compound is administered in combination with pegylated interferon alpha to patients with Kaposi's sarcoma.

In another embodiment, an immunomodulatory compound is administered in combination with fludarabine, carboplatin, and/or topotecan to patients with refractory or relapsed or high-risk acuted myelogenous leukemia.

In another embodiment, an immunomodulatory compound is administered in combination with liposomal daunorubicin,

US 8,648,095 B2

21

topotecan and/or cytarabine to patients with unfavorable karyotype acute myeloblastic leukemia.

In another embodiment, an immunomodulatory compound is administered in combination with gemcitabine and irinotecan to patients with non-small cell lung cancer. In one embodiment, an immunomodulatory compound is administered in combination with carboplatin and irinotecan to patients with non-small cell lung cancer. In one embodiment, an immunomodulatory compound is administered with doxetaxol to patients with non-small cell lung cancer who have been previously treated with carbo/VP 16 and radiotherapy.

In another embodiment, an immunomodulatory compound is administered in combination with carboplatin and/or taxotere, or in combination with carboplatin, paclitaxel and/or thoracic radiotherapy to patients with non-small cell lung cancer. In a specific embodiment, an immunomodulatory compound is administered in combination with taxotere to patients with stage IIIB or IV non-small cell lung cancer.

In another embodiment, an immunomodulatory compound of the invention is administered in combination with oblimersen (Genasense®) to patients with small cell lung cancer.

In another embodiment, an immunomodulatory compound is administered alone or in combination with a second active ingredient such as vinblastine or fludarabine to patients with various types of lymphoma, including, but not limited to, Hodgkin's lymphoma, non-Hodgkin's lymphoma, cutaneous T-Cell lymphoma, cutaneous B-Cell lymphoma, diffuse large B-Cell lymphoma or relapsed or refractory low grade follicular lymphoma.

In another embodiment, an immunomodulatory compound is administered in combination with taxotere, IL-2, IFN, GM-CSF, and/or dacarbazine to patients with various types or stages of melanoma.

In another embodiment, an immunomodulatory compound is administered alone or in combination with vinorelbine to patients with malignant mesothelioma, or stage IIIB non-small cell lung cancer with pleural implants or malignant pleural effusion mesothelioma syndrome.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of multiple myeloma in combination with dexamethasone, zoledronic acid, palmitronate, GM-CSF, biacin, vinblastine, melphalan, busulphan, cyclophosphamide, IFN, palmidronate, prednisone, bisphosphonate, celecoxib, arsenic trioxide, PEG INTRON-A, vincristine, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with relapsed or refractory multiple myeloma in combination with doxorubicin (Doxil®), vincristine and/or dexamethasone (Decadron®).

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of ovarian cancer such as peritoneal carcinoma, papillary serous carcinoma, refractory ovarian cancer or recurrent ovarian cancer, in combination with taxol, carboplatin, doxorubicin, gemcitabine, cisplatin, xeloda, paclitaxel, dexamethasone, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of prostate cancer, in combination with xeloda, 5 FU/LV, gemcitabine, irinotecan plus gemcitabine, cyclophosphamide, vincristine, dexamethasone, GM-CSF, celecoxib, taxotere, ganciclovir, paclitaxel, adriamycin, docetaxel, estramustine, Emcyt, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of

22

renal cell cancer, in combination with capecitabine, IFN, tamoxifen, IL-2, GM-CSF, Celebrex®, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of gynecologic, uterus or soft tissue sarcoma cancer in combination with IFN, a COX-2 inhibitor such as Celebrex®, and/or sulindac.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of solid tumors in combination with celebrex, etoposide, cyclophosphamide, docetaxel, apecitabine, IFN, tamoxifen, IL-2, GM-CSF, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with scleroderma or cutaneous vasculitis in combination with celebrex, etoposide, cyclophosphamide, docetaxel, apecitabine, IFN, tamoxifen, IL-2, GM-CSF, or a combination thereof.

This invention also encompasses a method of increasing the dosage of an anti-cancer drug or agent that can be safely and effectively administered to a patient, which comprises administering to a patient (e.g., a human) an immunomodulatory compound of the invention, or a pharmaceutically acceptable derivative, salt, solvate, clathrate, hydrate, or prodrug thereof. Patients that can benefit by this method are those likely to suffer from an adverse effect associated with anti-cancer drugs for treating a specific cancer of the skin, subcutaneous tissue, lymph nodes, brain, lung, liver, bone, intestine, colon, heart, pancreas, adrenal, kidney, prostate, breast, colorectal, or combinations thereof. The administration of an immunomodulatory compound of the invention alleviates or reduces adverse effects which are of such severity that it would otherwise limit the amount of anti-cancer drug.

In one embodiment, an immunomodulatory compound of the invention can be administered orally and daily in an amount of from about 0.1 to about 150 mg, and preferably from about 1 to about 50 mg, more preferably from about 2 to about 25 mg prior to, during, or after the occurrence of the adverse effect associated with the administration of an anti-cancer drug to a patient. In a particular embodiment, an immunomodulatory compound of the invention is administered in combination with specific agents such as heparin, aspirin, coumadin, or G-CSF to avoid adverse effects that are associated with anti-cancer drugs such as but not limited to neutropenia or thrombocytopenia.

In one embodiment, an immunomodulatory compound of the invention can be administered to patients with diseases and disorders associated with, or characterized by, undesired angiogenesis in combination with additional active ingredients including but not limited to anti-cancer drugs, anti-inflammatories, antihistamines, antibiotics, and steroids.

In another embodiment, this invention encompasses a method of treating, preventing and/or managing cancer, which comprises administering an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, in conjunction with (e.g. before, during, or after) conventional therapy including, but not limited to, surgery, immunotherapy, biological therapy, radiation therapy, or other non-drug based therapy presently used to treat, prevent or manage cancer. The combined use of the immunomodulatory compounds of the invention and conventional therapy may provide a unique treatment regimen that is unexpectedly effective in certain patients. Without being limited by theory, it is believed that immunomodulatory compounds of the invention may provide additive or synergistic effects when given concurrently with conventional therapy.

US 8,648,095 B2

23

As discussed elsewhere herein, the invention encompasses a method of reducing, treating and/or preventing adverse or undesired effects associated with conventional therapy including, but not limited to, surgery, chemotherapy, radiation therapy, hormonal therapy, biological therapy and immunotherapy. One or more immunomodulatory compounds of the invention and other active ingredient can be administered to a patient prior to, during, or after the occurrence of the adverse effect associated with conventional therapy.

In one embodiment, an immunomodulatory compound of the invention can be administered in an amount of from about 0.1 to about 150 mg, and preferably from about 1 to about 25 mg, more preferably from about 2 to about 10 mg orally and daily alone, or in combination with a second active agent disclosed herein (see, e.g., section 5.2), prior to, during, or after the use of conventional therapy.

In a specific embodiment of this method, an immunomodulatory compound of the invention and doxorubicin are administered to patients with non-small cell lung cancer who were previously treated with carbo/VP 16 and radiotherapy.

5.3.2 Use with Transplantation Therapy

Compounds of the invention can be used to reduce the risk of Graft Versus Host Disease (GVHD). Therefore, the invention encompasses a method of treating, preventing and/or managing cancer, which comprises administering the immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, in conjunction with transplantation therapy.

As those of ordinary skill in the art are aware, the treatment of cancer is often based on the stages and mechanism of the disease. For example, as inevitable leukemic transformation develops in certain stages of cancer, transplantation of peripheral blood stem cells, hematopoietic stem cell preparation or bone marrow may be necessary. The combined use of the immunomodulatory compound of the invention and transplantation therapy provides a unique and unexpected synergism. In particular, an immunomodulatory compound of the invention exhibits immunomodulatory activity that may provide additive or synergistic effects when given concurrently with transplantation therapy in patients with cancer.

An immunomodulatory compound of the invention can work in combination with transplantation therapy reducing complications associated with the invasive procedure of transplantation and risk of GVHD. This invention encompasses a method of treating, preventing and/or managing cancer which comprises administering to a patient (e.g., a human) an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, before, during, or after the transplantation of umbilical cord blood, placental blood, peripheral blood stem cell, hematopoietic stem cell preparation or bone marrow. Examples of stem cells suitable for use in the methods of the invention are disclosed in U.S. provisional patent application No. 60/372,348, filed Apr. 12, 2002 by R. Hariri et al., the entirety of which is incorporated herein by reference.

In one embodiment of this method, an immunomodulatory compound of the invention is administered to patients with multiple myeloma before, during, or after the transplantation of autologous peripheral blood progenitor cell.

In another embodiment, an immunomodulatory compound is administered to patients with relapsing multiple myeloma after the stem cell transplantation.

24

In another embodiment, an immunomodulatory compound and prednisone are administered as maintenance therapy to patients with multiple myeloma following the transplantation of autologous stem cell.

In another embodiment, an immunomodulatory compound and dexamethasone are administered as salvage therapy for low risk post transplantation to patients with multiple myeloma.

In another embodiment, an immunomodulatory compound and dexamethasone are administered as maintenance therapy to patients with multiple myeloma following the transplantation of autologous bone marrow.

In another embodiment, an immunomodulatory compound is administered following the administration of high dose of melphalan and the transplantation of autologous stem cell to patients with chemotherapy responsive multiple myeloma.

In another embodiment, an immunomodulatory compound and PEG INTRO-A are administered as maintenance therapy to patients with multiple myeloma following the transplantation of autologous CD34-selected peripheral stem cell.

In another embodiment, an immunomodulatory compound is administered with post transplant consolidation chemotherapy to patients with newly diagnosed multiple myeloma to evaluate anti-angiogenesis.

In another embodiment, an immunomodulatory compound and dexamethasone are administered as maintenance therapy after DCEP consolidation, following the treatment with high dose of melphalan and the transplantation of peripheral blood stem cell to 65 years of age or older patients with multiple myeloma.

5.3.3 Cycling Therapy

In certain embodiments, the prophylactic or therapeutic agents of the invention are cyclically administered to a patient. Cycling therapy involves the administration of an active agent for a period of time, followed by a rest for a period of time, and repeating this sequential administration. Cycling therapy can reduce the development of resistance to one or more of the therapies, avoid or reduce the side effects of one of the therapies, and/or improves the efficacy of the treatment.

Consequently, in one specific embodiment of the invention, an immunomodulatory compound of the invention is administered daily in a single or divided doses in a four to six week cycle with a rest period of about a week or two weeks. The invention further allows the frequency, number, and length of dosing cycles to be increased. Thus, another specific embodiment of the invention encompasses the administration of an immunomodulatory compound of the invention for more cycles than are typical when it is administered alone. In yet another specific embodiment of the invention, an immunomodulatory compound of the invention is administered for a greater number of cycles that would typically cause dose-limiting toxicity in a patient to whom a second active ingredient is not also being administered.

In one embodiment, an immunomodulatory compound of the invention is administered daily and continuously for three or four weeks at a dose of from about 0.1 to about 150 mg/d followed by a break of one or two weeks. Actimid™ is preferably administered daily and continuously at an initial dose of 0.1 to 5 mg/d with dose escalation (every week) by 1 to 10 mg/d to a maximum dose of 50 mg/d for as long as therapy is tolerated. In a particular embodiment, Revimid™ is administered in an amount of about 5, 10, or 25 mg/day, preferably in an amount of about 10 mg/day for three to four weeks, followed by one week or two weeks of rest in a four or six week cycle.

In one embodiment of the invention, an immunomodulatory compound of the invention and a second active ingredient are administered orally, with administration of an immunomodulatory compound of the invention occurring 30 to 60 minutes prior to a second active ingredient, during a cycle of four to six weeks. In another embodiment of the invention, the combination of an immunomodulatory compound of the invention and a second active ingredient is administered by intravenous infusion over about 90 minutes every cycle. In a specific embodiment, one cycle comprises the administration of from about 10 to about 25 mg/day of Revimid™ and from about 50 to about 200 mg/m²/day of a second active ingredient daily for three to four weeks and then one or two weeks of rest. In another specific embodiment, each cycle comprises the administration of from about 5 to about 10 mg/day of Actimid™ and from about 50 to about 200 mg/m²/day of a second active ingredient for 3 to 4 weeks followed by one or two weeks of rest. Typically, the number of cycles during which the combinatorial treatment is administered to a patient will be from about one to about 24 cycles, more typically from about two to about 16 cycles, and even more typically from about four to about three cycles.

5.4 Pharmaceutical Compositions and Dosage Forms

Pharmaceutical compositions can be used in the preparation of individual, single unit dosage forms. Pharmaceutical compositions and dosage forms of the invention comprise an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof. Pharmaceutical compositions and dosage forms of the invention can further comprise one or more excipients.

Pharmaceutical compositions and dosage forms of the invention can also comprise one or more additional active ingredients. Consequently, pharmaceutical compositions and dosage forms of the invention comprise the active ingredients disclosed herein (e.g., an immunomodulatory compound and a second active agent). Examples of optional second, or additional, active ingredients are disclosed herein (see, e.g., section 5.2).

Single unit dosage forms of the invention are suitable for oral, mucosal (e.g., nasal, sublingual, vaginal, buccal, or rectal), parenteral (e.g., subcutaneous, intravenous, bolus injection, intramuscular, or intraarterial), topical (e.g., eye drops or other ophthalmic preparations), transdermal or transcutaneous administration to a patient. Examples of dosage forms include, but are not limited to: tablets; caplets; capsules, such as soft elastic gelatin capsules; cachets; troches; lozenges; dispersions; suppositories; powders; aerosols (e.g., nasal sprays or inhalers); gels; liquid dosage forms suitable for oral or mucosal administration to a patient, including suspensions (e.g., aqueous or non-aqueous liquid suspensions, oil-in-water emulsions, or a water-in-oil liquid emulsions), solutions, and elixirs; liquid dosage forms suitable for parenteral administration to a patient; eye drops or other ophthalmic preparations suitable for topical administration; and sterile solids (e.g., crystalline or amorphous solids) that can be reconstituted to provide liquid dosage forms suitable for parenteral administration to a patient.

The composition, shape, and type of dosage forms of the invention will typically vary depending on their use. For example, a dosage form used in the acute treatment of a disease may contain larger amounts of one or more of the active ingredients it comprises than a dosage form used in the chronic treatment of the same disease. Similarly, a parenteral dosage form may contain smaller amounts of one or more of the active ingredients it comprises than an oral dosage form used to treat the same disease. These and other ways in which

specific dosage forms encompassed by this invention will vary from one another will be readily apparent to those skilled in the art. See, e.g., *Remington's Pharmaceutical Sciences*, 18th ed., Mack Publishing, Easton Pa. (1990).

Typical pharmaceutical compositions and dosage forms comprise one or more excipients. Suitable excipients are well known to those skilled in the art of pharmacy, and non-limiting examples of suitable excipients are provided herein. Whether a particular excipient is suitable for incorporation into a pharmaceutical composition or dosage form depends on a variety of factors well known in the art including, but not limited to, the way in which the dosage form will be administered to a patient. For example, oral dosage forms such as tablets may contain excipients not suited for use in parenteral dosage forms. The suitability of a particular excipient may also depend on the specific active ingredients in the dosage form. For example, the decomposition of some active ingredients may be accelerated by some excipients such as lactose, or when exposed to water. Active ingredients that comprise primary or secondary amines are particularly susceptible to such accelerated decomposition. Consequently, this invention encompasses pharmaceutical compositions and dosage forms that contain little, if any, lactose other mono- or disaccharides. As used herein, the term "lactose-free" means that the amount of lactose present, if any, is insufficient to substantially increase the degradation rate of an active ingredient.

Lactose-free compositions of the invention can comprise excipients that are well known in the art and are listed, for example, in the *U.S. Pharmacopeia* (USP) 25-NF20 (2002). In general, lactose-free compositions comprise active ingredients, a binder/filler, and a lubricant in pharmaceutically compatible and pharmaceutically acceptable amounts. Preferred lactose-free dosage forms comprise active ingredients, microcrystalline cellulose, pre-gelatinized starch, and magnesium stearate.

This invention further encompasses anhydrous pharmaceutical compositions and dosage forms comprising active ingredients, since water can facilitate the degradation of some compounds. For example, the addition of water (e.g., 5%) is widely accepted in the pharmaceutical arts as a means of simulating long-term storage in order to determine characteristics such as shelf-life or the stability of formulations over time. See, e.g., Jens T. Carstensen, *Drug Stability: Principles & Practice*, 2d. Ed., Marcel Dekker, NY, N.Y., 1995, pp. 379-80. In effect, water and heat accelerate the decomposition of some compounds. Thus, the effect of water on a formulation can be of great significance since moisture and/or humidity are commonly encountered during manufacture, handling, packaging, storage, shipment, and use of formulations.

Anhydrous pharmaceutical compositions and dosage forms of the invention can be prepared using anhydrous or low moisture containing ingredients and low moisture or low humidity conditions. Pharmaceutical compositions and dosage forms that comprise lactose and at least one active ingredient that comprises a primary or secondary amine are preferably anhydrous if substantial contact with moisture and/or humidity during manufacturing, packaging, and/or storage is expected.

An anhydrous pharmaceutical composition should be prepared and stored such that its anhydrous nature is maintained. Accordingly, anhydrous compositions are preferably packaged using materials known to prevent exposure to water such that they can be included in suitable formulary kits. Examples

of suitable packaging include, but are not limited to, hermetically sealed foils, plastics, unit dose containers (e.g., vials), blister packs, and strip packs.

The invention further encompasses pharmaceutical compositions and dosage forms that comprise one or more compounds that reduce the rate by which an active ingredient will decompose. Such compounds, which are referred to herein as “stabilizers,” include, but are not limited to, antioxidants such as ascorbic acid, pH buffers, or salt buffers.

Like the amounts and types of excipients, the amounts and specific types of active ingredients in a dosage form may differ depending on factors such as, but not limited to, the route by which it is to be administered to patients. However, typical dosage forms of the invention comprise an immunomodulatory compound of the invention or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof in an amount of from about 0.10 to about 150 mg. Typical dosage forms comprise an immunomodulatory compound of the invention or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof in an amount of about 0.1, 1, 2, 5, 7.5, 10, 12.5, 15, 17.5, 20, 25, 50, 100, 150 or 200 mg. In a particular embodiment, a preferred dosage form comprises 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™) in an amount of about 1, 2, 5, 10, 25 or 50 mg. In a specific embodiment, a preferred dosage form comprises 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) in an amount of about 5, 10, 25 or 50 mg. Typical dosage forms comprise the second active ingredient in an amount of 1 to about 1000 mg, from about 5 to about 500 mg, from about 10 to about 350 mg, or from about 50 to about 200 mg. Of course, the specific amount of the anti-cancer drug will depend on the specific agent used, the type of cancer being treated or managed, and the amount(s) of an immunomodulatory compound of the invention and any optional additional active agents concurrently administered to the patient.

5.4.1 Oral Dosage Forms

Pharmaceutical compositions of the invention that are suitable for oral administration can be presented as discrete dosage forms, such as, but are not limited to, tablets (e.g., chewable tablets), caplets, capsules, and liquids (e.g., flavored syrups). Such dosage forms contain predetermined amounts of active ingredients, and may be prepared by methods of pharmacy well known to those skilled in the art. See generally, *Remington's Pharmaceutical Sciences*, 18th ed., Mack Publishing, Easton Pa. (1990).

Typical oral dosage forms of the invention are prepared by combining the active ingredients in an intimate admixture with at least one excipient according to conventional pharmaceutical compounding techniques. Excipients can take a wide variety of forms depending on the form of preparation desired for administration. For example, excipients suitable for use in oral liquid or aerosol dosage forms include, but are not limited to, water, glycols, oils, alcohols, flavoring agents, preservatives, and coloring agents. Examples of excipients suitable for use in solid oral dosage forms (e.g., powders, tablets, capsules, and caplets) include, but are not limited to, starches, sugars, micro-crystalline cellulose, diluents, granulating agents, lubricants, binders, and disintegrating agents.

Because of their ease of administration, tablets and capsules represent the most advantageous oral dosage unit forms, in which case solid excipients are employed. If desired, tablets can be coated by standard aqueous or nonaqueous techniques. Such dosage forms can be prepared by any of the methods of pharmacy. In general, pharmaceutical compositions and dosage forms are prepared by uniformly and inti-

ately admixing the active ingredients with liquid carriers, finely divided solid carriers, or both, and then shaping the product into the desired presentation if necessary.

For example, a tablet can be prepared by compression or molding. Compressed tablets can be prepared by compressing in a suitable machine the active ingredients in a free-flowing form such as powder or granules, optionally mixed with an excipient. Molded tablets can be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

Examples of excipients that can be used in oral dosage forms of the invention include, but are not limited to, binders, fillers, disintegrants, and lubricants. Binders suitable for use in pharmaceutical compositions and dosage forms include, but are not limited to, corn starch, potato starch, or other starches, gelatin, natural and synthetic gums such as acacia, sodium alginate, alginic acid, other alginates, powdered tragacanth, guar gum, cellulose and its derivatives (e.g., ethyl cellulose, cellulose acetate, carboxymethyl cellulose calcium, sodium carboxymethyl cellulose), polyvinyl pyrrolidone, methyl cellulose, pre-gelatinized starch, hydroxypropyl methyl cellulose, (e.g., Nos. 2208, 2906, 2910), microcrystalline cellulose, and mixtures thereof.

Suitable forms of microcrystalline cellulose include, but are not limited to, the materials sold as AVICEL-PH-101, AVICEL-PH-103 AVICEL RC-581, AVICEL-PH-105 (available from FMC Corporation, American Viscose Division, Avicel Sales, Marcus Hook, Pa.), and mixtures thereof. A specific binder is a mixture of microcrystalline cellulose and sodium carboxymethyl cellulose sold as AVICEL RC-581. Suitable anhydrous or low moisture excipients or additives include AVICEL-PH-103™ and Starch 1500 LM.

Examples of fillers suitable for use in the pharmaceutical compositions and dosage forms disclosed herein include, but are not limited to, talc, calcium carbonate (e.g., granules or powder), microcrystalline cellulose, powdered cellulose, dextrates, kaolin, mannitol, silicic acid, sorbitol, starch, pre-gelatinized starch, and mixtures thereof. The binder or filler in pharmaceutical compositions of the invention is typically present in from about 50 to about 99 weight percent of the pharmaceutical composition or dosage form.

Disintegrants are used in the compositions of the invention to provide tablets that disintegrate when exposed to an aqueous environment. Tablets that contain too much disintegrant may disintegrate in storage, while those that contain too little may not disintegrate at a desired rate or under the desired conditions. Thus, a sufficient amount of disintegrant that is neither too much nor too little to detrimentally alter the release of the active ingredients should be used to form solid oral dosage forms of the invention. The amount of disintegrant used varies based upon the type of formulation, and is readily discernible to those of ordinary skill in the art. Typical pharmaceutical compositions comprise from about 0.5 to about 15 weight percent of disintegrant, preferably from about 1 to about 5 weight percent of disintegrant.

Disintegrants that can be used in pharmaceutical compositions and dosage forms of the invention include, but are not limited to, agar-agar, alginic acid, calcium carbonate, microcrystalline cellulose, croscarmellose sodium, crospovidone, polacrillin potassium, sodium starch glycolate, potato or tapioca starch, other starches, pre-gelatinized starch, other starches, clays, other alginates, other celluloses, gums, and mixtures thereof.

Lubricants that can be used in pharmaceutical compositions and dosage forms of the invention include, but are not limited to, calcium stearate, magnesium stearate, mineral oil, light mineral oil, glycerin, sorbitol, mannitol, polyethylene

US 8,648,095 B2

29

glycol, other glycols, stearic acid, sodium lauryl sulfate, talc, hydrogenated vegetable oil (e.g., peanut oil, cottonseed oil, sunflower oil, sesame oil, olive oil, corn oil, and soybean oil), zinc stearate, ethyl oleate, ethyl laureate, agar, and mixtures thereof. Additional lubricants include, for example, a syloid silica gel (AEROSIL200, manufactured by W.R. Grace Co. of Baltimore, Md.), a coagulated aerosol of synthetic silica (marketed by Degussa Co. of Plano, Tex.), CAB-O-SIL (a pyrogenic silicon dioxide product sold by Cabot Co. of Boston, Mass.), and mixtures thereof. If used at all, lubricants are typically used in an amount of less than about 1 weight percent of the pharmaceutical compositions or dosage forms into which they are incorporated.

A preferred solid oral dosage form of the invention comprises an immunomodulatory compound of the invention, anhydrous lactose, microcrystalline cellulose, polyvinylpyrrolidone, stearic acid, colloidal anhydrous silica, and gelatin.

5.4.2 Delayed Release Dosage Forms

Active ingredients of the invention can be administered by controlled release means or by delivery devices that are well known to those of ordinary skill in the art. Examples include, but are not limited to, those described in U.S. Pat. Nos. 3,845,770; 3,916,899; 3,536,809; 3,598,123; and 4,008,719, 5,674,533, 5,059,595, 5,591,767, 5,120,548, 5,073,543, 5,639,476, 5,354,556, and 5,733,566, each of which is incorporated herein by reference. Such dosage forms can be used to provide slow or controlled-release of one or more active ingredients using, for example, hydropropylmethyl cellulose, other polymer matrices, gels, permeable membranes, osmotic systems, multilayer coatings, microparticles, liposomes, microspheres, or a combination thereof to provide the desired release profile in varying proportions. Suitable controlled-release formulations known to those of ordinary skill in the art, including those described herein, can be readily selected for use with the active ingredients of the invention. The invention thus encompasses single unit dosage forms suitable for oral administration such as, but not limited to, tablets, capsules, gelcaps, and caplets that are adapted for controlled-release.

All controlled-release pharmaceutical products have a common goal of improving drug therapy over that achieved by their non-controlled counterparts. Ideally, the use of an optimally designed controlled-release preparation in medical treatment is characterized by a minimum of drug substance being employed to cure or control the condition in a minimum amount of time. Advantages of controlled-release formulations include extended activity of the drug, reduced dosage frequency, and increased patient compliance. In addition, controlled-release formulations can be used to affect the time of onset of action or other characteristics, such as blood levels of the drug, and can thus affect the occurrence of side (e.g., adverse) effects.

Most controlled-release formulations are designed to initially release an amount of drug (active ingredient) that promptly produces the desired therapeutic effect, and gradually and continually release of other amounts of drug to maintain this level of therapeutic or prophylactic effect over an extended period of time. In order to maintain this constant level of drug in the body, the drug must be released from the dosage form at a rate that will replace the amount of drug being metabolized and excreted from the body. Controlled-release of an active ingredient can be stimulated by various conditions including, but not limited to, pH, temperature, enzymes, water, or other physiological conditions or compounds.

30

5.4.3 Parenteral Dosage Forms

Parenteral dosage forms can be administered to patients by various routes including, but not limited to, subcutaneous, intravenous (including bolus injection), intramuscular, and intraarterial. Because their administration typically bypasses patients' natural defenses against contaminants, parenteral dosage forms are preferably sterile or capable of being sterilized prior to administration to a patient. Examples of parenteral dosage forms include, but are not limited to, solutions ready for injection, dry products ready to be dissolved or suspended in a pharmaceutically acceptable vehicle for injection, suspensions ready for injection, and emulsions.

Suitable vehicles that can be used to provide parenteral dosage forms of the invention are well known to those skilled in the art. Examples include, but are not limited to: Water for Injection USP; aqueous vehicles such as, but not limited to, Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, and Lactated Ringer's Injection; water-miscible vehicles such as, but not limited to, ethyl alcohol, polyethylene glycol, and polypropylene glycol; and non-aqueous vehicles such as, but not limited to, corn oil, cottonseed oil, peanut oil, sesame oil, ethyl oleate, isopropyl myristate, and benzyl benzoate.

Compounds that increase the solubility of one or more of the active ingredients disclosed herein can also be incorporated into the parenteral dosage forms of the invention. For example, cyclodextrin and its derivatives can be used to increase the solubility of an immunomodulatory compound of the invention and its derivatives. See, e.g., U.S. Pat. No. 5,134,127, which is incorporated herein by reference.

5.4.4 Topical and Mucosal Dosage Forms

Topical and mucosal dosage forms of the invention include, but are not limited to, sprays, aerosols, solutions, emulsions, suspensions, eye drops or other ophthalmic preparations, or other forms known to one of skill in the art. See, e.g., *Remington's Pharmaceutical Sciences*, 16th and 18th eds., Mack Publishing, Easton Pa. (1980 & 1990); and *Introduction to Pharmaceutical Dosage Forms*, 4th ed., Lea & Febiger, Philadelphia (1985). Dosage forms suitable for treating mucosal tissues within the oral cavity can be formulated as mouthwashes or as oral gels.

Suitable excipients (e.g., carriers and diluents) and other materials that can be used to provide topical and mucosal dosage forms encompassed by this invention are well known to those skilled in the pharmaceutical arts, and depend on the particular tissue to which a given pharmaceutical composition or dosage form will be applied. With that fact in mind, typical excipients include, but are not limited to, water, acetone, ethanol, ethylene glycol, propylene glycol, butane-1,3-diol, isopropyl myristate, isopropyl palmitate, mineral oil, and mixtures thereof to form solutions, emulsions or gels, which are non-toxic and pharmaceutically acceptable. Moisturizers or humectants can also be added to pharmaceutical compositions and dosage forms if desired. Examples of such additional ingredients are well known in the art. See, e.g., *Remington's Pharmaceutical Sciences*, 16th and 18th eds., Mack Publishing, Easton Pa. (1980 & 1990).

The pH of a pharmaceutical composition or dosage form may also be adjusted to improve delivery of one or more active ingredients. Similarly, the polarity of a solvent carrier, its ionic strength, or tonicity can be adjusted to improve delivery. Compounds such as stearates can also be added to pharmaceutical compositions or dosage forms to advantageously alter the hydrophilicity or lipophilicity of one or more active ingredients so as to improve delivery. In this regard, stearates can serve as a lipid vehicle for the formulation, as an emulsifying agent or surfactant, and as a delivery-

US 8,648,095 B2

31

enhancing or penetration-enhancing agent. Different salts, hydrates or solvates of the active ingredients can be used to further adjust the properties of the resulting composition.

5.4.5 Kits

Typically, active ingredients of the invention are preferably not administered to a patient at the same time or by the same route of administration. This invention therefore encompasses kits which, when used by the medical practitioner, can simplify the administration of appropriate amounts of active ingredients to a patient.

A typical kit of the invention comprises a dosage form of an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, pro-drug, or clathrate thereof. Kits encompassed by this invention can further comprise additional active ingredients such as oblimersen (Genasense®), melphalan, G-CSF, GM-CSF, EPO, topotecan, dacarbazine, irinotecan, taxotere, IFN, COX-2 inhibitor, pentoxifylline, ciprofloxacin, dexamethasone, IL2, IL8, IL18, Ara-C, vinorelbine, isotretinoin, 13 cis-retinoic acid, or a pharmacologically active mutant or derivative thereof, or a combination thereof. Examples of the additional active ingredients include, but are not limited to, those disclosed herein (see, e.g., section 5.2).

Kits of the invention can further comprise devices that are used to administer the active ingredients. Examples of such devices include, but are not limited to, syringes, drip bags, patches, and inhalers.

Kits of the invention can further comprise cells or blood for transplantation as well as pharmaceutically acceptable vehicles that can be used to administer one or more active ingredients. For example, if an active ingredient is provided in a solid form that must be reconstituted for parenteral administration, the kit can comprise a sealed container of a suitable vehicle in which the active ingredient can be dissolved to form a particulate-free sterile solution that is suitable for parenteral administration. Examples of pharmaceutically acceptable vehicles include, but are not limited to: Water for Injection USP; aqueous vehicles such as, but not limited to, Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, and Lactated Ringer's Injection; water-miscible vehicles such as, but not limited to, ethyl alcohol, polyethylene glycol, and polypropylene glycol; and non-aqueous vehicles such as, but not limited to, corn oil, cottonseed oil, peanut oil, sesame oil, ethyl oleate, isopropyl myristate, and benzyl benzoate.

6. EXAMPLES

Certain embodiments of the invention are illustrated by the following non-limiting examples.

6.1 Modulation of Cytokine Production

A series of non-clinical pharmacology and toxicology studies have been performed to support the clinical evaluation of an immunomodulatory compound of the invention in human subjects. These studies were performed in accordance with internationally recognized guidelines for study design and in compliance with the requirements of Good Laboratory Practice (GLP), unless otherwise noted.

Inhibition of TNF- α production following LPS-stimulation of human PBMC and human whole blood by 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™), 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione and thalidomide (Revimid™) was investigated in vitro (Muller et al., *Bioorg. Med. Chem. Lett.* 9:1625-1630, 1999). The IC₅₀'s of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione for inhibiting production of TNF- α following LPS-stimulation of PBMC and

32

human whole blood were ~24 nM (6.55 ng/mL) and ~25 nM (6.83 ng/mL), respectively. In vitro studies suggest a pharmacological activity profile for 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione that is similar to, but at least 200 times more potent than, thalidomide. In vitro studies have also demonstrated that concentrations of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione of 2.73 to 27.3 ng/mL (0.01 to 0.1 μ M) achieved 50% inhibition of the proliferation of MM.1S and Hs Sultan cells.

The IC₅₀'s of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione for inhibiting production of TNF- α following LPS-stimulation of PBMC and human whole blood were ~100 nM (25.9 ng/mL) and ~480 nM (103.6 ng/mL), respectively. Thalidomide, in contrast, had an IC₅₀ of ~194 μ M (50.2 μ g/mL) for inhibiting production of TNF- α following LPS-stimulation of PBMC. In vitro studies suggest a pharmacological activity profile for 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione that is similar to, but 50 to 2000 times more potent than, thalidomide. It has been shown that the compound is approximately 50-100 times more potent than thalidomide in stimulating the proliferation of T-cells following primary induction by T-cell receptor (TCR) activation. 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione is also approximately 50 to 100 times more potent than thalidomide in augmenting the production of IL-2 and IFN- γ following TCR activation of PBMC (IL-2) or T-cells (IFN- γ). In addition, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione exhibited dose-dependent inhibition of LPS-stimulated production of the pro-inflammatory cytokines TNF- α , IL-1 β , and IL-6 by PBMC while it increased production of the anti-inflammatory cytokine IL-10.

6.2 Inhibition of MM Cell Proliferation

The ability of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) and thalidomide for comparison to effect the proliferation of MM cell lines has been investigated in an in vitro study. Uptake [³H]-thymidine by different MM cell lines (MM.1S, Hs Sultan, U266 and RPMI-8226) was measured as an indicator of cell proliferation. Cells were incubated in the presence of compounds for 48 hours; [³H]-thymidine was included for the last 8 hours of the incubation period. Addition of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione to MM.1S and Hs Sultan cells resulted in 50% inhibition of cell proliferation at concentrations of 0.4 μ M and 1 μ M, respectively. In contrast, addition of thalidomide at concentrations up to 100 μ M resulted in only 15% and 20% inhibition of cell proliferation in MM.1S and Hs Sultan cells, respectively. These data are summarized in FIG. 1.

6.3 Toxicology Studies

The effects of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) on cardiovascular and respiratory function are investigated in anesthetized dogs. Two groups of Beagle dogs (2/sex/group) are used. One group receives three doses of vehicle only and the other receives three ascending doses of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (2, 10, and 20 mg/kg). In all cases, doses of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione or vehicle are successively administered via infusion through the jugular vein separated by intervals of at least 30 minutes.

The cardiovascular and respiratory changes induced by 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione are minimal at all doses when compared to the vehicle control group. The only statistically significant difference between the vehicle and treatment groups is a small increase in arterial blood pressure (from 94 mmHg to 101

mmHg) following administration of the low dose of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. This effect lasts approximately 15 minutes and is not seen at higher doses. Deviations in femoral blood flow, respiratory parameters, and Qtc interval are common to both the control and treated groups and are not considered treatment-related.

6.4 Cycling Therapy in Patients

In a specific embodiment, an immunomodulatory compound of the invention are cyclically administered to patients with cancer. Cycling therapy involves the administration of a first agent for a period of time, followed by a rest for a period of time and repeating this sequential administration. Cycling therapy can reduce the development of resistance to one or more of the therapies, avoid or reduce the side effects of one of the therapies, and/or improves the efficacy of the treatment.

In a specific embodiment, prophylactic or therapeutic agents are administered in a cycle of about 4 to 6 weeks, about once or twice every day. One cycle can comprise the administration of a therapeutic or prophylactic agent for three to four weeks and at least a week or two weeks of rest. The number of cycles administered is from about one to about 24 cycles, more typically from about two to about 16 cycles, and more typically from about four to about eight cycles.

For example, in a cycle of four weeks, on day 1, the administration of 25 mg/d of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione is started. On day 22, the administration of the compound is stopped for a week of rest. On day 29, the administration of 25 mg/d 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione is begun.

6.5 Clinical Studies in Patients

6.5.1 Treatment of Relapsed Multiple Myeloma

4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™) was administered to patients with relapsed/refractory multiple myeloma. The study was conducted in compliance with Good Clinical Practices. Patients were at least 18 years old, had been diagnosed with multiple myeloma (with paraprotein in serum and/or urine), and were considered refractory to treatment after at least two cycles of treatment, or have relapsed after two cycles of treatment.

Patients who have progressive disease, according to the Southwest Oncology Group (SWOG) criteria, on their prior regimen are considered treatment refractory. Relapse following remission is defined as >25% increase in M component from baseline levels; reappearance of the M paraprotein that had previously disappeared; or a definite increase in the size and number of lytic bone lesions recognized on radiographs. Patients may have had prior therapy with thalidomide, provided they were able to tolerate the treatment. A Zubrod performance status of 0 to 2 is required for all patients.

4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione is administered to patients at doses of 1, 2, 5, or 10 mg/day for up to four weeks; at each dose level, three patients are initially enrolled. Dosing occurs at approximately the same time each morning; all doses are administered in the fasted state (no eating for at least two hours prior to dosing and two hours after dosing). 4-(amino)-2-(2,6-dioxo(3-piperidyl))-

isoindoline-1,3-dione doses are administered in an ascending fashion such that patients in the first cohort receive the lowest dose of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (1 mg/day) and escalation to the next higher dose level occurs only following the establishment of safety and tolerability at the current dose. If one out of three patients at any dose level experience dose limiting toxicity (DLT), three additional patients are enrolled at that dose. If none of the three additional patients experience DLT, escalation to the next dose level occurs; dose escalations continue in a similar fashion until the MTD is established or the maximum daily dose (10 mg/day) is attained. However, if one of the three additional patients enrolled experiences DLT, the MTD has been reached. If two or more of the three additional patients enrolled experience DLT, the MTD is judged to have been exceeded and three additional patients are enrolled at the preceding dose level to confirm the MTD. Once the MTD has been identified, four additional patients are enrolled at that dose level so that a total of 10 patients is treated at the MTD.

Blood sampling for analysis of pharmacokinetic parameters is performed on Days 1 and 28 according to the following sampling schedule: pre-dose, 0.25, 0.5, 0.75, 1, 1.5, 2, 2.5, 3, 4, 6, 8, 10, 12, 18, and 24 hours post-dose. An additional blood sample is collected at each weekly visit for the determination of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione levels. Total urine collections are also made with urine pooled according to the following time intervals post-dose: 0 to 4, 4 to 8, 8 to 12, and 12 to 24 hours. Safety assessments are made by monitoring adverse events, vital signs, ECGs, clinical laboratory evaluations (blood chemistry, hematology, lymphocyte phenotyping, and urinalysis), and physical examination at specific times during the study.

Results of interim pharmacokinetic analyses obtained following single- and multiple-dose administration of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione to multiple myeloma patients are presented below in Tables 1 and 2. These data show that 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione was steadily absorbed at all dose levels in relapsed multiple myeloma patients. Maximum plasma concentrations occurred at a median T_{max} of between 2.5 and 2.8 hours post-dose at Day 1 and between 3 and 4 hours post-dose at Week 4. At all doses, plasma concentrations declined in a monophasic manner after reaching C_{max} . The start of the elimination phase occurred between 3 and 10 hours post-dose at Day 1 and Week 4, respectively.

These data also showed that after 4 weeks of dosing, 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione accumulated to a small extent (mean accumulation ratios ~1.02 to 1.52 and ~0.94 to 1.62 for C_{max} and $AUC_{(0-\tau)}$, respectively). There was almost a dose proportional increase in $AUC_{(0-\tau)}$ and C_{max} values with increasing dose. A five-fold higher dose of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione produced a 3.2- and 2.2-fold increase in C_{max} at Day 1 and Week 4, respectively. Similarly, a 5-fold increase in dose resulted in a 3.6- and 2.3-fold increase in $AUC_{(0-\tau)}$, at Day 1 and Week 4, respectively.

TABLE 1

| Pharmacokinetic parameters of Actimid™ in relapsed multiple myeloma patients | | | | |
|--|-------|--------------|--------------|---------------|
| Parameter | | 1 mg | 2 mg | 5 mg |
| | | (N = 6) | (N = 2) | (N = 3) |
| Day 1 | | | | |
| C_{max} | ng/mL | 15.03 (4.04) | 24.4* (12.1) | 48.56 (14.03) |
| t_{max} | h | 3.3 (2.6) | 2.7* (0.3) | 2.3 (0.3) |

US 8,648,095 B2

35

36

TABLE 1-continued

| Pharmacokinetic parameters of Actimid™ in relapsed multiple myeloma patients | | | | |
|--|-----------|----------------|----------------|-----------------|
| Parameter | | 1 mg | 2 mg | 5 mg |
| | | (N = 6) | (N = 2) | (N = 3) |
| Day 1 | | | | |
| AUC _(0-∞) | ng · h/mL | 152.90 (36.62) | 279.18 (51.10) | 593.10 (335.23) |
| AUC _(0-τ) | | 134.21 (27.14) | 249.57 (29.26) | 520.94 (267.32) |
| t _{1/2} | h | 7.3 (3.4) | 6.3 (1.4) | 6.5 (2.2) |
| CL/F | mL/min | 114.75 (29.20) | 121.43 (22.22) | 182.31 (117.06) |
| Vz/f | L | 69.55 (44.97) | 65.31 (2.80) | 87.24 (22.61) |

τ = 24 hours

N/A = not available

TABLE 2

| Pharmacokinetic parameters of Actimid™ following multiple oral doses(1, 2, and 5 mg/day) in relapsed multiple myeloma patients | | | | |
|--|-----------|-----------------|-----------------|-----------------|
| Parameter | | 1 mg | 2 mg | 5 mg |
| | | (N = 5) | (N = 2) | (N = 3) |
| Week 4 | | | | |
| C _{max} | ng/mL | 23.20 (7.48) | 30.05* (15.64) | 58.07 (38.08) |
| t _{max} | h | 3.6 (1.5) | 2.8* (0.3) | 5.0 (2.6) |
| AUC _(0-∞) | ng · h/mL | N/A | N/A | N/A |
| AUC _(0-τ) | | 239.31 (122.59) | 269.36 (186.34) | 597.24 (354.23) |
| t _{1/2} | h | 6.2* (0.6) | 7.7 (2.8) | 7.8 (4.0) |
| CL/F | mL/min | 87.85 (48.48) | 162.68 (112.54) | 207.50 (175.41) |
| Vz/f | L | 41.35* (8.84) | 95.04 (35.39) | 103.95 (27.25) |

τ = 24 hours

N/A = not available

*N = 3 patients

6.5.2 Treatment of Relapsed Multiple Myeloma

Two Phase 1 clinical studies of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) have been conducted to identify the maximum tolerated dose (MTD) in patients with refractory or relapsed multiple myeloma. These studies have also characterized the safety profile of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione when ascending doses of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione were given orally for up to 4 weeks. Patients started 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione treatment at 5 mg/day with subsequent escalation to 10, 25, and 50 mg/day. Patients were enrolled for 28 days at their assigned dose, with the option of extended treatment for those who did not exhibit disease progression or experience dose limiting toxicity (DLT). Patients were evaluated for adverse events at each visit and the severity of these events was graded according to the National Cancer Institute (NCI) Common Toxicity Criteria. Patients were discontinued if they experienced DLT (Grade 3 or greater non-hematological, or Grade 4 hematological toxicity).

In this study, 27 patients were enrolled. All patients had relapsed multiple myeloma and 18 (72%) were refractory to salvage therapy. Among these patients, 15 had undergone prior autologous stem cell transplantation and 16 patients had received prior thalidomide treatment. The median number of prior regimens was 3 (range 2 to 6).

Blood and urine samples were collected for analysis of pharmacokinetic parameters on Days 1 and 28. Blood samples were collected according to the following sampling schedule: pre-dose, 0.25, 0.5, 0.75, 1, 1.5, 2, 2.5, 3, 4, 6, 8, 10, 12, 18, and 24 hours post-dose. In addition, a blood sample was collected at each weekly clinic visit for 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione deter-

mination. Total urine was collected and pooled according to the following time intervals post-dose: 0 to 4, 4 to 8, 8 to 12, and 12 to 24 hours. Response to treatment was assessed by M-protein quantification (by immunoelectrophoresis) from serum and a 24-hour urine collection, with creatinine clearance and 24-hour protein calculations undertaken at screening, baseline, Weeks 2 and 4, and monthly thereafter (or upon early termination). Bone marrow aspirations and/or tissue biopsy are also performed at Months 3, 6 and 12 if a patient's paraprotein serum concentration or 24-hour urine protein excretion declined to the next lower level, based on best response criteria. Preliminary results for the 28-day treatment period are summarized below.

Preliminary pharmacokinetic analyses based on these two studies indicated that AUC and C_{max} values increase proportionally with dose following single and multiple doses in multiple myeloma patients (as was seen in healthy volunteers). Further, there was no evidence of accumulation with multiple dosing as single dose AUC_(0-∞) was comparable to multiple dose AUC_(0-τ) following the same dose of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione.

Similar to healthy volunteer studies, double peaks were observed. Exposure in multiple myeloma patients appeared to be slightly higher based on C_{max} and AUC values as compared to healthy male volunteers while clearance in multiple myeloma patients was lower than it was in healthy volunteers, consistent with their poorer renal function (both as a consequence of their age and their disease). Finally, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione half-life in patients was shorter than in healthy volunteers (mean 8 hours, ranging up to 17 hours).

In this study, the first cohort of 3 patients was treated for 28 days at 5 mg/day without any dose limiting toxicity (DLT). The second cohort of 3 patients subsequently commenced

US 8,648,095 B2

37

therapy at 10 mg/day. Patients in the second 10 mg/day of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione cohort tolerated treatment well.

6.5.3 Treatment of Solid Tumors

Study with 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) was conducted in patients with varying types of solid tumors, including malignant melanoma (13), carcinoma of the pancreas (2), carcinoid-unknown primary (1), renal carcinoma (1), breast carcinoma (1) and NSCLC (2). Patients received 5 mg/day 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione for seven days and are subsequently escalated every seven days to 10 mg/day, 25 mg/day, and 50 mg/day for a total of 4 weeks of treatment. Patients who, experienced clinical benefit were permitted to continue on treatment as Named Patients.

The study initially enrolled 20 patients and was subsequently amended to enroll 16 additional patients (adrenal carcinoma, NSCLC, malignant mesothelioma, breast cancer, malignant melanoma (8), renal cell cancer (4)) at a higher dose. The 16 additional patients were given weekly escalating doses of 25 mg/day, 50 mg/day, 75 mg/day, 100 mg/day, 125 mg/day, and 150 mg/day over a 6-week period with continuing treatment for an additional six weeks.

The study of Phase 1 study was designed to determine a maximum tolerated dose (MTD) of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione in patients with refractory solid tumors and/or lymphoma, as well as to characterize the pharmacokinetic and side effect profiles of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione in this patient population. The study design dictates that at least 3 patients must be enrolled at a dose level and have completed 28 days of treatment prior to enrollment of patients at the next higher dose level. Patients in the first cohort began dosing at 5 mg/day of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. Patients will be escalated to 10, 20, 25, and 30 mg/day provided there is no toxicity.

In this study, the MTD is defined as the highest dose level in which fewer than two of six patients treated did not experience Grade 3 or greater non-hematological toxicity or Grade 4 or greater hematological toxicity. If, at any given dose level in either study, one out of three patients experiences toxicity, three additional patients must be treated at that particular dose. If, however, two out of six patients experience DLT, the MTD is judged to have been exceeded. No further dose escalations are to occur and additional patients are to be enrolled at the previous dose level. The dose of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione administered is escalated until the MTD is achieved or the maximum daily dose of is reached.

No DLTs were reported in the initial group of 20 patients enrolled in the study. Thirteen of the original 20 trial patients, along with 2 non-trial patients, continued on treatment as named patients at doses up to 150 mg/day.

6.5.4 Treatment of Gliomas

This study was performed to find toxicity in patients with recurrent, high-grade gliomas. The study is designed such that patients are given increasingly higher doses of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione until a maximum tolerated dose (MTD) is established. The study also seeks to obtain preliminary toxicity information and pharmacokinetic data on 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione, as well as to develop exploratory data concerning surrogate end points of angiogenic activity in vivo using functional neuro-imaging studies, and in vitro assays of serum angiogenic peptides.

Patients enrolled in the first cohort receive 2.5 mg/m²/day for a 4-week cycle. During each 4-week cycle of therapy,

38

3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione administered once daily for 3 weeks followed by a week of rest. Patients who complete a treatment cycle may receive another cycle of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione treatment if two criteria are met. First, the patient must have stable disease or have experienced a partial response or complete response, or the patient is benefiting from the therapy with 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione as evidenced by a decrease in tumor-related symptoms such as neurological deficits. Second, the patient must have recovered from toxicity related to 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione which occurred in the prior cycle by Day 42 or sooner (28-day cycle plus limit of 2 weeks to recover) as evidenced by a return to Grade \leq 1 toxicity level. Patients who experience DLT in the previous cycle should have their dose modified. DLT is defined as a non-hematological event Grade \geq 3 toxicity or hematological event of Grade 4 toxicity thought to be related to the study medication. Patients who experience DLT in the first cycle and have no response to therapy are removed from the study.

3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione doses are subsequently escalated to 5, 8, 11, 15, and 20 mg/m²/day to a maximum total daily dose of 40 mg. Patients continue to receive 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione on a 4-week cycle per dose level until one of the off-study criteria are met.

Three patients are enrolled in each cohort. If at least one DLT occurs, three additional patients are added to the cohort at that particular dose level. If two DLTs occur, the MTD, defined as the dose at which fewer than one-third of patients at each dose level experiences DLT has been exceeded and four more patients are treated at the previous dose.

Patients who experience DLT during the first 4-week cycle are removed from the study, except if they have a response to therapy. For patients who have completed their first 4-week cycle of without DLT, but who subsequently experience Grade 3 or 4 hematological and/or nonhematological toxicity, treatment is suspended for a minimum of a week. If the toxicity resolves to <Grade 2 within three weeks, the patient is treated at two dose levels lower than the dose that caused the toxicity (or a 50% reduction if the patient was treated at the first or second dose level). Patients in whom Grade 3 or 4 toxicity does not resolve to <Grade 1 within three weeks, or those who have another Grade 3 toxicity at the reduced dose are removed from the study.

Pharmacokinetic sampling is performed prior the first dose of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Day 1) and 0.5, 1, 2, 4, 6, 8, 24, and 48 hours thereafter. Sampling is also conducted pre-dose on Days 7 and 21 and 0.5, 1, 2, 4, 6, 8, and 24 post-dose on Day 21 to evaluate steady-state 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione levels.

6.5.5 Treatment of Metastatic Melanoma

Patients with metastatic melanoma were started on 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) at 5 mg/day for seven days. The dose was then increased every seven days to 10 mg/day, 25 mg/day, and 50 mg/day, respectively, for a total of four weeks on therapy. Five of the 13 melanoma patients who were treated under this regimen either showed disease stabilization or a partial response in the first four weeks of treatment. Tumor response was seen in cutaneous and subcutaneous lesions (five patients), lymph nodes (two patients), and liver (one patient). The duration of response was approximately six months. The

US 8,648,095 B2

39

result suggests that the compound appears is a promising new anti-cancer agent and has both antiangiogenic and immunomodulatory properties.


6.5.6 Treatment of Relapsed or Refractory Multiple Myeloma

Patients with relapsed and refractory Dune-Salmon stage III multiple myeloma, who have either failed at least three previous regimens or presented with poor performance status, neutropenia or thrombocytopenia, are treated with up to four cycles of combination of melphalan (50 mg intravenously), an immunomodulatory compound of the invention (about 1 to 150 mg orally daily), and dexamethasone (40 mg/day orally on days 1 to 4) every four to six weeks. Maintenance treatment consisting of daily an immunomodulatory compound of the invention and monthly dexamethasone are continued until the disease progression. The therapy using an immunomodulatory compound of the invention in combination with melphalan and dexamethasone is highly active and generally tolerated in heavily pretreated multiple myeloma patients whose prognosis is otherwise poor.

The embodiments of the invention described above are intended to be merely exemplary, and those skilled in the art will recognize, or will be able to ascertain using no more than routine experimentation, numerous equivalents of specific compounds, materials, and procedures. All such equivalents are considered to be within the scope of the invention and are encompassed by the appended claims.

What is claimed is:

1. A method of treating multiple myeloma, which comprises cyclically administering to a patient having multiple myeloma: (a) about 1 to about 50 mg per day of a compound having the formula:



or a pharmaceutically acceptable salt, solvate or stereoisomer thereof for 21 consecutive days followed by seven consecutive days of rest from administration of said compound in a 28 day cycle, and (b) a therapeutically effective amount of dexamethasone.

2. The method of claim 1, wherein the multiple myeloma is newly diagnosed multiple myeloma, smoldering multiple myeloma, refractory multiple myeloma, relapsed multiple myeloma, or relapsed and refractory multiple myeloma.

3. The method of claim 1, wherein the compound is a pharmaceutically acceptable salt.

4. The method of claim 1, wherein the compound is a pharmaceutically acceptable stereoisomer.

5. The method of claim 4, wherein the stereoisomer is an enantiomerically pure R isomer.

6. The method of claim 4, wherein the stereoisomer is an enantiomerically pure S isomer.

40

7. The method of claim 1, which further comprises administering a therapeutically effective amount of an additional active agent.

8. The method of claim 7, wherein the additional active agent is melphalan, doxorubicin, vincristine, prednisone, cyclophosphamide, biacin, a proteasome inhibitor, or a combination thereof.

9. The method of claim 1, which further comprises autologous stem cell transplantation.

10. The method of claim 1, wherein the multiple myeloma is smoldering multiple myeloma.

11. The method of claim 1, wherein the multiple myeloma is refractory multiple myeloma.

12. The method of claim 1, wherein the multiple myeloma is relapsed multiple myeloma.

13. The method of claim 1, wherein the compound is administered in the form of a capsule or tablet.

14. The method of claim 1, wherein the compound is administered in a capsule of 5 mg, 10 mg, 15 mg or 25 mg.

15. The method of claim 1, wherein the compound is administered in an amount of about 25 mg per day.


16. The method of claim 1, wherein the compound is administered in an amount of 15 mg per day.

17. The method of claim 1, wherein the compound is administered in an amount of 10 mg per day.

18. The method of claim 1, wherein the compound is administered in an amount of 5 mg per day.

19. The method of claim 1, wherein the dexamethasone is administered in an amount of 40 mg per day on days 1-4 every 28 days.

20. The method of claim 1, wherein the compound is



and is not a pharmaceutically acceptable salt, solvate or stereoisomer hereof.

21. The method of claim 14, wherein the capsule comprises the compound, lactose anhydrous, microcrystalline cellulose, croscarmellose sodium and magnesium stearate.

22. The method of claim 1, wherein the dexamethasone is orally administered in an amount of 40 mg once daily on days 1, 8, 15 and 22 of each 28 day cycle.

23. The method of claim 1, wherein the multiple myeloma is newly diagnosed multiple myeloma.

24. The method of claim 1, wherein the compound is administered in a capsule in an amount of 1 mg to 50 mg.

25. The method of claim 24, wherein the capsule comprises the compound, lactose anhydrous, microcrystalline cellulose, croscarmellose sodium and magnesium stearate.

* * * * *

EXHIBIT F



US009101621B2

(12) **United States Patent**
Zeldis

(10) **Patent No.:** **US 9,101,621 B2**
(45) **Date of Patent:** ***Aug. 11, 2015**

(54) **METHODS FOR TREATING MULTIPLE MYELOMA WITH 3-(4-AMINO-1-OXO-1,3-DIHYDRO-ISOINDOL-2-YL)-PIPERIDINE-2,6-DIONE AFTER STEM CELL TRANSPLANTATION**

A61K 31/573 (2013.01); *A61K 31/675* (2013.01); *A61K 31/704* (2013.01); *A61K 31/7048* (2013.01); *A61K 35/12* (2013.01); *A61K 39/0011* (2013.01); *A61K 39/3955* (2013.01); *A61K 45/06* (2013.01)

(71) Applicant: **Celgene Corporation**, Summit, NJ (US)

(58) **Field of Classification Search**

(72) Inventor: **Jerome B. Zeldis**, Princeton, NJ (US)

CPC A61K 31/4035; A61K 31/425; A61K 31/445; A61K 31/454

(73) Assignee: **Celgene Corporation**, Summit, NJ (US)

USPC 514/323

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

See application file for complete search history.

This patent is subject to a terminal disclaimer.

(56) **References Cited**

U.S. PATENT DOCUMENTS

(21) Appl. No.: **14/255,211**

3,536,809 A 10/1970 Applezweig
3,598,123 A 8/1971 Zaffaroni et al.
3,845,770 A 11/1974 Theeuwes et al.
3,916,899 A 11/1975 Theeuwes et al.
4,008,719 A 2/1977 Theeuwes et al.
4,810,643 A 3/1989 Souza
4,999,291 A 3/1991 Souza

(22) Filed: **Apr. 17, 2014**

(65) **Prior Publication Data**

US 2014/0308344 A1 Oct. 16, 2014

(Continued)

Related U.S. Application Data

FOREIGN PATENT DOCUMENTS

(63) Continuation of application No. 14/201,069, filed on Mar. 7, 2014, which is a continuation of application No. 13/782,728, filed on Mar. 1, 2013, now Pat. No. 8,673,939, which is a continuation of application No. 13/488,888, filed on Jun. 5, 2012, now Pat. No. 8,648,095, which is a continuation of application No. 12/640,702, filed on Dec. 17, 2009, now Pat. No. 8,198,306, which is a continuation of application No. 10/438,213, filed on May 15, 2003, now Pat. No. 7,968,569.

JP 2002-513391 5/2002
WO WO 92/14455 9/1992

(Continued)

(60) Provisional application No. 60/380,842, filed on May 17, 2002, provisional application No. 60/424,600, filed on Nov. 6, 2002.

OTHER PUBLICATIONS

(51) **Int. Cl.**

A61K 31/4035 (2006.01)
A61K 31/425 (2006.01)
A61K 31/445 (2006.01)
A61K 31/454 (2006.01)
A61K 31/00 (2006.01)
A61K 31/40 (2006.01)
A61K 31/515 (2006.01)
A61K 45/06 (2006.01)
A61K 31/4439 (2006.01)
A61K 31/573 (2006.01)
A61K 31/198 (2006.01)
A61K 31/475 (2006.01)
A61K 31/675 (2006.01)
A61K 31/704 (2006.01)
A61K 31/7048 (2006.01)
A61K 35/12 (2015.01)
A61K 39/395 (2006.01)
A61K 39/00 (2006.01)

Bjorkstrand et al. (Leukemia&lymphoma (1994), 15, (3-4), 265-72).*

Canal et al. (Pathology Oncology Research, (1998), 4(3):171-178).*

Singhal et al. ((N Engl J Med (1999); 341(21):1565-1571).*

Stirling, D., (Seminars in Oncology, Dec. 2001, 28(6):602-606).*

U.S. Appl. No. 60/499,723, Markian.

U.S. Appl. No. 60/372,348, Hariri et al.

U.S. Appl. No. 10/732,867, D'Amato et al.

U.S. Appl. No. 09/545,654, D'Amato.

U.S. Appl. No. 09/287,377, D'Amato.

Carstensen, 1995, *Drug Stability: Principles & Practice*, 2nd. ed., Marcel Dekker, New York, NY pp. 379-380.

(Continued)

Primary Examiner — Jeffrey S Lundgren

Assistant Examiner — Chris Simmons

(74) *Attorney, Agent, or Firm* — Jones Day

(57) **ABSTRACT**

Methods of treating, preventing and/or managing cancer as well as and diseases and disorders associated with, or characterized by, undesired angiogenesis are disclosed. Specific methods encompass the administration of an immunomodulatory compound alone or in combination with a second active ingredient. The invention further relates to methods of reducing or avoiding adverse side effects associated with chemotherapy, radiation therapy, hormonal therapy, biological therapy or immunotherapy which comprise the administration of an immunomodulatory compound. Pharmaceutical compositions, single unit dosage forms, and kits suitable for use in methods of the invention are also disclosed.

(52) **U.S. Cl.**

CPC *A61K 31/454* (2013.01); *A61K 31/00* (2013.01); *A61K 31/198* (2013.01); *A61K 31/40* (2013.01); *A61K 31/4035* (2013.01); *A61K 31/425* (2013.01); *A61K 31/445* (2013.01); *A61K 31/4439* (2013.01); *A61K 31/475* (2013.01); *A61K 31/515* (2013.01);

US 9,101,621 B2

Page 2

(56)

References Cited

U.S. PATENT DOCUMENTS

| | | | | |
|--------------|------|---------|----------------------|---------|
| 5,059,595 | A | 10/1991 | Le Grazie | |
| 5,073,543 | A | 12/1991 | Marshall et al. | |
| 5,120,548 | A | 6/1992 | McClelland et al. | |
| 5,134,127 | A | 7/1992 | Stella et al. | |
| 5,229,496 | A | 7/1993 | Deeley et al. | |
| 5,354,556 | A | 10/1994 | Sparks et al. | |
| 5,385,901 | A | 1/1995 | Kaplan et al. | |
| 5,387,421 | A * | 2/1995 | Amidon et al. | 424/472 |
| 5,391,485 | A | 2/1995 | Deeley et al. | |
| 5,393,870 | A | 2/1995 | Deeley et al. | |
| 5,528,823 | A | 6/1996 | Rudy, Jr. et al. | |
| 5,580,755 | A | 12/1996 | Souza | |
| 5,591,767 | A | 1/1997 | Mohr et al. | |
| 5,593,990 | A | 1/1997 | D'Amato | |
| 5,629,327 | A | 5/1997 | D'Amato | |
| 5,635,517 | A * | 6/1997 | Muller et al. | 514/323 |
| 5,639,476 | A | 6/1997 | Oshlack et al. | |
| 5,674,533 | A | 10/1997 | Santus et al. | |
| 5,698,579 | A | 12/1997 | Muller | |
| 5,712,291 | A | 1/1998 | D'Amato | |
| 5,731,325 | A | 3/1998 | Andrulic, Jr. et al. | |
| 5,733,566 | A | 3/1998 | Lewis | |
| 5,798,368 | A | 8/1998 | Muller et al. | |
| 5,874,448 | A | 2/1999 | Muller et al. | |
| 5,877,200 | A | 3/1999 | Muller | |
| 5,929,117 | A | 7/1999 | Muller et al. | |
| 5,955,476 | A | 9/1999 | Muller et al. | |
| 6,020,358 | A | 2/2000 | Muller et al. | |
| 6,071,948 | A | 6/2000 | D'Amato | |
| 6,114,355 | A | 9/2000 | D'Amato | |
| 6,140,346 | A | 10/2000 | Andrulic, Jr. et al. | |
| 6,228,879 | B1 | 5/2001 | Green et al. | |
| 6,235,756 | B1 | 5/2001 | D'Amato | |
| 6,281,230 | B1 | 8/2001 | Muller et al. | |
| 6,316,471 | B1 | 11/2001 | Muller et al. | |
| 6,326,388 | B1 | 12/2001 | Man et al. | |
| 6,335,349 | B1 | 1/2002 | Muller et al. | |
| 6,380,239 | B1 | 4/2002 | Muller et al. | |
| 6,395,754 | B1 | 5/2002 | Muller et al. | |
| 6,403,613 | B1 | 6/2002 | Man et al. | |
| 6,420,414 | B1 | 7/2002 | D'Amato | |
| 6,458,810 | B1 | 10/2002 | Muller et al. | |
| 6,469,045 | B1 | 10/2002 | D'Amato | |
| 6,476,052 | B1 | 11/2002 | Muller et al. | |
| 6,518,298 | B2 | 2/2003 | Green et al. | |
| 6,555,554 | B2 | 4/2003 | Muller et al. | |
| 6,673,828 | B1 | 1/2004 | Green et al. | |
| 7,323,479 | B2 | 1/2008 | Zeldis | |
| 7,393,862 | B2 | 7/2008 | Zeldis | |
| 7,435,745 | B2 | 10/2008 | D'Amato | |
| 7,468,363 | B2 | 12/2008 | Zeldis | |
| 7,968,569 | B2 * | 6/2011 | Zeldis | 514/323 |
| 8,188,118 | B2 * | 5/2012 | Zeldis | 514/321 |
| 8,198,262 | B2 | 6/2012 | Zeldis | |
| 8,198,306 | B2 * | 6/2012 | Zeldis | 514/321 |
| 8,207,200 | B2 * | 6/2012 | Zeldis | 514/321 |
| 8,530,498 | B1 * | 9/2013 | Zeldis | 514/320 |
| 8,648,095 | B2 * | 2/2014 | Zeldis | 514/321 |
| 2001/0018445 | A1 | 8/2001 | Huang et al. | |
| 2001/0056114 | A1 | 12/2001 | D'Amato | |
| 2002/0035090 | A1 | 3/2002 | Zeldis et al. | |
| 2002/0045643 | A1 | 4/2002 | Muller et al. | |
| 2002/0052398 | A1 | 5/2002 | D'Amato | |
| 2002/0054899 | A1 | 5/2002 | Zeldis | |
| 2002/0061923 | A1 | 5/2002 | D'Amato | |
| 2002/0128228 | A1 | 9/2002 | Hwu | |
| 2002/0161023 | A1 | 10/2002 | D'Amato | |
| 2002/0173658 | A1 | 11/2002 | Muller et al. | |
| 2002/0183360 | A1 | 12/2002 | Muller et al. | |
| 2003/0013739 | A1 | 1/2003 | Masferrer et al. | |
| 2003/0028028 | A1 | 2/2003 | Man et al. | |
| 2003/0045552 | A1 | 3/2003 | Robarge et al. | |
| 2003/0069428 | A1 | 4/2003 | Muller et al. | |
| 2003/0096841 | A1 | 5/2003 | Robarge et al. | |
| 2003/0139451 | A1 | 7/2003 | Shah et al. | |

| | | | |
|--------------|----|---------|-------------------|
| 2003/0144325 | A1 | 7/2003 | Muller et al. |
| 2003/0181428 | A1 | 9/2003 | Green et al. |
| 2003/0187024 | A1 | 10/2003 | D'Amato |
| 2003/0191098 | A1 | 10/2003 | D'Amato |
| 2003/0235909 | A1 | 12/2003 | Hariri et al. |
| 2004/0029832 | A1 | 2/2004 | Zeldis |
| 2004/0077685 | A1 | 4/2004 | Figg et al. |
| 2004/0077686 | A1 | 4/2004 | Dannenberg et al. |
| 2004/0087546 | A1 | 5/2004 | Zeldis |
| 2004/0091455 | A1 | 5/2004 | Zeldis |
| 2004/0122052 | A1 | 6/2004 | Muller et al. |
| 2004/0266809 | A1 | 12/2004 | Emanuel et al. |
| 2008/0145368 | A1 | 6/2008 | Zeldis |
| 2008/0292583 | A1 | 11/2008 | Zeldis |
| 2009/0010877 | A1 | 1/2009 | Zeldis |
| 2009/0123416 | A1 | 5/2009 | Zeldis |
| 2010/0093683 | A1 | 4/2010 | Zeldis |
| 2010/0196369 | A1 | 8/2010 | Zeldis |
| 2010/0260719 | A1 | 10/2010 | Zeldis |
| 2012/0035145 | A1 | 2/2012 | Zeldis |
| 2012/0135042 | A1 | 5/2012 | Zeldis |
| 2013/0183381 | A1 | 7/2013 | Zeldis |

FOREIGN PATENT DOCUMENTS

| | | |
|----|----------------|---------|
| WO | WO 94/20085 | 9/1994 |
| WO | WO 98/03502 | 1/1998 |
| WO | WO 98/19649 | 5/1998 |
| WO | WO 98/54170 | 12/1998 |
| WO | WO 01/70275 | 9/2001 |
| WO | WO 01/87307 | 11/2001 |
| WO | WO 02/15926 | 2/2002 |
| WO | WO 02/059106 | 8/2002 |
| WO | WO 02/064083 | 8/2002 |
| WO | PCT/US03/11578 | 4/2003 |
| WO | WO 03/086373 | 10/2003 |

OTHER PUBLICATIONS

Corral et al., 1999, "Immunomodulation by thalidomide and thalidomide analogues," *Ann. Rheum. Dis.* 58(Suppl 1):1107-113.

Craig et al., 1967, "Potential anticancer agents. III. 2-phthalimidoaldehydes and derivatives," *Potential Anticancer Agents III* 10:1071-1073.

D'Amato et al., 2001, "Mechanism of action of thalidomide and 3-aminothalidomide in multiple myeloma," *Semin Oncol.* 28:597-601.

D'Amato et al., 1994, "Thalidomide is an Inhibitor of Angiogenesis", *Proc. Natl. Acad. Sci.* 91:4082-4085.

De et al., 1976, "Hansch analysis for some antineoplastic glutarimides," *J. Indian Chem. Soc. I.III:* 825-826.

De et al., 1976, "Possible antineoplastic agents: III. Synthesis of 6-alkyl-2-[4'-methoxyphthalimido] and 6-alkyl-3-[3'-4'-dimethoxyphenyl] glutarimides," *J. Indian Chem. Soc. I.III:*1122-1125.

Dredge et al., 2002, "Novel thalidomide analogues display anti-angiogenic activity independently of immunomodulatory effects," *Br. J. Cancer* 87(10):1166-1172.

Folkman et al., 1983, "Angiogenesis inhibition and tumor regression caused by heparin or a heparin fragment in the presence of cortisone," *Science* 221(4612):719-725.

Gershbein, 1991, "The thalidomide analog, EM 12, enhances 1,2-dimethylhydrazine-induction of rat colon adenocarcinomas," *Cancer Letters* 60: 129-133.

Grabstald et al., 1965, "Clinical experiences with thalidomide in patients with cancer," *Clinical Pharmacology and Therapeutics* 6:298-302.

Lentzsch et al., 2003, "Immunomodulatory analogs of thalidomide inhibit growth of Hs Sultan cells and angiogenesis in vivo," *Leukemia* 17(1):41-44.

Lentzsch et al., 2002, "S-3-amino-phthalimido-glutarimide inhibits angiogenesis and growth of B-cell neoplasias in mice", *Cancer Research* 62:2300-2305.

Miyachi et al., 1997, "Novel biological response modifiers: phthalimides with tumor necrosis factor-alpha production-regulating activity," *J. Med. Chem.* 40:2858-2865.

(56)

References Cited

OTHER PUBLICATIONS

- Muller et al., 1999, "Amino-substituted thalidomide analogs: potent inhibitors of TNF-alpha production," *Bioorg. Med. Chem. Lett.* 9(11):1625-1630.
- Muller et al., 1998, "Thalidomide analogs and PDE4 inhibition," *Bioorg. Med. Chem. Lett.* 8(19):2669-2674.
- Muller et al., 1996, "Structural modifications of thalidomide produce analogs with enhanced tumor necrosis factor inhibitory activity," *J. Med. Chem.* 39(17):3238-3240.
- Olson et al., 1965, "Thalidomide (N-phthaloylglutamimide) in the treatment of advanced cancer," *Clinical Pharmacology and Therapeutics* 6(3):292-297.
- Penichet et al., 2001, "Antibody-cytokine fusion proteins for the therapy of cancer," *J. Immunol. Methods* 248(1-2):91-101.
- Physician's Desk Reference*, 2002, 56th ed., pp. 1755-1760.
- Raza et al., 2001, "Thalidomide produces transfusion independence in long-standing refractory anemias of patients with myelodysplastic syndromes," *Blood* 98(4):958-965.
- Shah et al., 1999, "Synthesis and enantiomeric separation of 2-phthalimidino-glutaric acid analogues: potent inhibitors of tumor metastasis," *J. Med. Chem.* 42:3014-3017.
- Shibata et al., 1995, "N-alkylphthalimides: structural requirement of thalidomidial action on 12-O-tetradecanoylphorbol-13-acetate-induced tumor necrosis factor alpha production by human leukemia HL-60 cells," *Chem. Pharm. Bull.* 43(1):177-179.
- Shimazawa et al., 1999, "Antiangiogenic activity of tumor necrosis factor-alpha production regulators derived from thalidomide," *Biol. Pharm. Bull.* 22(2):224-226.
- Rubin et al., "Principles of Cancer Treatment-1", 12 ONCOIV 1, May 2003.
- Wilen et al., 1977, *Tetrahedron* 33:2725.
- Wilen, 1972, *Tables of Resolving Agents and Optical Resolutions*, E.L. Eliel, ed., Univ. of Notre Dame Press, Notre Dame, IN pp. 268.
- Wolff ed., 1995, *Burger's Medicinal Chemistry and Drug Discovery*, 5th ed., pp. 172-178, 949-982.
- N. Ake Jonnson, 1972, "Chemical Structure and Teratogenic Properties," *Acta Pharm.*, pp. 521-542.
- Alexanian et al., 2004, "VTD (Velcade, thalidomide, dexamethasone) as primary therapy for newly-diagnosed multiple myeloma," *Am. Soc. Hematol.* 46th Ann. Meeting Dec. 4-7, 2004, San Diego, CA Abstract #210.
- Anderson, 2000, "Thalidomide: Therapeutic potential in hematologic malignancies," *Seminars in Hematology* 37(1 Supp 3): 1-4.
- Attal et al., 2004, "Maintenance treatment with thalidomide after autologous transplantation for myeloma: First analysis of a prospective randomized study of the Intergroupe Francophone du Myelome (IFM 99 02)," *Am. Soc. Hematol.* 46th Ann. Meeting Dec. 4-7, 2004, San Diego, CA Abstract #535.
- Bernardeschi et al., 2003, *J. Exp. Clin. Cancer Res.* 22(4):129-133.
- Corral et al., 1999, "Differential cytokine modulation and T cell activation by two distinct classes of thalidomide analogues that are potent inhibitors of TNF-alpha," *J. Immunol.* 163(1):380-386.
- Davies et al., 2001, "Thalidomide and immunomodulatory derivatives augment natural killer cell cytotoxicity in multiple myeloma," *Blood* 98(1):210-216.
- Dimopoulos et al., 2004, "Primary treatment with pulsed melphalan, dexamethasone, thalidomide (MDT) for symptomatic patients with multiple myeloma ≥ 75 years of age," *Am. Soc. Hematol.* 46th Ann. Meeting Dec. 4-7, 2004, San Diego, CA Abstract #1482.
- Eisen et al., 2000, "Continuous low dose Thalidomide: a phase II study in advanced melanoma, renal cell, ovarian and breast cancer," *Br. J. Cancer* 82(4):812-817.
- Fakhouri et al., 2004, "Thalidomide in patients with multiple myeloma and renal failure," *Br. J. Haematol.* 125:90-102.
- Fenk et al., 2005, "Single-agent thalidomide for treatment of first relapse following high-dose chemotherapy in patients with multiple myeloma," *Leukemia* 19(1):156-159.
- Gupta et al., 2001, "Adherence of multiple myeloma cells to bone marrow stromal cells upregulates vascular endothelial growth factor secretion: therapeutic applications," *Leukemia* 15(12):1950-1961.
- Haslett et al., 2003, "Thalidomide and a thalidomide analogue drug costimulate virus-specific CD8+ T cells in vitro," *J. Infect. Dis.* 187(6):946-955.
- Hideshima et al., 2000, "Thalidomide and its analogs overcome drug resistance of human multiple myeloma cells to conventional therapy," *Blood* 96(9):2943-2950.
- Offidani et al., 2003, Thalidomide plus oral melphalan for advanced multiple myeloma: a phase II study. *Haematologica*. Dec. 2003;88(12):1432-1433.
- Palumbo et al., 2004, "A prospective randomized trial of oral melphalan prednisone, thalidomide (MPT) vs. oral melphalan, prednisone (MP): An interim analysis," *Am. Soc. Hematol.* 46th Ann. Meeting Dec. 4-7, 2004, San Diego, CA Abstract #207.
- Raje et al., 1999, "Thalidomide—a revival story," *N. Engl. J. Med.* 341(21):1606-1609.
- Rajkumar et al., 2004, "Thalidomide plus dexamethasone versus dexamethasone alone in newly diagnosed multiple myeloma (E1A00): Results of a phase III trial coordinated by the Eastern Cooperative Oncology Group," *Am. Soc. Hematol.* 46th Ann. Meeting Dec. 4-7, 2004, San Diego, CA Abstract #205.
- Rajkumar et al., 2000, "Prognostic value of bone marrow angiogenesis in multiple myeloma," *Clin. Cancer Res.* 6(8):3111-3116.
- Ribatti et al., 1999, "Bone marrow angiogenesis and mast cell density increase simultaneously with progression of human multiple myeloma," *Br. J. Cancer* 79(3-4):451-455.
- Singhal et al., 1999, Antitumor activity of thalidomide in refractory multiple myeloma, *N. Engl. J. Med.* 341(21):1565-1571.
- Steins et al., 2002, "Efficacy and safety of thalidomide in patients with acute myeloid leukemia," *Blood* 99(3):834-839.
- Vacca et al., 1999, "Bone marrow neovascularization, plasma cell angiogenic potential, and matrix metalloproteinase-2 secretion parallel progression of human multiple myeloma," *Blood* 93(9):3064-3073.
- Wohrer et al., 2004, "Effective treatment of primary plasma cell leukemia with thalidomide and dexamethasone—a case report," *Hematol. J.* 5(4):361-363.
- Bach, 1963, "Thalidomide in Cancer Chemotherapy," *The Lancet*, No. 1271, p. 71.
- Bach, 1963, "Studies on the Possible Anti-Neoplastic Effect of Thalidomide," *Acta Pathologica Et Microbiologica Scandinavica* 59:491-499.
- Chaundhry, 1966, *Cancer Research*, "Effect of Prednisolone and Thalidomide on Induced Submandibular Gland Tumors in Hamster," 26(part 1)1884-86.
- DiPaolo, 1963, "Effect of Thalidomide on a Variety of Transplantable Tumors," *Cancer Chemotherapy Reports* No. 29, p. 99-102.
- DiPaolo, 1963, "In vitro Test Systems for Cancer Chemotherapy. II. Correlation of in vitro Inhibition of Dehydrogenase and Growth with in vivo Inhibition of Ehrlich Ascites Tumor," *Proceedings of the Society for Experimental Biology & Medicine*, 114:384-387.
- DiPaolo, 1964, "Thalidomide: Effects on Ehrlich Ascites Tumor Cells in vitro" *Science* 144:1583.
- Mauad, 1963, "Clinical Improvements Obtained in Advanced Cancer Patients with Treatment with Thalidomide Associated with Hormones," *Anais Paulistas de Medicina e Cirurgia* 86:13-40.
- Roe and Mitchley, 1963, "Thalidomide and Neoplasia" *Nature* 200:1016-1017.
- Liu et al., "Phase I study of CC-5013 (Revimid), a thalidomide derivative, in patients with refractory metastatic cancer," *American Society of Clinical Oncology*, Abstract #927, 2003.
- Zangari et al., "Results of phase I study of CC-5013 for the treatment of multiple myeloma (MM) patients who relapse after high dose chemotherapy (HDCT)," *American Society of Hematology*, Abstract #3226, 2001.
- Zeldis et al., "Update on the evolution of the IMiD™," *International Society for Biological Therapy of Cancer*, Oral Abstract, 2003.
- Anderson, "Moving disease biology from the laboratory to the clinic," *Seminars in Oncology*, 2002 29:17-20.

US 9,101,621 B2

Page 4

(56)

References Cited

OTHER PUBLICATIONS

- Barlogie et al., "Total Therapy II (TTII) for newly diagnosed multiple myeloma (MM): preliminary data on feasibility and efficacy in the first 231 enrolled patients; comparison with predecessor trial total therapy I (TTI) (N=231)," *Blood*, Abstract # 2857, Dec. 7-11, 2001, *American Society of Hematology*.
- Barlogie et al., "High-dose therapy immunomodulatory drugs in multiple myeloma," *Seminars in Oncology*, 2002, 29 (6):26-33.
- Barlogie et al., "Introduction: Thalidomide and the IMiDs in multiple myeloma," *Seminars in Hematology*, 2003, 40 (4):1-2.
- Barlogie, "Thalidomide and CC-5013 in Multiple Myeloma: The University of Arkansas experience," *Seminars in Hematology*, 2003, 40 (4):33-38.
- Bartlett et al., "The evolution of thalidomide and its IMiD derivatives as anticancer agents," *Nature Reviews Cancer*, 2004, 4 (4):1-9.
- Bartlett et al., "Phase I study to determine the safety, tolerability and immunostimulatory activity of thalidomide analogue CC-5013 in patients with metastatic malignant melanoma and other advanced cancers," *British Journal of Cancer*, 2004, 90:955-961.
- Battagay, "Angiogenesis: mechanistic insights, neovascular diseases, and therapeutic prospects," *J. Mol. Med.*, 1995, 73:333-346.
- Baz et al., "Doxil (D), vincristine (V), reduced frequency dexamethasone (d) and revlimid (R) (DVD-R) results in a high response rate in patients with refractory multiple myeloma (RMM)," *Blood*, Abstract # 2559, *American Society of Hematology*, Dec. 10-13, 2005.
- Brennen et al., "Thalidomide and analogues: current proposed mechanisms and therapeutic usage," *Clinical Prostate Cancer*, 2004, 3 (1):54-61.
- Celgene Corporation, "Celgene advances immunomodulatory drug (IMiD™) clinical program," Press Release, Feb. 2000.
- Celgene Corporation, "Initial Phase I solid tumor data on Celgene's lead IMiD™, Revimid™," Press Release, Jun. 2001.
- Celgene Corporation, "Celgene Corporation receives orphan drug designation for Revimid™ for multiple myeloma," Press Release, Oct. 2001.
- Celgene Corporation, "Celgene Corporation announces third quarter results. Thalomid® (thalidomide) sales increase 24%. Prescriptions up 50%. Enhanced S.T.E.P.S.® launched. Pilot d-MPH data presented," Press Release, Oct. 2001.
- Celgene Corporation, "Celgene expands clinical development program for Revimid™. Five additional trials of Revimid initiated in hematological and solid tumor cancers," Press Release, Jun. 2002.
- Celgene Corporation, "Celgene Corporation announces third quarter results. Thalomid® (thalidomide) revenue increases 41% to \$30.5 million. Pivotal programs for Thalomid and Revimid™ finalized. Peer-reviewed publications of Thalomid and Revimid data. First JNK inhibitor advanced to Phase I clinical trial," Press Release, Oct. 2002.
- Celgene Corporation, "Blood reports Revimid™ has anti-tumor activity in patients with relapsed and refractory multiple myeloma," Press Release, Nov. 1, 2002.
- Celgene Corporation, "Celgene provides update on clinical pipeline. Celgene Announces first target indication for Actimid™, CC-8490. SelCID™ program to advance based on results from Phase I/II trial of CC-1088. First JNK inhibitor successfully completes phase I trial," Press Release, Jan. 2003.
- Celgene Corporation, "Celgene Corporation announces fourth quarter and full year results for 2002," Press Release, Jan. 2003.
- Celgene Corporation, "Celgene receives fast track status from FDA for Revimid™ in multiple myeloma," Press Release, Feb. 2003.
- Celgene Corporation, "Celgene receives fast track status from FDA for Revimid™ in myelodysplastic syndromes," Press Release, Apr. 2003.
- Celgene Corporation, "New Revimid™ clinical data shows potential as novel approach to treating myelodysplastic syndromes (MDS)," Press Release, May 2003.
- Celgene Corporation, "Celgene corporation reports strong operating performance in second quarter as total sales increase 100 percent and profits rise," Press Release, Jul. 2003.
- Celgene Corporation, "Celgene corporation reports record operating performance in third quarter as total revenue increases 117% and profits rise," Press Release, Oct. 2003.
- Celgene Corporation, "Celgene corporation advances Actimid™ (CC-4047) into phase II trial for prostate cancer," Press Release, Oct. 2003.
- Celgene Corporation, "Additional clinical data presented on Revimid™ in myelodysplastic syndromes at the American Society of Hematology 45th annual meeting," Press Release, Dec. 2003.
- Celgene Corporation, "Celgene corporation reviews 2003 achievements and announces 2004 financial outlook," Press Release, Jan. 2004.
- Celgene Corporation, "Revlimid™ receives orphan drug designation from the European commission for multiple myeloma," Press Release, Feb. 2004.
- Celgene Corporation, "Revlimid™ receives orphan drug designation from the European commission for myelodysplastic syndromes," Press Release, Mar. 2004.
- Celgene Corporation, "Celgene corporation reports record operating performance in first quarter with strong revenue growth and profits," Press Release, Apr. 2004.
- Celgene Corporation, "Celgene announces plans to stop phase III trials in melanoma due to lack of efficacy," Press Release, Apr. 2004.
- Dalglish, et al., "New thalidomide analogues; anti-cancer, anti-angiogenic and immunostimulatory," *British Journal of Cancer*, 2001, 85 (1):25.
- Dalglish et al., "Thalidomide analogues CC-5013 and CC-4047 induce T cell activation and IL-12 production in patients with both solid tumours and relapsed and refractory multiple myeloma," *British Journal of Cancer*, 2003, 88(Suppl 1), S25-S54.
- Davies et al., "Thalidomide (Thal) and immunomodulatory derivatives (IMiDs) augment natural killer (NK) cell cytotoxicity in multiple myeloma(MM)," Abstract # 3617, *American Society of Hematology*, Dec. 1-5, 2000.
- Davies et al., "Thalidomide (Thal) and immunomodulatory derivatives (IMiDs) augment natural killer (NK) cell cytotoxicity in multiple myeloma—MM," Abstract # P222, *VIIIth International Myeloma Workshop*, May 4-8, 2001.
- Dibbs et al., "Thalidomide and thalidomide analogs suppress TNF α secretion by myocytes," Abstract # 1284, *Circulation*, 1998.
- Dimopoulos et al., "Results of thalidomide and IMiDs in multiple myeloma," Abstract # P12.1.4, *International Multiple Myeloma Workshop*, May 23-27, 2003.
- Dimopoulos et al., "Treatment of plasma cell dyscrasias with thalidomide and its derivatives," *Journal of Clinical Oncology*, Dec. 1, 2003, 21 (23):4444-4454.
- Dimopoulos et al., "Study of lenalidomide plus dexamethasone versus dexamethasone alone in relapsed or refractory multiple myeloma (MM): Results of a phase 3 Study (MM-010)," Abstract # 6, *American Society of Hematology*, Dec. 10-13, 2005.
- Dredge et al., "A costimulatory thalidomide analog enhances the partial anti-tumor immunity of an autologous vaccination in a model of colorectal cancer," Abstract # 491, *American Association for Cancer Research*, Apr. 6-10, 2002.
- Dredge et al., "Adjuvants and the promotion of Th1-type cytokines in tumour immunotherapy," *Cancer Immunol. Immunother.*, 2002, 51:521-531.
- Dredge et al., "Immunological effects of thalidomide and its chemical and functional analogs," *Critical Reviews in Immunology*, 2002, 22 (5&6):425-437.
- Dredge et al., "Protective antitumor immunity induced by a costimulatory thalidomide analog in conjunction with whole tumor cell vaccination is mediated by increased Th1-type immunity¹," *The Journal of Immunology*, 2002, 168:4914-4919.
- Dredge et al., "Recent developments in antiangiogenic therapy," *Expert Opin. Biol. Ther.*, 2002, 2 (8):953-966.
- Dredge et al., "Angiogenesis inhibitors in cancer therapy," *Current Opinion in Investigational Drugs*, 2003, 4 (6):667-674.
- Dredge et al., "Thalidomide analogs as emerging anti-cancer drugs," *Anti-Cancer Drugs*, 2003, 14:331-335.
- Fickentscher et al., "Stereochemical properties and teratogenic activity of some tetrahydrothalimides," *Molecular Pharmacology*, 1976, 13:133-141.

(56)

References Cited

OTHER PUBLICATIONS

- Figg et al., "Inhibition of angiogenesis: treatment options for patients with metastatic prostate cancer," *Investigational New Drugs*, 2002, 20(2):183-194.
- Galustian et al., "Thalidomide-derived immunomodulatory drugs as therapeutic agents," *Expert Opin. Biol. Ther.*, 2004, 4 (12):1-8.
- Glaspay et al., "The potential role of thalidomide and thalidomide analogs in melanoma," *Clinical Advances in Hematology & Oncology*, 2004, 1-7.
- Gupta et al., "Adherence of multiple myeloma cells to bone marrow stromal cells upregulates vascular endothelial growth factor secretion: therapeutic applications," *Leukemia*, 2001, 15:1950-1961.
- Hayashi et al., "Mechanisms whereby immunomodulatory analogs of thalidomide augment autologous NK cell anti-myeloma immunity," *Blood*, Abstract #3219, Dec. 6-10, 2002, *American Society of Hematology*.
- He, W., et al., 1993, Abstract of papers, 206th American Chemical Society, Chicago, IL; Med. Chem., paper 216.
- Helm et al., "Comparative teratological investigation of compounds of structurally and pharmacologically related to thalidomide," *Arzneimittel Forschung/Drug Research*, 1981, 31 (1)941-949.
- Hernandez-Illizaliturri et al., "Addition of immunomodulatory drugs CC5013 or CC4047 to rituximab enhances anti-tumor activity in a severe combined immunodeficiency (SCID) mouse lymphoma model," Abstract # 235, *American Society of Hematology*, Dec. 6-9, 2003.
- Hideshima et al., "Thalidomide and its analogs overcome drug resistance of human multiple myeloma cells to conventional therapy," *Blood*, 2000, 96:2943-2950, *American Society of Hematology*.
- Hideshima et al., "Thalidomide (Thal) and its analogs overcome drug resistance of human multiple myeloma (MM) cells to conventional therapy," Abstract 1313, *American Society of Hematology*, Dec. 1-5, 2000.
- Hunt et al., "Markers of endothelial and haemostatic activation in the use of CC-4047, a structural analogue of thalidomide, in relapsed myeloma," *Blood*, Abstract # 3216, Dec. 6-10, 2002, *American Society of Hematology*.
- Hussein et al., "Doxil (D), vincristine (V), reduced frequency dexamethasone (d) and Revlimid (DVD-R) a phase I/II trial in advanced relapsed/refractory multiple myeloma (Rmm) patients," *Blood*, Abstract #208, *American Society of Hematology*, Dec. 4-7, 2004.
- Hwu et al., "Thalidomide and its analogues in the treatment of metastatic melanoma," *Chemotherapy Foundation Symposium*, Abstract #44, 2002.
- Kyle, "Current therapy of multiple myeloma," *Internal Medicine*, 2002, 41 (3)175-180.
- Kyle et al., "Multiple myeloma," *New England Journal of Medicine*, 2004, 351:1860-1873.
- Leblanc et al., "Immunomodulatory drug costimulates T cells via the B7-CD28 pathway," *Blood*, 2004, 103:1787-1790, *American Society of Hematology*.
- Lentzsch et al., "In vivo activity of thalidomide and immunomodulatory drugs against multiple myeloma," *VIIIth International Myeloma Workshop*, Abstract #P225, May 4-8, 2001.
- Lentzsch et al., "Immunomodulatory derivative of thalidomide (IMiD CC-4047) determine the lineage commitment of hematopoietic progenitors by down regulation of GATA-1 and modulation of cytokine secretion," Abstract # 3073, *American Society of Hematology*, Dec. 6-9, 2003.
- Lentzsch et al., "Immunomodulatory derivative of thalidomide (IMiD CC-4047) down regulates CAAT/enhancer-binding protein β (C/EBP β) in multiple myeloma (MM)," Abstract # 3456, *American Society of Hematology*, Dec. 6-9, 2003.
- Luzzio et al., "Thalidomide analogues: derivatives of an orphan drug with diverse biological activity," *Expert Opin. Ther. Patents*, 2004, 14 (2):215-229.
- Man et al., " α -Fluoro-substituted thalidomide analogues," *Bioorganic & Medicinal Chemistry Letters* 13, 2003, 3415-3417.
- Marriott et al., "Immunotherapeutic and antitumor potential of thalidomide analogues," *Expert Opin. Biol. Ther.*, 2001, 1 (4):1-8.
- Marriott et al., "New thalidomide analogues; anti-cancer, anti-angiogenic and immunostimulatory," *British Journal of Cancer*, 85:25, Jul. 6, 2001.
- Marriott et al., "Thalidomide and its analogues have distinct and opposing effects on TNF- α and TNFR2 during co-stimulation of both CD4⁺ and CD8⁺ T cells," *Clin. Exp. Immunol.*, 2002, 130:75-84.
- Marriott et al., "A novel subclass of thalidomide analogue with anti-solid tumor activity in which caspase-dependent apoptosis is associated with altered expression of bcl-2 family proteins¹," *Cancer Research*, 2003, 63:593-599.
- Marriott et al., "Thalidomide derived immunomodulatory drugs (IMiDs) as potential therapeutic agents," *Current Drug Targets—Immune, Endocrine & Metabolic Disorders*, 2003, 3:181-186.
- Masellis et al., "Changes in gene expression in bone marrow mesenchymal progenitor cells as a consequence of IMiD therapy in multiple myeloma patients," *Blood*, Abstract # 1548, Dec. 7-11, 2001, *American Society of Hematology*.
- McCarty, "Thalidomide may impede cell migration in primates by down-regulating integrin β -chains: potential therapeutic utility in solid malignancies, proliferative retinopathy, inflammatory disorders, neointimal hyperplasia, and osteoporosis," *Medical Hypotheses*, 1997, 49:123-131.
- Mitsiades et al., "Apoptotic signaling induced by immunomodulatory thalidomide analogs (Imids) in human multiple myeloma cells: therapeutic implications," Abstract # 3224, Dec. 7-11, 2001, *American Society of Hematology*.
- Mitsiades et al., "Apoptotic signaling induced by immunomodulatory thalidomide analogs in human multiple myeloma cells: therapeutic implications," *Blood*, 2002, 99:4525-4530, *American Society of Hematology*.
- Mitsiades et al., "CC-5013 Celgene," *Current Opinion in Investigational Drugs*, 2004, 5 (6):635-647.
- Moutouh et al., "Novel immunomodulatory drugs (IMiDs®): A potential, new therapy for β -hemoglobinopathies," Abstract # 3740, *American Society of Hematology*, Dec. 4-7, 2004.
- Patten et al., "The early use of the serum free light chain assay in patients with relapsed refractory myeloma receiving treatment with a thalidomide analogue (CC-4047)," Abstract # 1640, *American Society of Hematology*, Dec. 6-9, 2003.
- Payvandi et al., "Effects of a thalidomide analog on binding activity of transcription factors and cell cycle progression of multiple myeloma cell lines," *Blood*, Abstract #2487, Dec. 1-5, 2000, *American Society of Hematology*.
- Payvandi et al., "The thalidomide analogs IMiDs enhance expression of CD69 stimulatory receptor on natural killer cells," Abstract # 1793, *American Association for Cancer Research*, Mar. 24-28, 2001.
- Payvandi et al., "Thalidomide analogs IMiDs inhibit expression of cyclooxygenase-2 in multiple myeloma cell line and LPS stimulated PBMCs," *Blood*, Abstract # 2689, Dec. 7-11, 2001, *American Society of Hematology*.
- Payvandi et al., "Thalidomide and IMiDs inhibit microvessel formation from human arterial rings in the absence of human liver microsomes," *Blood*, Abstract # 5046, Dec. 6-10, 2002, *American Society of Hematology*.
- Payvandi et al., "CC-5013 inhibits the expression of adhesion molecules ICAM-1 and CD44 and prevents metastasis of B16 F10 mouse melanoma cells in an animal model," *American Society of Clinical Oncology*, Abstract # 992, 2003.
- Payvandi et al., "Immunomodulatory drugs inhibit expression of cyclooxygenase-2 from TNF- α , IL-1 β , and LPS-stimulated human PBMC in a partially IL-10-dependent manner," *Cellular Immunology*, 2004, 81-88.
- Raje et al., "Combination of the mTOR inhibitor rapamycin and CC-5013 has synergistic activity in multiple myeloma," *Blood*, Dec. 15, 2004, 104 (13)4188-4193.
- Rajkumar et al., "Combination therapy with lenalidomide plus dexamethasone (Rev/Dex) for newly diagnosed myeloma," *Blood*, Dec. 15, 2005, 106 (13)4050-4053.

US 9,101,621 B2

Page 6

(56) References Cited

OTHER PUBLICATIONS

- Richardson et al., "A Phase I study of oral CC5013, an immunomodulatory thalidomide (Thal) derivative, in patients with relapsed and refractory multiple myeloma (MM)," *Blood*, Abstract #3225, Dec. 7-11, 2001, *American Society of Hematology*.
- Richardson et al., "Immunomodulatory drug CC-5013 overcomes drug resistance and is well tolerated in patients with relapsed multiple myeloma," *Blood*, 2002 100:3063-3067, *American Society of Hematology*.
- Richardson et al., "A multi-center, randomized, phase 2 study to evaluate the efficacy and safety of 2 CDC-5013 dose regimens when used alone or in combination with dexamethasone (Dex) for the treatment of relapsed or refractory multiple myeloma (MM)," *Blood*, Abstract # 825, *American Society of Hematology*, Dec. 6-9, 2003.
- Richardson et al., "Immunomodulatory analogs of thalidomide: an emerging new therapy in myeloma," *Journal of Clinical Oncology*, 2004, 22(16) 3212-3214.
- Richardson et al., "A multicenter, single-arm, open-label study to evaluate the efficacy and safety of single-agent lenalidomide in patients with relapsed and refractory multiple myeloma; preliminary results," *10th International Myeloma Workshop*, Apr. 10-14, 2005.
- Richardson et al., "Novel biological therapies for the treatment of multiple myeloma," *Best Practice & Research Clinical Haematology*, 2005, 18 (4):619-634.
- Richardson et al., "A phase I trial of lenalidomide (Revlimid®) with bortezomib (Velcade®) in relapsed and refractory multiple myeloma," *Blood*, Abstract # 365, *American Society of Hematology*, Dec. 10-13, 2005.
- Rubin et al., "Principles of cancer treatment-1," 2003, 12 ONCO IV 1.
- Schafer et al., "Enhancement of cytokine production and AP-1 transcriptional activity in T cells by thalidomide-related immunomodulatory drugs," *Journal of Pharmacology and Experimental Therapeutics*, 2003, 305(3)1222-1232.
- Schey et al., "A phase I study of an immunomodulatory thalidomide analog, CC-4047, in relapsed or refractory multiple myeloma," *Journal of Clinical Oncology*, 2004, 22 (16):1-8.
- Schey et al., "A phase I study of an immunomodulatory thalidomide analogue (CC4047) in relapse/refractory multiple myeloma," *International Society for Experimental Hematology*, Abstract #248, 2002.
- Shaughnessy et al., "Global gene expression analysis shows loss of C-MYC and IL-6 receptor gene mRNA after exposure of myeloma to thalidomide and IMiD," Abstract # 2485, *The American Society of Hematology*, Dec. 1-5, 2000.
- Shire et al., "TNF- α inhibitors and rheumatoid arthritis," *Exp. Opin. Ther. Patents*, 1998, 8 (5):531-544.
- Sorbera et al., "CC-5013. Treatment of multiple myeloma. Treatment of Melanoma. Treatment of myelodysplastic syndrome. Angiogenesis inhibitor. TNF- α production inhibitor," *Drugs of the Future*, 2003, 28(5):425-431.
- Streetly et al., "Thalidomide analogue CC-4047 is effective in the treatment of patients with relapsed and refractory multiple myeloma (MM) and induces T-cell activation and IL-12 production," Abstract # 367, *International Multiple Myeloma Workshop*, May 23-27, 2003.
- Streetly et al., "Changes in neutrophil phenotype following the administration of CC-4047 (Actimid) to patients with multiple myeloma," Abstract # 2543, *American Society of Hematology*, Dec. 6-9, 2003.
- Streetly et al., "An update of the use and outcomes of the new immunomodulatory agent CC-4047 (Actimid) in patients with relapsed/refractory myeloma," Abstract #829, *American Society of Hematology*, Dec. 6-9, 2003.
- Teo et al., "A phase I, single-blind, placebo-controlled, ascending single oral dose, safety, tolerability and pharmacokinetic study of CDC-501, a novel immunomodulatory-oncologic agent, in healthy male subjects with a comparison of fed and fasted," *Clinical Pharmacology and Therapeutics*, 2002, 71 (2)93.
- Teo et al., "Chiral inversion of the second generation IMiD™ CC-4047 (Actimid™) in human plasma and phosphate-buffered saline," *Chirality*, 2003, 15:348-351.
- Thertulien et al., "Hybrid MEL/DT PACE autotransplant regimen for Multiple Myeloma (MM)—safety and efficacy data in pilot study of 15 patients," *Blood*, Abstract # 2869, *American Society of Hematology*, Dec. 7-11, 2001.
- Tohny et al., "A phase I study of oral CC-5013 (lenalidomide, Revlimid™), a thalidomide derivative, in patients with refractory metastatic cancer," *Clinical Prostate Cancer*, 2004, 2:241-243.
- Tricot et al., "Angiochemotherapy (ACT) for multiple myeloma (MM) with DT-PACE results in a high response rate, but in contrast to tandem transplants with melphalan does not affect durable disease control," *Blood*, Abstract # 3531, *American Society of Hematology*, Dec. 7-11, 2001.
- Tsenova et al., "Use of IMiD3, a thalidomide analog, as an adjunct to therapy for experimental tuberculous meningitis," *Antimicrobial Agents and Chemotherapy*, 2002, 46 (6)1887-1895.
- Weber, "Lenalidomide (CC-5013, Revlimid™) and other ImiDs," Abstract # PL5.02, *International Multiple Myeloma Workshop*, Apr. 10-14, 2005.
- Weber et al., "A multicenter, randomized, parallel-group, double-blind, placebo-controlled study of lenalidomide plus dexamethasone versus dexamethasone alone in previously treated subjects with multiple myeloma," Abstract # PO.738, *International Multiple Myeloma Workshop*, Apr. 10-14, 2005.
- Ye et al., "Novel IMiD drugs enhance expansion and regulate differentiation of human cord blood CD34+ cells with cytokines," *Blood*, Abstract #4099, *American Society of Hematology*, Dec. 6-10, 2002.
- Zangari et al., "Risk factors for deep vein thrombosis (DVT) in a large group of myeloma patients (Pts) treated with thalidomide (Thal): The Arkansas Experience," *Blood*, Abstract # 681, *American Society of Hematology*, Dec. 7-11, 2001.
- Zangari et al., "Revimid 25 mg (REV 25) x 20 versus 50 mg (REV 50) x 10 q 28 days with bridging of 5 mg x 10 versus 10 mg x 5 as post-transplant salvage therapy for multiple myeloma (MM)," *Blood*, Abstract # 1642, *American Society of Hematology*, Dec. 6-9, 2003.
- Zeldis et al., "Potential new therapeutics for Waldenström's macroglobulinemia," *Seminars in Oncology*, 2003, 30 (2):275-281.
- Zhang et al., "CC-5079, a novel microtubule and TNF- α inhibitor with anti-angiogenic and antimetastasis activity," Abstract # B012, *International Conference on Molecular Targets and Cancer Therapeutics*, Nov. 17-21, 2003.
- Anderson, "The Role of Immunomodulatory Drugs in Multiple Myeloma," *Seminars in Hematology*, vol. 40, No. 4, Suppl 4, 2003: pp. 23-32.
- Weber, "Thalidomide and Its Derivatives: New Promise for Multiple Myeloma," *Cancer Control*, vol. 10, No. 5, 375-383, 2003.
- Patt, Yehuda A.; Hassan, Manal M.; Lozano, Richard D.; Ellis, Lee M.; Peterson, J. Andrew; Waugh, Kimberly A.; *Durable Clinical Response of Refractory Hepatocellular Carcinoma to Orally Administered Thalidomide*. *American Journal of Clinical Oncology*, 2000.
- Richardson, Paul; Hideshima, Teru; Anderson, Kenneth; *Thalidomide: The Revival of a Drug with Therapeutic Promise in the Treatment of Cancer*; Principles & Practice of Oncology, vol. 15, No. 2, 2001.
- Thomas, Melodic; Doss, Deborah, *Thalidomide Nursing Roundtable Update*, Monograph, Sep. 2002.
- Richardson, Paul; Hideshima, Teru; Anderson, Kenneth; *Thalidomide: Emerging Role in Cancer Medicine*; Annual Review of Medicine, 2002.
- Berenson, J.R.; Bergsagel, P. L.; Munshi, N.; *Initiation and Maintenance of Multiple Myeloma*; Seminars in Hematology, vol. 36, No. 1, Supp. 3, Jan. 1999, pp. 9-13.
- Gollob, J.A.; Schinpper, C.P.; Orsini, E.; Murphy, E.; Daley, J.F.; Lazo, S.B.; Frank, D.A.; *Characterization of a Novel Subset of CD8 T Cells That Expands in patients Receiving Interleukin-12*, 02, Am. Soc. For Clin. Investigation, Inc., vol. 102, No. 3, Aug. 1998, pp. 561-575.
- Cavanagii, L.L.; Barnetson, R.S.; Basten, A.; Halliday, G.M.; *Dendritic Epidermal T-Cell Involvement in Induction of CD8+ T-Cell-Mediated Immunity Against an Ultraviolet Radiation-Induced Skin Tumor* Int. J. Cancer: 70, 98-105, 1997.

US 9,101,621 B2

Page 7

(56)

References Cited

OTHER PUBLICATIONS

- Thomas, D.A., Aguayo, A., Estey, E., Albitar, M., O'Brien, S., Giles, F.J., Beran, M., Cortes, J., Zeldis, J., Keating, M.J., Barlogie, B., Kantarjian, H.M., Thalidomide as anti-angiogenesis therapy (rx) in refractory or relapsed leukemia. Abstract #2269, American Society of Hematology, Dec. 3-7, 1999.
- Barlogie, B., Desikan, R., Munshi, N., Siegel, D., Mehta, J., Singhal, S., Anaissie, E., Single Course D.T. Pace Anti-Angiochemotherapy Effects CR in Plasma Cell Leukemia and Fulminant Multiple Myeloma (MM). Abstract #4180. American Society of Hematology, Dec. 4-9, 1998.
- Hideshima, T., Chauhan, D., Shima, Y., Noopur, R., Davies, F.E., Tai, Y., Treon, S.P., Lin, B.K., Schlossman, R.L., Richardson, P.G., Gupta, D., Muller, G.W., Stirling, D.I., Anderson, K.C., Thalidomide (THAL) and its Analogs Overcome Drug Resistance of Human Multiple Myeloma (MM) Cells to Conventional Therapy. Abstract #1313. American Society of Hematology, Dec. 1-5, 2000.
- Payvandi, F., Wu, L., Gupta, D., Hideshima, T., Haley, M., Muller, G., Chen, R., Anderson, K.C., Stirling, D., Effects of a Thalidomide Analog on Binding Activity of Transcription Factors and Cell Cycle Progression of Multiple Myeloma Cell Lines. Abstract #2487. American Society of Hematology, Dec. 1-5, 2000.
- Davies, F.E., Raje, N., Hideshima, T., Lentzsch, S., Young, G., Tai, Y., Lin, B.K., Podar, K., Chauhan, D., Treon, S.P., Gupta, D., Mitsiades, C., Mitsiades, N., Hayashi, T., Richardson, P.G., Schlossman, R.L., Muller, G.W., Stirling, D. I., Anderson, K.C., Thalidomide (THAL) and Immunomodulatory Derivatives (IMiDS) Augment Natural Killer (NK) Cell Cytotoxicity in Multiple Myeloma (MM). Abstract #3617. American Society of Hematology, Dec. 1-5, 2000.
- Hideshima, T., Chauhan, D., Castro, A., Hayashi, T., Mitsiades, C., Mitsiades, N., Akiyama, M., Richardson, P.G., Schlossman, R.L., Adams, J., Anderson, K.C., NF-KB as a Therapeutic Target in Multiple Myeloma (MM). Abstract #1581. American Society of Hematology, Dec. 7-11, 2001.
- Lentsch, S., Rogers, M., Leblanc, R., Birsner, A., Shah, J., Anderson K., D'Amato R., 3-Amino-Phthalimido-Glutarimide (S-3APG) Inhibits Angiogenesis and Growth in Drug Resistant Multiple Myeloma (MM) in vivo. Abstract #1976, American Society of Hematology, Dec. 7-11, 2001.
- Park, Y., Kim, S.A., Kim, C.J., Chung, J.H., Mechanism of the Effect of Thalidomide on Human Multiple Myeloma Cells. Abstract #2685. American Society of Clinical Oncology, May 12-17, 2001.
- Payvandi, F., Wu, L., Haley M., Gupta, D., Zhang, L., Schafer, P., Muller, G.W., Chen, R., Anderson, K.C., Stirling, D., Thalidomide Analogs IMiDS Inhibit Expression of Cyclooxygenase-2 in Multiple Myeloma Cell Line and LPS Stimulated PBMCs. Abstract #2689. American Society of Hematology, Dec. 7-11, 2001.
- Mitsiades, N., Mitsiades, C., Poulaki, V., Akiyama, M., Tai, Y., Lin, B., Hayashi, T., Catley, L., Hideshima, T., Chauhan, D., Treon, S.P., Anderson, K.C., Apoptotic Signaling Induced by Immunomodulatory Thalidomide Analogs (Imids) in Human Multiple Myeloma Cells; Therapeutic Implications. Abstract #3224. American Society of Hematology, Dec. 7-11, 2001.
- Richardson, P.G., Schlossman, R.L., Hideshima, T., Davies, F., Leblanc, R., Catley, L., Doss, D., Kelly, K.A., McKenney, M., Mechlowicz, J., Freeman, A., Deocampo, R., Rich, R., Ryou, J., Chauhan, D., Munshi, N., Weller, E., Zeldis, J., Anderson, K.C., A Phase I Study of Oral CC5013, an Immunomodulatory Thalidomide (Thal) Derivative, in Patients With Relapsed and Refractory Multiple Myeloma (MM). Abstract #3225. American Society of Hematology, Dec. 7-11, 2001.
- "Celgene drug promises activity in solid tumors," Marketletter, Jun. 18, 2001.
- Meregalli et al., "High-dose dexamethasone as first line therapy of multiple myeloma?" *Recenti Progressi in Medicina*, 1998, 89(1):18-20.
- Official Action in corresponding Canadian Application No. 2,476,983 dated Aug. 21, 2009.
- List, A., "New Approaches to the Treatment of Myelodysplasia," *The Oncologist*, 2002, 7(suppl. 1):39-49.
- Kurzrock, R., "Myelodysplastic syndrome overview," *Seminars in Hematology* (Abstract only), 2002, 39(3)(suppl. 2):18-25 Abstract only.
- Goerner, et al., "Morbidity and mortality of chronic GVHD after hematopoietic stem cell transplantation from HLA-identical siblings for patients with aplastic or refractory anemias," *Biology of Blood and Marrow Transplantation* (Abstract only), 2002, 8(1):47-56.
- Thomas, D., "Pilot studies of Thalidomide in Acute Myelogenous Leukemia, Myelodysplastic Syndromes, and Myeloproliferative Disorders," *Seminars in Hematology*, 2000, 37(1)(suppl. 3):26-34.
- Zorat, F. et al., "The clinical and biological effects of thalidomide in patients with myelodysplastic syndromes," *British Journal of Haematology*, 2001, 115:881-894.
- Official Action dated Feb. 10, 2009 in JP Application No. 2004-545192. (English translation provided.)
- Teramura, M., Men-ekiyokusei Ryouhou, *Current Therapy*, 2000, 18(5):140-144 (in Japanese).
- Kon-nichi no Chiryu Shishin, 1997 [Pocket Edition], Igaku Shoin, 1997, 513-514 (in Japanese).
- Okamoto, T., Kotsuzukeisei Shoukougun to Men-eki Ijo, Bessatsu Nihon Rinsho, Syndrome Series for each area, No. 22, Blood Syndromes III, Nihon Rinsho, 213-216 (in Japanese), Oct. 1998.
- Merck Manual, 17th ed. Japanese version, 1999, 951-952.
- Notice of Allowance from U.S. Appl. No. 11/096,155 dated Jan. 12, 2010.
- Rajkumar et al., "Combination therapy with thalidomide plus dexamethasone for newly diagnosed multiple myeloma," *American Society of Hematology*, 43rd Annual Meeting, Dec. 7-11, 2001, Abstract #3525.
- Scheffler et al., "Safety and pharmacokinetics of CDC-501, a novel immunomodulatory-oncologic agent, after single then multiple, oral 100 mg twice daily doses," *American Society for Clinical Pharmacology and Therapeutics*, Mar. 24-27, 2002, Abstract #WP111-63.
- Marriott et al., "Thalidomide analogue CDC-501 is safe and well tolerated by patients with end stage cancer and shows evidence of clinical responses and extensive immune activation," *Br. J. Cancer*, 2002, 86(Suppl. 1):Abst 64.
- Kast, R.E., "Evidence of a mechanism by which etanercept increased TNF-alpha in multiple myeloma: New insights into the biology of TNF-alpha giving new treatment opportunities—the role of burpion," *Leukemia Research*, 2005, 29:1459-1463.
- Tsimberidou, A. et al., "Pilot study of recombinant human soluble tumor necrosis factor (TNF) receptor (p75) fusion protein (TNFR:Fc;Enbrel) in patients with refractory multiple myeloma: increase in plasma TNF α levels during treatment," *Leukemia Research*, 2003, 27:375-380.
- Dimopoulos, et al., "Long-term follow-up on overall survival from the MM-009 and MM-010 phase III trials of lenalidomide plus dexamethasone in patients with relapsed or refractory multiple myeloma," *Leukemia*, 2009, 1-6.
- Hideshima, T., et al., "A review of lenalidomide in combination with dexamethasone for the treatment of multiple myeloma," *Therapeutics and Clinical Risk Management*, 2008, 4(1):129-136.
- Wang, M., et al., "Lenalidomide plus dexamethasone is more effective than dexamethasone alone in patients with relapsed or refractory multiple myeloma regardless of prior thalidomide exposure," *Blood*, 2008, 112(12):4445-4451.
- Gandhi, A., et al., "Dexamethasone Synergizes with Lenalidomide to Inhibit Multiple Myeloma Tumor Growth, But Reduces Lenalidomide-Induced Immunomodulation of T and NK Cell Function," *Current Cancer Drug Targets*, 2010, 10(1):1-13.
- Gay, F. et al., "Lenalidomide plus dexamethasone versus thalidomide plus dexamethasone in newly diagnosed multiple myeloma: a comparative analysis of 411 patients," *Blood*, 2010, 115(97):1343-150.
- Richardson, P. et al., "Thalidomide in multiple myeloma," *Biomed Pharmacother*, 2002, 56:115-28.
- Swartz, G. et al., "Pre-clinical evaluation of ENMD-0995: A thalidomide analog with activity against multiple myeloma and solid tumors," *Cell and Tumor Biology*, 2002, 43:181-182, Abstract# 910.
- Mazucco, R., "Angiogenesis and Anti-angiogenesis Therapeutics," *IDrugs*, 2002, 5(4): 320-322.
- Worker, C., "JP Morgan Hambrecht & Quist—20th Annual Healthcare Conference," *IDrugs*, 2002, 5(2):113-116.

US 9,101,621 B2

Page 8

(56) References Cited

OTHER PUBLICATIONS

Treston, A. et al., "Pre-Clinical Evaluation of a Thalidomide Analog with Activity Against Multiple Myeloma and Solid Tumors—ENMD-0995 (S-(-)-3-(3-amino-phthalimido)-glutarimide)," *Blood*, 2002, 100(11):816a, Abstract #3225.

Mazucco, R. and Williams, L., "Immunotherapy, chemoprevention and angiogenesis," *IDrugs*, 2002, 5(5):408-411.

Fernandes, P., "Anti-Cancer Drug Discovery and Development Summit," *IDrugs*, 2002, 5(8):757-764.

Notification letter dated Aug. 30, 2010 from Natco Pharma Limited to Celgene Corporation re: Notification pursuant to § 505(j)(2)(B) of the Federal Food, Drug and Cosmetic Act.

Complaint for Patent Infringement filed on Oct. 8, 2010 by Celgene Corporation in the U.S. District Court, District of New Jersey against Natco Pharma Limited.

Answer to Complaint filed on Nov. 18, 2010 by Nalco Pharma Limited in the U.S. District Court, District of New Jersey.

Grosshans, E. and Illy, G., "Thalidomide Therapy for Inflammatory Dermatoses," *International Journal of Dermatology*, 1984, 23(9):598-602.

Krenn, M. et al., "Improvements in Solubility and Stability of Thalidomide upon Complexation with Hydropropyl-β-Cyclodextrin," *Journal of Pharmaceutical Sciences*, 1992, 81(7):685-689.

Schmahl, H. J. et al., "Pharmacokinetics of the Teratogenic and Nonteratogenic Thalidomide Analogs EM12 and Supidimide in the Rat and Marmoset Monkey," in *Pharmacokinetics in Teratogenesis*, CRC Press, 1987, vol. I, Ch. 12, pp. 181-192.

Schumacher, H. et al., "The Teratogenic Activity of a Thalidomide Analogue, EM₁₂, in Rabbits, Rats, and Monkeys," *Teratology*, 1971, 5:233-240.

Smith, R. et al., "Studies on the Relationship Between the Chemical Structure and Embryotoxic Activity of Thalidomide and Related Compounds," in *A Symposium on Embryopathic Activity of Drugs*, J. & A. Churchill Ltd., 1965, Session 6, pp. 194-209.

Sheskin, J. and Sagher, F., "Trials with Thalidomide Derivatives in Leprosy Reactions," *Leprosy Review*, 1968, 39(4):203-205.

Sheskin, J., "Study with Nine Thalidomide Derivatives in the Leprosy Reaction," *Pharmacology and Therapeutics*, 1978, 17:82-84.

Raje, N. and Anderson, K., "Thalidomide and immunomodulatory drugs as cancer therapy," *Current Opinions in Oncology*, 2002, 14:635-640.

Kumar, S. et al., "Thalidomide as an anti-cancer agent," *J. Cell. Mod. Med.*, 2002, 6(2):160-174.

Singhal, S. and Mehta, J., "Thalidomide in Cancer," *BioDrugs*, 2001, 15(3):163-172.

Notice of Opposition to EP 1 505 973 filed by Synthon B.V. on Nov. 30, 2010.

Notice of Opposition to EP 1 505 973 filed by Strawman Limited on Dec. 1, 2010.

Samson, D. et al., "Infusion of Vincristine and Doxorubicin with Oral Dexamethasone as First-Line Therapy for Multiple Myeloma," *The Lancet*, 1989, 334(8668):882-885.

Barlogie, B. et al., "Effective Treatment of Advanced Multiple Myeloma Refractory to Alkylating Agents," *N. Engl. J. Med.*, 1984, 310(21):1353-1356.

Dimopoulos, M. et al., "Thalidomide and dexamethasone combination for refractory multiple myeloma," *Annals of Oncology*, 2001, 12:991-995.

Zangari, M., et al., "Thrombogenic activity of doxorubicin in myeloma patients receiving thalidomide: implications for therapy," *Blood*, 2002, 100:1168-1171.

List, A. et al., "High Erythropoietic Remitting Activity of the Immunomodulatory Thalidomide Analog, CC5013, in Patients with Myelodysplastic Syndrome (MDS)," Abstract #353, *Blood*, 2002, 100(11):96a.

Mufti, G. et al., "Myelodysplastic Syndrome," *American Society of Hematology*, 2003, pp. 176-199.

Extracts from drug databases: retrieved from <http://www.nextbio.com/b/search/ov/IMiD3%20cpd> on Nov. 26, 2010 and <http://pubchem.ncbi.nlm.nih.gov/summary/summary.cgi?cid=216326> on Nov. 26, 2010.

Stockdale, 1998, *Medicine*, Rubenstein and Federman, eds., vol. 3, Ch. 12, Sections IV and X.

"List of Approved Oncology Drugs with Approved Indications," <http://www.accessdata.fda.gov/scripts/cder/onctools/druglist.cfm>, last accessed on Sep. 11, 2008.

Office Action mailed Jun. 18, 2008, U.S. Appl. No. 11/325,954.

Gamberi et al., "Overall Safety and Treatment Duration in Lenalidomide (LEN)-, Thalidomide (THAL)-, and Bortezomib (BORT)-Treated Patients (Pts) within the European Post-Approval Safety Study (EU PASS) of Relapsed/Refractory Multiple Myeloma (RRMM)," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #4068.

Korde et al., "Phase II Clinical and Correlative Study of Carfilzomib, Lenalidomide, and Dexamethasone (CRd) in Newly Diagnosed Multiple Myeloma (MM) Patients", presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #732.

Kumar et al., "A Phase 1/2 Study of Weekly MLN9708, an Investigational Oral Proteasome Inhibitor, in Combination with Lenalidomide and Dexamethasone in Patients with Previously Untreated Multiple Myeloma (MM)", presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #332.

Palumbo et al., "Pomalidomide Cyclophosphamide and Prednisone (PCP) Treatment for Relapsed/Refractory Multiple Myeloma", presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #446.

Richardson et al., "A Phase 2 Study of Elotuzumab (Elo) in Combination with Lenalidomide and Low-Dose Dexamethasone (Ld) in Patients (pts) with Relapsed/Refractory Multiple Myeloma (R/R MM): Updated Results", presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #202.

Sacchi et al., "A Phase I/II Study of Bendamustine, Low-Dose Dexamethasone, and Lenalidomide (BdL) for the Treatment of Patients with Relapsed Multiple Myeloma", presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #1851.

Sonneveld et al., "Escalated Dose Bortezomib Once Weekly Combined with Lenalidomide and Dexamethasone (eVRD) Followed by Lenalidomide Maintenance in First Relapse of Multiple Myeloma (MM), the HOVON 86 Phase 2 Trial", presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #1853.

Suvannasankha et al., "A Phase I/II Trial Combining High-Dose Lenalidomide with Melphalan and Autologous Transplant for Multiple Myeloma: A Report of the Phase I Dose-Finding Study", presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #3146.

Mark et al., "ClAPD (Clarithromycin, Pomalidomide, Dexamethasone) Therapy in Relapsed or Refractory Multiple Myeloma", presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #77.

Lacy et al., "Pomalidomide Plus Low-Dose Dexamethasone (Pom/Dex) in Relapsed Myeloma: Long Term Follow up and Factors Predicting Outcome in 345 Patients," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #201.

Jagannath et al., "Pomalidomide (POM) with Low-Dose Dexamethasone (LoDex) in Patients (Pts) with Relapsed and Refractory Multiple Myeloma Who Have Received Prior Therapy with Lenalidomide (LEN) and Bortezomib (BORT): Updated Phase 2 Results and Age Subgroup Analysis," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #450.

US 9,101,621 B2

Page 9

(56)

References Cited

OTHER PUBLICATIONS

Richardson et al., "MM-005: A Phase 1, Multicenter, Open-Label, Dose-Escalation Study to Determine the Maximum Tolerated Dose for the Combination of Pomalidomide, Bortezomib, and Low-Dose Dexamethasone in Subjects with Relapsed or Refractory Multiple Myeloma," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #727.

Leleu et al., "Prolonged Overall Survival with Pomalidomide and Dexamethasone in Myeloma Characterized with End Stage Disease," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #2961.

Berenson et al., "A Phase 1/2 Study of Pomalidomide, Dexamethasone and Pegylated Liposomal Doxorubicin for Patients with Relapsed/Refractory Multiple Myeloma," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012., Abstract #2979.

Lonial et al., "Improvement in Clinical Benefit Parameters with Pomalidomide (POM) in Combination with Low-Dose Dexamethasone (LoDex) in Patients with Relapsed and Refractory Multiple Myeloma (RRMM): Results From a Phase 2 Study," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #4052.

Vij et al., "Pomalidomide (POM) with Low-Dose Dexamethasone (LoDex) in Patients with Relapsed and Refractory Multiple Myeloma (RRMM): Outcomes Based on Prior Treatment Exposure," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #4070.

Richardson et al., "Treatment Outcomes with Pomalidomide (POM) in Combination with Low-Dose Dexamethasone (LoDex) in Patients with Relapsed and Refractory Multiple Myeloma (RRMM) and Del(17p13) and/or t(4;14) (p16;q32) Cytogenic Abnormalities Who Have Received Prior Therapy with Lenalidomide (LEN) and Bortezomib (BORT)," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #4053.

Dimopoulos et al., "Pomalidomide in Combination with Low-Dose Dexamethasone: Demonstrates a Significant Progression Free Survival and Overall Survival Advantage, in Relapsed/Refractory MM: A Phase 3, Multicenter, Randomized, Open-Label Study," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #LBA-6.

Shastri et al., "A Phase II Study of Low-Dose Pomalidomide (0.5mg/day) and Prednisone Combination Therapy in Patients with Myelofibrosis and Significant Anemia," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #1728.

Shah et al., "A Multi-Center Phase I/II Trial of Carfilzomib and Pomalidomide with Dexamethasone (Car-Pom-d) in Patients with Relapsed/Refractory Multiple Myeloma," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #74.

Office Action in corresponding CN Application No. 201110256752.0 dated Feb. 8, 2013.

Stirling, D., "Thalidomide: A Novel Template for Anticancer Drugs," *Seminars in Oncology*, Dec. 2001, 28(6):602-606.

Celgene Press Release, "Celgene Will Discontinue Phase III Origin® Trial in Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukemia," published on Celgene Newsroom, <http://newsroom.celgene.com> on Jul. 18, 2013 at 7:30 am EDT.

Mateos, M.-V., Ph.D. et al., "Lenalidomide plus Dexamethasone for High-Risk Smoldering Multiple Myeloma," *New England Journal of Medicine*, Aug. 2013, 369(5):438-447.

English translation of Japanese IP High Court decision in Application No. JP 2004-505051, dated Apr. 11, 2013.

Jagannath, S. et al., "Pomalidomide (POM) with or without low-dose dexamethasone (LoDEX) in patients (Pts) with relapsed and refractory multiple myeloma (RRMM): MM-002 phase II age subgroup analysis," *J Clin Oncol* 31, 2013 (suppl; abstr 8532).

Siegel, D. et al., "Long-term safety and efficacy of pomalidomide (POM) with or without low-dose dexamethasone (LoDEX) in

relapsed and refractory multiple myeloma (RRMM) patients enrolled in the MM-002 phase II trial," *J Clin Oncol* 31, 2013 (suppl; abstr 8588).

Richardson, P.G. et al., A Phase 1/2 Multi-Center, Randomized, Open Label Dose Escalation Study to Determine the Maximum Tolerated Dose (MTD), Safety, and Efficacy of Pomalidomide (POM) Alone or in Combination with Low-Dose Dexamethasone (DEX) in Patients (PTS) with Relapsed and Refractory Multiple Myeloma (RRMM) Who Have Received Prior Treatment (TX) That Includes Lenalidomide (LEN) and Bortezomib (BORT), *Haematologica*, 2001; 96(s1):S31, Abstract O-12, *13th International Myeloma Workshop*, Paris, France—May 3-6, 2011.

MacNeil, J.S., "Pomalidomide Picks Up Where Both Earlier IMiDs Stop Working," *The Oncology Report*, Mar./Apr. 2010, p. 34.

Alexanian, R. et al., "Primary Dexamethasone Treatment of Multiple Myeloma," *Blood*, 1992, 80(4):887-890.

Berenson, J. R. et al., "Long-Term Pamidronate Treatment of Advanced Multiple Myeloma Patients Reduces Skeletal Events," *Journal of Clinical Oncology*, 1998, 16(2):593-602.

Attal, M. et al., "Lenalidomide Maintenance after Stem-Cell Transplantation for Multiple Myeloma", *N Engl J Med*, 2012, 366(19):1782-1791.

McCarthy, P.L., et al., "Lenalidomide after Stem-Cell Transplantation for Multiple Myeloma", *N Engl J Med*, 2012, 366(19):1770-1781.

Fifth Amended Complaint for Patent Infringement, dated May 6, 2013, *Celgene Corp. v. NATCO Pharma Ltd.*, D.N.J. (10-cv-5197). Amended Answer, Affirmative Defense and Counterclaims to Plaintiff's Fifth Amended Complaint, dated Aug. 1, 2013, *Celgene Corp. v. NATCO Pharma Ltd.*, D.N.J. (10-cv-5197).

Reply to Counterclaims to Fifth Amended Complaint, Jun. 10, 2013, *Celgene Corp. v. NATCO Pharma Ltd.*, D.N.J. (10-cv-5197).

Celgene's Opening Markman Brief, dated Oct. 21, 2013, *Celgene Corp. v. NATCO Pharma Ltd.*, D.N.J. (10-cv-5197).

Markman Opinion, dated May 27, 2014, *Celgene Corp. v. NATCO Pharma Ltd.*, D.N.J. (10-cv-5197).

European Patent Office's Decision revoking European Patent No. EP-B-1 505 973, dated Feb. 25, 2013.

Office Action in U.S. Appl. No. 13/276,867 dated Mar. 3, 2015.

Office Action in corresponding Japanese Patent Application No. 2012-273326 dated May 13, 2014.

Office Action in corresponding Canadian Patent Application No. 2,855,359 dated Aug. 14, 2014.

"A Study to Compare MPR With MP in Newly Diagnosed Multiple Myeloma Subjects 65 Years Old or Older," available at <http://clinicaltrials.gov/show/NCT00405756>, last accessed on Jul. 21, 2014.

Durie B., "VIIIth International Myeloma Workshop Meeting Report from Banff, Canada May 4-8, 2001", *Myeloma Today*.

Gertz et al., "Melphalan and Prednisone," *Myeloma Today*, 1999, 3(6):4 and 14-15.

Gregory, W. et al., "Combination Chemotherapy Versus Melphalan and Prednisolone in the Treatment of Multiple Myeloma: An Overview of Published Trials," *J of Clinical Oncology*, 1992, 10(2):334-342.

Kyle et al., "Therapeutic Application of Thalidomide in Multiple Myeloma", *Seminars in Oncology* (2001) 28, 583-587.

Reiman T. et al., "Meeting Synopsis, VIII International Myeloma Workshop, Banff Springs Hotel, Banff, Alberta, Canada, May 4-8, 2001", *European Journal of Haematology*, 2001, 67: 199-202.

Riet, I., "Angiogenesis in Multiple Myeloma," The Comprehensive Guide to Banff: Understanding the VIIIth International Myeloma Workshop, p. 12-13, May 5-8, 2001.

Weerdt et al., "Continuous low-dose cyclophosphamide-prednisone is effective and well tolerated in patients with advanced multiple myeloma," *The Netherlands Journal of Medicine*, Aug. 2001, 59(2):50-56.

Zaidi et al. "Multiple myeloma: an old disease with new hope for the future." *CA Cancer J Clin.* 2001 51(5):273-85.



* cited by examiner

U.S. Patent

Aug. 11, 2015

US 9,101,621 B2

Effects of Revimid™ and Thalidomide on MM Cell Proliferation



US 9,101,621 B2

1

**METHODS FOR TREATING MULTIPLE
MYELOMA WITH 3-(4-AMINO-1-OXO-1,3-
DIHYDRO-ISOINDOL-2-YL)-PIPERIDINE-
2,6-DIONE AFTER STEM CELL
TRANSPLANTATION**

This application is a continuation of U.S. patent application Ser. No. 14/201,069, filed Mar. 7, 2014, which is a continuation of U.S. patent application Ser. No. 13/782,728, filed Mar. 1, 2013, now U.S. Pat. No. 8,673,939, which is a continuation of U.S. patent application Ser. No. 13/488,888, filed Jun. 5, 2012, now U.S. Pat. No. 8,648,095, which is a continuation of U.S. patent application Ser. No. 12/640,702, filed Dec. 17, 2009, now U.S. Pat. No. 8,198,306, which is a continuation application of U.S. patent application Ser. No. 10/438,213, filed May 15, 2003, now U.S. Pat. No. 7,968,569, which claims the benefit of U.S. provisional application nos. 60/380,842, filed May 17, 2002, and 60/424,600, filed Nov. 6, 2002, the entireties of which are incorporated herein by reference.

1. FIELD OF THE INVENTION

This invention relates to methods of treating, preventing and/or managing specific cancers, and other diseases including, but not limited to, those associated with, or characterized by, undesired angiogenesis, by the administration of one or more immunomodulatory compounds alone or in combination with other therapeutics. In particular, the invention encompasses the use of specific combinations, or "cocktails," of drugs and other therapy, e.g., radiation to treat these specific cancers, including those refractory to conventional therapy. The invention also relates to pharmaceutical compositions and dosing regimens.

2. BACKGROUND OF THE INVENTION

2.1 Pathobiology of Cancer and Other Diseases

Cancer is characterized primarily by an increase in the number of abnormal cells derived from a given normal tissue, invasion of adjacent tissues by these abnormal cells, or lymphatic or blood-borne spread of malignant cells to regional lymph nodes and to distant sites (metastasis). Clinical data and molecular biologic studies indicate that cancer is a multistep process that begins with minor preneoplastic changes, which may under certain conditions progress to neoplasia. The neoplastic lesion may evolve clonally and develop an increasing capacity for invasion, growth, metastasis, and heterogeneity, especially under conditions in which the neoplastic cells escape the host's immune surveillance. Roitt, I., Brostoff, J and Kale, D., *Immunology*, 17.1-17.12 (3rd ed., Mosby, St. Louis, Mo., 1993).

There is an enormous variety of cancers which are described in detail in the medical literature. Examples includes cancer of the lung, colon, rectum, prostate, breast, brain, and intestine. The incidence of cancer continues to climb as the general population ages, as new cancers develop, and as susceptible populations (e.g., people infected with AIDS or excessively exposed to sunlight) grow. A tremendous demand therefore exists for new methods and compositions that can be used to treat patients with cancer.

Many types of cancers are associated with new blood vessel formation, a process known as angiogenesis. Several of the mechanisms involved in tumor-induced angiogenesis have been elucidated. The most direct of these mechanisms is the secretion by the tumor cells of cytokines with angiogenic

2

properties. Examples of these cytokines include acidic and basic fibroblastic growth factor (a,b-FGF), angiogenin, vascular endothelial growth factor (VEGF), and TNF- α . Alternatively, tumor cells can release angiogenic peptides through the production of proteases and the subsequent breakdown of the extracellular matrix where some cytokines are stored (e.g., b-FGF). Angiogenesis can also be induced indirectly through the recruitment of inflammatory cells (particularly macrophages) and their subsequent release of angiogenic cytokines (e.g., TNF- α , bFGF).

A variety of other diseases and disorders are also associated with, or characterized by, undesired angiogenesis. For example, enhanced or unregulated angiogenesis has been implicated in a number of diseases and medical conditions including, but not limited to, ocular neovascular diseases, choroidal neovascular diseases, retina neovascular diseases, rubeosis (neovascularization of the angle), viral diseases, genetic diseases, inflammatory diseases, allergic diseases, and autoimmune diseases. Examples of such diseases and conditions include, but are not limited to: diabetic retinopathy; retinopathy of prematurity; corneal graft rejection; neovascular glaucoma; retrolental fibroplasia; and proliferative vitreoretinopathy.

Accordingly, compounds that can control angiogenesis or inhibit the production of certain cytokines, including TNF- α , may be useful in the treatment and prevention of various diseases and conditions.

2.2 Methods of Treating Cancer

Current cancer therapy may involve surgery, chemotherapy, hormonal therapy and/or radiation treatment to eradicate neoplastic cells in a patient (see, for example, Stockdale, 1998, *Medicine*, vol. 3, Rubenstein and Federman, eds., Chapter 12, Section IV). Recently, cancer therapy could also involve biological therapy or immunotherapy. All of these approaches pose significant drawbacks for the patient. Surgery, for example, may be contraindicated due to the health of a patient or may be unacceptable to the patient. Additionally, surgery may not completely remove neoplastic tissue. Radiation therapy is only effective when the neoplastic tissue exhibits a higher sensitivity to radiation than normal tissue. Radiation therapy can also often elicit serious side effects. Hormonal therapy is rarely given as a single agent. Although hormonal therapy can be effective, it is often used to prevent or delay recurrence of cancer after other treatments have removed the majority of cancer cells. Biological therapies and immunotherapies are limited in number and may produce side effects such as rashes or swellings, flu-like symptoms, including fever, chills and fatigue, digestive tract problems or allergic reactions.

With respect to chemotherapy, there are a variety of chemotherapeutic agents available for treatment of cancer. A majority of cancer chemotherapeutics act by inhibiting DNA synthesis, either directly, or indirectly by inhibiting the biosynthesis of deoxyribonucleotide triphosphate precursors, to prevent DNA replication and concomitant cell division. Gilman et al., *Goodman and Gilman's: The Pharmacological Basis of Therapeutics*, Tenth Ed. (McGraw Hill, New York).

Despite availability of a variety of chemotherapeutic agents, chemotherapy has many drawbacks. Stockdale, *Medicine*, vol. 3, Rubenstein and Federman, eds., ch. 12, sect. 10, 1998. Almost all chemotherapeutic agents are toxic, and chemotherapy causes significant, and often dangerous side effects including severe nausea, bone marrow depression, and immunosuppression. Additionally, even with administration of combinations of chemotherapeutic agents, many tumor

US 9,101,621 B2

3

cells are resistant or develop resistance to the chemotherapeutic agents. In fact, those cells resistant to the particular chemotherapeutic agents used in the treatment protocol often prove to be resistant to other drugs, even if those agents act by different mechanism from those of the drugs used in the specific treatment. This phenomenon is referred to as pleiotropic drug or multidrug resistance. Because of the drug resistance, many cancers prove refractory to standard chemotherapeutic treatment protocols.

Other diseases or conditions associated with, or characterized by, undesired angiogenesis are also difficult to treat. However, some compounds such as protamine, heparin and steroids have been proposed to be useful in the treatment of certain specific diseases. Taylor et al., *Nature* 297:307 (1982); Folkman et al., *Science* 221:719 (1983); and U.S. Pat. Nos. 5,001,116 and 4,994,443. Thalidomide and certain derivatives of it have also been proposed for the treatment of such diseases and conditions. U.S. Pat. Nos. 5,593,990, 5,629,327, 5,712,291, 6,071,948 and 6,114,355 to D'Amato.

Still, there is a significant need for safe and effective methods of treating, preventing and managing cancer and other diseases and conditions, particularly for diseases that are refractory to standard treatments, such as surgery, radiation therapy, chemotherapy and hormonal therapy, while reducing or avoiding the toxicities and/or side effects associated with the conventional therapies.

2.3 IMIDS™

A number of studies have been conducted with the aim of providing compounds that can safely and effectively be used to treat diseases associated with abnormal production of TNF- α . See, e.g., Marriott, J. B., et al., *Expert Opin. Biol. Ther.* 1(4):1-8 (2001); G. W. Muller, et al., *Journal of Medicinal Chemistry* 39(17): 3238-3240 (1996); and G. W. Muller, et al., *Bioorganic & Medicinal Chemistry Letters* 8: 2669-2674 (1998). Some studies have focused on a group of compounds selected for their capacity to potently inhibit TNF- α production by LPS stimulated PBMC. L. G. Corral, et al., *Ann. Rheum. Dis.* 58:(Suppl I) 1107-1113 (1999). These compounds, which are referred to as IMiDs™ (Celgene Corporation) or Immunomodulatory Drugs, show not only potent inhibition of TNF- α but also marked inhibition of LPS induced monocyte IL1 β and IL12 production. LPS induced IL6 is also inhibited by immunomodulatory compounds, albeit partially. These compounds are potent stimulators of LPS induced IL10. Id. Particular examples of IMiD™s include, but are not limited to, the substituted 2-(2,6-dioxopiperidin-3-yl)phthalimides and substituted 2-(2,6-dioxopiperidin-3-yl)-1-oxoisindoles described in U.S. Pat. Nos. 6,281,230 and 6,316,471, both to G. W. Muller, et al.

3. SUMMARY OF THE INVENTION

This invention encompasses methods of treating and preventing certain types of cancer, including primary and metastatic cancer, as well as cancers that are refractory or resistant to conventional chemotherapy. The methods comprise administering to a patient in need of such treatment or prevention a therapeutically or prophylactically effective amount of an immunomodulatory compound, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof. The invention also encompasses methods of managing certain cancers (e.g., preventing or prolonging their recurrence, or lengthening the time of remission) which comprise administering to a patient in need of such management a prophylactically effective amount of an

4

immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

In particular methods of the invention, an immunomodulatory compound is administered in combination with a therapy conventionally used to treat, prevent or manage cancer. Examples of such conventional therapies include, but are not limited to, surgery, chemotherapy, radiation therapy, hormonal therapy, biological therapy and immunotherapy.

This invention also encompasses methods of treating, managing or preventing diseases and disorders other than cancer that are associated with, or characterized by, undesired angiogenesis, which comprise administering to a patient in need of such treatment, management or prevention a therapeutically or prophylactically effective amount of an immunomodulatory compound, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

In other methods of the invention, an immunomodulatory compound is administered in combination with a therapy conventionally used to treat, prevent or manage diseases or disorders associated with, or characterized by, undesired angiogenesis. Examples of such conventional therapies include, but are not limited to, surgery, chemotherapy, radiation therapy, hormonal therapy, biological therapy and immunotherapy.

This invention encompasses pharmaceutical compositions, single unit dosage forms, dosing regimens and kits which comprise an immunomodulatory compound, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, and a second, or additional, active agent. Second active agents include specific combinations, or "cocktails," of drugs.

4. BRIEF DESCRIPTION OF FIGURE

FIG. 1 shows a comparison of the effects of 3-(4-amino-1-oxo-1,3-dihydro-isindol-2-yl)-piperidine-2,6-dione (Revimid™) and thalidomide in inhibiting the proliferation of multiple myeloma (MM) cell lines in an in vitro study. The uptake of [³H]-thymidine by different MM cell lines (MM.1S, Hs Sultan, U266 and RPMI-8226) was measured as an indicator of the cell proliferation.

5. DETAILED DESCRIPTION OF THE INVENTION

A first embodiment of the invention encompasses methods of treating, managing, or preventing cancer which comprises administering to a patient in need of such treatment or prevention a therapeutically or prophylactically effective amount of an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

In particular methods encompassed by this embodiment, the immunomodulatory compound is administered in combination with another drug ("second active agent") or method of treating, managing, or preventing cancer. Second active agents include small molecules and large molecules (e.g., proteins and antibodies), examples of which are provided herein, as well as stem cells. Methods, or therapies, that can be used in combination with the administration of the immunomodulatory compound include, but are not limited to, surgery, blood transfusions, immunotherapy, biological therapy, radiation therapy, and other non-drug based therapies presently used to treat, prevent or manage cancer.

Another embodiment of the invention encompasses methods of treating, managing or preventing diseases and disorders

ders other than cancer that are characterized by undesired angiogenesis. These methods comprise the administration of a therapeutically or prophylactically effective amount of an immunomodulatory compound, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

Examples of diseases and disorders associated with, or characterized by, undesired angiogenesis include, but are not limited to, inflammatory diseases, autoimmune diseases, viral diseases, genetic diseases, allergic diseases, bacterial diseases, ocular neovascular diseases, choroidal neovascular diseases, retina neovascular diseases, and rubeosis (neovascularization of the angle).

In particular methods encompassed by this embodiment, the immunomodulatory compound is administered in combination with a second active agent or method of treating, managing, or preventing the disease or condition. Second active agents include small molecules and large molecules (e.g., proteins and antibodies), examples of which are provided herein, as well as stem cells. Methods, or therapies, that can be used in combination with the administration of the immunomodulatory compound include, but are not limited to, surgery, blood transfusions, immunotherapy, biological therapy, radiation therapy, and other non-drug based therapies presently used to treat, prevent or manage disease and conditions associated with, or characterized by, undesired angiogenesis.

The invention also encompasses pharmaceutical compositions (e.g., single unit dosage forms) that can be used in methods disclosed herein. Particular pharmaceutical compositions comprise an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, and a second active agent.

5.1 Immunomodulatory Compounds

Compounds used in the invention include immunomodulatory compounds that are racemic, stereomerically enriched or stereomerically pure, and pharmaceutically acceptable salts, solvates, hydrates, stereoisomers, clathrates, and prodrugs thereof. Preferred compounds used in the invention are small organic molecules having a molecular weight less than about 1,000 g/mol, and are not proteins, peptides, oligonucleotides, oligosaccharides or other macromolecules.

As used herein and unless otherwise indicated, the terms "immunomodulatory compounds" and "IMiDs™" (Celgene Corporation) encompasses small organic molecules that markedly inhibit TNF- α , LPS induced monocyte IL1 β and IL12, and partially inhibit IL6 production. Specific immunomodulatory compounds are discussed below.


TNF- α is an inflammatory cytokine produced by macrophages and monocytes during acute inflammation. TNF- α is responsible for a diverse range of signaling events within cells. TNF- α may play a pathological role in cancer. Without being limited by theory, one of the biological effects exerted by the immunomodulatory compounds of the invention is the reduction of synthesis of TNF- α . Immunomodulatory compounds of the invention enhance the degradation of TNF- α mRNA.

Further, without being limited by theory, immunomodulatory compounds used in the invention may also be potent co-stimulators of T cells and increase cell proliferation dramatically in a dose dependent manner. Immunomodulatory compounds of the invention may also have a greater co-stimulatory effect on the CD8+ T cell subset than on the

CD4+ T cell subset. In addition, the compounds preferably have anti-inflammatory properties, and efficiently co-stimulate T cells.

Specific examples of immunomodulatory compounds of the invention, include, but are not limited to, cyano and carboxy derivatives of substituted styrenes such as those disclosed in U.S. Pat. No. 5,929,117; 1-oxo-2-(2,6-dioxo-3-fluoropiperidin-3-yl) isoindolines and 1,3-dioxo-2-(2,6-dioxo-3-fluoropiperidine-3-yl) isoindolines such as those described in U.S. Pat. No. 5,874,448; the tetra substituted 2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolines described in U.S. Pat. No. 5,798,368; 1-oxo and 1,3-dioxo-2-(2,6-dioxopiperidin-3-yl)isoindolines (e.g., 4-methyl derivatives of thalidomide and EM-12), including, but not limited to, those disclosed in U.S. Pat. No. 5,635,517; and a class of non-polypeptide cyclic amides disclosed in U.S. Pat. Nos. 5,698,579 and 5,877,200; analogs and derivatives of thalidomide, including hydrolysis products, metabolites, derivatives and precursors of thalidomide, such as those described in U.S. Pat. Nos. 5,593,990, 5,629,327, and 6,071,948 to D'Amato; aminothalidomide, as well as analogs, hydrolysis products, metabolites, derivatives and precursors of aminothalidomide, and substituted 2-(2,6-dioxopiperidin-3-yl)phthalimides and substituted 2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindoles such as those described in U.S. Pat. Nos. 6,281,230 and 6,316,471; isoindole-imide compounds such as those described in U.S. patent application Ser. No. 09/972,487 filed on Oct. 5, 2001, U.S. patent application Ser. No. 10/032,286 filed on Dec. 21, 2001, and International Application No. PCT/US01/50401 (International Publication No. WO 02/059106). The entireties of each of the patents and patent applications identified herein are incorporated herein by reference. Immunomodulatory compounds of the invention do not include thalidomide.

Other specific immunomodulatory compounds of the invention include, but are not limited to, 1-oxo- and 1,3-dioxo-2-(2,6-dioxopiperidin-3-yl)isoindolines substituted with amino in the benzo ring as described in U.S. Pat. No. 5,635,517 which is incorporated herein by reference. These compounds have the structure I:



in which one of X and Y is C=O, the other of X and Y is C=O or CH₂, and R² is hydrogen or lower alkyl, in particular methyl. Specific immunomodulatory compounds include, but are not limited to:


1-oxo-2-(2,6-dioxopiperidin-3-yl)-4-aminoisoindoline;
1-oxo-2-(2,6-dioxopiperidin-3-yl)-5-aminoisoindoline;
1-oxo-2-(2,6-dioxopiperidin-3-yl)-6-aminoisoindoline;
1-oxo-2-(2,6-dioxopiperidin-3-yl)-7-aminoisoindoline;
1,3-dioxo-2-(2,6-dioxopiperidin-3-yl)-4-aminoisoindoline;
and
1,3-dioxo-2-(2,6-dioxopiperidin-3-yl)-5-aminoisoindoline.

Other specific immunomodulatory compounds of the invention belong to a class of substituted 2-(2,6-dioxopiperidin-3-yl)phthalimides and substituted 2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindoles, such as those described in U.S. Pat. Nos. 6,281,230; 6,316,471; 6,335,349; and 6,476,052, and International Patent Application No. PCT/US97/13375 (In-

US 9,101,621 B2


7

ternational Publication No. WO 98/03502), each of which is incorporated herein by reference. Compounds representative of this class are of the formulas:



wherein R^1 is hydrogen or methyl. In a separate embodiment, the invention encompasses the use of enantiomerically pure forms (e.g. optically pure (R) or (S) enantiomers) of these compounds.

Still other specific immunomodulatory compounds of the invention belong to a class of isoindole-imides disclosed in U.S. patent application Ser. Nos. 10/032,286 and 09/972,487, and International Application No. PCT/US01/50401 (International Publication No. WO 02/059106), each of which are incorporated herein by reference. Representative compounds are of formula II:



and pharmaceutically acceptable salts, hydrates, solvates, clathrates, enantiomers, diastereomers, racemates, and mixtures of stereoisomers thereof, wherein:

one of X and Y is C=O and the other is CH₂ or C=O;
 R^1 is H, (C₁-C₈)alkyl, (C₃-C₇)cycloalkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, benzyl, aryl, (C₀-C₄)alkyl-(C₁-C₆)heterocycloalkyl, (C₀-C₄)alkyl-(C₂-C₅)heteroaryl, C(O)R³, C(S)R³, C(O)OR⁴, (C₁-C₈)alkyl-N(R⁶)₂, (C₁-C₈)alkyl-OR⁵, (C₁-C₈)alkyl-C(O)OR⁵, C(O)NHR³, C(S)NHR³, C(O)NR³R³, C(S)NR³R³ or (C₁-C₈)alkyl-O(CO)R⁵;

R^2 is H, F, benzyl, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, or (C₂-C₈)alkynyl;

R^3 and R^3 are independently (C₁-C₈)alkyl, (C₃-C₇)cycloalkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, benzyl, aryl, (C₀-

8

C₄)alkyl-(C₁-C₆)heterocycloalkyl, (C₀-C₄)alkyl-(C₂-C₅)heteroaryl, (C₀-C₈)alkyl-N(R⁶)₂, (C₁-C₈)alkyl-OR⁵, (C₁-C₈)alkyl-C(O)OR⁵, (C₁-C₈)alkyl-O(CO)R⁵, or C(O)OR⁵;

R^4 is (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, (C₁-C₄)alkyl-OR⁵, benzyl, aryl, (C₀-C₄)alkyl-(C₁-C₆)heterocycloalkyl, or (C₀-C₄)alkyl-(C₂-C₅)heteroaryl;

R^5 is (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, benzyl, aryl, or (C₂-C₅)heteroaryl;

each occurrence of R^6 is independently H, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, benzyl, aryl, (C₂-C₅)heteroaryl, or (C₀-C₈)alkyl-C(O)O— R^5 or the R^6 groups can join to form a heterocycloalkyl group;

n is 0 or 1; and

* represents a chiral-carbon center.

15 In specific compounds of formula II, when n is 0 then R^1 is (C₃-C₇)cycloalkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, benzyl, aryl, (C₀-C₄)alkyl-(C₁-C₆)heterocycloalkyl, (C₀-C₄)alkyl-(C₂-C₅)heteroaryl, C(O)R³, C(O)OR⁴, (C₁-C₈)alkyl-N(R⁶)₂, (C₁-C₈)alkyl-OR⁵, (C₁-C₈)alkyl-C(O)OR⁵, C(S)NHR³, or (C₁-C₈)alkyl-O(CO)R⁵;


R^2 is H or (C₁-C₈)alkyl; and

R^3 is (C₁-C₈)alkyl, (C₃-C₇)cycloalkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, benzyl, aryl, (C₀-C₄)alkyl-(C₁-C₆)heterocycloalkyl, (C₀-C₄)alkyl-(C₂-C₅)heteroaryl, (C₅-C₈)alkyl-N(R⁶)₂; (C₀-C₈)alkyl-NH—C(O)O— R^5 ; (C₁-C₈)alkyl-OR⁵, (C₁-C₈)alkyl-C(O)OR⁵, (C₁-C₈)alkyl-O(CO)R⁵, or C(O)OR⁵; and the other variables have the same definitions.


In other specific compounds of formula II, R^2 is H or (C₁-C₄)alkyl.

30 In other specific compounds of formula II, R^1 is (C₁-C₈)alkyl or benzyl.

In other specific compounds of formula II, R^1 is H, (C₁-C₈)alkyl, benzyl, CH₂OCH₃, CH₂CH₂OCH₃, or



40 In another embodiment of the compounds of formula II, R^1 is



wherein Q is O or S, and each occurrence of R^7 is independently H, (C₁-C₈)alkyl, benzyl, CH₂OCH₃, or CH₂CH₂OCH₃.

55 In other specific compounds of formula II, R^1 is C(O)R³.
 In other specific compounds of formula II, R^3 is (C₀-C₄)alkyl-(C₂-C₅)heteroaryl, (C₁-C₈)alkyl, aryl, or (C₀-C₄)alkyl-OR⁵.

In other specific compounds of formula II, heteroaryl is pyridyl, furyl, or thienyl.

In other specific compounds of formula II, R^1 is C(O)OR⁴.


In other specific compounds of formula II, the H of C(O)NHC(O) can be replaced with (C₁-C₄)alkyl, aryl, or benzyl.

65 Still other specific immunomodulatory compounds of the invention belong to a class of isoindole-imides disclosed in U.S. patent application Ser. No. 09/781,179, International Publication No. WO 98/54170, and U.S. Pat. No. 6,395,754,

US 9,101,621 B2

9

each of which are incorporated herein by reference. Representative compounds are of formula III:



and pharmaceutically acceptable salts, hydrates, solvates, clathrates, enantiomers, diastereomers, racemates, and mixtures of stereoisomers thereof, wherein:

one of X and Y is C=O and the other is CH₂ or C=O;

R is H or CH₂OCOR';

(i) each of R¹, R², R³, or R⁴, independently of the others, is halo, alkyl of 1 to 4 carbon atoms, or alkoxy of 1 to 4 carbon atoms or (ii) one of R¹, R², R³, or R⁴ is nitro or —NHR⁵ and the remaining of R¹, R², R³, or R⁴ are hydrogen;

R⁵ is hydrogen or alkyl of 1 to 8 carbons

R⁶ hydrogen, alkyl of 1 to 8 carbon atoms, benzo, chloro, or fluoro;

R' is R⁷—CHR¹⁰—N(R⁸R⁹);


R⁷ is m-phenylene or p-phenylene or —(C_nH_{2n})— in which n has a value of 0 to 4;

each of R⁸ and R⁹ taken independently of the other is hydrogen or alkyl of 1 to 8 carbon atoms, or R⁸ and R⁹ taken together are tetramethylene, pentamethylene, hexamethylene, or —CH₂CH₂[X]X₁CH₂CH₂— in which [X]X₁ is —O—, —S—, or —NH—;


R¹⁰ is hydrogen, alkyl of 1 to 8 carbon atoms, or phenyl; and

* represents a chiral-carbon center.

The most preferred immunomodulatory compounds of the invention are 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione and 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. The compounds can be obtained via standard, synthetic methods (see e.g., U.S. Pat. No. 5,635, 517, incorporated herein by reference). The compounds are available from Celgene Corporation, Warren, N.J. 4-(Amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (ACTIMID™) has the following chemical structure:



The compound 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (REVIMID™) has the following chemical structure:



Compounds of the invention can either be commercially purchased or prepared according to the methods described in the patents or patent publications disclosed herein. Further, optically pure compounds can be asymmetrically synthesized

10

or resolved using known resolving agents or chiral columns as well as other standard synthetic organic chemistry techniques.

III

As used herein and unless otherwise indicated, the term “pharmaceutically acceptable salt” encompasses non-toxic acid and base addition salts of the compound to which the term refers. Acceptable non-toxic acid addition salts include those derived from organic and inorganic acids or bases known in the art, which include, for example, hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid, methanesulphonic acid, acetic acid, tartaric acid, lactic acid, succinic acid, citric acid, malic acid, maleic acid, sorbic acid, aconitic acid, salicylic acid, phthalic acid, embolic acid, enanthic acid, and the like.

Compounds that are acidic in nature are capable of forming salts with various pharmaceutically acceptable bases. The bases that can be used to prepare pharmaceutically acceptable base addition salts of such acidic compounds are those that form non-toxic base addition salts, i.e., salts containing pharmacologically acceptable cations such as, but not limited to, alkali metal or alkaline earth metal salts and the calcium, magnesium, sodium or potassium salts in particular. Suitable organic bases include, but are not limited to, N,N-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumine (N-methylglucamine), lysine, and procaine.

As used herein and unless otherwise indicated, the term “prodrug” means a derivative of a compound that can hydrolyze, oxidize, or otherwise react under biological conditions (in vitro or in vivo) to provide the compound. Examples of prodrugs include, but are not limited to, derivatives of immunomodulatory compounds of the invention that comprise biohydrolyzable moieties such as biohydrolyzable amides, biohydrolyzable esters, biohydrolyzable carbamates, biohydrolyzable carbonates, biohydrolyzable ureides, and biohydrolyzable phosphate analogues. Other examples of prodrugs include derivatives of immunomodulatory compounds of the invention that comprise —NO, —NO₂, —ONO, or —ONO₂ moieties. Prodrugs can typically be prepared using well-known methods, such as those described in 1 *Burger's Medicinal Chemistry and Drug Discovery*, 172-178, 949-982 (Manfred E. Wolff ed., 5th ed. 1995), and *Design of Prodrugs* (H. Bundgaard ed., Elsevier, New York 1985).

As used herein and unless otherwise indicated, the terms “biohydrolyzable amide,” “biohydrolyzable ester,” “biohydrolyzable carbamate,” “biohydrolyzable carbonate,” “biohydrolyzable ureide,” “biohydrolyzable phosphate” mean an amide, ester, carbamate, carbonate, ureide, or phosphate, respectively, of a compound that either: 1) does not interfere with the biological activity of the compound but can confer upon that compound advantageous properties in vivo, such as uptake, duration of action, or onset of action; or 2) is biologically inactive but is converted in vivo to the biologically active compound.

Examples of biohydrolyzable esters include, but are not limited to, lower alkyl esters, lower acyloxyalkyl esters (such as acetoxyethyl, acetoxyethyl, aminocarbonyloxyethyl, pivaloyloxyethyl, and pivaloyloxyethyl esters), lactonyl esters (such as phthalidyl and thiophthalidyl esters), lower alkoxyacyloxyalkyl esters (such as methoxycarbonyloxyethyl, ethoxycarbonyloxyethyl and isopropoxycarbonyloxyethyl esters), alkoxyalkyl esters, choline esters, and acylamino alkyl esters (such as acetamidomethyl esters). Examples of biohydrolyzable amides include, but are not limited to, lower alkyl amides, α-amino acid amides, alkoxyacyl amides, and alkylaminoalkylcarbonyl amides. Examples of biohydrolyzable carbamates

include, but are not limited to, lower alkylamines, substituted ethylenediamines, amino acids, hydroxyalkylamines, heterocyclic and heteroaromatic amines, and polyether amines.

Various immunomodulatory compounds of the invention contain one or more chiral centers, and can exist as racemic mixtures of enantiomers or mixtures of diastereomers. This invention encompasses the use of stereomerically pure forms of such compounds, as well as the use of mixtures of those forms. For example, mixtures comprising equal or unequal amounts of the enantiomers of a particular immunomodulatory compounds of the invention may be used in methods and compositions of the invention. These isomers may be asymmetrically synthesized or resolved using standard techniques such as chiral columns or chiral resolving agents. See, e.g., Jacques, J., et al., *Enantiomers, Racemates and Resolutions* (Wiley-Interscience, New York, 1981); Wilen, S. H., et al., *Tetrahedron* 33:2725 (1977); Eliel, E. L., *Stereochemistry of Carbon Compounds* (McGraw-Hill, NY, 1962); and Wilen, S. H., *Tables of Resolving Agents and Optical Resolutions* p. 268 (E. L. Eliel, Ed., Univ. of Notre Dame Press, Notre Dame, Ind., 1972).

As used herein and unless otherwise indicated, the term "stereomerically pure" means a composition that comprises one stereoisomer of a compound and is substantially free of other stereoisomers of that compound. For example, a stereomerically pure composition of a compound having one chiral center will be substantially free of the opposite enantiomer of the compound. A stereomerically pure composition of a compound having two chiral centers will be substantially free of other diastereomers of the compound. A typical stereomerically pure compound comprises greater than about 80% by weight of one stereoisomer of the compound and less than about 20% by weight of other stereoisomers of the compound, more preferably greater than about 90% by weight of one stereoisomer of the compound and less than about 10% by weight of the other stereoisomers of the compound, even more preferably greater than about 95% by weight of one stereoisomer of the compound and less than about 5% by weight of the other stereoisomers of the compound, and most preferably greater than about 97% by weight of one stereoisomer of the compound and less than about 3% by weight of the other stereoisomers of the compound. As used herein and unless otherwise indicated, the term "stereomerically enriched" means a composition that comprises greater than about 60% by weight of one stereoisomer of a compound, preferably greater than about 70% by weight, more preferably greater than about 80% by weight of one stereoisomer of a compound. As used herein and unless otherwise indicated, the term "enantiomerically pure" means a stereomerically pure composition of a compound having one chiral center. Similarly, the term "stereomerically enriched" means a stereomerically enriched composition of a compound having one chiral center.

It should be noted that if there is a discrepancy between a depicted structure and a name given that structure, the depicted structure is to be accorded more weight. In addition, if the stereochemistry of a structure or a portion of a structure is not indicated with, for example, bold or dashed lines, the structure or portion of the structure is to be interpreted as encompassing all stereoisomers of it.

5.2 Second Active Agents

Immunomodulatory compounds can be combined with other pharmacologically active compounds ("second active agents") in methods and compositions of the invention. It is believed that certain combinations work synergistically in the

treatment of particular types of cancer and certain diseases and conditions associated with, or characterized by, undesired angiogenesis. Immunomodulatory compounds can also work to alleviate adverse effects associated with certain second active agents, and some second active agents can be used to alleviate adverse effects associated with immunomodulatory compounds.

One or more second active ingredients or agents can be used in the methods and compositions of the invention together with an immunomodulatory compound. Second active agents can be large molecules (e.g., proteins) or small molecules (e.g., synthetic inorganic, organometallic, or organic molecules).

Examples of large molecule active agents include, but are not limited to, hematopoietic growth factors, cytokines, and monoclonal and polyclonal antibodies. Typical large molecule active agents are biological molecules, such as naturally occurring or artificially made proteins. Proteins that are particularly useful in this invention include proteins that stimulate the survival and/or proliferation of hematopoietic precursor cells and immunologically active poietic cells in vitro or in vivo. Others stimulate the division and differentiation of committed erythroid progenitors in cells in vitro or in vivo. Particular proteins include, but are not limited to: interleukins, such as IL-2 (including recombinant IL-II ("rIL2")) and canarypox TL-2), IL-10, IL-12, and IL-18; interferons, such as interferon alfa-2a, interferon alfa-2b, interferon alfa-n1, interferon alfa-n3, interferon beta-I a, and interferon gamma-I b; GM-CSF and GM-CSF; and EPO.

Particular proteins that can be used in the methods and compositions of the invention include, but are not limited to: filgrastim, which is sold in the United States under the trade name Neupogen® (Amgen, Thousand Oaks, Calif.); sargramostim, which is sold in the United States under the trade name Leukine® (Immunex, Seattle, Wash.); and recombinant EPO, which is sold in the United States under the trade name Epogen® (Amgen, Thousand Oaks, Calif.).

Recombinant and mutated forms of GM-CSF can be prepared as described in U.S. Pat. Nos. 5,391,485; 5,393,870; and 5,229,496; all of which are incorporated herein by reference. Recombinant and mutated forms of G-CSF can be prepared as described in U.S. Pat. Nos. 4,810,643; 4,999,291; 5,528,823; and 5,580,755; all of which are incorporated herein by reference.

This invention encompasses the use of native, naturally occurring, and recombinant proteins. The invention further encompasses mutants and derivatives (e.g., modified forms) of naturally occurring proteins that exhibit, in vivo, at least some of the pharmacological activity of the proteins upon which they are based. Examples of mutants include, but are not limited to, proteins that have one or more amino acid residues that differ from the corresponding residues in the naturally occurring forms of the proteins. Also encompassed by the term "mutants" are proteins that lack carbohydrate moieties normally present in their naturally occurring forms (e.g., nonglycosylated forms). Examples of derivatives include, but are not limited to, pegylated derivatives and fusion proteins, such as proteins formed by fusing IgG1 or IgG3 to the protein or active portion of the protein of interest. See, e.g., Penichet, M. L. and Morrison, S. L., *J. Immunol. Methods* 248:91-101 (2001).

Antibodies that can be used in combination with compounds of the invention include monoclonal and polyclonal antibodies. Examples of antibodies include, but are not limited to, trastuzumab (Herceptin®), rituximab (Rituxan®), bevacizumab (Avastin™), pertuzumab (Omnitarg™), tositumomab (Bexxar®), edrecolomab (Panorex®), and G250.

US 9,101,621 B2

13

Compounds of the invention can also be combined with, or used in combination with, anti-TNF- α antibodies.

Large molecule active agents may be administered in the form of anti-cancer vaccines. For example, vaccines that secrete, or cause the secretion of, cytokines such as IL-2, G-CSF, and GM-CSF can be used in the methods, pharmaceutical compositions, and kits of the invention. See, e.g., Emens, L. A., et al., *Curr. Opin. Mol. Ther.* 3(1):77-84 (2001).

In one embodiment of the invention, the large molecule active agent reduces, eliminates, or prevents an adverse effect associated with the administration of an immunomodulatory compound. Depending on the particular immunomodulatory compound and the disease or disorder being treated, adverse effects can include, but are not limited to, drowsiness and somnolence, dizziness and orthostatic hypotension, neutropenia, infections that result from neutropenia, increased HIV-viral load, bradycardia, Stevens-Johnson Syndrome and toxic epidermal necrolysis, and seizures (e.g., grand mal convulsions). A specific adverse effect is neutropenia.

Second active agents that are small molecules can also be used to alleviate adverse effects associated with the administration of an immunomodulatory compound. However, like some large molecules, many are believed to be capable of providing a synergistic effect when administered with (e.g., before, after or simultaneously) an immunomodulatory compound. Examples of small molecule second active agents include, but are not limited to, anti-cancer agents, antibiotics, immunosuppressive agents, and steroids.

Examples of anti-cancer agents include, but are not limited to: acivicin; aclarubicin; acodazole hydrochloride; acronine; adozelesin; aldesleukin; altretamine; ambomycin; ametranone acetate; amsacrine; anastrozole; anthramycin; asparaginase; asperlin; azacitidine; azetepa; azotomycin; batimastat; benzodepa; bicalutamide; bisantrene hydrochloride; bisnafide dimesylate; bizelesin; bleomycin sulfate; brequinar sodium; bropirimine; busulfan; cactinomycin; calusterone; caracemide; carbetimer; carboplatin; carmustine; carubicin hydrochloride; carzelesin; cedefingol; celecoxib (COX-2 inhibitor); chlorambucil; cirolemycin; cisplatin; cladribine; crisnatol mesylate; cyclophosphamide; cytarabine; dacarbazine; dactinomycin; daunorubicin hydrochloride; decitabine; dexormaplatin; dezaguanine; dezaguanine mesylate; diaziquone; docetaxel; doxorubicin; doxorubicin hydrochloride; droloxifene; droloxifene citrate; dromostanolone propionate; diazomycin; edatrexate; eflornithine hydrochloride; elsamitucin; enloplatin; enpromate; epiropropidine; epirubicin hydrochloride; erbulozole; esorubicin hydrochloride; estramustine; estramustine phosphate sodium; etanidazole; etoposide; etoposide phosphate; etoprine; fadrozole hydrochloride; fazarabine; fenretinide; floxuridine; fludarabine phosphate; fluorouracil; fluorocitabine; fosquidone; fostriecin sodium; gemcitabine; gemcitabine hydrochloride; hydroxyurea; idarubicin hydrochloride; ifosfamide; ilmofofosine; iroplatin; irinotecan; irinotecan hydrochloride; lanreotide acetate; letrozole; leuprolide acetate; liarozole hydrochloride; lometrexol sodium; lomustine; losoxantrone hydrochloride; masoprocol; maytansine; mechlorethamine hydrochloride; megestrol acetate; melengestrol acetate; melphalan; menogaril; mercaptopurine; methotrexate; methotrexate sodium; metoprine; meturedepa; mitidomide; mitocarcin; mitocromin; mitogillin; mitomycin; mitomycin; mitosper; mitotane; mitoxantrone hydrochloride; mycophenolic acid; nocodazole; nogalamycin; ormaplatin; oxisuran; paclitaxel; pegaspargase; peliomycin; pentamustine; peplomycin sulfate; perfosfamide; pipobroman; pipsulfan; piroxantrone hydrochloride;

14

plicamycin; plomestane; porfimer sodium; porfiromycin; prednimustine; procarbazine hydrochloride; puromycin; puromycin hydrochloride; pyrazofurin; riboprine; safingol; safingol hydrochloride; semustine; simtrazene; sparfosate sodium; sparsomycin; spirogermanium hydrochloride; spiro-mustine; spiroplatin; streptonigrin; streptozocin; sulofenur; talisomycin; tecogalan sodium; taxotere; tegafur; teloxantrone hydrochloride; temoporfin; teniposide; teroxirone; testolactone; thiamiprine; thioguanine; thiotepa; tiazofurin; tirapazamine; toremifene citrate; trestolone acetate; tricitribine phosphate; trimetrexate; trimetrexate glucuronate; triptorelin; tubulozole hydrochloride; uracil mustard; uredepa; vaporeotide; verteporfin; vinblastine sulfate; vincristine sulfate; vindesine; vindesine sulfate; vinepidine sulfate; vinyglycinate sulfate; vinleurosine sulfate; vinorelbine tartrate; vinrosidine sulfate; vinzolidine sulfate; vorozole; zeniplatin; zinostatin; and zorubicin hydrochloride.

Other anti-cancer drugs include, but are not limited to: 20-epi-1,25 dihydroxyvitamin D3; 5-ethynyluracil; abiraterone; aclarubicin; acylfulvene; adecypenol; adozelesin; aldesleukin; ALL-TK antagonists; altretamine; ambamustine; amidox; amifostine; aminolevulinic acid; amrubicin; amsacrine; anagrelide; anastrozole; andrographolide; angiogenesis inhibitors; antagonist D; antagonist G; antarelix; antidorsalizing morphogenetic protein-1; antiandrogen, prostatic carcinoma; antiestrogen; antineoplaston; antisense oligonucleotides; aphidicolin glycinate; apoptosis gene modulators; apoptosis regulators; apurinic acid; ara-CDP-DL-PTBA; arginine deaminase; asulacrine; atamestane; atrimustine; axinastatin 1; axinastatin 2; axinastatin 3; azasctron; azatoxin; azatyrosine; baccatin ITT derivatives; balanol; batimastat; BCR/ABL antagonists; benzochlorins; benzoylstauroporine; beta lactam derivatives; beta-alethine; betaclamycin B; betulinic acid; bFGF inhibitor; bicalutamide; bisantrene; bisaziridinylspermine; bisnafide; bistratene A; bizelesin; breflate; bropirimine; budotitane; buthionine sulfoximine; calcipotriol; calphostin C; camptothecin derivatives; capecitabine; carboxamide-amino-triazole; carboxamidotriazole; CaRest M3; CARN 700; cartilage derived inhibitor; carzelesin; casein kinase inhibitors (ICOS); castanospermine; cecropin B; cetrorelix; chlorins; chloroquinoline sulfonamide; cicaprost; cis-porphyrin; cladribine; clomifene analogues; clotrimazole; collismycin A; collismycin B; combretastatin A4; combretastatin analogue; conagein; crambescidin 816; crisnatol; cryptophycin 8; cryptophycin A derivatives; curacin A; cyclopentantraquinones; cycloplatin; cypemycin; cytarabine ocfosfate; cytolytic factor; cytosatin; dacliximab; decitabine; dehydrididemnin B; deslorelin; dexamethasone; dexifosfamide; dexrazoxane; dexverapamil; diaziquone; didemnin B; didox; diethylnorspermine; dihydro-5-azacytidine; dihydrotaxol, 9-; dioxamycin; diphenyl spiro-mustine; docetaxel; docosanol; dolasetron; doxifluridine; doxorubicin; droloxifene; dronabinol; duocarmycin SA; ebselen; ecomustine; edelfosine; edrecolomab; eflornithine; elemene; emitofur; epirubicin; epristeride; estramustine analogue; estrogen agonists; estrogen antagonists; etanidazole; etoposide phosphate; exemestane; fadrozole; fazarabine; fenretinide; filgrastim; finasteride; flavopiridol; flezelastine; fluasterone; fludarabine; fluorodaunorubicin hydrochloride; forfenimex; formestane; fostriecin; fotemustine; gadolinium texaphyrin; gallium nitrate; galocitabine; ganirelix; gelatinase inhibitors; gemcitabine; glutathione inhibitors; hepsulfam; heregulin; hexamethylene bisacetamide; hypericin; ibandron acid; idarubicin; idoxifene; idramantone; ilmofofosine; ilomastat; imatinib (e.g., Gleevec®), imiquimod; immunostimulant peptides; insulin-like growth factor-1 receptor inhibitor; interferon

agonists; interferons; interleukins; iobenguane; iododoxorubicin; ipomeanol, 4-; iroplact; irsogladine; isobengazole; isohomohalicondrin B; itasetron; jasplakinolide; kahalalide F; lamellarin-N triacetate; lanreotide; leinamycin; lenograstim; lentinan sulfate; leptolstatin; letrozole; leukemia inhibiting factor; leukocyte alpha interferon; leuprolide+estrogen+progesterone; leuprorelin; levamisole; liarozole; linear polyamine analogue; lipophilic disaccharide peptide; lipophilic platinum compounds; lissoclinamide 7; lobaplatin; lombricine; lometrexol; lomidamine; losoxantrone; lxoribine; lurtotecan; lutetium texaphyrin; lysofylline; lytic peptides; maitansine; mannostatin A; marimastat; masoprocol; maspin; matrilysin inhibitors; matrix metalloproteinase inhibitors; menogaryl; merbarone; meterelin; methioninase; metoclopramide; MIF inhibitor; mifepristone; miltefosine; mirimostim; mitoguanine; mitolactol; mitomycin analogues; mitonafide; mitotoxin fibroblast growth factor-saporin; mitoxantrone; mofarotene; molgramostim; Erbitux, human chorionic gonadotrophin; monophosphoryl lipid A+myobacterium cell wall sk; mopidamol; mustard anticancer agent; mycaperoxide B; mycobacterial cell wall extract; myriaporone; N-acetyldinaline; N-substituted benzamides; nafarelin; nagrestip; naloxone+pentazocine; napavin; naphterpin; narograstim; nedaplatin; nemorubicin; neridronic acid; nilutamide; nisamycin; nitric oxide modulators; nitroxide antioxidant; nitrullyn; oblimersen (Genasense®); O⁶-benzylguanine; octreotide; okicenone; oligonucleotides; onapristone; ondansetron; ondansetron; oracin; oral cytokine inducer; ormaplatin; osaterone; oxaliplatin; oxaunomycin; paclitaxel; paclitaxel analogues; paclitaxel derivatives; palauamine; palmitoylrhizoxin; pamidronic acid; panaxytriol; panomifene; parabactin; pazelliptine; pegaspargase; peldesine; pentosan polysulfate sodium; pentostatin; pentrozole; perflubron; perfosfamide; perillyl alcohol; phenazinomycin; phenylacetate; phosphatase inhibitors; picibanil; pilocarpine hydrochloride; pirarubicin; piritrexim; placetin A; placetin B; plasminogen activator inhibitor; platinum complex; platinum compounds; platinum-triamine complex; porfimer sodium; porfiromycin; prednisone; propyl bis-acridone; prostaglandin J2; proteasome inhibitors; protein A-based immune modulator; protein kinase C inhibitor; protein kinase C inhibitors, microalgal; protein tyrosine phosphatase inhibitors; purine nucleoside phosphorylase inhibitors; purpurins; pyrazoloacridine; pyridoxylated hemoglobin polyoxyethylene conjugate; raf antagonists; raltitrexed; ramosetron; ras farnesyl protein transferase inhibitors; ras inhibitors; ras-GAP inhibitor; retelliptine demethylated; rhenium Re 186 etidronate; rhizoxin; ribozymes; RII retinamide; rohitukine; romurtide; roquinimex; rubiginone B1; ruboxyl; safingol; saintopin; SarCNU; sarcophytol A; sargramostim; Sdi 1 mimetics; semustine; senescence derived inhibitor 1; sense oligonucleotides; signal transduction inhibitors; sizofiran; sobuzoxane; sodium borocaptate; sodium phenylacetate; solverol; somatomedin binding protein; sonermin; sparfosic acid; spicamycin D; spiromustine; splenopentin; spongistatin 1; squalamine; stipiamide; stromelysin inhibitors; sulfinosine; superactive vasoactive intestinal peptide antagonist; suradista; suramin; swainsonine; tallimustine; tamoxifen methiodide; tauromustine; tazarotene; tecogalan sodium; tegafur; tellurapyrylium; telomerase inhibitors; temoporfin; teniposide; tetrachlorodecaoxide; tetrazomine; thaliblastine; thiocoraline; thrombopoietin; thrombopoietin mimetic; thymalfasin; thymopoietin receptor agonist; thymotrinan; thyroid stimulating hormone; tin ethyl etiopurpurin; tirapazamine; titanocnc bichloride; topsentin; toremifene; translation inhibitors; tretinoin; triacetyluridine; triciribine; trimetrexate; triptorelin; tropisetron; turosteride; tyrosine

kinase inhibitors; tyrphostins; UBC inhibitors; ubenimex; urogenital sinus-derived growth inhibitory factor; urokinase receptor antagonists; vapreotide; variolin B; velaresol; veramine; verdins; verteporfin; vinorelbine; vinxaltine; vitaxin; vorozole; zanoterone; zeniplatin; zilascorb; and zinostatin stimalamer.

Specific second active agents include, but are not limited to, oblimersen (Genasense®), remicade, docetaxel, celecoxib, melphalan, dexamethasone (Decadron®), steroids, gemcitabine, cisplatin, temozolomide, etoposide, cyclophosphamide, temodar, carboplatin, procarbazine, gliadel, tamoxifen, topotecan, methotrexate, Arisa®, taxol, taxotere, fluorouracil, leucovorin, irinotecan, xeloda, CPT-11, interferon alpha, pegylated interferon alpha (e.g., PEG INTRON-A), capecitabine, cisplatin, thiotepa, fludarabine, carboplatin, liposomal daunorubicin, cytarabine, doxetaxol, paclitaxel, vinblastine, IL-2, GM-CSF, dacarbazine, vinorelbine, zoledronic acid, palmitronate, biaxin, busulphan, prednisone, bisphosphonate, arsenic trioxide, vincristine, doxorubicin (Doxil®), paclitaxel, ganciclovir, adriamycin, estramustine sodium phosphate (Emcyt®), sulindac, and etoposide.

5.3 Methods of Treatments and Prevention

Methods of this invention encompass methods of treating, preventing and/or managing various types of cancer and diseases and disorders associated with, or characterized by, undesired angiogenesis. As used herein, unless otherwise specified, the term “treating” refers to the administration of a compound of the invention or other additional active agent after the onset of symptoms of the particular disease or disorder. As used herein, unless otherwise specified, the term “preventing” refers to the administration prior to the onset of symptoms, particularly to patients at risk of cancer, and other diseases and disorders associated with, or characterized by, undesired angiogenesis. The term “prevention” includes the inhibition of a symptom of the particular disease or disorder. Patients with familial history of cancer and diseases and disorders associated with, or characterized by, undesired angiogenesis are preferred candidates for preventive regimens. As used herein and unless otherwise indicated, the term “managing” encompasses preventing the recurrence of the particular disease or disorder in a patient who had suffered from it, and/or lengthening the time a patient who had suffered from the disease or disorder remains in remission.

As used herein, the term “cancer” includes, but is not limited to, solid tumors and blood born tumors. The term “cancer” refers to disease of skin tissues, organs, blood, and vessels, including, but not limited to, cancers of the bladder, bone or blood, brain, breast, cervix, chest, colon, endometrium, esophagus, eye, head, kidney, liver, lymph nodes, lung, mouth, neck, ovaries, pancreas, prostate, rectum, stomach, testis, throat, and uterus. Specific cancers include, but are not limited to, advanced malignancy, amyloidosis, neuroblastoma, meningioma, hemangiopericytoma, multiple brain metastase, glioblastoma multiforms, glioblastoma, brain stem glioma, poor prognosis malignant brain tumor, malignant glioma, recurrent malignant glioma, anaplastic astrocytoma, anaplastic oligodendroglioma, neuroendocrine tumor, rectal adenocarcinoma, Dukes C & D colorectal cancer, unresectable colorectal carcinoma, metastatic hepatocellular carcinoma, Kaposi’s sarcoma, karotype acute myeloblastic leukemia, Hodgkin’s lymphoma, non-Hodgkin’s lymphoma, cutaneous T-Cell lymphoma, cutaneous B-Cell lymphoma, diffuse large B-Cell lymphoma, low grade follicular lymphoma, malignant melanoma, malignant mesothelioma, malignant pleural effusion mesothelioma syndrome,

peritoneal carcinoma, papillary serous carcinoma, gynecologic sarcoma, soft tissue sarcoma, scleroderma, cutaneous vasculitis, Langerhans cell histiocytosis, leiomyosarcoma, fibrodysplasia ossificans progressive, hormone refractory prostate cancer, resected high-risk soft tissue sarcoma, unresectable hepatocellular carcinoma, Waldenstrom's macroglobulinemia, smoldering myeloma, indolent myeloma, fallopian tube cancer, androgen independent prostate cancer, androgen dependent stage IV non-metastatic prostate cancer, hormone-insensitive prostate cancer, chemotherapy-insensitive prostate cancer, papillary thyroid carcinoma, follicular thyroid carcinoma, medullary thyroid carcinoma, and leiomyoma. In a specific embodiment, the cancer is metastatic. In another embodiment, the cancer is refractory or resistance to chemotherapy or radiation; in particular, refractory to thalidomide.

As used herein to refer to diseases and conditions other than cancer, the terms "diseases or disorders associated with, or characterized by, undesired angiogenesis," "diseases or disorders associated with undesired angiogenesis," and "diseases or disorders characterized by undesired angiogenesis" refer to diseases, disorders and conditions that are caused, mediated or attended by undesired, unwanted or uncontrolled angiogenesis, including, but not limited to, inflammatory diseases, autoimmune diseases, genetic diseases, allergic diseases, bacterial diseases, ocular neovascular diseases, choroidal neovascular diseases, and retina neovascular diseases.

Examples of such diseases or disorders associated with undesired angiogenesis include, but are not limited to, diabetic retinopathy, retinopathy of prematurity, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, proliferative vitreoretinopathy, trachoma, myopia, optic pits, epidemic keratoconjunctivitis, atopic keratitis, superior limbic keratitis, pterygium keratitis sicca, sjogrens, acne rosacea, phlyctenulosis, syphilis, lipid degeneration, bacterial ulcer, fungal ulcer, Herpes simplex infection, Herpes zoster infection, protozoan infection, Kaposi sarcoma, Mooren ulcer, Terrien's marginal degeneration, mariginal keratolysis, rheumatoid arthritis, systemic lupus, polyarteritis, trauma, Wegeners sarcoidosis, Scleritis, Steven's Johnson disease, periphigoid radial keratotomy, sickle cell anemia, sarcoid, pseudoxanthoma elasticum, Pagets disease, vein occlusion, artery occlusion, carotid obstructive disease, chronic uveitis, chronic vitritis, Lyme's disease, Eales disease, Behcet's disease, retinitis, choroiditis, presumed ocular histoplasmosis, Bests disease, Stargarts disease, pars planitis, chronic retinal detachment, hyperviscosity syndromes, toxoplasmosis, rubeosis, sarcodiosis, sclerosis, soriatis, psoriasis, primary sclerosing cholangitis, proctitis, primary biliary srosis, idiopathic pulmonary fibrosis, and alcoholic hepatitis.

In specific embodiments of the invention, diseases or disorders associated with undesired angiogenesis do not include congestive heart failure, cardiomyopathy, pulmonary edema, endotoxin-mediated septic shock, acute viral myocarditis, cardiac allograft rejection, myocardial infarction, HIV, hepatitis, adult respiratory distress syndrome, bone-resorption disease, chronic obstructive pulmonary diseases, chronic pulmonary inflammatory disease, dermatitis, cystic fibrosis, septic shock, sepsis, endotoxic shock, hemodynamic shock, sepsis syndrome, post ischemic reperfusion injury, meningitis, psoriasis, fibrotic disease, cachexia, graft rejection, rheumatoid spondylitis, osteoporosis, Crohn's disease, ulcerative colitis, inflammatory-bowel disease, multiple sclerosis, systemic lupus erythrematosus, erythema nodosum leprosum in leprosy, radiation damage, asthma, hyperoxic alveolar injury, malaria, mycobacterial infection, and opportunistic infections resulting from HIV.

This invention encompasses methods of treating patients who have been previously treated for cancer or diseases or disorders associated with, or characterized by, undesired angiogenesis, but are non-responsive to standard therapies, as well as those who have not previously been treated. The invention also encompasses methods of treating patients regardless of patient's age, although some diseases or disorders are more common in certain age groups. The invention further encompasses methods of treating patients who have undergone surgery in an attempt to treat the disease or condition at issue, as well as those who have not. Because patients with cancer and diseases and disorders characterized by undesired angiogenesis have heterogenous clinical manifestations and varying clinical outcomes, the treatment given to a patient may vary, depending on his/her prognosis. The skilled clinician will be able to readily determine without undue experimentation specific secondary agents, types of surgery, and types of non-drug based standard therapy that can be effectively used to treat an individual patient with cancer and other diseases or disorders.

Methods encompassed by this invention comprise administering one or more immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, to a patient (e.g., a human) suffering, or likely to suffer, from cancer or a disease or disorder mediated by undesired angiogenesis.

In one embodiment of the invention, an immunomodulatory compound of the invention can be administered orally and in single or divided daily doses in an amount of from about 0.10 to about 150 mg/day. In a particular embodiment, 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™) may be administered in an amount of from about 0.1 to about 1 mg per day, or alternatively from about 0.1 to about 5 mg every other day. In a preferred embodiment, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) may be administered in an amount of from about 5 to 25 mg per day, or alternatively from about 10 to about 50 mg every other day.

In a specific embodiment, 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™) may be administered in an amount of about 1, 2, or 5 mg per day to patients with relapsed multiple myeloma. In a particular embodiment, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) may be administered initially in an amount of 5 mg/day and the dose can be escalated every week to 10, 20, 25, 30 and 50 mg/day. In a specific embodiment, Revimid™ can be administered in an amount of up to about 30 mg/day to patients with solid tumor. In a particular embodiment, Revimid™ can be administered in an amount of up to about 40 mg/day to patients with glioma.

5.3.1 Combination Therapy with a Second Active Agent

Specific methods of the invention comprise administering an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, in combination with one or more second active agents, and/or in combination with radiation therapy, blood transfusions, or surgery. Examples of immunomodulatory compounds of the invention are disclosed herein (see, e.g., section 5.1). Examples of second active agents are also disclosed herein (see, e.g., section 5.2).

Administration of the immunomodulatory compounds and the second active agents to a patient can occur simultaneously or sequentially by the same or different routes of administra-

tion. The suitability of a particular route of administration employed for a particular active agent will depend on the active agent itself (e.g., whether it can be administered orally without decomposing prior to entering the blood stream) and the disease being treated. A preferred route of administration

for an immunomodulatory compound of the invention is orally. Preferred routes of administration for the second active agents or ingredients of the invention are known to those of ordinary skill in the art. See, e.g., *Physicians' Desk Reference*, 1755-1760 (56th ed., 2002).

In one embodiment of the invention, the second active agent is administered intravenously or subcutaneously and once or twice daily in an amount of from about 1 to about 1000 mg, from about 5 to about 500 mg, from about 10 to about 350 mg, or from about 50 to about 200 mg. The specific amount of the second active agent will depend on the specific agent used, the type of disease being treated or managed, the severity and stage of disease, and the amount(s) of immunomodulatory compounds of the invention and any optional additional active agents concurrently administered to the patient. In a particular embodiment, the second active agent is oblimersen (Genasense®), GM-CSF, G-CSF, EPO, taxotere, irinotecan, dacarbazine, transretinoic acid, topotecan, pentoxifylline, ciprofloxacin, dexamethasone, vincristine, doxorubicin, COX-2 inhibitor, IL2, IL8, IL18, IFN, Ara-C, vinorelbine, or a combination thereof.

In a particular embodiment, GM-CSF, G-CSF or EPO is administered subcutaneously during about five days in a four or six week cycle in an amount of from about 1 to about 750 mg/m²/day, preferably in an amount of from about 25 to about 500 mg/m²/day, more preferably in an amount of from about 50 to about 250 mg/m²/day, and most preferably in an amount of from about 50 to about 200 mg/m²/day. In a certain embodiment, GM-CSF may be administered in an amount of from about 60 to about 500 mcg/m² intravenously over 2 hours, or from about 5 to about 12 mcg/m²/day subcutaneously. In a specific embodiment, G-CSF may be administered subcutaneously in an amount of about 1 mcg/kg/day initially and can be adjusted depending on rise of total granulocyte counts. The maintenance dose of G-CSF may be administered in an amount of about 300 (in smaller patients) or 480 mcg subcutaneously. In a certain embodiment, EPO may be administered subcutaneously in an amount of 10,000 Unit 3 times per week.

In another embodiment, Revimid™ in an amount of about 25 mg/d and dacarbazine in an amount of about from 200 to 1,000 mg/m²/d are administered to patients with metastatic malignant melanoma. In a specific embodiment, Revimid™ is administered in an amount of from about 5 to about 25 mg/d to patients with metastatic malignant melanoma whose disease has progressed on treatment with dacarbazine, IL-2 or IFN. In a specific embodiment, Revimid™ is administered to patients with relapsed or refractory multiple myeloma in an amount of about 15 mg/d twice a day or about 30 mg/d four times a day in a combination with dexamethasone.

In another embodiment, an immunomodulatory compound is administered with melphalan and dexamethasone to patients with amyloidosis. In a specific embodiment, an immunomodulatory compound of the invention and steroids can be administered to patients with amyloidosis.

In another embodiment, an immunomodulatory compound is administered with gemcitabine and cisplatin to patients with locally advanced or metastatic transitional cell bladder cancer.

In another embodiment, an immunomodulatory compound is administered in combination with a second active ingredient as follows: temozolomide to pediatric patients with

relapsed or progressive brain tumors or recurrent neuroblastoma; celecoxib, etoposide and cyclophosphamide for relapsed or progressive CNS cancer; temodar to patients with recurrent or progressive meningioma, malignant meningioma, hemangiopericytoma, multiple brain metastases, relapsed brain tumors, or newly diagnosed glioblastoma multiformis; irinotecan to patients with recurrent glioblastoma; carboplatin to pediatric patients with brain stem glioma; procarbazine to pediatric patients with progressive malignant gliomas; cyclophosphamide to patients with poor prognosis malignant brain tumors, newly diagnosed or recurrent glioblastoma multiformis; Gliadel® for high grade recurrent malignant gliomas; temozolomide and tamoxifen for anaplastic astrocytoma; or topotecan for gliomas, glioblastoma, anaplastic astrocytoma or anaplastic oligodendroglioma.

In another embodiment, an immunomodulatory compound is administered with methotrexate and cyclophosphamide to patients with metastatic breast cancer.

In another embodiment, an immunomodulatory compound is administered with temozolomide to patients with neuroendocrine tumors.

In another embodiment, an immunomodulatory compound is administered with gemcitabine to patients with recurrent or metastatic head or neck cancer. In another embodiment, an immunomodulatory compound is administered with gemcitabine to patients with pancreatic cancer.

In another embodiment, an immunomodulatory compound is administered to patients with colon cancer in combination with Arisa®, taxol and/or taxotere.

In another embodiment, an immunomodulatory compound is administered with capecitabine to patients with refractory colorectal cancer or patients who fail first line therapy or have poor performance in colon or rectal adenocarcinoma.

In another embodiment, an immunomodulatory compound is administered in combination with fluorouracil, leucovorin, and irinotecan to patients with Dukes C & D colorectal cancer or to patients who have been previously treated for metastatic colorectal cancer.

In another embodiment, an immunomodulatory compound is administered to patients with refractory colorectal cancer in combination with capecitabine, xeloda, and/or CPT-11.

In another embodiment, an immunomodulatory compound of the invention is administered with capecitabine and irinotecan to patients with refractory colorectal cancer or to patients with unresectable or metastatic colorectal carcinoma.

In another embodiment, an immunomodulatory compound is administered alone or in combination with interferon alpha or capecitabine to patients with unresectable or metastatic hepatocellular carcinoma; or with cisplatin and thiotepa to patients with primary or metastatic liver cancer.

In another embodiment, an immunomodulatory compound is administered in combination with pegylated interferon alpha to patients with Kaposi's sarcoma.

In another embodiment, an immunomodulatory compound is administered in combination with fludarabine, carboplatin, and/or topotecan to patients with refractory or relapsed or high-risk acute myelogenous leukemia.

In another embodiment, an immunomodulatory compound is administered in combination with liposomal daunorubicin, topotecan and/or cytarabine to patients with unfavorable karyotype acute myeloblastic leukemia.

In another embodiment, an immunomodulatory compound is administered in combination with gemcitabine and irinotecan to patients with non-small cell lung cancer. In one embodiment, an immunomodulatory compound is administered in combination with carboplatin and irinotecan to

US 9,101,621 B2

21

patients with non-small cell lung cancer. In one embodiment, an immunomodulatory compound is administered with doxetaxol to patients with non-small cell lung cancer who have been previously treated with carbo/VP 16 and radiotherapy.

In another embodiment, an immunomodulatory compound is administered in combination with carboplatin and/or taxotere, or in combination with carboplatin, paclitaxel and/or thoracic radiotherapy to patients with non-small cell lung cancer. In a specific embodiment, an immunomodulatory compound is administered in combination with taxotere to patients with stage IIIB or TV non-small cell lung cancer.

In another embodiment, an immunomodulatory compound of the invention is administered in combination with oblimersen (Genasense®) to patients with small cell lung cancer.

In another embodiment, an immunomodulatory compound is administered alone or in combination with a second active ingredient such as vinblastine or fludarabine to patients with various types of lymphoma, including, but not limited to, Hodgkin's lymphoma, non-Hodgkin's lymphoma, cutaneous T-Cell lymphoma, cutaneous B-Cell lymphoma, diffuse large B-Cell lymphoma or relapsed or refractory low grade follicular lymphoma.

In another embodiment, an immunomodulatory compound is administered in combination with taxotere, IL-2, IFN, GM-CSF, and/or dacarbazine to patients with various types or stages of melanoma.

In another embodiment, an immunomodulatory compound is administered alone or in combination with vinorelbine to patients with malignant mesothelioma, or stage IIIB non-small cell lung cancer with pleural implants or malignant pleural effusion mesothelioma syndrome.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of multiple myeloma in combination with dexamethasone, zoledronic acid, palmitronate, GM-CSF, biacin, vinblastine, melphalan, busulphan, cyclophosphamide, IFN, palmitronate, prednisone, bisphosphonate, celecoxib, arsenic trioxide, PEG INTRON-A, vincristine, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with relapsed or refractory multiple myeloma in combination with doxorubicin (Doxil®), vincristine and/or dexamethasone (Decadron®).

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of ovarian cancer such as peritoneal carcinoma, papillary serous carcinoma, refractory ovarian cancer or recurrent ovarian cancer, in combination with taxol, carboplatin, doxorubicin, gemcitabine, cisplatin, xeloda, paclitaxel, dexamethasone, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of prostate cancer, in combination with xeloda, 5 FU/LV, gemcitabine, irinotecan plus gemcitabine, cyclophosphamide, vincristine, dexamethasone, GM-CSF, celecoxib, taxotere, ganciclovir, paclitaxel, adriamycin, docetaxel, estramustine, Emcyt, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of renal cell cancer, in combination with capecitabine, IFN, tamoxifen, IL-2, GM-CSF, Celebrex®, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of gynecologic, uterus or soft tissue sarcoma cancer in combination with IFN, a COX-2 inhibitor such as Celebrex®, and/or sulindac.

22

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of solid tumors in combination with celebrex, etoposide, cyclophosphamide, docetaxel, apicitabine, IFN, tamoxifen, IL-2, GM-CSF, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with scleroderma or cutaneous vasculitis in combination with celebrex, etoposide, cyclophosphamide, docetaxel, apicitabine, IFN, tamoxifen, IL-2, GM-CSF, or a combination thereof.

This invention also encompasses a method of increasing the dosage of an anti-cancer drug or agent that can be safely and effectively administered to a patient, which comprises administering to a patient (e.g., a human) an immunomodulatory compound of the invention, or a pharmaceutically acceptable derivative, salt, solvate, clathrate, hydrate, or prodrug thereof. Patients that can benefit by this method are those likely to suffer from an adverse effect associated with anti-cancer drugs for treating a specific cancer of the skin, subcutaneous tissue, lymph nodes, brain, lung, liver, bone, intestine, colon, heart, pancreas, adrenal, kidney, prostate, breast, colorectal, or combinations thereof. The administration of an immunomodulatory compound of the invention alleviates or reduces adverse effects which are of such severity that it would otherwise limit the amount of anti-cancer drug.

In one embodiment, an immunomodulatory compound of the invention can be administered orally and daily in an amount of from about 0.1 to about 150 mg, and preferably from about 1 to about 50 mg, more preferably from about 2 to about 25 mg prior to, during, or after the occurrence of the adverse effect associated with the administration of an anti-cancer drug to a patient. In a particular embodiment, an immunomodulatory compound of the invention is administered in combination with specific agents such as heparin, aspirin, coumadin, or G-CSF to avoid adverse effects that are associated with anti-cancer drugs such as but not limited to neutropenia or thrombocytopenia.

In one embodiment, an immunomodulatory compound of the invention can be administered to patients with diseases and disorders associated with, or characterized by, undesired angiogenesis in combination with additional active ingredients including but not limited to anti-cancer drugs, anti-inflammatories, antihistamines, antibiotics, and steroids.

In another embodiment, this invention encompasses a method of treating, preventing and/or managing cancer, which comprises administering an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, in conjunction with (e.g. before, during, or after) conventional therapy including, but not limited to, surgery, immunotherapy, biological therapy, radiation therapy, or other non-drug based therapy presently used to treat, prevent or manage cancer. The combined use of the immunomodulatory compounds of the invention and conventional therapy may provide a unique treatment regimen that is unexpectedly effective in certain patients. Without being limited by theory, it is believed that immunomodulatory compounds of the invention may provide additive or synergistic effects when given concurrently with conventional therapy.

As discussed elsewhere herein, the invention encompasses a method of reducing, treating and/or preventing adverse or undesired effects associated with conventional therapy including, but not limited to, surgery, chemotherapy, radiation therapy, hormonal therapy, biological therapy and immunotherapy. One or more immunomodulatory compounds of the invention and other active ingredient can be administered

to a patient prior to, during, or after the occurrence of the adverse effect associated with conventional therapy.

In one embodiment, an immunomodulatory compound of the invention can be administered in an amount of from about 0.1 to about 150 mg, and preferably from about 1 to about 25 mg, more preferably from about 2 to about 10 mg orally and daily alone, or in combination with a second active agent disclosed herein (see, e.g., section 5.2), prior to, during, or after the use of conventional therapy.

In a specific embodiment of this method, an immunomodulatory compound of the invention and doxorubicin are administered to patients with non-small cell lung cancer who were previously treated with carboplatin and radiotherapy.

5.3.2 Use with Transplantation Therapy

Compounds of the invention can be used to reduce the risk of Graft Versus Host Disease (GVHD). Therefore, the invention encompasses a method of treating, preventing and/or managing cancer, which comprises administering the immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, in conjunction with transplantation therapy.

As those of ordinary skill in the art are aware, the treatment of cancer is often based on the stages and mechanism of the disease. For example, as inevitable leukemic transformation develops in certain stages of cancer, transplantation of peripheral blood stem cells, hematopoietic stem cell preparation or bone marrow may be necessary. The combined use of the immunomodulatory compound of the invention and transplantation therapy provides a unique and unexpected synergism. In particular, an immunomodulatory compound of the invention exhibits immunomodulatory activity that may provide additive or synergistic effects when given concurrently with transplantation therapy in patients with cancer.

An immunomodulatory compound of the invention can work in combination with transplantation therapy reducing complications associated with the invasive procedure of transplantation and risk of GVHD. This invention encompasses a method of treating, preventing and/or managing cancer which comprises administering to a patient (e.g., a human) an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, before, during, or after the transplantation of umbilical cord blood, placental blood, peripheral blood stem cell, hematopoietic stem cell preparation or bone marrow. Examples of stem cells suitable for use in the methods of the invention are disclosed in U.S. provisional patent application No. 60/372,348, filed Apr. 12, 2002 by R. Hariri et al., the entirety of which is incorporated herein by reference.

In one embodiment of this method, an immunomodulatory compound of the invention is administered to patients with multiple myeloma before, during, or after the transplantation of autologous peripheral blood progenitor cell.

In another embodiment, an immunomodulatory compound is administered to patients with relapsing multiple myeloma after the stem cell transplantation.

In another embodiment, an immunomodulatory compound and prednisone are administered as maintenance therapy to patients with multiple myeloma following the transplantation of autologous stem cell.

In another embodiment, an immunomodulatory compound and dexamethasone are administered as salvage therapy for low risk post transplantation to patients with multiple myeloma.

In another embodiment, an immunomodulatory compound and dexamethasone are administered as maintenance therapy to patients with multiple myeloma following the transplantation of autologous bone marrow.

In another embodiment, an immunomodulatory compound is administered following the administration of high dose of melphalan and the transplantation of autologous stem cell to patients with chemotherapy responsive multiple myeloma.

In another embodiment, an immunomodulatory compound and PEG INTRO-A are administered as maintenance therapy to patients with multiple myeloma following the transplantation of autologous CD34-selected peripheral stem cell.

In another embodiment, an immunomodulatory compound is administered with post transplant consolidation chemotherapy to patients with newly diagnosed multiple myeloma to evaluate anti-angiogenesis.

In another embodiment, an immunomodulatory compound and dexamethasone are administered as maintenance therapy after DCEP consolidation, following the treatment with high dose of melphalan and the transplantation of peripheral blood stem cell to 65 years of age or older patients with multiple myeloma.

5.3.3 Cycling Therapy

In certain embodiments, the prophylactic or therapeutic agents of the invention are cyclically administered to a patient. Cycling therapy involves the administration of an active agent for a period of time, followed by a rest for a period of time, and repeating this sequential administration. Cycling therapy can reduce the development of resistance to one or more of the therapies, avoid or reduce the side effects of one of the therapies, and/or improves the efficacy of the treatment.

Consequently, in one specific embodiment of the invention, an immunomodulatory compound of the invention is administered daily in a single or divided doses in a four to six week cycle with a rest period of about a week or two weeks. The invention further allows the frequency, number, and length of dosing cycles to be increased. Thus, another specific embodiment of the invention encompasses the administration of an immunomodulatory compound of the invention for more cycles than are typical when it is administered alone. In yet another specific embodiment of the invention, an immunomodulatory compound of the invention is administered for a greater number of cycles that would typically cause dose-limiting toxicity in a patient to whom a second active ingredient is not also being administered.

In one embodiment, an immunomodulatory compound of the invention is administered daily and continuously for three or four weeks at a dose of from about 0.1 to about 150 mg/d followed by a break of one or two weeks. Actimid™ is preferably administered daily and continuously at an initial dose of 0.1 to 5 mg/d with dose escalation (every week) by 1 to 10 mg/d to a maximum dose of 50 mg/d for as long as therapy is tolerated. In a particular embodiment, Revimid™ is administered in an amount of about 5, 10, or 25 mg/day, preferably in an amount of about 10 mg/day for three to four weeks, followed by one week or two weeks of rest in a four or six week cycle.

In one embodiment of the invention, an immunomodulatory compound of the invention and a second active ingredient are administered orally, with administration of an immunomodulatory compound of the invention occurring 30 to 60 minutes prior to a second active ingredient, during a cycle of four to six weeks. In another embodiment of the invention, the combination of an immunomodulatory compound of the

US 9,101,621 B2

25

invention and a second active ingredient is administered by intravenous infusion over about 90 minutes every cycle. In a specific embodiment, one cycle comprises the administration of from about 10 to about 25 mg/day of Revimid™ and from about 50 to about 200 mg/m²/day of a second active ingredient daily for three to four weeks and then one or two weeks of rest. In another specific embodiment, each cycle comprises the administration of from about 5 to about 10 mg/day of Actimid™ and from about 50 to about 200 mg/m²/day of a second active ingredient for 3 to 4 weeks followed by one or two weeks of rest. Typically, the number of cycles during which the combinatorial treatment is administered to a patient will be from about one to about 24 cycles, more typically from about two to about 16 cycles, and even more typically from about four to about three cycles.

5.4 Pharmaceutical Compositions and Dosage Forms

Pharmaceutical compositions can be used in the preparation of individual, single unit dosage forms. Pharmaceutical compositions and dosage forms of the invention comprise an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof. Pharmaceutical compositions and dosage forms of the invention can further comprise one or more excipients.

Pharmaceutical compositions and dosage forms of the invention can also comprise one or more additional active ingredients. Consequently, pharmaceutical compositions and dosage forms of the invention comprise the active ingredients disclosed herein (e.g., an immunomodulatory compound and a second active agent). Examples of optional second, or additional, active ingredients are disclosed herein (see, e.g., section 5.2).

Single unit dosage forms of the invention are suitable for oral, mucosal (e.g., nasal, sublingual, vaginal, buccal, or rectal), parenteral (e.g., subcutaneous, intravenous, bolus injection, intramuscular, or intraarterial), topical (e.g., eye drops or other ophthalmic preparations), transdermal or transcutaneous administration to a patient. Examples of dosage forms include, but are not limited to: tablets; caplets; capsules, such as soft elastic gelatin capsules; cachets; troches; lozenges; dispersions; suppositories; powders; aerosols (e.g., nasal sprays or inhalers); gels; liquid dosage forms suitable for oral or mucosal administration to a patient, including suspensions (e.g., aqueous or non-aqueous liquid suspensions, oil-in-water emulsions, or a water-in-oil liquid emulsions), solutions, and elixirs; liquid dosage forms suitable for parenteral administration to a patient; eye drops or other ophthalmic preparations suitable for topical administration; and sterile solids (e.g., crystalline or amorphous solids) that can be reconstituted to provide liquid dosage forms suitable for parenteral administration to a patient.

The composition, shape, and type of dosage forms of the invention will typically vary depending on their use. For example, a dosage form used in the acute treatment of a disease may contain larger amounts of one or more of the active ingredients it comprises than a dosage form used in the chronic treatment of the same disease. Similarly, a parenteral dosage form may contain smaller amounts of one or more of the active ingredients it comprises than an oral dosage form used to treat the same disease. These and other ways in which specific dosage forms encompassed by this invention will vary from one another will be readily apparent to those skilled in the art. See, e.g., *Remington's Pharmaceutical Sciences*, 18th ed., Mack Publishing, Easton Pa. (1990).

26

Typical pharmaceutical compositions and dosage forms comprise one or more excipients. Suitable excipients are well known to those skilled in the art of pharmacy, and non-limiting examples of suitable excipients are provided herein.

Whether a particular excipient is suitable for incorporation into a pharmaceutical composition or dosage form depends on a variety of factors well known in the art including, but not limited to, the way in which the dosage form will be administered to a patient. For example, oral dosage forms such as tablets may contain excipients not suited for use in parenteral dosage forms. The suitability of a particular excipient may also depend on the specific active ingredients in the dosage form. For example, the decomposition of some active ingredients may be accelerated by some excipients such as lactose, or when exposed to water. Active ingredients that comprise primary or secondary amines are particularly susceptible to such accelerated decomposition. Consequently, this invention encompasses pharmaceutical compositions and dosage forms that contain little, if any, lactose other mono- or disaccharides. As used herein, the term "lactose-free" means that the amount of lactose present, if any, is insufficient to substantially increase the degradation rate of an active ingredient.

Lactose-free compositions of the invention can comprise excipients that are well known in the art and are listed, for example, in the *U.S. Pharmacopeia* (USP) 25-NF20 (2002). In general, lactose-free compositions comprise active ingredients, a binder/filler, and a lubricant in pharmaceutically compatible and pharmaceutically acceptable amounts. Preferred lactose-free dosage forms comprise active ingredients, microcrystalline cellulose, pre-gelatinized starch, and magnesium stearate.

This invention further encompasses anhydrous pharmaceutical compositions and dosage forms comprising active ingredients, since water can facilitate the degradation of some compounds. For example, the addition of water (e.g., 5%) is widely accepted in the pharmaceutical arts as a means of simulating long-term storage in order to determine characteristics such as shelf-life or the stability of formulations over time. See, e.g., Jens T. Carstensen, *Drug Stability: Principles & Practice*, 2d. Ed., Marcel Dekker, NY, N.Y., 1995, pp. 379-80. In effect, water and heat accelerate the decomposition of some compounds. Thus, the effect of water on a formulation can be of great significance since moisture and/or humidity are commonly encountered during manufacture, handling, packaging, storage, shipment, and use of formulations.

Anhydrous pharmaceutical compositions and dosage forms of the invention can be prepared using anhydrous or low moisture containing ingredients and low moisture or low humidity conditions. Pharmaceutical compositions and dosage forms that comprise lactose and at least one active ingredient that comprises a primary or secondary amine are preferably anhydrous if substantial contact with moisture and/or humidity during manufacturing, packaging, and/or storage is expected.

An anhydrous pharmaceutical composition should be prepared and stored such that its anhydrous nature is maintained. Accordingly, anhydrous compositions are preferably packaged using materials known to prevent exposure to water such that they can be included in suitable formulary kits. Examples of suitable packaging include, but are not limited to, hermetically sealed foils, plastics, unit dose containers (e.g., vials), blister packs, and strip packs.

The invention further encompasses pharmaceutical compositions and dosage forms that comprise one or more compounds that reduce the rate by which an active ingredient will

decompose. Such compounds, which are referred to herein as “stabilizers,” include, but are not limited to, antioxidants such as ascorbic acid, pH buffers, or salt buffers.

Like the amounts and types of excipients, the amounts and specific types of active ingredients in a dosage form may differ depending on factors such as, but not limited to, the route by which it is to be administered to patients. However, typical dosage forms of the invention comprise an immunomodulatory compound of the invention or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof in an amount of from about 0.10 to about 150 mg. Typical dosage forms comprise an immunomodulatory compound of the invention or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof in an amount of about 0.1, 1, 2, 5, 7.5, 10, 12.5, 15, 17.5, 20, 25, 50, 100, 150 or 200 mg. In a particular embodiment, a preferred dosage form comprises 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™) in an amount of about 1, 2, 5, 10, 25 or 50 mg. In a specific embodiment, a preferred dosage form comprises 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) in an amount of about 5, 10, 25 or 50 mg. Typical dosage forms comprise the second active ingredient in an amount of 1 to about 1000 mg, from about 5 to about 500 mg, from about 10 to about 350 mg, or from about 50 to about 200 mg. Of course, the specific amount of the anti-cancer drug will depend on the specific agent used, the type of cancer being treated or managed, and the amount(s) of an immunomodulatory compound of the invention and any optional additional active agents concurrently administered to the patient.

5.4.1 Oral Dosage Forms

Pharmaceutical compositions of the invention that are suitable for oral administration can be presented as discrete dosage forms, such as, but are not limited to, tablets (e.g., chewable tablets), caplets, capsules, and liquids (e.g., flavored syrups). Such dosage forms contain predetermined amounts of active ingredients, and may be prepared by methods of pharmacy well known to those skilled in the art. See generally, *Remington's Pharmaceutical Sciences*, 18th ed., Mack Publishing, Easton Pa. (1990).

Typical oral dosage forms of the invention are prepared by combining the active ingredients in an intimate admixture with at least one excipient according to conventional pharmaceutical compounding techniques. Excipients can take a wide variety of forms depending on the form of preparation desired for administration. For example, excipients suitable for use in oral liquid or aerosol dosage forms include, but are not limited to, water, glycols, oils, alcohols, flavoring agents, preservatives, and coloring agents. Examples of excipients suitable for use in solid oral dosage forms (e.g., powders, tablets, capsules, and caplets) include, but are not limited to, starches, sugars, micro-crystalline cellulose, diluents, granulating agents, lubricants, binders, and disintegrating agents.

Because of their ease of administration, tablets and capsules represent the most advantageous oral dosage unit forms, in which case solid excipients are employed. If desired, tablets can be coated by standard aqueous or nonaqueous techniques. Such dosage forms can be prepared by any of the methods of pharmacy. In general, pharmaceutical compositions and dosage forms are prepared by uniformly and intimately admixing the active ingredients with liquid carriers, finely divided solid carriers, or both, and then shaping the product into the desired presentation if necessary.

For example, a tablet can be prepared by compression or molding. Compressed tablets can be prepared by compressing in a suitable machine the active ingredients in a free-flowing form such as powder or granules, optionally mixed with an excipient. Molded tablets can be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

Examples of excipients that can be used in oral dosage forms of the invention include, but are not limited to, binders, fillers, disintegrants, and lubricants. Binders suitable for use in pharmaceutical compositions and dosage forms include, but are not limited to, corn starch, potato starch, or other starches, gelatin, natural and synthetic gums such as acacia, sodium alginate, alginic acid, other alginates, powdered tragacanth, guar gum, cellulose and its derivatives (e.g., ethyl cellulose, cellulose acetate, carboxymethyl cellulose calcium, sodium carboxymethyl cellulose), polyvinyl pyrrolidone, methyl cellulose, pre-gelatinized starch, hydroxypropyl methyl cellulose, (e.g., Nos. 2208, 2906, 2910), microcrystalline cellulose, and mixtures thereof.

Suitable forms of microcrystalline cellulose include, but are not limited to, the materials sold as AVICEL-PH-101, AVICEL-PH-103 AVICEL RC-581, AVICEL-PH-105 (available from FMC Corporation, American Viscose Division, Avicel Sales, Marcus Hook, Pa.), and mixtures thereof. A specific binder is a mixture of microcrystalline cellulose and sodium carboxymethyl cellulose sold as AVICEL RC-581. Suitable anhydrous or low moisture excipients or additives include AVICEL-PH-103™ and Starch 1500 LM.

Examples of fillers suitable for use in the pharmaceutical compositions and dosage forms disclosed herein include, but are not limited to, talc, calcium carbonate (e.g., granules or powder), microcrystalline cellulose, powdered cellulose, dextrates, kaolin, mannitol, silicic acid, sorbitol, starch, pre-gelatinized starch, and mixtures thereof. The binder or filler in pharmaceutical compositions of the invention is typically present in from about 50 to about 99 weight percent of the pharmaceutical composition or dosage form.

Disintegrants are used in the compositions of the invention to provide tablets that disintegrate when exposed to an aqueous environment. Tablets that contain too much disintegrant may disintegrate in storage, while those that contain too little may not disintegrate at a desired rate or under the desired conditions. Thus, a sufficient amount of disintegrant that is neither too much nor too little to detrimentally alter the release of the active ingredients should be used to form solid oral dosage forms of the invention. The amount of disintegrant used varies based upon the type of formulation, and is readily discernible to those of ordinary skill in the art. Typical pharmaceutical compositions comprise from about 0.5 to about 15 weight percent of disintegrant, preferably from about 1 to about 5 weight percent of disintegrant.

Disintegrants that can be used in pharmaceutical compositions and dosage forms of the invention include, but are not limited to, agar-agar, alginic acid, calcium carbonate, microcrystalline cellulose, croscarmellose sodium, crospovidone, polacrillin potassium, sodium starch glycolate, potato or tapioca starch, other starches, pre-gelatinized starch, other starches, clays, other alginates, other celluloses, gums, and mixtures thereof.

Lubricants that can be used in pharmaceutical compositions and dosage forms of the invention include, but are not limited to, calcium stearate, magnesium stearate, mineral oil, light mineral oil, glycerin, sorbitol, mannitol, polyethylene glycol, other glycols, stearic acid, sodium lauryl sulfate, talc, hydrogenated vegetable oil (e.g., peanut oil, cottonseed oil, sunflower oil, sesame oil, olive oil, corn oil, and soybean oil),

zinc stearate, ethyl oleate, ethyl laureate, agar, and mixtures thereof. Additional lubricants include, for example, a syloid silica gel (AEROSIL200, manufactured by W.R. Grace Co. of Baltimore, Md.), a coagulated aerosol of synthetic silica (marketed by Degussa Co. of Plano, Tex.), CAB-O-SIL (a pyrogenic silicon dioxide product sold by Cabot Co. of Boston, Mass.), and mixtures thereof. If used at all, lubricants are typically used in an amount of less than about 1 weight percent of the pharmaceutical compositions or dosage forms into which they are incorporated.

A preferred solid oral dosage form of the invention comprises an immunomodulatory compound of the invention, anhydrous lactose, microcrystalline cellulose, polyvinylpyrrolidone, stearic acid, colloidal anhydrous silica, and gelatin.

5.4.2 Delayed Release Dosage Forms

Active ingredients of the invention can be administered by controlled release means or by delivery devices that are well known to those of ordinary skill in the art. Examples include, but are not limited to, those described in U.S. Pat. Nos. 3,845,770; 3,916,899; 3,536,809; 3,598,123; and 4,008,719, 5,674,533, 5,059,595, 5,591,767, 5,120,548, 5,073,543, 5,639,476, 5,354,556, and 5,733,566, each of which is incorporated herein by reference. Such dosage forms can be used to provide slow or controlled-release of one or more active ingredients using, for example, hydropropylmethyl cellulose, other polymer matrices, gels, permeable membranes, osmotic systems, multilayer coatings, microparticles, liposomes, microspheres, or a combination thereof to provide the desired release profile in varying proportions. Suitable controlled-release formulations known to those of ordinary skill in the art, including those described herein, can be readily selected for use with the active ingredients of the invention. The invention thus encompasses single unit dosage forms suitable for oral administration such as, but not limited to, tablets, capsules, gelpcaps, and caplets that are adapted for controlled-release.

All controlled-release pharmaceutical products have a common goal of improving drug therapy over that achieved by their non-controlled counterparts. Ideally, the use of an optimally designed controlled-release preparation in medical treatment is characterized by a minimum of drug substance being employed to cure or control the condition in a minimum amount of time. Advantages of controlled-release formulations include extended activity of the drug, reduced dosage frequency, and increased patient compliance. In addition, controlled-release formulations can be used to affect the time of onset of action or other characteristics, such as blood levels of the drug, and can thus affect the occurrence of side (e.g., adverse) effects.

Most controlled-release formulations are designed to initially release an amount of drug (active ingredient) that promptly produces the desired therapeutic effect, and gradually and continually release of other amounts of drug to maintain this level of therapeutic or prophylactic effect over an extended period of time. In order to maintain this constant level of drug in the body, the drug must be released from the dosage form at a rate that will replace the amount of drug being metabolized and excreted from the body. Controlled-release of an active ingredient can be stimulated by various conditions including, but not limited to, pH, temperature, enzymes, water, or other physiological conditions or compounds.

5.4.3 Parenteral Dosage Forms

Parenteral dosage forms can be administered to patients by various routes including, but not limited to, subcutaneous,

intravenous (including bolus injection), intramuscular, and intraarterial. Because their administration typically bypasses patients' natural defenses against contaminants, parenteral dosage forms are preferably sterile or capable of being sterilized prior to administration to a patient. Examples of parenteral dosage forms include, but are not limited to, solutions ready for injection, dry products ready to be dissolved or suspended in a pharmaceutically acceptable vehicle for injection, suspensions ready for injection, and emulsions.

Suitable vehicles that can be used to provide parenteral dosage forms of the invention are well known to those skilled in the art. Examples include, but are not limited to: Water for Injection USP; aqueous vehicles such as, but not limited to, Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, and Lactated Ringer's Injection; water-miscible vehicles such as, but not limited to, ethyl alcohol, polyethylene glycol, and polypropylene glycol; and non-aqueous vehicles such as, but not limited to, corn oil, cottonseed oil, peanut oil, sesame oil, ethyl oleate, isopropyl myristate, and benzyl benzoate.

Compounds that increase the solubility of one or more of the active ingredients disclosed herein can also be incorporated into the parenteral dosage forms of the invention. For example, cyclodextrin and its derivatives can be used to increase the solubility of an immunomodulatory compound of the invention and its derivatives. See, e.g., U.S. Pat. No. 5,134,127, which is incorporated herein by reference.

5.4.4 Topical and Mucosal Dosage Forms

Topical and mucosal dosage forms of the invention include, but are not limited to, sprays, aerosols, solutions, emulsions, suspensions, eye drops or other ophthalmic preparations, or other forms known to one of skill in the art. See, e.g., *Remington's Pharmaceutical Sciences*, 16th and 18th eds., Mack Publishing, Easton Pa. (1980 & 1990); and *Introduction to Pharmaceutical Dosage Forms*, 4th ed., Lea & Febiger, Philadelphia (1985). Dosage forms suitable for treating mucosal tissues within the oral cavity can be formulated as mouthwashes or as oral gels.

Suitable excipients (e.g., carriers and diluents) and other materials that can be used to provide topical and mucosal dosage forms encompassed by this invention are well known to those skilled in the pharmaceutical arts, and depend on the particular tissue to which a given pharmaceutical composition or dosage form will be applied. With that fact in mind, typical excipients include, but are not limited to, water, acetone, ethanol, ethylene glycol, propylene glycol, butane-1,3-diol, isopropyl myristate, isopropyl palmitate, mineral oil, and mixtures thereof to form solutions, emulsions or gels, which are non-toxic and pharmaceutically acceptable. Moisturizers or humectants can also be added to pharmaceutical compositions and dosage forms if desired. Examples of such additional ingredients are well known in the art. See, e.g., *Remington's Pharmaceutical Sciences*, 16th and 18th eds., Mack Publishing, Easton Pa. (1980 & 1990).

The pH of a pharmaceutical composition or dosage form may also be adjusted to improve delivery of one or more active ingredients. Similarly, the polarity of a solvent carrier, its ionic strength, or tonicity can be adjusted to improve delivery. Compounds such as stearates can also be added to pharmaceutical compositions or dosage forms to advantageously alter the hydrophilicity or lipophilicity of one or more active ingredients so as to improve delivery. In this regard, stearates can serve as a lipid vehicle for the formulation, as an emulsifying agent or surfactant, and as a delivery-enhancing or penetration-enhancing agent. Different salts,

hydrates or solvates of the active ingredients can be used to further adjust the properties of the resulting composition.

5.4.5 Kits

Typically, active ingredients of the invention are preferably not administered to a patient at the same time or by the same route of administration. This invention therefore encompasses kits which, when used by the medical practitioner, can simplify the administration of appropriate amounts of active ingredients to a patient.

A typical kit of the invention comprises a dosage form of an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, pro-drug, or clathrate thereof. Kits encompassed by this invention can further comprise additional active ingredients such as oblimersen (Genasense®), melphalan, G-CSF, GM-CSF, EPO, topotecan, dacarbazine, irinotecan, taxotere, IFN, COX-2 inhibitor, pentoxifylline, ciprofloxacin, dexamethasone, IL2, IL8, IL18, Ara-C, vinorelbine, isotretinoin, 13 cis-retinoic acid, or a pharmacologically active mutant or derivative thereof, or a combination thereof. Examples of the additional active ingredients include, but are not limited to, those disclosed herein (see, e.g., section 5.2).

Kits of the invention can further comprise devices that are used to administer the active ingredients. Examples of such devices include, but are not limited to, syringes, drip bags, patches, and inhalers.

Kits of the invention can further comprise cells or blood for transplantation as well as pharmaceutically acceptable vehicles that can be used to administer one or more active ingredients. For example, if an active ingredient is provided in a solid form that must be reconstituted for parenteral administration, the kit can comprise a sealed container of a suitable vehicle in which the active ingredient can be dissolved to form a particulate-free sterile solution that is suitable for parenteral administration. Examples of pharmaceutically acceptable vehicles include, but are not limited to: Water for Injection USP; aqueous vehicles such as, but not limited to, Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, and Lactated Ringer's Injection; water-miscible vehicles such as, but not limited to, ethyl alcohol, polyethylene glycol, and polypropylene glycol; and non-aqueous vehicles such as, but not limited to, corn oil, cottonseed oil, peanut oil, sesame oil, ethyl oleate, isopropyl myristate, and benzyl benzoate.

6. EXAMPLES

Certain embodiments of the invention are illustrated by the following non-limiting examples.

6.1 Modulation of Cytokine Production

A series of non-clinical pharmacology and toxicology studies have been performed to support the clinical evaluation of an immunomodulatory compound of the invention in human subjects. These studies were performed in accordance with internationally recognized guidelines for study design and in compliance with the requirements of Good Laboratory Practice (GLP), unless otherwise noted.

Inhibition of TNF- α production following LPS-stimulation of human PBMC and human whole blood by 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™), 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione and thalidomide (Revimid™) was investigated in vitro (Muller et al., *Bioorg. Med. Chem. Lett.* 9:1625-1630, 1999). The IC₅₀'s of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione for inhibiting production of TNF- α following LPS-stimulation of PBMC and

human whole blood were ~24 nM (6.55 ng/mL) and ~25 nM (6.83 ng/mL), respectively. In vitro studies suggest a pharmacological activity profile for 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione that is similar to, but at least 200 times more potent than, thalidomide. In vitro studies have also demonstrated that concentrations of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione of 2.73 to 27.3 ng/mL (0.01 to 0.1 μ M) achieved 50% inhibition of the proliferation of MM.1S and Hs Sultan cells.

The IC₅₀'s of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione for inhibiting production of TNF- α following LPS-stimulation of PBMC and human whole blood were ~100 nM (25.9 ng/mL) and ~480 nM (103.6 ng/mL), respectively. Thalidomide, in contrast, had an IC₅₀ of ~194 μ M (50.2 μ g/mL) for inhibiting production of TNF- α following LPS-stimulation of PBMC. In vitro studies suggest a pharmacological activity profile for 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione that is similar to, but 50 to 2000 times more potent than, thalidomide. It has been shown that the compound is approximately 50-100 times more potent than thalidomide in stimulating the proliferation of T-cells following primary induction by T-cell receptor (TCR) activation. 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione is also approximately 50 to 100 times more potent than thalidomide in augmenting the production of IL-2 and IFN- γ following TCR activation of PBMC (IL-2) or T-cells (IFN- γ). In addition, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione exhibited dose-dependent inhibition of LPS-stimulated production of the pro-inflammatory cytokines TNF- α , IL-1 β , and IL-6 by PBMC while it increased production of the anti-inflammatory cytokine IL-10.

6.2 Inhibition of MM Cell Proliferation

The ability of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) and thalidomide for comparison to effect the proliferation of MM cell lines has been investigated in an in vitro study. Uptake [³H]-thymidine by different MM cell lines (MM.1S, Hs Sultan, U266 and RPMI-8226) was measured as an indicator of cell proliferation. Cells were incubated in the presence of compounds for 48 hours; [³H]-thymidine was included for the last 8 hours of the incubation period. Addition of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione to MM.1S and Hs Sultan cells resulted in 50% inhibition of cell proliferation at concentrations of 0.4 μ M and 1 μ M, respectively. In contrast, addition of thalidomide at concentrations up to 100 μ M resulted in only 15% and 20% inhibition of cell proliferation in MM.1S and Hs Sultan cells, respectively. These data are summarized in FIG. 1.

6.3 Toxicology Studies

The effects of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) on cardiovascular and respiratory function are investigated in anesthetized dogs. Two groups of Beagle dogs (2/sex/group) are used. One group receives three doses of vehicle only and the other receives three ascending doses of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (2, 10, and 20 mg/kg). In all cases, doses of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione or vehicle are successively administered via infusion through the jugular vein separated by intervals of at least 30 minutes.

The cardiovascular and respiratory changes induced by 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione are minimal at all doses when compared to the vehicle control group. The only statistically significant difference between the vehicle and treatment groups is a small

increase in arterial blood pressure (from 94 mmHg to 101 mmHg) following administration of the low dose of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. This effect lasts approximately 15 minutes and is not seen at higher doses. Deviations in femoral blood flow, respiratory parameters, and Qtc interval are common to both the control and treated groups and are not considered treatment-related.

6.4 Cycling Therapy in Patients

In a specific embodiment, an immunomodulatory compound of the invention are cyclically administered to patients with cancer. Cycling therapy involves the administration of a first agent for a period of time, followed by a rest for a period of time and repeating this sequential administration. Cycling therapy can reduce the development of resistance to one or more of the therapies, avoid or reduce the side effects of one of the therapies, and/or improves the efficacy of the treatment.

In a specific embodiment, prophylactic or therapeutic agents are administered in a cycle of about 4 to 6 weeks, about once or twice every day. One cycle can comprise the administration of a therapeutic on prophylactic agent for three to four weeks and at least a week or two weeks of rest. The number of cycles administered is from about one to about 24 cycles, more typically from about two to about 16 cycles, and more typically from about four to about eight cycles.

For example, in a cycle of four weeks, on day 1, the administration of 25 mg/d of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione is started. On day 22, the administration of the compound is stopped for a week of rest. On day 29, the administration of 25 mg/d 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidin-2,6-dione is begun.

6.5 Clinical Studies in Patients

6.5.1 Treatment of Relapsed Multiple Myeloma

4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™) was administered to patients with relapsed/refractory multiple myeloma. The study was conducted in compliance with Good Clinical Practices. Patients were at least 18 years old, had been diagnosed with multiple myeloma (with paraprotein in serum and/or urine), and were considered refractory to treatment after at least two cycles of treatment, or have relapsed after two cycles of treatment.

Patients who have progressive disease, according to the Southwest Oncology Group (SWOG) criteria, on their prior regimen are considered treatment refractory. Relapse following remission is defined as >25% increase in M component from baseline levels; reappearance of the M paraprotein that had previously disappeared; or a definite increase in the size and number of lytic bone lesions recognized on radiographs. Patients may have had prior therapy with thalidomide, provided they were able to tolerate the treatment. A Zubrod performance status of 0 to 2 is required for all patients.

4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione is administered to patients at doses of 1, 2, 5, or 10 mg/day for up to four weeks; at each dose level, three patients are initially enrolled. Dosing occurs at approximately the same time each morning; all doses are administered in the fasted state (no eating for at least two hours prior to dosing and two hours after dosing). 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione doses are administered in an ascending fashion such that patients in the first cohort receive the lowest dose of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (1 mg/day) and escalation to the next higher dose

level occurs only following the establishment of safety and tolerability at the current dose. If one out of three patients at any dose level experience dose limiting toxicity (DLT), three additional patients are enrolled at that dose. If none of the three additional patients experience DLT, escalation to the next dose level occurs; dose escalations continue in a similar fashion until the MTD is established or the maximum daily dose (10 mg/day) is attained. However, if one of the three additional patients enrolled experiences DLT, the MTD has been reached. If two or more of the three additional patients enrolled experience DLT, the MTD is judged to have been exceeded and three additional patients are enrolled at the preceding dose level to confirm the MTD. Once the MTD has been identified, four additional patients are enrolled at that dose level so that a total of 10 patients is treated at the MTD.

Blood sampling for analysis of pharmacokinetic parameters is performed on Days 1 and 28 according to the following sampling schedule: pre-dose, 0.25, 0.5, 0.75, 1, 1.5, 2, 2.5, 3, 4, 6, 8, 10, 12, 18, and 24 hours post-dose. An additional blood sample is collected at each weekly visit for the determination of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione levels. Total urine collections are also made with urine pooled according to the following time intervals post-dose: 0 to 4, 4 to 8, 8 to 12, and 12 to 24 hours. Safety assessments are made by monitoring adverse events, vital signs, ECGs, clinical laboratory evaluations (blood chemistry, hematology, lymphocyte phenotyping, and urinalysis), and physical examination at specific times during the study.

Results of interim pharmacokinetic analyses obtained following single- and multiple-dose administration of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione to multiple myeloma patients are presented below in Tables 1 and 2. These data show that 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione was steadily absorbed at all dose levels in relapsed multiple myeloma patients. Maximum plasma concentrations occurred at a median T_{max} of between 2.5 and 2.8 hours post-dose at Day 1 and between 3 and 4 hours post-dose at Week 4. At all doses, plasma concentrations declined in a monophasic manner after reaching C_{max} . The start of the elimination phase occurred between 3 and 10 hours post-dose at Day 1 and Week 4, respectively.

These data also showed that after 4 weeks of dosing, 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione accumulated to a small extent (mean accumulation ratios ~1.02 to 1.52 and ~0.94 to 1.62 for C_{max} and $AUC_{(0-\tau)}$, respectively). There was almost a dose proportional increase in $AUC_{(0-\tau)}$ and C_{max} values with increasing dose. A five-fold higher dose of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione produced a 3.2- and 2.2-fold increase in C_{max} at Day 1 and Week 4, respectively. Similarly, a 5-fold increase in dose resulted in a 3.6- and 2.3-fold increase in $AUC_{(0-\tau)}$, at Day 1 and Week 4, respectively.

TABLE 1

| Pharmacokinetic parameters of Actimid™ in relapsed multiple myeloma patients | | | | |
|---|-----------|-------------------|-------------------|--------------------|
| Parameter | | 1 mg | 2 mg | 5 mg |
| | | (N = 6) | (N = 2) | (N = 3) |
| Day 1 | | | | |
| C_{max} | ng/mL | 15.03 (4.04) | 24.4* (12.1) | 48.56 (14.03) |
| t_{max} | h | 3.3 (2.6) | 2.7* (0.3) | 2.3 (0.3) |
| $AUC_{(0-\infty)}$ | ng · h/mL | 152.90 (36.62) | 279.18 (51.10) | 593.10 (335.23) |

TABLE 1-continued

| Pharmacokinetic parameters of Actimid™ in relapsed multiple myeloma patients | | | | |
|---|--------|-------------------|-------------------|--------------------|
| Parameter | | 1 mg (N = 6) | 2 mg (N = 2) | 5 mg (N = 3) |
| AUC _(0-τ) | | 134.21 (27.14) | 249.57 (29.26) | 520.94 (267.32) |
| t _{1/2} | h | 7.3 (3.4) | 6.3 (1.4) | 6.5 (2.2) |
| CL/F | mL/min | 114.75 (29.20) | 121.43 (22.22) | 182.31 (117.06) |
| Vz/f | L | 69.55 (44.97) | 65.31 (2.80) | 87.24 (22.61) |

τ = 24 hours

N/A = not available

TABLE 2

| Pharmacokinetic parameters of Actimid™ following multiple oral doses (1, 2, and 5 mg/day) in relapsed multiple myeloma patients | | | | |
|--|-----------|--------------------|--------------------|--------------------|
| Parameter | | 1 mg (N = 5) | 2 mg (N = 2) | 5 mg (N = 3) |
| Week 4 | | | | |
| C _{max} | ng/mL | 23.20 (7.48) | 30.05* (15.64) | 58.07 (38.08) |
| t _{max} | h | 3.6 (1.5) | 2.8* (0.3) | 5.0 (2.6) |
| AUC _(0-∞) | ng · h/mL | N/A | N/A | N/A |
| AUC _(0-τ) | | 239.31 (122.59) | 269.36 (186.34) | 597.24 (354.23) |
| t _{1/2} | h | 6.2* (0.6) | 7.7 (2.8) | 7.8 (4.0) |
| CL/F | mL/min | 87.85 (48.48) | 162.68 (112.54) | 207.50 (175.41) |
| Vz/f | L | 41.35* (8.84) | 95.04 (35.39) | 103.95 (27.25) |

τ = 24 hours

N/A = not available

*N = 3 patients

6.5.2 Treatment of Relapsed Multiple Myeloma

Two Phase 1 clinical studies of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) have been conducted to identify the maximum tolerated dose (MTD) in patients with refractory or relapsed multiple myeloma. These studies have also characterized the safety profile of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione when ascending doses of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione were given orally for up to 4 weeks. Patients started 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione treatment at 5 mg/day with subsequent escalation to 10, 25, and 50 mg/day. Patients were enrolled for 28 days at their assigned dose, with the option of extended treatment for those who did not exhibit disease progression or experience dose limiting toxicity (DLT). Patients were evaluated for adverse events at each visit and the severity of these events was graded according to the National Cancer Institute (NCI) Common Toxicity Criteria. Patients were discontinued if they experienced DLT (Grade 3 or greater non-hematological, or Grade 4 hematological toxicity).

In this study, 27 patients were enrolled. All patients had relapsed multiple myeloma and 18 (72%) were refractory to salvage therapy. Among these patients, 15 had undergone prior autologous stem cell transplantation and 16 patients had

received prior thalidomide treatment. The median number of prior regimens was 3 (range 2 to 6).

Blood and urine samples were collected for analysis of pharmacokinetic parameters on Days 1 and 28. Blood samples were collected according to the following sampling schedule: pre-dose, 0.25, 0.5, 0.75, 1, 1.5, 2, 2.5, 3, 4, 6, 8, 10, 12, 18, and 24 hours post-dose. In addition, a blood sample was collected at each weekly clinic visit for 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione determination. Total urine was collected and pooled according to the following time intervals post-dose: 0 to 4, 4 to 8, 8 to 12, and 12 to 24 hours. Response to treatment was assessed by M-protein quantification (by immunoelectrophoresis) from serum and a 24-hour urine collection, with creatinine clearance and 24-hour protein calculations undertaken at screening, baseline, Weeks 2 and 4, and monthly thereafter (or upon early termination). Bone marrow aspirations and/or tissue biopsy are also performed at Months 3, 6 and 12 if a patient's paraprotein serum concentration or 24-hour urine protein excretion declined to the next lower level, based on best response criteria. Preliminary results for the 28-day treatment period are summarized below.

Preliminary pharmacokinetic analyses based on these two studies indicated that AUC and C_{max} values increase proportionally with dose following single and multiple doses in multiple myeloma patients (as was seen in healthy volunteers). Further, there was no evidence of accumulation with multiple dosing as single dose AUC_(0-∞) was comparable to multiple dose AUC_(0-τ) following the same dose of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. Similar to healthy volunteer studies, double peaks were observed. Exposure in multiple myeloma patients appeared to be slightly higher based on C_{max} and AUC values as compared to healthy male volunteers while clearance in multiple myeloma patients was lower than it was in healthy volunteers, consistent with their poorer renal function (both as a consequence of their age and their disease). Finally, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione half-life in patients was shorter than in healthy volunteers (mean 8 hours, ranging up to 17 hours).

In this study, the first cohort of 3 patients was treated for 28 days at 5 mg/day without any dose limiting toxicity (DLT). The second cohort of 3 patients subsequently commenced therapy at 10 mg/day. Patients in the second 10 mg/day of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione cohort tolerated treatment well.

6.5.3 Treatment of Solid Tumors

Study with 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) was conducted in patients with varying types of solid tumors, including malignant melanoma (13), carcinoma of the pancreas (2), carcinoid-unknown primary (1), renal carcinoma (1), breast carcinoma (1) and NSCLC (2). Patients received 5 mg/day 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione for seven days and are subsequently escalated every seven days to 10 mg/day, 25 mg/day, and 50 mg/day for a total of 4 weeks of treatment. Patients who, experienced clinical benefit were permitted to continue on treatment as Named Patients.

The study initially enrolled 20 patients and was subsequently amended to enroll 16 additional patients (adrenal carcinoma, NSCLC, malignant mesothelioma, breast cancer, malignant melanoma (8), renal cell cancer (4)) at a higher dose. The 16 additional patients were given weekly escalating doses of 25 mg/day, 50 mg/day, 75 mg/day, 100 mg/day, 125

US 9,101,621 B2

37

mg/day, and 150 mg/day over a 6-week period with continuing treatment for an additional six weeks.

The study of Phase 1 study was designed to determine a maximum tolerated dose (MTD) of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione in patients with refractory solid tumors and/or lymphoma, as well as to characterize the pharmacokinetic and side effect profiles of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione in this patient population. The study design dictates that at least 3 patients must be enrolled at a dose level and have completed 28 days of treatment prior to enrollment of patients at the next higher dose level. Patients in the first cohort began dosing at 5 mg/day of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. Patients will be escalated to 10, 20, 25, and 30 mg/day provided there is no toxicity.

In this study, the MTD is defined as the highest dose level in which fewer than two of six patients treated did not experience Grade 3 or greater non-hematological toxicity or Grade 4 or greater hematological toxicity. If, at any given dose level in either study, one out of three patients experiences toxicity, three additional patients must be treated at that particular dose. If, however, two out of six patients experience DLT, the MTD is judged to have been exceeded. No further dose escalations are to occur and additional patients are to be enrolled at the previous dose level. The dose of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione administered is escalated until the MTD is achieved or the maximum daily dose of is reached.

No DLTs were reported in the initial group of 20 patients enrolled in the study. Thirteen of the original 20 trial patients, along with 2 non-trial patients, continued on treatment as named patients at doses up to 150 mg/day.

6.5.4 Treatment of Gliomas

This study was performed to find toxicity in patients with recurrent, high-grade gliomas. The study is designed such that patients are given increasingly higher doses of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione until a maximum tolerated dose (MTD) is established. The study also seeks to obtain preliminary toxicity information and pharmacokinetic data on 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione, as well as to develop exploratory data concerning surrogate end points of angiogenic activity in vivo using functional neuro-imaging studies, and in vitro assays of scrum angiogenic peptides.

Patients enrolled in the first cohort receive 2.5 mg/m²/day for a 4-week cycle. During each 4-week cycle of therapy, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione administered once daily for 3 weeks followed by a week of rest. Patients who complete a treatment cycle may receive another cycle of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione treatment if two criteria are met. First, the patient must have stable disease or have experienced a partial response or complete response, or the patient is benefiting from the therapy with 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione as evidenced by a decrease in tumor-related symptoms such as neurological deficits. Second, the patient must have recovered from toxicity related to 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione which occurred in the prior cycle by Day 42 or sooner (28-day cycle plus limit of 2 weeks to recover) as evidenced by a return to Grade \leq 1 toxicity level. Patients who experience DLT in the previous cycle should have their dose modified. DLT is defined as a non-hematological event Grade \geq 3 toxicity or hematological event of Grade 4 toxicity thought to be related to the study

38

medication. Patients who experience DLT in the first cycle and have no response to therapy are removed from the study.

3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione doses are subsequently escalated to 5, 8, 11, 15, and 20 mg/m²/day to a maximum total daily dose of 40 mg. Patients continue to receive 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione on a 4-week cycle per dose level until one of the off-study criteria are met.

Three patients are enrolled in each cohort. If at least one DLT occurs, three additional patients are added to the cohort at that particular dose level. If two DLTs occur, the MTD, defined as the dose at which fewer than one-third of patients at each dose level experiences DLT has been exceeded and four more patients are treated at the previous dose.

Patients who experience DLT during the first 4-week cycle are removed from the study, except if they have a response to therapy. For patients who have completed their first 4-week cycle of without DLT, but who subsequently experience Grade 3 or 4 hematological and/or nonhematological toxicity, treatment is suspended for a minimum of a week. If the toxicity resolves to <Grade 2 within three weeks, the patient is treated at two dose levels lower than the dose that caused the toxicity (or a 50% reduction if the patient was treated at the first or second dose level). Patients in whom Grade 3 or 4 toxicity does not resolve to <Grade 1 within three weeks, or those who have another Grade 3 toxicity at the reduced dose are removed from the study.

Pharmacokinetic sampling is performed prior the first dose of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Day 1) and 0.5, 1, 2, 4, 6, 8, 24, and 48 hours thereafter. Sampling is also conducted pre-dose on Days 7 and 21 and 0.5, 1, 2, 4, 6, 8, and 24 post-dose on Day 21 to evaluate steady-state 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione levels.

6.5.5 Treatment of Metastatic Melanoma

Patients with metastatic melanoma were started on 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revmid™) at 5 mg/day for seven days. The dose was then increased every seven days to 10 mg/day, 25 mg/day, and 50 mg/day, respectively, for a total of four weeks on therapy. Five of the 13 melanoma patients who were treated under this regimen either showed disease stabilization or a partial response in the first four weeks of treatment. Tumor response was seen in cutaneous and subcutaneous lesions (five patients), lymph nodes (two patients), and liver (one patient). The duration of response was approximately six months. The result suggests that the compound appears is a promising new anti-cancer agent and has both antiangiogenic and immunomodulatory properties.

6.5.6 Treatment of Relapsed or Refractory Multiple Myeloma

Patients with relapsed and refractory Dune-Salmon stage III multiple myeloma, who have either failed at least three previous regimens or presented with poor performance status, neutropenia or thrombocytopenia, are treated with up to four cycles of combination of melphalan (50 mg intravenously), an immunomodulatory compound of the invention (about 1 to 150 mg orally daily), and dexamethasone (40 mg/day orally on days 1 to 4) every four to six weeks. Maintenance treatment consisting of daily an immunomodulatory compound of the invention and monthly dexamethasone are continued until the disease progression. The therapy using an immunomodulatory compound of the invention in combination with mel-

US 9,101,621 B2


39

phalan and dexamethasone is highly active and generally tolerated in heavily pretreated multiple myeloma patients whose prognosis is otherwise poor.

The embodiments of the invention described above are intended to be merely exemplary, and those skilled in the art will recognize, or will be able to ascertain using no more than routine experimentation, numerous equivalents of specific compounds, materials, and procedures. All such equivalents are considered to be within the scope of the invention and are encompassed by the appended claims.

What is claimed is:

1. A method of treating multiple myeloma, which comprises administering to a patient having multiple myeloma about 1 to about 50 mg per day of a compound having the formula:



or a pharmaceutically acceptable salt, solvate or stereoisomer thereof, wherein the patient has previously received stem cell transplantation.

2. The method of claim 1, wherein the multiple myeloma is relapsed, refractory, or relapsed and refractory multiple myeloma.

3. The method of claim 1, wherein the compound is administered for 21 consecutive days followed by seven consecutive days of rest in a 28 day cycle.

4. The method of claim 1, wherein the compound is administered every day.

5. The method of claim 1, wherein the compound is administered orally.

40

6. The method of claim 1, wherein the compound is administered in the form of a capsule or tablet.

7. The method of claim 1, wherein the compound is administered in a capsule of 2.5 mg, 5 mg, 10 mg, 15 mg, 20 mg or 25 mg.

8. The method of claim 6, wherein the capsule comprises the compound, lactose anhydrous, microcrystalline cellulose, croscarmellose sodium and magnesium stearate.

9. The method of claim 1, wherein the compound is administered in an amount of about 25 mg per day.

10. The method of claim 1, wherein the compound is administered in an amount of about 15 mg per day.

11. The method of claim 1, wherein the compound is administered in an amount of about 10 mg per day.

12. The method of claim 1, wherein the compound is administered in an amount of about 5 mg per day.

13. The method of claim 3, wherein the compound is administered in an amount of about 25 mg per day.

14. The method of claim 4, wherein the compound is administered in an amount of about 5 mg, 10 mg or 15 mg per day.

15. The method of claim 1, wherein the stem cell transplantation is autologous stem cell transplantation.

16. The method of claim 1, wherein the stem cell transplantation is hematopoietic stem cell transplantation.

17. The method of claim 1, wherein the stem cell transplantation is peripheral blood stem cell transplantation.

18. The method of claim 1, wherein the compound is administered in an amount of about 2.5 mg per day.

19. The method of claim 3, wherein the compound is administered in an amount of about 10 mg per day.

20. The method of claim 1, wherein the compound is administered in an amount of about 20 mg per day.

21. The method of claim 1, wherein the compound is administered in a capsule containing from about 1 to about 50 mg of the compound.

* * * * *

EXHIBIT G



US009101622B2

(12) **United States Patent**
Zeldis

(10) **Patent No.:** **US 9,101,622 B2**
(45) **Date of Patent:** ***Aug. 11, 2015**

(54) **METHODS FOR TREATING NEWLY DIAGNOSED MULTIPLE MYELOMA 3-(4-AMINO-1-OXO-1,3-DIHYDRO-ISOINDOL-2-YL)-PIPERIDINE-2,6-DIONE IN COMBINATION WITH DEXAMETHASONE**

(71) Applicant: **Celgene Corporation**, Summit, NJ (US)

(72) Inventor: **Jerome B. Zeldis**, Princeton, NJ (US)

(73) Assignee: **Celgene Corporation**, Summit, NJ (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **14/482,745**

(22) Filed: **Sep. 10, 2014**

(65) **Prior Publication Data**

US 2014/0377293 A1 Dec. 25, 2014

Related U.S. Application Data

(63) Continuation of application No. 14/255,211, filed on Apr. 17, 2014, which is a continuation of application No. 14/201,069, filed on Mar. 7, 2014, which is a continuation of application No. 13/782,728, filed on Mar. 1, 2013, now Pat. No. 8,673,939, which is a continuation of application No. 13/488,888, filed on Jun. 5, 2012, now Pat. No. 8,648,095, which is a continuation of application No. 12/640,702, filed on Dec. 17, 2009, now Pat. No. 8,198,306, which is a continuation of application No. 10/438,213, filed on May 15, 2003, now Pat. No. 7,968,569.

(60) Provisional application No. 60/380,842, filed on May 17, 2002, provisional application No. 60/424,600, filed on Nov. 6, 2002.

(51) **Int. Cl.**

A61K 31/4035 (2006.01)
A61K 31/425 (2006.01)
A61K 31/445 (2006.01)
A61K 31/454 (2006.01)
A61K 31/00 (2006.01)
A61K 31/40 (2006.01)
A61K 31/515 (2006.01)
A61K 45/06 (2006.01)
A61K 31/4439 (2006.01)
A61K 31/573 (2006.01)
A61K 31/198 (2006.01)
A61K 31/475 (2006.01)
A61K 31/675 (2006.01)
A61K 31/704 (2006.01)
A61K 31/7048 (2006.01)
A61K 35/12 (2015.01)
A61K 39/395 (2006.01)
A61K 39/00 (2006.01)

(52) **U.S. Cl.**

CPC **A61K 31/454** (2013.01); **A61K 31/00** (2013.01); **A61K 31/198** (2013.01); **A61K 31/40** (2013.01); **A61K 31/4035** (2013.01); **A61K 31/425** (2013.01); **A61K 31/445** (2013.01); **A61K 31/4439** (2013.01); **A61K 31/475** (2013.01); **A61K 31/515** (2013.01); **A61K 31/573** (2013.01); **A61K 31/675** (2013.01); **A61K 31/704** (2013.01); **A61K 31/7048** (2013.01); **A61K 35/12** (2013.01); **A61K 39/0011** (2013.01); **A61K 39/3955** (2013.01); **A61K 45/06** (2013.01)

(58) **Field of Classification Search**

CPC **A61K 31/4035**; **A61K 31/425**; **A61K 31/445**; **A61K 31/454**
USPC **514/323**
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

| | | |
|-------------|---------|------------------|
| 3,536,809 A | 10/1970 | Applezweig |
| 3,598,123 A | 8/1971 | Zaffaroni et al. |
| 3,845,770 A | 11/1974 | Theeuwes et al. |
| 3,916,899 A | 11/1975 | Theeuwes et al. |
| 4,008,719 A | 2/1977 | Theeuwes et al. |

(Continued)

FOREIGN PATENT DOCUMENTS

| | | |
|----|----------------|---------|
| JP | 2002-513391 | 5/2002 |
| WO | WO 92/14455 | 9/1992 |
| WO | WO 94/20085 | 9/1994 |
| WO | WO 98/03502 | 1/1998 |
| WO | WO 98/19649 | 5/1998 |
| WO | WO 98/54170 | 12/1998 |
| WO | WO 01/70275 | 9/2001 |
| WO | WO 01/87307 | 11/2001 |
| WO | WO 02/15926 | 2/2002 |
| WO | WO 02/059106 | 8/2002 |
| WO | WO 02/064083 | 8/2002 |
| WO | PCT/US03/11578 | 4/2003 |
| WO | WO 03/086373 | 10/2003 |

OTHER PUBLICATIONS

Zeldis, Jerome B. U.S. Appl. No. 14/255,211, filed Sep. 10, 2014.*

(Continued)

Primary Examiner — Jeffrey S Lundgren

Assistant Examiner — Chris Simmons

(74) *Attorney, Agent, or Firm* — Jones Day

(57) **ABSTRACT**

Methods of treating, preventing and/or managing cancer as well as and diseases and disorders associated with, or characterized by, undesired angiogenesis are disclosed. Specific methods encompass the administration of an immunomodulatory compound alone or in combination with a second active ingredient. The invention further relates to methods of reducing or avoiding adverse side effects associated with chemotherapy, radiation therapy, hormonal therapy, biological therapy or immunotherapy which comprise the administration of an immunomodulatory compound. Pharmaceutical compositions, single unit dosage forms, and kits suitable for use in methods of the invention are also disclosed.

25 Claims, 1 Drawing Sheet

US 9,101,622 B2

Page 2

(56)

References Cited

U.S. PATENT DOCUMENTS

- | | | | | | | | |
|--------------|------|---------|----------------------|--------------|----|---------|-------------------|
| 4,810,643 | A | 3/1989 | Souza | 2002/0183360 | A1 | 12/2002 | Muller et al. |
| 4,999,291 | A | 3/1991 | Souza | 2003/0013739 | A1 | 1/2003 | Masferrer et al. |
| 5,059,595 | A | 10/1991 | Le Grazie | 2003/0028028 | A1 | 2/2003 | Man et al. |
| 5,073,543 | A | 12/1991 | Marshall et al. | 2003/0045552 | A1 | 3/2003 | Robarge et al. |
| 5,120,548 | A | 6/1992 | McClelland et al. | 2003/0069428 | A1 | 4/2003 | Muller et al. |
| 5,134,127 | A | 7/1992 | Stella et al. | 2003/0096841 | A1 | 5/2003 | Robarge et al. |
| 5,229,496 | A | 7/1993 | Deeley et al. | 2003/0139451 | A1 | 7/2003 | Shah et al. |
| 5,354,556 | A | 10/1994 | Sparks et al. | 2003/0144325 | A1 | 7/2003 | Muller et al. |
| 5,385,901 | A | 1/1995 | Kaplan et al. | 2003/0181428 | A1 | 9/2003 | Green et al. |
| 5,391,485 | A | 2/1995 | Deeley et al. | 2003/0187024 | A1 | 10/2003 | D'Amato |
| 5,393,870 | A | 2/1995 | Deeley et al. | 2003/0191098 | A1 | 10/2003 | D'Amato |
| 5,528,823 | A | 6/1996 | Rudy, Jr. et al. | 2003/0235909 | A1 | 12/2003 | Hariri et al. |
| 5,580,755 | A | 12/1996 | Souza | 2004/0029832 | A1 | 2/2004 | Zeldis |
| 5,591,767 | A | 1/1997 | Mohr et al. | 2004/0077685 | A1 | 4/2004 | Figg et al. |
| 5,593,990 | A | 1/1997 | D'Amato | 2004/0077686 | A1 | 4/2004 | Dannenberg et al. |
| 5,629,327 | A | 5/1997 | D'Amato | 2004/0087546 | A1 | 5/2004 | Zeldis |
| 5,635,517 | A | 6/1997 | Muller et al. | 2004/0091455 | A1 | 5/2004 | Zeldis |
| 5,639,476 | A | 6/1997 | Oshlack et al. | 2004/0122052 | A1 | 6/2004 | Muller et al. |
| 5,674,533 | A | 10/1997 | Santus et al. | 2004/0266809 | A1 | 12/2004 | Emanuel et al. |
| 5,698,579 | A | 12/1997 | Muller | 2009/0010877 | A1 | 1/2009 | Zeldis |
| 5,712,291 | A | 1/1998 | D'Amato | 2009/0123416 | A1 | 5/2009 | Zeldis |
| 5,731,325 | A | 3/1998 | Andrulis, Jr. et al. | 2012/0035145 | A1 | 2/2012 | Zeldis |
| 5,733,566 | A | 3/1998 | Lewis | 2014/0186404 | A1 | 7/2014 | Zeldis |
| 5,798,368 | A | 8/1998 | Muller et al. | | | | |
| 5,874,448 | A | 2/1999 | Muller et al. | | | | |
| 5,877,200 | A | 3/1999 | Muller | | | | |
| 5,929,117 | A | 7/1999 | Muller et al. | | | | |
| 5,955,476 | A | 9/1999 | Muller et al. | | | | |
| 6,020,358 | A | 2/2000 | Muller et al. | | | | |
| 6,071,948 | A | 6/2000 | D'Amato | | | | |
| 6,114,355 | A | 9/2000 | D'Amato | | | | |
| 6,140,346 | A | 10/2000 | Andrulis, Jr. et al. | | | | |
| 6,228,879 | B1 | 5/2001 | Green et al. | | | | |
| 6,235,756 | B1 | 5/2001 | D'Amato | | | | |
| 6,281,230 | B1 | 8/2001 | Muller et al. | | | | |
| 6,316,471 | B1 | 11/2001 | Muller et al. | | | | |
| 6,326,388 | B1 | 12/2001 | Man et al. | | | | |
| 6,335,349 | B1 | 1/2002 | Muller et al. | | | | |
| 6,380,239 | B1 | 4/2002 | Muller et al. | | | | |
| 6,395,754 | B1 | 5/2002 | Muller et al. | | | | |
| 6,403,613 | B1 | 6/2002 | Man et al. | | | | |
| 6,420,414 | B1 | 7/2002 | D'Amato | | | | |
| 6,458,810 | B1 | 10/2002 | Muller et al. | | | | |
| 6,469,045 | B1 | 10/2002 | D'Amato | | | | |
| 6,476,052 | B1 | 11/2002 | Muller et al. | | | | |
| 6,518,298 | B2 | 2/2003 | Green et al. | | | | |
| 6,555,554 | B2 | 4/2003 | Muller et al. | | | | |
| 6,673,828 | B1 | 1/2004 | Green et al. | | | | |
| 7,323,479 | B2 | 1/2008 | Zeldis | | | | |
| 7,393,862 | B2 | 7/2008 | Zeldis | | | | |
| 7,435,745 | B2 | 10/2008 | D'Amato | | | | |
| 7,468,363 | B2 | 12/2008 | Zeldis | | | | |
| 7,968,569 | B2 * | 6/2011 | Zeldis | 514/323 | | | |
| 8,188,118 | B2 * | 5/2012 | Zeldis | 514/321 | | | |
| 8,198,262 | B2 | 6/2012 | Zeldis | | | | |
| 8,198,306 | B2 * | 6/2012 | Zeldis | 514/321 | | | |
| 8,207,200 | B2 * | 6/2012 | Zeldis | 514/321 | | | |
| 8,410,136 | B2 | 4/2013 | Zeldis | | | | |
| 8,440,194 | B2 | 5/2013 | Zeldis | | | | |
| 8,492,406 | B2 | 7/2013 | Zeldis | | | | |
| 8,530,498 | B1 * | 9/2013 | Zeldis | 514/320 | | | |
| 8,632,787 | B2 | 1/2014 | Zeldis | | | | |
| 8,648,095 | B2 * | 2/2014 | Zeldis | 514/321 | | | |
| 8,673,939 | B2 | 3/2014 | Zeldis | | | | |
| 8,759,375 | B2 | 6/2014 | Zeldis | | | | |
| 2001/0018445 | A1 | 8/2001 | Huang et al. | | | | |
| 2001/0056114 | A1 | 12/2001 | D'Amato | | | | |
| 2002/0035090 | A1 | 3/2002 | Zeldis et al. | | | | |
| 2002/0045643 | A1 | 4/2002 | Muller et al. | | | | |
| 2002/0052398 | A1 | 5/2002 | D'Amato | | | | |
| 2002/0054899 | A1 | 5/2002 | Zeldis | | | | |
| 2002/0061923 | A1 | 5/2002 | D'Amato | | | | |
| 2002/0128228 | A1 | 9/2002 | Hwu | | | | |
| 2002/0161023 | A1 | 10/2002 | D'Amato | | | | |
| 2002/0173658 | A1 | 11/2002 | Muller et al. | | | | |

OTHER PUBLICATIONS

- U.S. Appl. No. 60/499,723, Markian.
U.S. Appl. No. 60/372,348, Hariri et al.
U.S. Appl. No. 10/732,867, D'Amato et al.
U.S. Appl. No. 09/545,654, D'Amato.
U.S. Appl. No. 09/287,377, D'Amato.
Carstensen, 1995, *Drug Stability: Principles & Practice*, 2nd ed., Marcel Dekker, New York, NY pp. 379-380.
Corral et al., 1999, "Immunomodulation by thalidomide and thalidomide analogues," *Ann. Rheum. Dis.* 58(Suppl 1):1107-1113.
Craig et al., 1967, "Potential anticancer agents. III. 2-phthalimidoaldehydes and derivatives," *Potential Anticancer Agents III* 10:1071-1073.
D'Amato et al., 2001, "Mechanism of action of thalidomide and 3-aminothalidomide in multiple myeloma," *Semin Oncol.* 28:597-601.
D'Amato et al., 1994, "Thalidomide is an Inhibitor of Angiogenesis," *Proc. Natl. Acad. Sci.* 91:4082-4085.
De et al., 1976, "Hansch analysis for some antineoplastic glutarimides," *J. Indian Chem. Soc.* I.III: 825-826.
De et al., 1976, "Possible antineoplastic agents: III. Synthesis of 6-alkyl-2-[4'-methoxyphthalimido] and 6-alkyl-3-[3'-4'-dimethoxyphenyl] glutarimides," *J. Indian Chem. Soc.* I.III: 1122-1125.
Dredge et al., 2002, "Novel thalidomide analogues display anti-angiogenic activity independently of immunomodulatory effects," *Br. J. Cancer* 87(10):1166-1172.
Folkman et al., 1983, "Angiogenesis inhibition and tumor regression caused by heparin or a heparin fragment in the presence of cortisone," *Science* 221(4612):719-725.
Gershbein, 1991, "The thalidomide analog, EM 12, enhances 1,2-dimethylhydrazine-induction of rat colon adenocarcinomas," *Cancer Letters* 60: 129-133.
Grabstald et al., 1965, "Clinical experiences with thalidomide in patients with cancer," *Clinical Pharmacology and Therapeutics* 6:298-302.
Lentzsch et al., 2003, "Immunomodulatory analogs of thalidomide inhibit growth of Hs Sultan cells and angiogenesis in vivo," *Leukemia* 17(1):41-44.
Lentzsch et al., 2002, "S-3-amino-phthalimido-glutarimide inhibits angiogenesis and growth of B-cell neoplasias in mice", *Cancer Research* 62:2300-2305.
Miyachi et al., 1997, "Novel biological response modifiers: phthalimides with tumor necrosis factor-alpha production-regulating activity," *J. Med. Chem.* 40:2858-2865.
Muller et al., 1999, "Amino-substituted thalidomide analogs: potent inhibitors of TNF-alpha production," *Bioorg. Med. Chem. Lett.* 9(11):1625-1630.
Muller et al., 1998, "Thalidomide analogs and PDE4 inhibition," *Bioorg. Med. Chem. Lett.* 8(19):2669-2674.

US 9,101,622 B2

Page 3

(56)

References Cited

OTHER PUBLICATIONS

- Muller et al., 1996, "Structural modifications of thalidomide produce analogs with enhanced tumor necrosis factor inhibitory activity," *J. Med. Chem.* 39(17):3238-3240.
- Olson et al., 1965, "Thalidomide (N-phthaloylglutamimide) in the treatment of advanced cancer," *Clinical Pharmacology and Therapeutics* 6(3):292-297.
- Penichet et al., 2001, "Antibody-cytokine fusion proteins for the therapy of cancer," *J. Immunol. Methods* 248(1-2):91-101.
- Physician's Desk Reference*, 2002, 56th ed., pp. 1755-1760.
- Raza et al., 2001, "Thalidomide produces transfusion independence in long-standing refractory anemias of patients with myelodysplastic syndromes," *Blood* 98(4):958-965.
- Shah et al., 1999, "Synthesis and enantiomeric separation of 2-phthalimidino-glutaric acid analogues: potent inhibitors of tumor metastasis," *J. Med. Chem.* 42:3014-3017.
- Shibata et al., 1995, "N-alkylphthalimides: structural requirement of thalidomide action on 12-O-tetradecanoylphorbol-13-acetate-induced tumor necrosis factor production by human leukemia HL-60 cells," *Chem. Pharm. Bull.* 43(1):177-179.
- Shimazawa et al., 1999, "Antiangiogenic activity of tumor necrosis factor-alpha production regulators derived from thalidomide," *Biol. Pharm. Bull.* 22(2):224-226.
- Rubin et al., "Principles of Cancer Treatment—1", 12 ONCO IV 1, May 2003.
- Wilens et al., 1977, *Tetrahedron* 33:2725.
- Wilens, 1972, *Tables of Resolving Agents and Optical Resolutions*, E.L. Eliel, ed., Univ. of Notre Dame Press, Notre Dame, IN pp. 268.
- Wolff ed., 1995, *Burger's Medicinal Chemistry and Drug Discovery*, 5th ed., pp. 172-178, 949-982.
- N. Ake Jonsson, 1972, "Chemical Structure and Teratogenic Properties," *Acta Pharm.*, pp. 521-542.
- Alexanian et al., 2004, "VTD (Velcade, thalidomide, dexamethasone) as primary therapy for newly-diagnosed multiple myeloma," *Am. Soc. Hematol. 46th Ann. Meeting Dec. 4-7, 2004, San Diego, CA Abstract #210.*
- Anderson, 2000, "Thalidomide: Therapeutic potential in hematologic malignancies," *Seminars in Hematology* 37(1 Supp 3): 1-4.
- Attal et al., 2004, "Maintenance treatment with thalidomide after autologous transplantation for myeloma: First analysis of a prospective randomized study of the Intergroupe Francophone du Myelome (IFM 99 02)," *Am. Soc. Hematol. 46th Ann. Meeting Dec. 4-7, 2004, San Diego, CA Abstract #535.*
- Bemardeschi et al., 2003, *J. Exp. Clin. Cancer Res.* 22(4):129-133.
- Corral et al., 1999, "Differential cytokine modulation and T cell activation by two distinct classes of thalidomide analogues that are potent inhibitors of TNF-alpha," *J. Immunol.* 163(1):380-386.
- Davies et al., 2001, "Thalidomide and immunomodulatory derivatives augment natural killer cell cytotoxicity in multiple myeloma," *Blood* 98(1):210-216.
- Dimopoulos et al., 2004, "Primary treatment with pulsed melphalan, dexamethasone, thalidomide (MDT) for symptomatic patients with multiple myeloma ≥ 75 years of age," *Am. Soc. Hematol. 46th Ann. Meeting Dec. 4-7, 2004, San Diego, CA Abstract #1482.*
- Eisen et al., 2000, "Continuous low dose Thalidomide: a phase II study in advanced melanoma, renal cell, ovarian and breast cancer," *Br. J. Cancer* 82(4):812-817.
- Fakhouri et al., 2004, "Thalidomide in patients with multiple myeloma and renal failure," *Br. J. Haematol.* 125:90-102.
- Fenk et al., 2005, "Single-agent thalidomide for treatment of first relapse following high-dose chemotherapy in patients with multiple myeloma," *Leukemia* 19(1):156-159.
- Gupta et al., 2001, "Adherence of multiple myeloma cells to bone marrow stromal cells upregulates vascular endothelial growth factor secretion: therapeutic applications," *Leukemia* 15(12):1950-1961.
- Haslett et al., 2003, "Thalidomide and a thalidomide analogue drug costimulate virus-specific CD8+ T cells in vitro," *J. Infect. Dis.* 187(6):946-955.
- Hideshima et al., 2000, "Thalidomide and its analogs overcome drug resistance of human multiple myeloma cells to conventional therapy," *Blood* 96(9):2943-2950.
- Offidani et al., 2003, "Thalidomide plus oral melphalan for advanced multiple myeloma: a phase II study. *Haematologica*. Dec. 2003;88(12):1432-1433.
- Palumbo et al., 2004, "A prospective randomized trial of oral melphalan prednisone, thalidomide (MPT) vs. oral melphalan, prednisone (MP): an interim analysis," *Am. Soc. Hematol. 46th Ann. Meeting Dec. 4-7, 2004, San Diego, CA Abstract #207.*
- Raje et al., 1999, "Thalidomide—a revival story," *N. Engl. J. Med.* 341(21):1606-1609.
- Rajkumar et al., 2004, "Thalidomide plus dexamethasone versus dexamethasone alone in newly diagnosed multiple myeloma (E1A00): Results of a phase III trial coordinated by the Eastern Cooperative Oncology Group," *Am. Soc. Hematol. 46th Ann. Meeting Dec. 4-7, 2004, San Diego, CA Abstract #205.*
- Rajkumar et al., 2000, "Prognostic value of bone marrow angiogenesis in multiple myeloma," *Clin. Cancer Res.* 6(8):3111-3116.
- Ribatti et al., 1999, "Bone marrow angiogenesis and mast cell density increase simultaneously with progression of human multiple myeloma," *Br. J. Cancer* 79(3-4):451-455.
- Singhal et al., 1999, "Antitumor activity of thalidomide in refractory multiple myeloma," *N. Engl. J. Med.* 341(21):1565-1571.
- Steins et al., 2002, "Efficacy and safety of thalidomide in patients with acute myeloid leukemia," *Blood* 99(3):834-839.
- Vacca et al., 1999, "Bone marrow neovascularization, plasma cell angiogenic potential, and matrix metalloproteinase-2 secretion parallel progression of human multiple myeloma," *Blood* 93(9):3064-3073.
- Wohrer et al., 2004, "Effective treatment of primary plasma cell leukemia with thalidomide and dexamethasone—a case report," *Hematol. J.* 5(4):361-363.
- Bach, 1963, "Thalidomide in Cancer Chemotherapy," *The Lancet*, No. 1271, p. 71.
- Bach, 1963, "Studies on the Possible Anti-Neoplastic Effect of Thalidomide," *Acta Pathologica Et Microbiologica Scandinavica* 59:491-499.
- Chandhry, 1966, *Cancer Research*, "Effect of Prednisolone and Thalidomide on Induced Submandibular Gland Tumors in Hamster," 26(part 1):1884-86.
- DiPaolo, 1963, "Effect of Thalidomide on a Variety of Transplantable Tumors," *Cancer Chemotherapy Reports* No. 29, p. 99-102.
- DiPaolo, 1963, "In vitro Test Systems for Cancer Chemotherapy. II. Correlation of in vitro Inhibition of Dehydrogenase and Growth with in vivo Inhibition of Ehrlich Ascites Tumor," *Proceedings of the Society for Experimental Biology & Medicine*, 114:384-387.
- DiPaolo, 1964, "Thalidomide: Effects on Ehrlich Ascites Tumor Cells in vitro" *Science* 144:1583.
- Maud, 1963, "Clinical Improvements Obtained in Advanced Cancer Patients with Treatment with Thalidomide Associated with Hormones," *Anais Paulistas de Medicina e Cirurgia* 86:13-40.
- Roe and Mitchley, 1963, "Thalidomide and Neoplasia" *Nature* 200:1016-1017.
- Liu et al., "Phase I study of CC-5013 (Revimid), a thalidomide derivative, in patients with refractory metastatic cancer," *American Society of Clinical Oncology*, Abstract #927, 2003.
- Zangari et al., "Results of phase I study of CC-5013 for the treatment of multiple myeloma (MM) patients who relapse after high dose chemotherapy (HDCT)," *American Society of Hematology, Abstract #3226*, 2001.
- Zeldis et al., "Update on the evolution of the IMiD™," *International Society for Biological Therapy of Cancer, Oral Abstract*, 2003.
- Anderson, "Moving disease biology from the laboratory to the clinic," *Seminars in Oncology*, 2002 29:17-20.
- Barlogie et al., "Total Therapy II (TTII) for newly diagnosed multiple myeloma (MM): preliminary data on feasibility and efficacy in the first 231 enrolled patients; comparison with predecessor trial total therapy I (TTI) (N=231)," *Blood, Abstract # 2857*, Dec. 7-11, 2001, *American Society of Hematology*.
- Barlogie et al., "High-dose therapy immunomodulatory drugs in multiple myeloma," *Seminars in Oncology*, 2002, 29 (6):26-33.

(56) **References Cited**

OTHER PUBLICATIONS

Barlogie et al., "Introduction: Thalidomide and the IMiDs in multiple myeloma," *Seminars in Hematology*, 2003, 40 (4):1-2.

Barlogie, "Thalidomide and CC-5013 in Multiple Myeloma: The University of Arkansas experience," *Seminars in Hematology*, 2003, 40 (4):33-38.

Bartlett et al., "The evolution of thalidomide and its IMiD derivatives as anticancer agents," *Nature Reviews Cancer*, 2004, 4 (4):1-9.

Bartlett et al., "Phase I study to determine the safety, tolerability and immunostimulatory activity of thalidomide analogue CC-5013 in patients with metastatic malignant melanoma and other advanced cancers," *British Journal of Cancer*, 2004, 90:955-961.

Battegay, "Angiogenesis: mechanistic insights, neovascular diseases, and therapeutic prospects," *J. Mol. Med.*, 1995, 73:333-346.

Baz et al., "Doxil (D), vincristine (V), reduced frequency dexamethasone (d) and revlimid (R) (DVd-R) results in a high response rate in patients with refractory multiple myeloma (RMM)," *Blood, Abstract # 2559, American Society of Hematology*, Dec. 10-13, 2005.

Brennen et al., "Thalidomide and analogues: current proposed mechanisms and therapeutic usage," *Clinical Prostate Cancer*, 2004, 3 (1):54-61.

Celgene Corporation, "Celgene advances immunomodulatory drug (IMiD™) clinical program," Press Release, Feb. 2000.

Celgene Corporation, "Initial Phase I solid tumor data on Celgene's lead IMiD™, Revimid™," Press Release, Jun. 2001.

Celgene Corporation, "Celgene Corporation receives orphan drug designation for Revimid™ for multiple myeloma," Press Release, Oct. 2001.

Celgene Corporation, "Celgene Corporation announces third quarter results. Thalomid® (thalidomide) sales increase 24%. Prescriptions up 50%. Enhanced S.T.E.P.S.® launched. Pilot d-MPH data presented," Press Release, Oct. 2001.

Celgene Corporation, "Celgene expands clinical development program for Revimid™. Five additional trials of Revimid initiated in hematological and solid tumor cancers," Press Release, Jun. 2002.

Celgene Corporation, "Celgene Corporation announces third quarter results. Thalomid® (thalidomide) revenue increases 41% to \$30.5 million. Pivotal programs for Thalomid and Revimid™ finalized. Peer-reviewed publications of Thalomid and Revimid data. First JNK inhibitor advanced to Phase I clinical trial," Press Release, Oct. 2002.

Celgene Corporation, "Blood reports Revimid™ has anti-tumor activity in patients with relapsed and refractory multiple myeloma," Press Release, Nov. 1, 2002.

Celgene Corporation, "Celgene provides update on clinical pipeline. Celgene Announces first target indication for Actimid™, CC-8490. SelCID™ program to advance based on results from Phase I/II trial of CC-1088. First JNK inhibitor successfully completes phase I trial," Press Release, Jan. 2003.

Celgene Corporation, "Celgene Corporation announces fourth quarter and full year results for 2002," Press Release, Jan. 2003.

Celgene Corporation, "Celgene receives fast track status from FDA for Revimid™ in multiple myeloma," Press Release, Feb. 2003.

Celgene Corporation, "Celgene receives fast track status from FDA for Revimid™ in myelodysplastic syndromes," Press Release, Apr. 2003.

Celgene Corporation, "New Revimid™ clinical data shows potential as novel approach to treating myelodysplastic syndromes (MDS)," Press Release, May 2003.

Celgene Corporation, "Celgene corporation reports strong operating performance in second quarter as total sales increase 100 percent and profits rise," Press Release, Jul. 2003.

Celgene Corporation, "Celgene corporation reports record operating performance in third quarter as total revenue increases 117% and profits rise," Press Release, Oct. 2003.

Celgene Corporation, "Celgene corporation advances Actimid™ (CC-4047) into phase II trial for prostate cancer," Press Release, Oct. 2003.

Celgene Corporation, "Additional clinical data presented on Revimid™ in myelodysplastic syndromes at the American Society of Hematology 45th annual meeting," Press Release, Dec. 2003.

Celgene Corporation, "Celgene corporation reviews 2003 achievements and announces 2004 financial outlook," Press Release, Jan. 2004.

Celgene Corporation, "Revlimid™ receives orphan drug designation from the European commission for multiple myeloma," Press Release, Feb. 2004.

Celgene Corporation, "Revlimid™ receives orphan drug designation from the European commission for myelodysplastic syndromes," Press Release, Mar. 2004.

Celgene Corporation, "Celgene corporation reports record operating performance in first quarter with strong revenue growth and profits," Press Release, Apr. 2004.

Celgene Corporation, "Celgene announces plans to stop phase III trials in melanoma due to lack of efficacy," Press Release, Apr. 2004.

Dalgleish, et al., "New thalidomide analogues; anti-cancer, anti-angiogenic and immunostimulatory," *British Journal of Cancer*, 2001, 85 (1)25.

Dalgleish et al., "Thalidomide analogues CC-5013 and CC-4047 induce T cell activation and IL-12 production in patients with both solid tumours and relapsed and refractory multiple myeloma," *British Journal of Cancer*, 2003, 88(Suppl 1), S25-S54.

Davies et al., "Thalidomide (Thal) and immunomodulatory derivatives (IMiDs) augment natural killer (NK) cell cytotoxicity in multiple myeloma(MM)," Abstract # 3617, *American Society of Hematology*, Dec. 1-5, 2000.

Davies et al., "Thalidomide (Thal) and immunomodulatory derivatives (IMiDs) augment natural killer (NK) cell cytotoxicity in multiple myeloma ~MM," Abstract # P222, *VIIIth International Myeloma Workshop*, May 4-8, 2001.

Dibbs et al., "Thalidomide and thalidomide analogs suppress TNF α secretion by myocytes," Abstract # 1284, *Circulation*, 1998.

Dimopoulos et al., "Results of thalidomide and IMiDs in multiple myeloma," Abstract # P12.1.4, *International Multiple Myeloma Workshop*, May 23-27, 2003.

Dimopoulos et al., "Treatment of plasma cell dyscrasias with thalidomide and its derivatives," *Journal of Clinical Oncology*, Dec. 1, 2003, 21 (23)4444-4454.

Dimopoulos et al., "Study of lenalidomide plus dexamethasone versus dexamethasone alone in relapsed or refractory multiple myeloma (MM): Results of a phase 3 Study (MM-010)," Abstract # 6, *American Society of Hematology*, Dec. 10-13, 2005.

Dredge et al., "A costimulatory thalidomide analog enhances the partial anti-tumor immunity of an autologous vaccination in a model of colorectal cancer," Abstract # 491, *American Association for Cancer Research*, Apr. 6-10, 2002.

Dredge et al., "Adjuvants and the promotion of Th1-type cytokines in tumour immunotherapy," *Cancer Immunol. Immunother.*, 2002, 51:521-531.

Dredge et al., "Immunological effects of thalidomide and its chemical and functional analogs," *Critical Reviews in Immunology*, 2002, 22 (5&6):425-437.

Dredge et al., "Protective antitumor immunity induced by a costimulatory thalidomide analog in conjunction with whole tumor cell vaccination is mediated by increased Th1-type immunity¹," *The Journal of Immunology*, 2002, 168:4914-4919.

Dredge et al., "Recent developments in antiangiogenic therapy," *Expert Opin. Biol. Ther.*, 2002, 2 (8):953-966.

Dredge et al., "Angiogenesis inhibitors in cancer therapy," *Current Opinion in Investigational Drugs*, 2003, 4 (6):667-674.

Dredge et al., "Thalidomide analogs as emerging anti-cancer drugs," *Anti-Cancer Drugs*, 2003, 14:331-335.

Fickentscher et al., "Stereochemical properties and teratogenic activity of some tetrahydrophthalimides," *Molecular Pharmacology*, 1976, 13:133-141.

Figg et al., "Inhibition of angiogenesis: treatment options for patients with metastatic prostate cancer," *Investigational New Drugs*, 2002, 20(2):183-194.

Galustian et al., "Thalidomide-derived immunomodulatory drugs as therapeutic agents," *Expert Opin. Biol. Ther.*, 2004, 4 (12):1-8.

(56)

References Cited

OTHER PUBLICATIONS

- Glaspy et al., "The potential role of thalidomide and thalidomide analogs in melanoma," *Clinical Advances in Hematology & Oncology*, 2004, 1-7.
- Gupta et al., "Adherence of multiple myeloma cells to bone marrow stromal cells upregulates vascular endothelial growth factor secretion: therapeutic applications," *Leukemia*, 2001, 15:1950-1961.
- Hayashi et al., "Mechanisms whereby immunomodulatory analogs of thalidomide augment autologous NK cell anti-myeloma immunity," *Blood, Abstract #3219*, Dec. 6-10, 2002, *American Society of Hematology*.
- He, W., et al., 1993, Abstract of papers, 206th American Chemical Society, Chicago, IL; Med. Chem., paper 216.
- Helm et al., "Comparative teratological investigation of compounds of structurally and pharmacologically related to thalidomide," *Arzneimittel Forschung/Drug Research*, 1981, 31 (I)941-949.
- Hernandez-Illizalitur et al., "Addition of immunomodulatory drugs CC5013 or CC4047 to rituximab enhances anti-tumor activity in a severe combined immunodeficiency (SCID) mouse lymphoma model," *Abstract # 235*, *American Society of Hematology*, Dec. 6-9, 2003.
- Hideshima et al., "Thalidomide and its analogs overcome drug resistance of human multiple myeloma cells to conventional therapy," *Blood*, 2000, 96:2943-2950, *American Society of Hematology*.
- Hideshima et al., "Thalidomide (Thal) and its analogs overcome drug resistance of human multiple myeloma (MM) cells to conventional therapy," *Abstract 1313*, *American Society of Hematology*, Dec. 1-5, 2000.
- Hunt et al., "Markers of endothelial and haemostatic activation in the use of CC-4047, a structural analogue of thalidomide, in relapsed myeloma," *Blood, Abstract # 3216*, Dec. 6-10, 2002, *American Society of Hematology*.
- Hussein et al., "Doxil (D), vincristine (V), reduced frequency dexamethasone (d) and Revlimid (DVd-R) a phase I/II trial in advanced relapsed/refractory multiple myeloma (Rmm) patients," *Blood, Abstract #208*, *American Society of Hematology*, Dec. 4-7, 2004.
- Hwu et al., "Thalidomide and its analogues in the treatment of metastatic melanoma," *Chemotherapy Foundation Symposium, Abstract #44*, 2002.
- Kyle, "Current therapy of multiple myeloma," *Internal Medicine*, 2002, 41 (3)175-180.
- Kyle et al., "Multiple myeloma," *New England Journal of Medicine*, 2004, 351:1860-1873.
- Leblanc et al., "Immunomodulatory drug costimulates T cells via the B7-CD28 pathway," *Blood*, 2004, 103:1787-1790, *American Society of Hematology*.
- Lentzsch et al., "In vivo activity of thalidomide and immunomodulatory drugs against multiple myeloma," *VIIIth International Myeloma Workshop, Abstract #P225*, May 4-8, 2001.
- Lentzsch et al., "Immunomodulatory derivative of thalidomide (IMiD CC-4047) determine the lineage commitment of hematopoietic progenitors by down regulation of GATA-1 and modulation of cytokine secretion," *Abstract # 3073*, *American Society of Hematology*, Dec. 6-9, 2003.
- Lentzsch et al., "Immunomodulatory derivative of thalidomide (IMiD CC-4047) down regulates CAAT/enhancer-binding protein β (C/EBP β) in multiple myeloma (MM)," *Abstract # 3456*, *American Society of Hematology*, Dec. 6-9, 2003.
- Luzzio et al., "Thalidomide analogues: derivatives of an orphan drug with diverse biological activity," *Expert Opin. Ther. Patents*, 2004, 14 (2):215-229.
- Man et al., " α -Fluoro-substituted thalidomide analogues," *Bioorganic & Medicinal Chemistry Letters* 13, 2003, 3415-3417.
- Marriott et al., "Immunotherapeutic and antitumor potential of thalidomide analogues," *Expert Opin. Biol. Ther.*, 2001, 1 (4):1-8.
- Marriott et al., "New thalidomide analogues; anti-cancer, anti-angiogenic and immunostimulatory," *British Journal of Cancer*, 85:25, Jul. 6, 2001.
- Marriott et al., "Thalidomide and its analogues have distinct and opposing effects on TNF- α and TNFR2 during co-stimulation of both CD4⁺ and CD8⁺ T cells," *Clin. Exp. Immunol.*, 2002, 130:75-84.
- Marriott et al., "A novel subclass of thalidomide analogue with anti-solid tumor activity in which caspase-dependent apoptosis is associated with altered expression of bcl-2 family proteins," *Cancer Research*, 2003, 63:593-599.
- Marriott et al., "Thalidomide derived immunomodulatory drugs (IMiDs) as potential therapeutic agents," *Current Drug Targets—Immune, Endocrine & Metabolic Disorders*, 2003, 3:181-186.
- Masellis et al., "Changes in gene expression in bone marrow mesenchymal progenitor cells as a consequence of IMiD therapy in multiple myeloma patients," *Blood, Abstract # 1548*, Dec. 7-11, 2001, *American Society of Hematology*.
- McCarty, "Thalidomide may impede cell migration in primates by down-regulating integrin β -chains: potential therapeutic utility in solid malignancies, proliferative retinopathy, inflammatory disorders, neointimal hyperplasia, and osteoporosis," *Medical Hypotheses*, 1997, 49:123-131.
- Mitsiades et al., "Apoptotic signaling induced by immunomodulatory thalidomide analogs (Imids) in human multiple myeloma cells: therapeutic implications," *Abstract # 3224*, Dec. 7-11, 2001, *American Society of Hematology*.
- Mitsiades et al., "Apoptotic signaling induced by immunomodulatory thalidomide analogs in human multiple myeloma cells: therapeutic implications," *Blood*, 2002, 99:4525-4530, *American Society of Hematology*.
- Mitsiades et al., "CC-5013 Celgene," *Current Opinion in Investigational Drugs*, 2004, 5 (6):635-647.
- Moutouh et al., "Novel immunomodulatory drugs (IMiDs®): A potential, new therapy for β -hemoglobinopathies," *Abstract # 3740*, *American Society of Hematology*, Dec. 4-7, 2004.
- Patten et al., "The early use of the serum free light chain assay in patients with relapsed refractory myeloma receiving treatment with a thalidomide analogue (CC-4047)," *Abstract # 1640*, *American Society of Hematology*, Dec. 6-9, 2003.
- Payvandi et al., "Effects of a thalidomide analog on binding activity of transcription factors and cell cycle progression of multiple myeloma cell lines," *Blood, Abstract #2487*, Dec. 1-5, 2000, *American Society of Hematology*.
- Payvandi et al., "The thalidomide analogs IMiDs enhance expression of CD69 stimulatory receptor on natural killer cells," *Abstract # 1793*, *American Association for Cancer Research*, Mar. 24-28, 2001.
- Payvandi et al., "Thalidomide analogs IMiDs inhibit expression of cyclooxygenase-2 in multiple myeloma cell line and LPS stimulated PBMCs," *Blood, Abstract # 2689*, Dec. 7-11, 2001, *American Society of Hematology*.
- Payvandi et al., "Thalidomide and IMiDs inhibit microvessel formation from human arterial rings in the absence of human liver microsomes," *Blood, Abstract # 5046*, Dec. 6-10, 2002, *American Society of Hematology*.
- Payvandi et al., "CC-5013 inhibits the expression of adhesion molecules ICAM-1 and CD44 and prevents metastasis of B16 F10 mouse melanoma cells in an animal model," *American Society of Clinical Oncology, Abstract # 992*, 2003.
- Payvandi et al., "Immunomodulatory drugs inhibit expression of cyclooxygenase-2 from TNF- α , IL-1 β , and LPS-stimulated human PBMC in a partially IL-10-dependent manner," *Cellular Immunology*, 2004, 81-88.
- Raje et al., "Combination of the mTOR inhibitor rapamycin and CC-5013 has synergistic activity in multiple myeloma," *Blood*, Dec. 15, 2004, 104 (13)4188-4193.
- Rajkumar et al., "Combination therapy with lenalidomide plus dexamethasone (Rev/Dex) for newly diagnosed myeloma," *Blood*, Dec. 15, 2005, 106 (13)4050-4053.
- Richardson et al., "A Phase I study of oral CC5013, an immunomodulatory thalidomide (Thal) derivative, in patients with relapsed and refractory multiple myeloma (MM)," *Blood, Abstract #3225*, Dec. 7-11, 2001, *American Society of Hematology*.
- Richardson et al., "Immunomodulatory drug CC-5013 overcomes drug resistance and is well tolerated in patients with relapsed multiple myeloma," *Blood*, 2002 100:3063-3067, *American Society of Hematology*.

US 9,101,622 B2

Page 6

(56)

References Cited

OTHER PUBLICATIONS

- Richardson et al., "A multi-center, randomized, phase 2 study to evaluate the efficacy and safety of 2 CDC-5013 dose regimens when used alone or in combination with dexamethasone (Dex) for the treatment of relapsed or refractory multiple myeloma (MM)," *Blood, Abstract # 825, American Society of Hematology*, Dec. 6-9, 2003.
- Richardson et al., "Immunomodulatory analogs of thalidomide: an emerging new therapy in myeloma," *Journal of Clinical Oncology*, 2004, 22(16) 3212-3214.
- Richardson et al., "A multicenter, single-arm, open-label study to evaluate the efficacy and safety of single-agent lenalidomide in patients with relapsed and refractory multiple myeloma; preliminary results," *10th International Myeloma Workshop*, Apr. 10-14, 2005.
- Richardson et al., "Novel biological therapies for the treatment of multiple myeloma," *Best Practice & Research Clinical Haematology*, 2005, 18 (4):619-634.
- Richardson et al., "A phase I trial of lenalidomide (Revlimid®) with bortezomib (Velcade®) in relapsed and refractory multiple myeloma," *Blood, Abstract # 365, American Society of Hematology*, Dec. 10-13, 2005.
- Rubin et al., "Principles of cancer treatment-1," 2003, 12 ONCO IV 1.
- Schafer et al., "Enhancement of cytokine production and AP-1 transcriptional activity in T cells by thalidomide-related immunomodulatory drugs," *Journal of Pharmacology and Experimental Therapeutics*, 2003, 305(3)1222-1232.
- Schey et al., "A phase I study of an immunomodulatory thalidomide analog, CC-4047, in relapsed or refractory multiple myeloma," *Journal of Clinical Oncology*, 2004, 22 (16):1-8.
- Schey et al., "A phase I study of an immunomodulatory thalidomide analogue (CC4047) in relapse/refractory multiple myeloma," *International Society for Experimental Hematology, Abstract #248*, 2002.
- Shaughnessy et al., "Global gene expression analysis shows loss of C-MYC and IL-6 receptor gene mRNA after exposure of myeloma to thalidomide and IMiD," *Abstract # 2485, The American Society of Hematology*, Dec. 1-5, 2000.
- Shire et al., "TNF- α inhibitors and rheumatoid arthritis," *Exp. Opin. Ther. Patents*, 1998, 8 (5):531-544.
- Sorbera et al., "CC-5013. Treatment of multiple myeloma. Treatment of Melanoma. Treatment of myelodysplastic syndrome. Angiogenesis inhibitor. TNF- α production inhibitor," *Drugs of the Future*, 2003, 28(5):425-431.
- Streety et al., "Thalidomide analogue CC-4047 is effective in the treatment of patients with relapsed and refractory multiple myeloma (MM) and induces T-cell activation and IL-12 production," *Abstract # 367, International Multiple Myeloma Workshop*, May 23-27, 2003.
- Streety et al., "Changes in neutrophil phenotype following the administration of CC-4047 (Actimid) to patients with multiple myeloma," *Abstract # 2543, American Society of Hematology*, Dec. 6-9, 2003.
- Streety et al., "An update of the use and outcomes of the new immunomodulatory agent CC-4047 (Actimid) in patients with relapsed/refractory myeloma," *Abstract #829, American Society of Hematology*, Dec. 6-9, 2003.
- Teo et al., "A phase I, single-blind, placebo-controlled, ascending single oral dose, safety, tolerability and pharmacokinetic study of CDC-501, a novel immunomodulatory-oncologic agent, in healthy male subjects with a comparison of fed and fasted," *Clinical Pharmacology and Therapeutics*, 2002, 71 (2)93.
- Teo et al., "Chiral inversion of the second generation IMiD™ CC-4047 (Actimid™) in human plasma and phosphate-buffered saline," *Chirality*, 2003, 15:348-351.
- Theutulien et al., "Hybrid MEL/DT PACE autotransplant regimen for Multiple Myeloma (MM)—safety and efficacy data in pilot study of 15 patients," *Blood, Abstract # 2869, American Society of Hematology*, Dec. 7-11, 2001.
- Tohny et al., "A phase I study of oral CC-5013 (lenalidomide, Revlimid™), a thalidomide derivative, in patients with refractory metastatic cancer," *Clinical Prostate Cancer*, 2004, 2:241-243.
- Tricot et al., "Angiochemotherapy (ACT) for multiple myeloma (MM) with DT-PACE results in a high response rate, but in contrast to tandem transplants with melphalan does not affect durable disease control," *Blood, Abstract # 3531, American Society of Hematology*, Dec. 7-11, 2001.
- Tsenova et al., "Use of IMiD3, a thalidomide analog, as an adjunct to therapy for experimental tuberculous meningitis," *Antimicrobial Agents and Chemotherapy*, 2002, 46 (6)1887-1895.
- Weber, "Lenalidomide (CC-5013, Revlimid™) and other ImiDs," *Abstract # PL5.02, International Multiple Myeloma Workshop*, Apr. 10-14, 2005.
- Weber et al., "A multicenter, randomized, parallel-group, double-blind, placebo-controlled study of lenalidomide plus dexamethasone versus dexamethasone alone in previously treated subjects with multiple myeloma," *Abstract # PO.738, International Multiple Myeloma Workshop*, Apr. 10-14, 2005.
- Ye et al., "Novel IMiD drugs enhance expansion and regulate differentiation of human cord blood CD34+ cells with cytokines," *Blood, Abstract #4099, American Society of Hematology*, Dec. 6-10, 2002.
- Zangari et al., "Risk factors for deep vein thrombosis (DVT) in a large group of myeloma patients (Pts) treated with thalidomide (Thal): The Arkansas Experience," *Blood, Abstract # 681, American Society of Hematology*, Dec. 7-11, 2001.
- Zangari et al., "Revimid 25 mg (REV 25) x 20 versus 50 mg (REV 50) x 10 q 28 days with bridging of 5 mg x 10 versus 10 mg x 5 as post-transplant salvage therapy for multiple myeloma (MM)," *Blood, Abstract # 1642, American Society of Hematology*, Dec. 6-9, 2003.
- Zeldis et al., "Potential new therapeutics for Waldenstrom's macroglobulinemia," *Seminars in Oncology*, 2003, 30 (2):275-281.
- Zhang et al., "CC-5079, a novel microtubule and TNF- α inhibitor with anti-angiogenic and antimetastasis activity," *Abstract # B012, International Conference on Molecular Targets and Cancer Therapeutics*, Nov. 17-21, 2003.
- Anderson, "The Role of Immunomodulatory Drugs in Multiple Myeloma," *Seminars in Hematology*, vol. 40, No. 4, Suppl 4, 2003: pp. 23-32.
- Weber, "Thalidomide and Its Derivatives: New Promise for Multiple Myeloma," *Cancer Control*, vol. 10, No. 5, 375-383, 2003.
- Patt, Yehuda A.; Iiassan, Manal M.; Lozano, Richard D.; Ellis, Lee M.; Peterson, J. Andrew; Waugh, Kimberly A.; Durable Clinical Response of Refractory Hepatocellular Carcinoma to Orally Administered Thalidomide. *American Journal of Clinical Oncology*, 2000.
- Richardson, Paul; Hideshima, Tem; Anderson, Kenneth; Thalidomide: The Revival of a Drug with Therapeutic Promise in the Treatment of Cancer; *Principles & Practice of Oncology*, vol. 15, No. 2, 2001.
- Thomas, Melodic; Doss, Deborah, Thalidomide Nursing Roundtable Update, Monograph, Sep. 2002.
- Richardson, Paul; Hideshima, Teru; Anderson, Kenneth; Thalidomide: Emerging Role in Cancer Medicine; *Annual Review of Medicine*, 2002.
- Berenson, J.R.; Bergsagel, P. L.; Munshi, N.; Initiation and Maintenance of Multiple Myeloma; *Seminars in Hematology*, vol. 36, No. 1, Supp. 3, Jan. 1999, pp. 9-13.
- Gollob, J.A.; Schinpper, C.P.; Orsini, E.; Murphy, E.; Daley, J.F.; Lazo, S.B.; Frank, D.A.; Characterization of a Novel Subset of CD8 T Cells That Expands in patients Receiving Interleukin-12, 02, *Am. Soc. For Clin. Investigation, Inc.*, vol. 102, No. 3, Aug. 1998, pp. 561-575.
- Cavanagh, L.L.; Barnetson, R.S.; Basten, A.; Halliday, G.M.; Dendritic Epidermal T-Cell Involvement in Induction of CD8+ T-Cell-Mediated Immunity Against an Ultraviolet Radiation-Induced Skin Tumor *Int. J. Cancer*: 70, 98-105, 1997.
- Thomas, D.A., Aguayo, A., Estey, E., Albitar, M., O'Brien, S., Giles, F.J., Beran, M., Cortes, J., Zeldis, J., Keating, M.J., Barlogie, B., Kantarjian, H.M., Thalidomide as anti-angiogenesis therapy (rx) in refractory or relapsed leukemia. *Abstract #2269, American Society of Hematology*, Dec. 3-7, 1999.
- Barlogie, B., Desikan, R., Munshi, N., Siegel, D., Mehta, J., Singhal, S., Anaissie, E., Single Course D.T. Pace Anti-Angiochemotherapy Effects CR in Plasma Cell Leukemia and Fulminant Multiple Myeloma (MM). *Abstract #4180, American Society of Hematology*, Dec. 4-9, 1998.

(56)

References Cited

OTHER PUBLICATIONS

- Hideshima, T., Chauhan, D., Shima, Y., Noopur, R., Davies, F.E., Tai, Y., Treon, S.P., Lin, B.K., Schlossman, R.L., Richardson, P.G., Gupta, D., Muller, G.W., Stirling, D.I., Anderson, K.C., Thalidome (THAL) and its Analogs Overcome Drug Resistance of Human Multiple Myeloma (MM) Cells to Conventional Therapy. Abstract #1313. American Society of Hematology, Dec. 1-5, 2000.
- Payvandi, F., Wu, L., Gupta, D., Hideshima, T., Haley, M., Muller, G., Chen, R., Anderson, K.C., Stirling, D., Effects of a Thalidomide Analog on Binding Activity of Transcription Factors and Cell Cycle Progression of Multiple Myeloma Cell Lines. Abstract #2487. American Society of Hematology, Dec. 1-5, 2000.
- Davies, F.E., Rajc, N., Hideshima, T., Lentzsch, S., Young, G., Tai, Y., Lin, B.K., Podar, K., Chauhan, D., Treon, S.P., Gupta, D., Mitsiades, C., Mitsiades, N., Hayashi, T., Richardson, P.G., Schlossman, R.L., Muller, G.W., Stirling, D. I., Anderson, K.C., Thalidomide (THAL) and Immunomodulatory Derivatives (IMiDS) Augment Natural Killer (NK) Cell Cytotoxicity in Multiple Myeloma (MM). Abstract #3617. American Society of Hematology, Dec. 15, 2000.
- Hideshima, T., Chauhan, D., Castro, A., Hayashi, T., Mitsiades, C., Mitsiades, N., Akiyama, M., Richardson, P.G., Schlossman, R.L., Adams, J., Anderson, K.C., NF-KB as a Therapeutic Target in Multiple Myeloma (MM). Abstract #1581. American Society of Hematology, Dec. 7-11, 2001.
- Lentzsch, S., Rogers, M., Leblanc, R., Birsner, A., Shah, J., Anderson K., D'Amato R., 3-Amino-Phthalimido-Glutarimide (S-3APG) Inhibits Angiogenesis and Growth in Drug Resistant Multiple Myeloma (MM) in vivo. Abstract #1976, American Society of Hematology, Dec. 7-11, 2001.
- Park, Y., Kim, S.A., Kim, C.J., Chung, J.H., Mechanism of the Effect of Thalidomide on Human Multiple Myeloma Cells. Abstract #2685. American Society of Clinical Oncology, May 12-17, 2001.
- Payvandi, F., Wu, L., Haley M., Gupta, D., Zhang, L., Schafer, P., Muller, G.W., Chen, R., Anderson, K.C., Stirling, D., Thalidomide Analogs IMiDS Inhibit Expression of Cyclooxygenase-2 in Multiple Myeloma. Cell Line and LPS Stimulated PBMCs. Abstract #2689. American Society of Hematology, Dec. 7-11, 2001.
- Mitsiades, N., Mitsiades, C., Poulaki, V., Akiyama, M., Tai, Y., Lin, B., Hayashi, T., Catley, L., Hideshima, T., Chauhan, D., Treon, S.P., Anderson, K.C., Apoptotic Signaling Induced by Immunomodulatory Thalidomide Analogs (Imids) in Human Multiple Myeloma Cells; Therapeutic Implications. Abstract #3224. American Society of Hematology, Dec. 7-11, 2001.
- Richardson, P.G., Schlossman, R.L., Hideshima, T., Davies, F., Leblanc, R., Catley, L., Doss, D., Kelly, K.A., Mckenney, M., Mechlowicz, J., Freeman, A., Deocampo, R., Rich, R., Ryoo, J., Chauhan, D., Munshi, N., Weller, E., Zeldis, J., Anderson, K.C., A Phase I Study of Oral CC5013, an Immunomodulatory Thalidomide (Thal) Derivative, in Patients With Relapsed and Refractory Multiple Myeloma (MM). Abstract #3225. American Society of Hematology, Dec. 7-11, 2001.
- "Celgene drug promises activity in solid tumors," Marketletter, Jun. 18, 2001.
- Meregalli et al., "High-dose dexamethasone as first line therapy of multiple myeloma?," *Recenti Progressi in Medicina*, 1998, 89(1):18-20.
- Official Action in corresponding Canadian Application No. 2,476,983 dated Aug. 21, 2009.
- List, A., "New Approaches to the Treatment of Myelodysplasia," *The Oncologist*, 2002, 7(suppl. 1):39-49.
- Kurzrock, R., "Myelodysplastic syndrome overview," *Seminars in Hematology* (Abstract only), 2002, 39(3)(suppl. 2):18-25 Abstract only.
- Goerner, et al., "Morbidity and mortality of chronic GVHD after hematopoietic stem cell transplantation from HLA-identical siblings for patients with aplastic or refractory anemias," *Biology of Blood and Marrow Transplantation* (Abstract only), 2002, 8(1):47-56.
- Thomas, D., "Pilot studies of Thalidomide in Acute Myelogenous Leukemia, Myelodysplastic Syndromes, and Myeloproliferative Disorders," *Seminars in Hematology*, 2000, 37(1)(suppl. 3):26-34.
- Zorat, F. et al., "The clinical and biological effects of thalidomide in patients with myelodysplastic syndromes," *British Journal of Haematology*, 2001, 115:881-894.
- Official Action dated Feb. 10, 2009 in JP Application No. 2004-545192. (English translation provided).
- Teramura, M., Men-ekiyokusei Ryouhou, *Current Therapy*, 2000, 18(5):140-144 (in Japanese).
- Kon-nichi no Chiryuu Shishin, 1997 [Pocket Edition], Igaku Shoin, 1997, 513-514 (in Japanese).
- Okamoto, T., Kotsuzuikeisei Shoukougun to Men-eki Ijo, Bessatsu Nihon Rinsho, Syndrome Series for each area, No. 22, Blood Syndromes III, Nihon Rinsho, 213-216 (in Japanese), Oct. 1998.
- Merck Manual, 17th ed. Japanese version, 1999, 951-952.
- Notice of Allowance from U.S. Appl. No. 11/096,155 dated Jan. 12, 2010.
- Rajkumar et al., "Combination therapy with thalidomide plus dexamethasone for newly diagnosed multiple myeloma," *American Society of Hematology*, 43rd Annual Meeting, Dec. 7-11, 2001, Abstract #3525.
- Scheffler et al., "Safety and pharmacokinetics of CDC-501, a novel immunomodulatory-oncologic agent, after single then multiple, oral 100 mg twice daily doses," *American Society for Clinical Pharmacology and Therapeutics*, Mar. 24-27, 2002, Abstract #WP111-63.
- Marriott et al., "Thalidomide analogue CDC-501 is safe and well tolerated by patients with end stage cancer and shows evidence of clinical responses and extensive immune activation," *Br. J. Cancer*, 2002, 86(Suppl. 1):Abst 6.4.
- Kast, R.E., "Evidence of a mechanism by which etanercept increased TNF-alpha in multiple myeloma: New insights into the biology of TNF-alpha giving new treatment opportunities—the role of burproion," *Leukemia Research*, 2005, 29:1459-1463.
- Tsimberidou, A. et al., "Pilot study of recombinant human soluble tumor necrosis factor (TNF) receptor (p75) fusion protein (TNFR:Fc:Enbrel) in patients with refractory multiple myeloma: increase in plasma TNE α levels during treatment," *Leukemia Research*, 2003, 27:375-380.
- Dimopoulos, et al., "Long-term follow-up on overall survival from the MM-009 and MM-010 phase III trials of lenalidomide plus dexamethasone in patients with relapsed or refractory multiple myeloma," *Leukemia*, 2009, 1-6.
- Hideshima, T., et al., "A review of lenalidomide in combination with dexathasone for the treatment of multiple myeloma," *Therapeutics and Clinical Risk Management*, 2008, 4(1):129-136.
- Wang, M., et al., "Lenalidomide plus dexamethasone is more effective than dexamethasone alone in patients with relapsed or refractory multiple myeloma regardless of prior thalidomide exposure," *Blood*, 2008, 112(12):4445-4451.
- Gandhi, A., et al., "Dexamethasone Synergizes with Lenalidomide to Inhibit Multiple Myeloma Tumor Growth, But Reduces Lenalidomide-Induced Immunomodulation of T and NK Cell Function," *Current Cancer Drug Targets*, 2010, 10(1):1-13.
- Gay, F. et al., "Lenalidomide plus dexamethasone versus thalidomide plus dexamethasone in newly diagnosed multiple myeloma: a comparative analysis of 411 patients," *Blood*, 2010, 115(97):1343-150.
- Richardson, P. et al., "Thalidomide in multiple myeloma," *Biomed Pharmacother*, 2002, 56:115-28.
- Swartz, G. et al., "Pre-clinical evaluation of ENMD-0995: A thalidomide analog with activity against multiple myeloma and solid tumors," *Cell and Tumor Biology*, 2002, 43:181-182, Abstract# 910.
- Mazucco, R., "Angiogenesis and Anti-angiogenesis Therapeutics," *IDrugs*, 2002, 5(4): 320-322.
- Worker, C., "JP Morgan Hambrecht & Quist—20th Annual Healthcare Conference," *IDrugs*, 2002, 5(2):113-116.
- Treston, A. et al., "Pre-Clinical Evaluation of a Thalidomide Analog with Activity Against Multiple Myeloma and Solid Tumors—ENMD-0995 (S(-)-3-(3-amino-phthalimido)-glutarimide)," *Blood*, 2002, 100(11):816a, Abstract #3225.
- Mazucco, R. and Williams, L., "Immunotherapy, chemoprevention and angiogenesis," *IDrugs*, 2002, 5(5):408-411.
- Fernandes, P., "Anti-Cancer Drug Discovery and Development Summit," *IDrugs*, 2002, 5(8):757-764.

US 9,101,622 B2

Page 8

(56) **References Cited**

OTHER PUBLICATIONS

Notification letter dated Aug. 30, 2010 from Natco Pharma Limited to Celgene Corporation re: Notification pursuant to § 505(j)(2)(B) of the Federal Food, Drug and Cosmetic Act.

Complaint for Patent Infringement filed on Oct. 8, 2010 by Celgene Corporation in the U.S. District Court, District of New Jersey against Natco Pharma Limited.

Answer to Complaint filed on Nov. 18, 2010 by Natco Pharma Limited in the U.S. District Court, District of New Jersey.

Grosshans, E. and Illy, G., "Thalidomide Therapy for Inflammatory Dermatoses," *International Journal of Dermatology*, 1984, 23(9):598-602.

Krenn, M. et al., "Improvements in Solubility and Stability of Thalidomide upon Complexation with Hydropropyl- β -Cyclodextrin," *Journal of Pharmaceutical Sciences*, 1992, 81(7):685-689.

Schmahl, H. J. et al., "Pharmacokinetics of the Teratogenic and Nonteratogenic Thalidomide Analogs EM 12 and Supidimide in the Rat and Marmoset Monkey", in *Pharmacokinetics in Teratogenesis*, CRC Press, 1987, vol. I, Ch. 12, pp. 181-192.

Schumacher, H. et al., "The Teratogenic Activity of a Thalidomide Analogue, EM₁₂, in Rabbits, Rats, and Monkeys," *Teratology*, 1971, 5:233-240.

Smith, R. et al., "Studies on the Relationship Between the Chemical Structure and Embryotoxic Activity of Thalidomide and Related Compounds," in *A Symposium on Embryopathic Activity of Drugs*, J. & A. Churchill Ltd., 1965, Session 6, pp. 194-209.

Sheskin, J. and Sagher, F., "Trials with Thalidomide Derivatives in Leprosy Reactions," *Leprosy Review*, 1968, 39(4):203-205.

Sheskin, J., "Study with Nine Thalidomide Derivatives in the Lepra Reaction," *Pharmacology and Therapeutics*, 1978, 17:82-84.

Raje, N. and Anderson, K., "Thalidomide and immunomodulatory drugs as cancer therapy," *Current Opinions in Oncology*, 2002, 14:635-640.

Kumar, S. et al., "Thalidomide as an anti-cancer agent," *J. Cell. Mod. Med.*, 2002, 6(2):160-174.

Singhal, S. and Mehta, J., "Thalidomide in Cancer," *BioDrugs*, 2001, 15(3):163-172.

Notice of Opposition to EP 1 505 973 filed by Synthron B.V. on Nov. 30, 2010.

Notice of Opposition to EP 1 505 973 filed by Strawman Limited on Dec. 1, 2010.

Samson, D. et al., "Infusion of Vincristine and Doxorubicin with Oral Dexamethasone as First-Line Therapy for Multiple Myeloma," *The Lancet*, 1989, 334(8668):882-885.

Barlogie, B. et al., "Effective Treatment of Advanced Multiple Myeloma Refractory to Alkylating Agents," *N. Engl. J. Med.*, 1984, 310(21):1353-1356.

Dimopoulos, M. et al., "Thalidomide and dexamethasone combination for refractory multiple myeloma," *Annals of Oncology*, 2001, 12:991-995.

Zangari, M., et al., "Thrombogenic activity of doxorubicin in myeloma patients receiving thalidomide: implications for therapy," *Blood*, 2002, 100:1168-1171.

List, A. et al., "High Erythropoietic Remitting Activity of the Immunomodulatory Thalidomide Analog, CC5013, in Patients with Myelodysplastic Syndrome (MDS)," Abstract #353, *Blood*, 2002, 100(11):96a.

Mufti, G. et al., "Myelodysplastic Syndrome," *American Society of Hematology*, 2003, pp. 176-199.

Extracts from drug databases: retrieved from <http://www.nextbio.com/b/search/ov/IMI03%20cpd> on Nov. 26, 2010 and <http://pubchem.ncbi.nlm.nih.gov/summary/summary.cgi?cid=216326> on Nov. 26, 2010.

Stockdale, 1998, *Medicine*, Rubenstein and Federman, eds., vol. 3, Ch. 12, Sections IV and X.

"List of Approved Oncology Drugs with Approved Indications," <http://www.accessdata.fda.gov/scripts/cder/onctools/druglist.cfm>, last accessed on Sep. 11, 2008.

Office Action mailed Jun. 18, 2008, U.S. Appl. No. 11/325,954.

Gamberi et al., "Overall Safety and Treatment Duration in Lenalidomide (LEN)-, Thalidomide (THAL)-, and Bortezomib (BORT)-Treated Patients (Pts) within the European Post-Approval Safety Study (EU PASS) of Relapsed/Refractory Multiple Myeloma (RRMM)", presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #4068.

Korde et al., "Phase II Clinical and Correlative Study of Carfilzomib, Lenalidomide, and Dexamethasone (CRd) in Newly Diagnosed Multiple Myeloma (MM) Patients", presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #732.

Kumar et al., "A Phase I/2 Study of Weekly MLN9708, an Investigational Oral Proteasome Inhibitor, in Combination with Lenalidomide and Dexamethasone in Patients with Previously Untreated Multiple Myeloma (MM)", presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #332.

Palumbo et al., "Pomalidomide Cyclophosphamide and Prednisone (PCP) Treatment for Relapsed/Refractory Multiple Myeloma", presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #446.

Richardson et al., "A Phase 2 Study of Elotuzumab (Elo) in Combination with Lenalidomide and Low-Dose Dexamethasone (Ld) in Patients (pts) with Relapsed/Refractory Multiple Myeloma (R/R MM): Updated Results", presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #202.

Sacchi et al., "A Phase I/II Study of Bendamustine, Low-Dose Dexamethasone, and Lenalidomide (BdL) for the Treatment of Patients with Relapsed Multiple Myeloma", presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #1851.

Sonneveld et al., "Escalated Dose Bortezomib Once Weekly Combined with Lenalidomide and Dexamethasone (eVRD) Followed by Lenalidomide Maintenance in First Relapse of Multiple Myeloma (MM), the HOVON 86 Phase 2 Trial", presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #1853.

Suvannasankha et al., "A Phase I/II Trial Combining High-Dose Lenalidomide with Melphalan and Autologous Transplant for Multiple Myeloma: A Report of the Phase I Dose-Finding Study", presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #3146.

Mark et al., "ClAPD (Clarithromycin, Pomalidomide, Dexamethasone) Therapy in Relapsed or Refractory Multiple Myeloma", presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #77.

Lacy et al., "Pomalidomide Plus Low-Dose Dexamethasone (Pom/Dex) in Relapsed Myeloma: Long Term Follow up and Factors Predicting Outcome in 345 Patients," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #201.

Jagannath et al., "Pomalidomide (POM) with Low-Dose Dexamethasone (LoDex) in Patients (Pts) with Relapsed and Refractory Multiple Myeloma Who Have Received Prior Therapy with Lenalidomide (LEN) and Bortezomib (BORT): Updated Phase 2 Results and Age Subgroup Analysis," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #450.

Richardson et al., "MM-005: A Phase 1, Multicenter, Open-Label, Dose-Escalation Study to Determine the Maximum Tolerated Dose for the Combination of Pomalidomide, Bortezomib, and Low-Dose Dexamethasone in Subjects with Relapsed or Refractory Multiple Myeloma," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #727.

Leleu et al., "Prolonged Overall Survival with Pomalidomide and Dexamethasone in Myeloma Characterized with End Stage Disease," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #2961.

Berenson et al., "A Phase 1/2 Study of Pomalidomide, Dexamethasone and Pegylated Liposomal Doxorubicin for Patients with Relapsed/Refractory Multiple Myeloma," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #2979.

US 9,101,622 B2

Page 9

(56)

References Cited

OTHER PUBLICATIONS

- Lonial et al., "Improvement in Clinical Benefit Parameters with Pomalidomide (POM) in Combination with Low-Dose Dexamethasone (LoDex) in Patients with Relapsed and Refractory Multiple Myeloma (RRMM): Results From a Phase 2 Study," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #4052.
- Vij et al., "Pomalidomide (POM) with Low-Dose Dexamethasone (LoDex) in Patients with Relapsed and Refractory Multiple Myeloma (RRMM): Outcomes Based on Prior Treatment Exposure," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #4070.
- Richardson et al., "Treatment Outcomes with Pomalidomide (POM) in Combination with Low-Dose Dexamethasone (LoDex) in Patients with Relapsed and Refractory Multiple Myeloma (RRMM) and Del(17p13) and/or t(4;14) (p16;q32) Cytogenic Abnormalities Who Have Received Prior Therapy with Lenalidomide (LEN) and Bortezomib (BORT)", presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #4053.
- Dimopoulos et al., "Pomalidomide in Combination with Low-Dose Dexamethasone: Demonstrates a Significant Progression Free Survival and Overall Survival Advantage, in Relapsed/Refractory MM: A Phase 3, Multicenter, Randomized, Open-Label Study," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #LBA-6.
- Shastri et al., "A Phase II Study of Low-Dose Pomalidomide (0.5mg/day) and Prednisone Combination Therapy in Patients with Myelofibrosis and Significant Anemia," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #1728.
- Shah et al., "A Multi-Center Phase I/II Trial of Carfilzomib and Pomalidomide with Dexamethasone (Car-Pom-d) in Patients with Relapsed/Refractory Multiple Myeloma," presented at 54th ASH Annual Meeting and Exposition, Atlanta, Georgia, Dec. 8-11, 2012, Abstract #74.
- Office Action in corresponding CN Application No. 201110256752.0 dated Feb. 8, 2013.
- Stirling, D., "Thalidomide: A Novel Template for Anticancer Drugs," *Seminars in Oncology*, Dec. 2001, 28(6):602-606**.
- Celgene Press Release, "Celgene Will Discontinue Phase III Origin® Trial in Previously Untreated Elderly Patients with B-Cell Chronic Lymphocytic Leukemia," published on Celgene Newsroom, <http://newsroom.celgene.com> on Jul. 18, 2013 at 7:30 am EDT.
- Mateos, M.-V., Ph.D. et al., "Lenalidomide plus Dexamethasone for High-Risk Smoldering Multiple Myeloma," *New England Journal of Medicine*, Aug. 2013, 369(5):438-447.
- English translation of Japanese IP High Court decision in Application No. JP 2004-505051, dated Apr. 11, 2013.
- Jagannath, S. et al., "Pomalidomide (POM) with or without low-dose dexamethasone (LoDEX) in patients (Pts) with relapsed and refractory multiple myeloma (RRMM): MM-002 phase II age subgroup analysis," *J Clin Oncol* 31, 2013 (suppl; abstr 8532).
- Siegel, D. et al., "Long-term safety and efficacy of pomalidomide (POM) with or without low-dose dexamethasone (LoDEX) in relapsed and refractory multiple myeloma (RRMM) patients enrolled in the MM-002 phase II trial," *J Clin Oncol* 31, 2013 (suppl; abstr 8588).
- Richardson, P.G. et al., A Phase 1/2 Multi-Center, Randomized, Open Label Dose Escalation Study to Determine the Maximum Tolerated Dose (MTD), Safety, and Efficacy of Pomalidomide (POM) Alone or in Combination with Low-Dose Dexamethasone (DEX) in Patients (PTS) with Relapsed and Refractory Multiple Myeloma (RRMM) Who Have Received Prior Treatment (TX) That Includes Lenalidomide (LEN) and Bortezomib (BORT), *Haematologica*, 2001; 96(s1):S31, Abstract 0-12, 13th International Myeloma Workshop, Paris, France—May 3-6, 2011.
- MacNeil, J.S., "Pomalidomide Picks Up Where Both Earlier IMiDs Stop Working," *The Oncology Report*, Mar./Apr. 2010, p. 34.
- Alexanian, R. et al., "Primary Dexamethasone Treatment of Multiple Myeloma," *Blood*, 1992, 80(4):887-890.
- Berenson, J. R. et al., "Long-Term Pamidronate Treatment of Advanced Multiple Myeloma Patients Reduces Skeletal Events," *Journal of Clinical Oncology*, 1998, 16(2):593-602.
- Attal, M. et al., "Lenalidomide Maintenance after Stem-Cell Transplantation for Multiple Myeloma", *N Engl J Med*, 2012, 366(19):1782-1791.
- McCarthy, P.L., et al., "Lenalidomide after Stem-Cell Transplantation for Multiple Myeloma", *N Engl J Med*, 2012, 366(19):1770-1781.
- Fifth Amended Complaint for Patent Infringement, dated May 6, 2013, *Celgene Corp. v. NATCO Pharma Ltd.*, D.N.J. (10-cv-5197).
- Amended Answer, Affirmative Defense and Counterclaims to Plaintiff's Fifth Amended Complaint, dated Aug. 1, 2013, *Celgene Corp. v. NATCO Pharma Ltd.*, D.N.J. (10-cv-5197).
- Reply to Counterclaims to Fifth Amended Complaint, Jun. 10, 2013, *Celgene Corp. v. NATCO Pharma Ltd.*, D.N.J. (10-cv-5197).
- Celgene's Opening Markman Brief, dated Oct. 21, 2013, *Celgene Corp. v. NATCO Pharma Ltd.*, D.N.J. (10-cv-5197).
- Markman Opinion, dated May 27, 2014, *Celgene Corp. v. NATCO Pharma Ltd.*, D.N.J. (10-cv-5197).
- European Patent Office's Decision revoking European Patent No. EP-B-1 505 973, dated Feb. 25, 2013.
- Office Action in corresponding Japanese Patent Application No. 2012-273326 dated May 13, 2014.
- Office Action in corresponding Canadian Patent Application No. 2,855,359 dated Aug. 14, 2014.
- Office Action in U.S. Appl. No. 13/276,867 dated Mar. 3, 2015.
- "A Study to Compare MPR With MP in Newly Diagnosed Multiple Myeloma Subjects 65 Years Old or Older," available at <http://clinicaltrials.gov/show/NCT00405756>, last accessed on Jul. 21, 2014.
- Durie B., "VIIIth International Myeloma Workshop Meeting Report from Banff, Canada May 4th-8th, 2001", *Myeloma Today*.
- Gertz et al., "Melphalan and Prednisone." *Myeloma Today*, 1999, 3(6):4 and 14-15.
- Gregory, W. et al., "Combination Chemotherapy Versus Melphalan and Prednisolone in the Treatment of Multiple Myeloma: An Overview of Published Trials," *J of Clinical Oncology*, 1992, 10(2):334-342.
- Kyle et al., "Therapeutic Application of Thalidomide in Multiple Myeloma", *Seminars in Oncology* (2001) 28, 583-587.
- Reiman T. et al., "Meeting Synopsis, VIII International Myeloma Workshop, Banff Springs Hotel, Banff, Alberta, Canada, May 4-8, 2001", *European Journal of Haematology*, 2001, 67: 199-202.
- Riet, I., "Angiogenesis in Multiple Myeloma." The Comprehensive Guide to Banff: Understanding the VIIIth International Myeloma Workshop, p. 12-13, May 5-8, 2001.
- Weerd et al., "Continuous low-dose cyclophosphamide-prednisone is effective and well tolerated in patients with advanced multiple myeloma," *The Netherlands Journal of Medicine*, Aug. 2001, 59(2):50-56.
- Zaidi et al. "Multiple myeloma: an old disease with new hope for the future." *CA Cancer J Clin*. 2001 51(5):273-85.



* cited by examiner

U.S. Patent

Aug. 11, 2015

US 9,101,622 B2

Effects of Revimid™ and Thalidomide on MM Cell Proliferation



US 9,101,622 B2

1

**METHODS FOR TREATING NEWLY
DIAGNOSED MULTIPLE MYELOMA
3-(4-AMINO-1-OXO-1,3-DIHYDRO-ISOINDOL-
2-YL)-PIPERIDINE-2,6-DIONE IN
COMBINATION WITH DEXAMETHASONE**

This application is a continuation of U.S. patent application Ser. No. 14/255,211, filed Apr. 17, 2014, which is continuation of U.S. patent application Ser. No. 14/201,069, filed Mar. 7, 2014, which is a continuation of U.S. patent application Ser. No. 13/782,728, filed Mar. 1, 2013, now U.S. Pat. No. 8,673,939, which is a continuation of U.S. patent application Ser. No. 13/488,888, filed Jun. 5, 2012, now U.S. Pat. No. 8,648,095, which is a continuation of U.S. patent application Ser. No. 12/640,702, filed Dec. 17, 2009, now U.S. Pat. No. 8,198,306, which is a continuation of U.S. patent application Ser. No. 10/438,213, filed May 15, 2003, now U.S. Pat. No. 7,968,569, which claims the benefit of U.S. provisional application No. 60/380,842, filed May 17, 2002, and 60/424,600, filed Nov. 6, 2002, the entireties of which are incorporated herein by reference.

1. FIELD OF THE INVENTION

This invention relates to methods of treating, preventing and/or managing specific cancers, and other diseases including, but not limited to, those associated with, or characterized by, undesired angiogenesis, by the administration of one or more immunomodulatory compounds alone or in combination with other therapeutics. In particular, the invention encompasses the use of specific combinations, or “cocktails,” of drugs and other therapy, e.g., radiation to treat these specific cancers, including those refractory to conventional therapy. The invention also relates to pharmaceutical compositions and dosing regimens.

2. BACKGROUND OF THE INVENTION

2.1 Pathobiology of Cancer and Other Diseases

Cancer is characterized primarily by an increase in the number of abnormal cells derived from a given normal tissue, invasion of adjacent tissues by these abnormal cells, or lymphatic or blood-borne spread of malignant cells to regional lymph nodes and to distant sites (metastasis). Clinical data and molecular biologic studies indicate that cancer is a multistep process that begins with minor preneoplastic changes, which may under certain conditions progress to neoplasia. The neoplastic lesion may evolve clonally and develop an increasing capacity for invasion, growth, metastasis, and heterogeneity, especially under conditions in which the neoplastic cells escape the host's immune surveillance. Roitt, I., Brostoff, J and Kale, D., *Immunology*, 17.1-17.12 (3rd ed., Mosby, St. Louis, Mo., 1993).

There is an enormous variety of cancers which are described in detail in the medical literature. Examples includes cancer of the lung, colon, rectum, prostate, breast, brain, and intestine. The incidence of cancer continues to climb as the general population ages, as new cancers develop, and as susceptible populations (e.g., people infected with AIDS or excessively exposed to sunlight) grow. A tremendous demand therefore exists for new methods and compositions that can be used to treat patients with cancer.

Many types of cancers are associated with new blood vessel formation, a process known as angiogenesis. Several of the mechanisms involved in tumor-induced angiogenesis have been elucidated. The most direct of these mechanisms is the secretion by the tumor cells of cytokines with angiogenic

2

properties. Examples of these cytokines include acidic and basic fibroblastic growth factor (a,b-FGF), angiogenin, vascular endothelial growth factor (VEGF), and TNF- α . Alternatively, tumor cells can release angiogenic peptides through the production of proteases and the subsequent breakdown of the extracellular matrix where some cytokines are stored (e.g., b-FGF). Angiogenesis can also be induced indirectly through the recruitment of inflammatory cells (particularly macrophages) and their subsequent release of angiogenic cytokines (e.g., TNF- α , bFGF).

A variety of other diseases and disorders are also associated with, or characterized by, undesired angiogenesis. For example, enhanced or unregulated angiogenesis has been implicated in a number of diseases and medical conditions including, but not limited to, ocular neovascular diseases, choroidal neovascular diseases, retina neovascular diseases, rubeosis (neovascularization of the angle), viral diseases, genetic diseases, inflammatory diseases, allergic diseases, and autoimmune diseases. Examples of such diseases and conditions include, but are not limited to: diabetic retinopathy; retinopathy of prematurity; corneal graft rejection; neovascular glaucoma; retrolental fibroplasia; and proliferative vitreoretinopathy.

Accordingly, compounds that can control angiogenesis or inhibit the production of certain cytokines, including TNF- α , may be useful in the treatment and prevention of various diseases and conditions.

2.2 Methods of Treating Cancer

Current cancer therapy may involve surgery, chemotherapy, hormonal therapy and/or radiation treatment to eradicate neoplastic cells in a patient (see, for example, Stockdale, 1998, *Medicine*, vol. 3, Rubenstein and Federman, eds., Chapter 12, Section IV). Recently, cancer therapy could also involve biological therapy or immunotherapy. All of these approaches pose significant drawbacks for the patient. Surgery, for example, may be contraindicated due to the health of a patient or may be unacceptable to the patient. Additionally, surgery may not completely remove neoplastic tissue. Radiation therapy is only effective when the neoplastic tissue exhibits a higher sensitivity to radiation than normal tissue. Radiation therapy can also often elicit serious side effects. Hormonal therapy is rarely given as a single agent. Although hormonal therapy can be effective, it is often used to prevent or delay recurrence of cancer after other treatments have removed the majority of cancer cells. Biological therapies and immunotherapies are limited in number and may produce side effects such as rashes or swellings, flu-like symptoms, including fever, chills and fatigue, digestive tract problems or allergic reactions.

With respect to chemotherapy, there are a variety of chemotherapeutic agents available for treatment of cancer. A majority of cancer chemotherapeutics act by inhibiting DNA synthesis, either directly, or indirectly by inhibiting the biosynthesis of deoxyribonucleotide triphosphate precursors, to prevent DNA replication and concomitant cell division. Gilman et al., *Goodman and Gilman's: The Pharmacological Basis of Therapeutics*, Tenth Ed. (McGraw Hill, New York).

Despite availability of a variety of chemotherapeutic agents, chemotherapy has many drawbacks. Stockdale, *Medicine*, vol. 3, Rubenstein and Federman, eds., ch. 12, sect. 10, 1998. Almost all chemotherapeutic agents are toxic, and chemotherapy causes significant, and often dangerous side effects including severe nausea, bone marrow depression, and immunosuppression. Additionally, even with administration of combinations of chemotherapeutic agents, many tumor cells are resistant or develop resistance to the chemotherapeutic agents. In fact, those cells resistant to the particular

US 9,101,622 B2

3

chemotherapeutic agents used in the treatment protocol often prove to be resistant to other drugs, even if those agents act by different mechanism from those of the drugs used in the specific treatment. This phenomenon is referred to as pleiotropic drug or multidrug resistance. Because of the drug resistance, many cancers prove refractory to standard chemotherapeutic treatment protocols.

Other diseases or conditions associated with, or characterized by, undesired angiogenesis are also difficult to treat. However, some compounds such as protamine, heparin and steroids have been proposed to be useful in the treatment of certain specific diseases. Taylor et al., *Nature* 297:307 (1982); Folkman et al., *Science* 221:719 (1983); and U.S. Pat. Nos. 5,001,116 and 4,994,443. Thalidomide and certain derivatives of it have also been proposed for the treatment of such diseases and conditions. U.S. Pat. Nos. 5,593,990, 5,629,327, 5,712,291, 6,071,948 and 6,114,355 to D'Amato.

Still, there is a significant need for safe and effective methods of treating, preventing and managing cancer and other diseases and conditions, particularly for diseases that are refractory to standard treatments, such as surgery, radiation therapy, chemotherapy and hormonal therapy, while reducing or avoiding the toxicities and/or side effects associated with the conventional therapies.

2.3 IMiDS™

A number of studies have been conducted with the aim of providing compounds that can safely and effectively be used to treat diseases associated with abnormal production of TNF- α . See, e.g., Marriott, J. B., et al., *Expert Opin. Biol. Ther.* 1(4):1-8 (2001); G. W. Muller, et al., *Journal of Medicinal Chemistry* 39(17): 3238-3240 (1996); and G. W. Muller, et al., *Bioorganic & Medicinal Chemistry Letters* 8: 2669-2674 (1998). Some studies have focused on a group of compounds selected for their capacity to potently inhibit TNF- α production by LPS stimulated PBMC. L. G. Corral, et al., *Ann. Rheum. Dis.* 58:(Suppl I) 1107-1113 (1999). These compounds, which are referred to as IMiDS™ (Celgene Corporation) or Immunomodulatory Drugs, show not only potent inhibition of TNF- α but also marked inhibition of LPS induced monocyte IL1 β and IL12 production. LPS induced IL6 is also inhibited by immunomodulatory compounds, albeit partially. These compounds are potent stimulators of LPS induced IL10. Id. Particular examples of IMiDS™s include, but are not limited to, the substituted 2-(2,6-dioxopiperidin-3-yl) phthalimides and substituted 2-(2,6-dioxopiperidin-3-yl)-1-oxoisindoles described in U.S. Pat. Nos. 6,281,230 and 6,316,471, both to G. W. Muller, et al.

3. SUMMARY OF THE INVENTION

This invention encompasses methods of treating and preventing certain types of cancer, including primary and metastatic cancer, as well as cancers that are refractory or resistant to conventional chemotherapy. The methods comprise administering to a patient in need of such treatment or prevention a therapeutically or prophylactically effective amount of an immunomodulatory compound, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof. The invention also encompasses methods of managing certain cancers (e.g., preventing or prolonging their recurrence, or lengthening the time of remission) which comprise administering to a patient in need of such management a prophylactically effective amount of an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

4

In particular methods of the invention, an immunomodulatory compound is administered in combination with a therapy conventionally used to treat, prevent or manage cancer. Examples of such conventional therapies include, but are not limited to, surgery, chemotherapy, radiation therapy, hormonal therapy, biological therapy and immunotherapy.

This invention also encompasses methods of treating, managing or preventing diseases and disorders other than cancer that are associated with, or characterized by, undesired angiogenesis, which comprise administering to a patient in need of such treatment, management or prevention a therapeutically or prophylactically effective amount of an immunomodulatory compound, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

In other methods of the invention, an immunomodulatory compound is administered in combination with a therapy conventionally used to treat, prevent or manage diseases or disorders associated with, or characterized by, undesired angiogenesis. Examples of such conventional therapies include, but are not limited to, surgery, chemotherapy, radiation therapy, hormonal therapy, biological therapy and immunotherapy.

This invention encompasses pharmaceutical compositions, single unit dosage forms, dosing regimens and kits which comprise an immunomodulatory compound, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, and a second, or additional, active agent. Second active agents include specific combinations, or "cocktails," of drugs.

4. BRIEF DESCRIPTION OF FIGURE

FIG. 1 shows a comparison of the effects of 3-(4-amino-1-oxo-1,3-dihydro-isindol-2-yl)-piperidine-2,6-dione (Revimid™) and thalidomide in inhibiting the proliferation of multiple myeloma (MM) cell lines in an in vitro study. The uptake of [³H]-thymidine by different MM cell lines (MM.1S, Hs Sultan, U266 and RPMI-8226) was measured as an indicator of the cell proliferation.

5. DETAILED DESCRIPTION OF THE INVENTION

A first embodiment of the invention encompasses methods of treating, managing, or preventing cancer which comprises administering to a patient in need of such treatment or prevention a therapeutically or prophylactically effective amount of an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

In particular methods encompassed by this embodiment, the immunomodulatory compound is administered in combination with another drug ("second active agent") or method of treating, managing, or preventing cancer. Second active agents include small molecules and large molecules (e.g., proteins and antibodies), examples of which are provided herein, as well as stem cells. Methods, or therapies, that can be used in combination with the administration of the immunomodulatory compound include, but are not limited to, surgery, blood transfusions, immunotherapy, biological therapy, radiation therapy, and other non-drug based therapies presently used to treat, prevent or manage cancer.

Another embodiment of the invention encompasses methods of treating, managing or preventing diseases and disorders other than cancer that are characterized by undesired angiogenesis. These methods comprise the administration of a therapeutically or prophylactically effective amount of an

US 9,101,622 B2

5

immunomodulatory compound, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof.

Examples of diseases and disorders associated with, or characterized by, undesired angiogenesis include, but are not limited to, inflammatory diseases, autoimmune diseases, viral diseases, genetic diseases, allergic diseases, bacterial diseases, ocular neovascular diseases, choroidal neovascular diseases, retina neovascular diseases, and rubeosis (neovascularization of the angle).

In particular methods encompassed by this embodiment, the immunomodulatory compound is administered in combination with a second active agent or method of treating, managing, or preventing the disease or condition. Second active agents include small molecules and large molecules (e.g., proteins and antibodies), examples of which are provided herein, as well as stem cells. Methods, or therapies, that can be used in combination with the administration of the immunomodulatory compound include, but are not limited to, surgery, blood transfusions, immunotherapy, biological therapy, radiation therapy, and other non-drug based therapies presently used to treat, prevent or manage disease and conditions associated with, or characterized by, undesired angiogenesis.

The invention also encompasses pharmaceutical compositions (e.g., single unit dosage forms) that can be used in methods disclosed herein. Particular pharmaceutical compositions comprise an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, and a second active agent.

5.1 Immunomodulatory Compounds

Compounds used in the invention include immunomodulatory compounds that are racemic, stereomerically enriched or stereomerically pure, and pharmaceutically acceptable salts, solvates, hydrates, stereoisomers, clathrates, and prodrugs thereof. Preferred compounds used in the invention are small organic molecules having a molecular weight less than about 1,000 g/mol, and are not proteins, peptides, oligonucleotides, oligosaccharides or other macromolecules.

As used herein and unless otherwise indicated, the terms "immunomodulatory compounds" and "IMiDs™" (Celgene Corporation) encompasses small organic molecules that markedly inhibit TNF- α , LPS induced monocyte IL1 β and IL12, and partially inhibit IL6 production. Specific immunomodulatory compounds are discussed below.

TNF- α is an inflammatory cytokine produced by macrophages and monocytes during acute inflammation. TNF- α is responsible for a diverse range of signaling events within cells. TNF- α may play a pathological role in cancer. Without being limited by theory, one of the biological effects exerted by the immunomodulatory compounds of the invention is the reduction of synthesis of TNF- α . Immunomodulatory compounds of the invention enhance the degradation of TNF- α mRNA.


Further, without being limited by theory, immunomodulatory compounds used in the invention may also be potent co-stimulators of T cells and increase cell proliferation dramatically in a dose dependent manner. Immunomodulatory compounds of the invention may also have a greater co-stimulatory effect on the CD8+ T cell subset than on the CD4+ T cell subset. In addition, the compounds preferably have anti-inflammatory properties, and efficiently co-stimulate T cells.

Specific examples of immunomodulatory compounds of the invention, include, but are not limited to, cyano and carboxy derivatives of substituted styrenes such as those disclosed in U.S. Pat. No. 5,929,117; 1-oxo-2-(2,6-dioxo-3-

6

fluoropiperidin-3-yl) isoindolines and 1,3-dioxo-2-(2,6-dioxo-3-fluoropiperidin-3-yl) isoindolines such as those described in U.S. Pat. No. 5,874,448; the tetra substituted 2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindolines described in U.S. Pat. No. 5,798,368; 1-oxo and 1,3-dioxo-2-(2,6-dioxopiperidin-3-yl) isoindolines (e.g., 4-methyl derivatives of thalidomide and EM-12), including, but not limited to, those disclosed in U.S. Pat. No. 5,635,517; and a class of non-polypeptide cyclic amides disclosed in U.S. Pat. Nos. 5,698,579 and 5,877,200; analogs and derivatives of thalidomide, including hydrolysis products, metabolites, derivatives and precursors of thalidomide, such as those described in U.S. Pat. Nos. 5,593,990, 5,629,327, and 6,071,948 to D'Amato; aminothalidomide, as well as analogs, hydrolysis products, metabolites, derivatives and precursors of aminothalidomide, and substituted 2-(2,6-dioxopiperidin-3-yl) phthalimides and substituted 2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindoles such as those described in U.S. Pat. Nos. 6,281,230 and 6,316,471; isoindole-imide compounds such as those described in U.S. patent application Ser. No. 09/972,487 filed on Oct. 5, 2001, U.S. patent application Ser. No. 10/032,286 filed on Dec. 21, 2001, and International Application No. PCT/US01/50401 (International Publication No. WO 02/059106). The entireties of each of the patents and patent applications identified herein are incorporated herein by reference. Immunomodulatory compounds of the invention do not include thalidomide.

Other specific immunomodulatory compounds of the invention include, but are not limited to, 1-oxo- and 1,3-dioxo-2-(2,6-dioxopiperidin-3-yl) isoindolines substituted with amino in the benzo ring as described in U.S. Pat. No. 5,635,517 which is incorporated herein by reference. These compounds have the structure I:




in which one of X and Y is C=O, the other of X and Y is C=O or CH₂, and R² is hydrogen or lower alkyl, in particular methyl. Specific immunomodulatory compounds include, but are not limited to:

- 1-oxo-2-(2,6-dioxopiperidin-3-yl)-4-aminoisoindoline;
- 1-oxo-2-(2,6-dioxopiperidin-3-yl)-5-aminoisoindoline;
- 1-oxo-2-(2,6-dioxopiperidin-3-yl)-6-aminoisoindoline;
- 1-oxo-2-(2,6-dioxopiperidin-3-yl)-7-aminoisoindoline;
- 1,3-dioxo-2-(2,6-dioxopiperidin-3-yl)-4-aminoisoindoline; and
- 1,3-dioxo-2-(2,6-dioxopiperidin-3-yl)-5-aminoisoindoline.

Other specific immunomodulatory compounds of the invention belong to a class of substituted 2-(2,6-dioxopiperidin-3-yl) phthalimides and substituted 2-(2,6-dioxopiperidin-3-yl)-1-oxoisoindoles, such as those described in U.S. Pat. Nos. 6,281,230; 6,316,471; 6,335,349; and 6,476,052, and International Patent Application No. PCT/US97/13375 (International Publication No. WO 98/03502), each of which is incorporated herein by reference. Compounds representative of this class are of the formulas:


US 9,101,622 B2

7



wherein R^1 is hydrogen or methyl. In a separate embodiment, the invention encompasses the use of enantiomerically pure forms (e.g. optically pure (R) or (S) enantiomers) of these compounds.

Still other specific immunomodulatory compounds of the invention belong to a class of isoindole-imides disclosed in U.S. patent application Ser. Nos. 10/032,286 and 09/972,487, and International Application No. PCT/US01/50401 (International Publication No. WO 02/059106), each of which are incorporated herein by reference. Representative compounds are of formula II:



and pharmaceutically acceptable salts, hydrates, solvates, clathrates, enantiomers, diastereomers, racemates, and mixtures of stereoisomers thereof, wherein:

one of X and Y is $C=O$ and the other is CH_2 or $C=O$;

R^1 is H, (C_1-C_8) alkyl, (C_3-C_7) cycloalkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, benzyl, aryl, (C_0-C_4) alkyl- (C_1-C_6) heterocycloalkyl, (C_0-C_4) alkyl- (C_2-C_5) heteroaryl, $C(O)R^3$, $C(S)R^3$, $C(O)OR^4$, (C_1-C_8) alkyl- $N(R^6)_2$, (C_1-C_8) alkyl- OR^5 , (C_1-C_8) alkyl- $C(O)OR^5$, $C(O)NHR^3$, $C(S)NHR^3$, $C(O)NR^3R^3$, $C(S)NR^3R^3$ or (C_1-C_8) alkyl- $O(CO)R^5$;

R^2 is H, F, benzyl, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, or (C_2-C_8) alkynyl;

R^3 and R^3' are independently (C_1-C_8) alkyl, (C_3-C_7) cycloalkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, benzyl, aryl, $(C_0-$

8

$C_4)$ alkyl- (C_1-C_6) heterocycloalkyl, (C_0-C_4) alkyl- (C_2-C_5) heteroaryl, (C_0-C_8) alkyl- $N(R^6)_2$, (C_1-C_8) alkyl- OR^5 , (C_1-C_8) alkyl- $C(O)OR^5$, (C_1-C_8) alkyl- $O(CO)R^5$, or $C(O)OR^5$;

R^4 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, (C_1-C_4) alkyl- OR^5 , benzyl, aryl, (C_0-C_4) alkyl- (C_1-C_6) heterocycloalkyl, or (C_0-C_4) alkyl- (C_2-C_5) heteroaryl;

R^5 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, benzyl, aryl, or (C_2-C_5) heteroaryl;

each occurrence of R^6 is independently H, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, benzyl, aryl, (C_2-C_5) heteroaryl, or (C_0-C_8) alkyl- $C(O)OR^5$ or the R^6 groups can join to form a heterocycloalkyl group;

n is 0 or 1; and

* represents a chiral-carbon center.

In specific compounds of formula II, when n is 0 then R^1 is (C_3-C_7) cycloalkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, benzyl, aryl, (C_0-C_4) alkyl- (C_1-C_6) heterocycloalkyl, (C_0-C_4) alkyl- (C_2-C_5) heteroaryl, $C(O)R^3$, $C(O)OR^4$, (C_1-C_8) alkyl- $N(R^6)_2$, (C_1-C_8) alkyl- OR^5 , (C_1-C_8) alkyl- $C(O)OR^5$, $C(S)NHR^3$, or (C_1-C_8) alkyl- $O(CO)R^5$;


R^2 is H or (C_1-C_8) alkyl; and

R^3 is (C_1-C_8) alkyl, (C_3-C_7) cycloalkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, benzyl, aryl, (C_0-C_4) alkyl- (C_1-C_6) heterocycloalkyl, (C_0-C_4) alkyl- (C_2-C_5) heteroaryl, (C_5-C_8) alkyl- $N(R^6)_2$; (C_0-C_8) alkyl- $NH-C(O)OR^5$; (C_1-C_8) alkyl- OR^5 , (C_1-C_8) alkyl- $C(O)OR^5$, (C_1-C_8) alkyl- $O(CO)R^5$, or $C(O)OR^5$; and the other variables have the same definitions.


In other specific compounds of formula II, R^2 is H or (C_1-C_4) alkyl.

In other specific compounds of formula II, R^1 is (C_1-C_8) alkyl or benzyl.

In other specific compounds of formula II, R^1 is H, (C_1-C_8) alkyl, benzyl, CH_2OCH_3 , $CH_2CH_2OCH_3$, or



In another embodiment of the compounds of formula II, R^1 is



wherein Q is O or S, and each occurrence of R^7 is independently H, (C_1-C_8) alkyl, benzyl, CH_2OCH_3 , or $CH_2CH_2OCH_3$.

In other specific compounds of formula II, R^1 is $C(O)R^3$.

In other specific compounds of formula II, R^3 is (C_0-C_4) alkyl- (C_2-C_5) heteroaryl, (C_1-C_8) alkyl, aryl, or (C_0-C_4) alkyl- OR^5 .

In other specific compounds of formula II, heteroaryl is pyridyl, furyl, or thienyl.


In other specific compounds of formula II, R^1 is $C(O)OR^4$.

In other specific compounds of formula II, the H of $C(O)NHC(O)$ can be replaced with (C_1-C_4) alkyl, aryl, or benzyl.

US 9,101,622 B2

9

Still other specific immunomodulatory compounds of the invention belong to a class of isoindole-imides disclosed in U.S. patent application Ser. No. 09/781,179, International Publication No. WO 98/54170, and U.S. Pat. No. 6,395,754, each of which are incorporated herein by reference. Representative compounds are of formula III:



III

and pharmaceutically acceptable salts, hydrates, solvates, clathrates, enantiomers, diastereomers, racemates, and mixtures of stereoisomers thereof, wherein:

one of X and Y is C=O and the other is CH₂ or C=O;

R is H or CH₂OCOR';

(i) each of R¹, R², R³, or R⁴, independently of the others, is halo, alkyl of 1 to 4 carbon atoms, or alkoxy of 1 to 4 carbon atoms or (ii) one of R¹, R², R³, or R⁴ is nitro or —NHR⁵ and the remaining of R¹, R², R³, or R⁴ are hydrogen;

R⁵ is hydrogen or alkyl of 1 to 8 carbons

R⁶ hydrogen, alkyl of 1 to 8 carbon atoms, benzo, chloro, or fluoro;

R⁷ is R⁷—CHR¹⁰—N(R⁸R⁹);


R⁷ is m-phenylene or p-phenylene or —(C_nH_{2n})— in which n has a value of 0 to 4;

each of R⁸ and R⁹ taken independently of the other is hydrogen or alkyl of 1 to 8 carbon atoms, or R⁸ and R⁹ taken together are tetramethylene, pentamethylene, hexamethylene, or —CH₂CH₂[X]X₁CH₂CH₂— in which [X]X₁ is —O—, —S—, or —NH—;

R¹⁰ is hydrogen, alkyl of 1 to 8 carbon atoms, or phenyl; and


* represents a chiral-carbon center.

The most preferred immunomodulatory compounds of the invention are 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione and 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. The compounds can be obtained via standard, synthetic methods (see e.g., U.S. Pat. No. 5,635,517, incorporated herein by reference). The compounds are available from Celgene Corporation, Warren, N.J. 4-(Amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (ACTIMID™) has the following chemical structure:



The compound 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (REVIMID™) has the following chemical structure:

10



5

Compounds of the invention can either be commercially purchased or prepared according to the methods described in the patents or patent publications disclosed herein. Further, optically pure compounds can be asymmetrically synthesized or resolved using known resolving agents or chiral columns as well as other standard synthetic organic chemistry techniques.

As used herein and unless otherwise indicated, the term “pharmaceutically acceptable salt” encompasses non-toxic acid and base addition salts of the compound to which the term refers. Acceptable non-toxic acid addition salts include those derived from organic and inorganic acids or bases known in the art, which include, for example, hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid, methanesulphonic acid, acetic acid, tartaric acid, lactic acid, succinic acid, citric acid, malic acid, maleic acid, sorbic acid, aconitic acid, salicylic acid, phthalic acid, embolic acid, enanthic acid, and the like.

Compounds that are acidic in nature are capable of forming salts with various pharmaceutically acceptable bases. The bases that can be used to prepare pharmaceutically acceptable base addition salts of such acidic compounds are those that form non-toxic base addition salts, i.e., salts containing pharmacologically acceptable cations such as, but not limited to, alkali metal or alkaline earth metal salts and the calcium, magnesium, sodium or potassium salts in particular. Suitable organic bases include, but are not limited to, N,N-dibenzylethylenediamine, chlorprocaine, choline, diethanolamine, ethylenediamine, meglumaine (N-methylglucamine), lysine, and procaine.

As used herein and unless otherwise indicated, the term “prodrug” means a derivative of a compound that can hydrolyze, oxidize, or otherwise react under biological conditions (in vitro or in vivo) to provide the compound. Examples of prodrugs include, but are not limited to, derivatives of immunomodulatory compounds of the invention that comprise biohydrolyzable moieties such as biohydrolyzable amides, biohydrolyzable esters, biohydrolyzable carbamates, biohydrolyzable carbonates, biohydrolyzable ureides, and biohydrolyzable phosphate analogues. Other examples of prodrugs include derivatives of immunomodulatory compounds of the invention that comprise —NO, —NO₂, —ONO, or —ONO₂ moieties. Prodrugs can typically be prepared using well-known methods, such as those described in 1 *Burger's Medicinal Chemistry and Drug Discovery*, 172-178, 949-982 (Manfred E. Wolff ed., 5th ed. 1995), and *Design of Prodrugs* (H. Bundgaard ed., Elsevier, N.Y. 1985).

As used herein and unless otherwise indicated, the terms “biohydrolyzable amide,” “biohydrolyzable ester,” “biohydrolyzable carbamate,” “biohydrolyzable carbonate,” “biohydrolyzable ureide,” “biohydrolyzable phosphate” mean an amide, ester, carbamate, carbonate, ureide, or phosphate, respectively, of a compound that either: 1) does not interfere with the biological activity of the compound but can confer upon that compound advantageous properties in vivo, such as uptake, duration of action, or onset of action; or 2) is biologically inactive but is converted in vivo to the biologically

55

60

65

active compound. Examples of biohydrolyzable esters include, but are not limited to, lower alkyl esters, lower acyloxyalkyl esters (such as acetoxyethyl, acetoxyethyl, aminocarbonyloxymethyl, pivaloyloxymethyl, and pivaloyloxyethyl esters), lactonyl esters (such as phthalidyl and thiophthalidyl esters), lower alkoxyacyloxyalkyl esters (such as methoxycarbonyloxymethyl, ethoxycarbonyloxyethyl and isopropoxycarbonyloxyethyl esters), alkoxyalkyl esters, choline esters, and acylamino alkyl esters (such as acetamidomethyl esters). Examples of biohydrolyzable amides include, but are not limited to, lower alkyl amides, α -amino acid amides, alkoxyacyl amides, and alkylaminoalkylcarbonyl amides. Examples of biohydrolyzable carbamates include, but are not limited to, lower alkylamines, substituted ethylenediamines, amino acids, hydroxyalkylamines, heterocyclic and heteroaromatic amines, and polyether amines.

Various immunomodulatory compounds of the invention contain one or more chiral centers, and can exist as racemic mixtures of enantiomers or mixtures of diastereomers. This invention encompasses the use of stereomerically pure forms of such compounds, as well as the use of mixtures of those forms. For example, mixtures comprising equal or unequal amounts of the enantiomers of a particular immunomodulatory compounds of the invention may be used in methods and compositions of the invention. These isomers may be asymmetrically synthesized or resolved using standard techniques such as chiral columns or chiral resolving agents. See, e.g., Jacques, J., et al., *Enantiomers, Racemates and Resolutions* (Wiley-Interscience, New York, 1981); Wilen, S. H., et al., *Tetrahedron* 33:2725 (1977); Eliel, E. L., *Stereochemistry of Carbon Compounds* (McGraw-Hill, NY, 1962); and Wilen, S. H., *Tables of Resolving Agents and Optical Resolutions* p. 268 (E. L. Eliel, Ed., Univ. of Notre Dame Press, Notre Dame, Ind., 1972).

As used herein and unless otherwise indicated, the term "stereomerically pure" means a composition that comprises one stereoisomer of a compound and is substantially free of other stereoisomers of that compound. For example, a stereomerically pure composition of a compound having one chiral center will be substantially free of the opposite enantiomer of the compound. A stereomerically pure composition of a compound having two chiral centers will be substantially free of other diastereomers of the compound. A typical stereomerically pure compound comprises greater than about 80% by weight of one stereoisomer of the compound and less than about 20% by weight of other stereoisomers of the compound, more preferably greater than about 90% by weight of one stereoisomer of the compound and less than about 10% by weight of the other stereoisomers of the compound, even more preferably greater than about 95% by weight of one stereoisomer of the compound and less than about 5% by weight of the other stereoisomers of the compound, and most preferably greater than about 97% by weight of one stereoisomer of the compound and less than about 3% by weight of the other stereoisomers of the compound. As used herein and unless otherwise indicated, the term "stereomerically enriched" means a composition that comprises greater than about 60% by weight of one stereoisomer of a compound, preferably greater than about 70% by weight, more preferably greater than about 80% by weight of one stereoisomer of a compound. As used herein and unless otherwise indicated, the term "enantiomerically pure" means a stereomerically pure composition of a compound having one chiral center. Similarly, the term "stereomerically enriched" means a stereomerically enriched composition of a compound having one chiral center.

It should be noted that if there is a discrepancy between a depicted structure and a name given that structure, the depicted structure is to be accorded more weight. In addition, if the stereochemistry of a structure or a portion of a structure is not indicated with, for example, bold or dashed lines, the structure or portion of the structure is to be interpreted as encompassing all stereoisomers of it.

5.2 Second Active Agents

Immunomodulatory compounds can be combined with other pharmacologically active compounds ("second active agents") in methods and compositions of the invention. It is believed that certain combinations work synergistically in the treatment of particular types of cancer and certain diseases and conditions associated with, or characterized by, undesired angiogenesis. Immunomodulatory compounds can also work to alleviate adverse effects associated with certain second active agents, and some second active agents can be used to alleviate adverse effects associated with immunomodulatory compounds.

One or more second active ingredients or agents can be used in the methods and compositions of the invention together with an immunomodulatory compound. Second active agents can be large molecules (e.g., proteins) or small molecules (e.g., synthetic inorganic, organometallic, or organic molecules).

Examples of large molecule active agents include, but are not limited to, hematopoietic growth factors, cytokines, and monoclonal and polyclonal antibodies. Typical large molecule active agents are biological molecules, such as naturally occurring or artificially made proteins. Proteins that are particularly useful in this invention include proteins that stimulate the survival and/or proliferation of hematopoietic precursor cells and immunologically active poietic cells in vitro or in vivo. Others stimulate the division and differentiation of committed erythroid progenitors in cells in vitro or in vivo. Particular proteins include, but are not limited to: interleukins, such as IL-2 (including recombinant IL-II ("rIL.2") and canarypox IL-2), IL-10, IL-12, and IL-18; interferons, such as interferon alfa-2a, interferon alfa-2b, interferon alfa-n1, interferon alfa-n3, interferon beta-1 a, and interferon gamma-I b; GM-CSF and GM-CSF; and EPO.

Particular proteins that can be used in the methods and compositions of the invention include, but are not limited to: filgrastim, which is sold in the United States under the trade name Neupogen® (Amgen, Thousand Oaks, Calif.); sargramostim, which is sold in the United States under the trade name Leukine® (Immunex, Seattle, Wash.); and recombinant EPO, which is sold in the United States under the trade name Epogen® (Amgen, Thousand Oaks, Calif.).

Recombinant and mutated forms of GM-CSF can be prepared as described in U.S. Pat. Nos. 5,391,485; 5,393,870; and 5,229,496; all of which are incorporated herein by reference. Recombinant and mutated forms of G-CSF can be prepared as described in U.S. Pat. Nos. 4,810,643; 4,999,291; 5,528,823; and 5,580,755; all of which are incorporated herein by reference.

This invention encompasses the use of native, naturally occurring, and recombinant proteins. The invention further encompasses mutants and derivatives (e.g., modified forms) of naturally occurring proteins that exhibit, in vivo, at least some of the pharmacological activity of the proteins upon which they are based. Examples of mutants include, but are not limited to, proteins that have one or more amino acid residues that differ from the corresponding residues in the naturally occurring forms of the proteins. Also encompassed by the term "mutants" are proteins that lack carbohydrate moieties normally present in their naturally occurring forms

US 9,101,622 B2

13

(e.g., nonglycosylated forms). Examples of derivatives include, but are not limited to, pegylated derivatives and fusion proteins, such as proteins formed by fusing IgG1 or IgG3 to the protein or active portion of the protein of interest. See, e.g., Penichet, M. L. and Morrison, S. L., *J. Immunol. Methods* 248:91-101 (2001).

Antibodies that can be used in combination with compounds of the invention include monoclonal and polyclonal antibodies. Examples of antibodies include, but are not limited to, trastuzumab (Herceptin®), rituximab (Rituxan®), bevacizumab (Avastin™), pertuzumab (Omnitarg™), tositumomab (Bexxar®), edrelomab (Panorex®), and G250. Compounds of the invention can also be combined with, or used in combination with, anti-TNF- α antibodies.

Large molecule active agents may be administered in the form of anti-cancer vaccines. For example, vaccines that secrete, or cause the secretion of, cytokines such as IL-2, G-CSF, and GM-CSF can be used in the methods, pharmaceutical compositions, and kits of the invention. See, e.g., Emens, L. A., et al., *Curr. Opin. Mol. Ther.* 3(1):77-84 (2001).

In one embodiment of the invention, the large molecule active agent reduces, eliminates, or prevents an adverse effect associated with the administration of an immunomodulatory compound. Depending on the particular immunomodulatory compound and the disease or disorder being treated, adverse effects can include, but are not limited to, drowsiness and somnolence, dizziness and orthostatic hypotension, neutropenia, infections that result from neutropenia, increased HIV-viral load, bradycardia, Stevens-Johnson Syndrome and toxic epidermal necrolysis, and seizures (e.g., grand mal convulsions). A specific adverse effect is neutropenia.

Second active agents that are small molecules can also be used to alleviate adverse effects associated with the administration of an immunomodulatory compound. However, like some large molecules, many are believed to be capable of providing a synergistic effect when administered with (e.g., before, after or simultaneously) an immunomodulatory compound. Examples of small molecule second active agents include, but are not limited to, anti-cancer agents, antibiotics, immunosuppressive agents, and steroids.

Examples of anti-cancer agents include, but are not limited to: acivicin; aclarubicin; acodazole hydrochloride; acronine; adozelesin; aldesleukin; altretamine; ambomycin; ametantrone acetate; amsacrine; anastrozole; anthramycin; asparaginase; asperlin; azacitidine; azetepa; azotomycin; batimastat; benzodepa; bicalutamide; bisantrene hydrochloride; bisnafide dimesylate; bizelesin; bleomycin sulfate; brequinar sodium; bropirimine; busulfan; cactinomycin; calusterone; caracemide; carbetimer; carboplatin; carmustine; carubicin hydrochloride; carzelesin; cedefingol; celecoxib (COX-2 inhibitor); chlorambucil; cirolemycin; cisplatin; cladribine; crisnatol mesylate; cyclophosphamide; cytarabine; dacarbazine; dactinomycin; daunorubicin hydrochloride; decitabine; dexormaplatin; dezaguanine; dezaguanine mesylate; diaziquone; docetaxel; doxorubicin; doxorubicin hydrochloride; droloxifene; droloxifene citrate; dromostanolone propionate; duazomycin; edatrexate; eflornithine hydrochloride; elsamitucin; enloplatin; enpromate; epiropropidine; epirubicin hydrochloride; erbulozole; esorubicin hydrochloride; estramustine; estramustine phosphate sodium; etanidazole; etoposide; etoposide phosphate; etoprine; fadrozole hydrochloride; fazarabine; fenretinide; floxuridine; fludarabine phosphate; fluorouracil; fluorocytidine; fosquidone; fostriecin sodium; gemcitabine; gemcitabine hydrochloride; hydroxyurea; idarubicin hydrochloride; ifosfamide; ilmofofosine; iproplatin; irinotecan; irinotecan

14

hydrochloride; lanreotide acetate; letrozole; leuprolide acetate; liarozole hydrochloride; lometrexol sodium; lomustine; losoxantrone hydrochloride; masoprocol; maytansine; mechlorethamine hydrochloride; megestrol acetate; melengestrol acetate; melphalan; menogaril; mercaptopurine; methotrexate; methotrexate sodium; metoprine; meturedepa; mitindomide; mitocarcin; mitocromin; mitogillin; mitomalcin; mitomycin; mitosper; mitotane; mitoxantrone hydrochloride; mycophenolic acid; nocodazole; nogalamycin; ormaplatin; oxisuran; paclitaxel; pegaspargase; peliomycin; pentamustine; peplomycin sulfate; perfosfamide; pipobroman; piposulfan; piroxantrone hydrochloride; plicamycin; plomestane; porfimer sodium; porfiromycin; prednimustine; procarbazine hydrochloride; puromycin; puromycin hydrochloride; pyrazofurin; riboprine; safingol; safingol hydrochloride; semustine; simtrazene; sparfosate sodium; sparsomycin; spirogermanium hydrochloride; spiro-mustine; spiroplatin; streptonigrin; streptozocin; sulofenur; talisomycin; tecogalan sodium; taxotere; tegafur; teloxantrone hydrochloride; temoporfin; teniposide; teroxirone; testolactone; thiamiprine; thioguanine; thiotepa; tiazofurin; tirapazamine; toremifene citrate; tretolone acetate; triciribine phosphate; trimetrexate; trimetrexate glucuronate; triptorelin; tubulozole hydrochloride; uracil mustard; uredepa; vaporeotide; verteporfin; vinblastine sulfate; vincristine sulfate; vindesine; vindesine sulfate; vinepidine sulfate; vinglycinate sulfate; vinleurosine sulfate; vinorelbine tartrate; vinrosidine sulfate; vinzolidine sulfate; vorozole; zeniplatin; zinostatin; and zorubicin hydrochloride.

Other anti-cancer drugs include, but are not limited to: 20-epi-1,25 dihydroxyvitamin D3; 5-ethynyluracil; abiraterone; aclarubicin; acylfulvene; adecypenol; adozelesin; aldesleukin; ALL-TK antagonists; altretamine; ambamustine; amidox; amifostine; aminolevulinic acid; amrubicin; amsacrine; anagrelide; anastrozole; andrographolide; angiogenesis inhibitors; antagonist D; antagonist G; antarelix; antidorsalizing morphogenetic protein-1; antiandrogen, prostatic carcinoma; antiestrogen; antineoplaston; antisense oligonucleotides; aphidicolin glycinate; apoptosis gene modulators; apoptosis regulators; apurinic acid; ara-CDP-DL-PTBA; arginine deaminase; asulacrine; atamcstane; atrimustine; axinastatin 1; axinastatin 2; axinastatin 3; azasetron; azatoxin; azatyrosine; baccatin ITT derivatives; balanol; batimastat; BCR/ABL antagonists; benzochlorins; benzoylstauroporine; beta lactam derivatives; beta-alethine; betaclamycin B; betulinic acid; bFGF inhibitor; bicalutamide; bisantrene; bisaziridinylspermine; bisnafide; bistratene A; bizelesin; breflate; bropirimine; budotitane; buthionine sulfoximine; calcipotriol; calphostin C; camptothecin derivatives; capecitabine; carboxamide-amino-triazole; carboxamidotriazole; CaRest M3; CARN 700; cartilage derived inhibitor; carzelesin; casein kinase inhibitors (ICOS); castanospermine; cecropin B; cetrorelix; chlorins; chloroquinoline sulfonamide; cicaprost; cis-porphyrin; cladribine; clomifene analogues; clotrimazole; collismycin A; collismycin B; combretastatin A4; combretastatin analogue; conagein; crambescidin 816; crisnatol; cryptophycin 8; cryptophycin A derivatives; curacin A; cyclopentantraquinones; cycloplatin; cypemycin; cytarabine ocfosfate; cytolytic factor; cytostatin; dacliximab; decitabine; dehydridemnin B; deslorelin; dexamethasone; dexifosfamide; dexrazoxane; dexverapamil; diaziquone; didemnin B; didox; diethylnor-spermine; dihydro-5-azacytidine; dihydrotaxol, 9-; dioxamycin; diphenyl spiromustine; docetaxel; docosanol; dolasetron; doxifluridine; doxorubicin; droloxifene; dronabinol; duocarmycin SA; ebselen; ecomustine; edelfosine; edrecolomab; eflornithine; elemene; emitefur; epirubicin; epristeride;

US 9,101,622 B2

15

estramustine analogue; estrogen agonists; estrogen antagonists; etanidazole; etoposide phosphate; exemestane; fadrozole; fazarabine; fenretinide; filgrastim; finasteride; flavopiridol; flezelastine; fluasterone; fludarabine; fluorodaunorubicin hydrochloride; forfenimex; formestane; fostriecin; fotemustine; gadolinium texaphyrin; gallium nitrate; galocitabine; ganirelix; gelatinase inhibitors; gemcitabine; glutathione inhibitors; hepsulfam; heregulin; hexamethylene bisacetamide; hypericin; ibandronic acid; idarubicin; idoxifene; idramantone; ilmofosine; ilomastat; imatinib (e.g., Gleevec®), imiquimod; immunostimulant peptides; insulin-like growth factor-1 receptor inhibitor; interferon agonists; interferons; interleukins; iobenguane; iododoxorubicin; ipomeanol, 4-; iroplact; irsogladine; isobengazole; isohomohalicondrin B; itasetron; jasplakinolide; kahalalide F; lamellarin-N triacetate; lanreotide; leinamycin; lenograstim; lentinan sulfate; leptolstatin; letrozole; leukemia inhibiting factor; leukocyte alpha interferon; leuprolide+estrogen+progesterone; leuprorelin; levamisole; liarozole; linear polyamine analogue; lipophilic disaccharide peptide; lipophilic platinum compounds; lissoclinamide 7; lobaplatin; lombricine; lometrexol; lonidamine; losoxantrone; loxoribine; lurtotecan; lutetium texaphyrin; lysofylline; lytic peptides; maitansine; mannostatin A; marimastat; masoprocol; maspin; matrilysin inhibitors; matrix metalloproteinase inhibitors; mcnogaril; merbarone; meterclin; methioninase; metoclopramide; MIF inhibitor; mifepristone; miltefosine; mirimostim; mitoguanzone; mitolactol; mitomycin analogues; mitonafide; mitotoxin fibroblast growth factor-saporin; mitoxantrone; mofarotene; molgramostim; Erbitux, human chorionic gonadotrophin; monophosphoryl lipid A+mycobacterium cell wall sk; mopidamol; mustard anticancer agent; mycaperoxide B; mycobacterial cell wall extract; myriaporone; N-acetyldinaline; N-substituted benzamides; nafarelin; nagrestip; naloxone+pentazocine; napavin; naphterpin; narograstim; nedaplatin; nemorubicin; neridronic acid; nilutamide; nisamycin; nitric oxide modulators; nitroxide antioxidant; nitrullyn; oblimersen (Genasense®); O⁶-benzylguanine; octreotide; okicenone; oligonucleotides; onapristone; ondansetron; ondansetron; oracin; oral cytokine inducer; ormaplatin; osaterone; oxaliplatin; oxaunomycin; paclitaxel; paclitaxel analogues; paclitaxel derivatives; palauamine; palmitoylrhizoxin; pamidronic acid; panaxytriol; panomifene; parabactin; pazelliptine; pegaspargase; peldesine; pentosan polysulfate sodium; pentostatin; pentozole; perflubron; perfosfamide; perillyl alcohol; phenazinomycin; phenylacetate; phosphatase inhibitors; picibanil; pilocarpine hydrochloride; pirarubicin; piritrexim; placetin A; placetin B; plasminogen activator inhibitor; platinum complex; platinum compounds; platinum-triamine complex; porrimer sodium; porfirromycin; prednisone; propyl bis-acridone; prostaglandin J2; proteasome inhibitors; protein A-based immune modulator; protein kinase C inhibitor; protein kinase C inhibitors, microalgal; protein tyrosine phosphatase inhibitors; purine nucleoside phosphorylase inhibitors; purpurins; pyrazoloacridine; pyridoxylated hemoglobin polyoxyethylene conjugate; raf antagonists; raltitrexed; ramosestron; ras farnesyl protein transferase inhibitors; ras inhibitors; ras-GAP inhibitor; retelliptine demethylated; rhenium Re 186 etidronate; rhizoxin; ribozymes; RII retinamide; rohitukine; romurtide; roquinimex; rubiginone B1; ruboxyl; safingol; saintopin; SarCNU; sarcophytol A; sargramostim; Sdi 1 mimetics; semustine; senescence derived inhibitor 1; sense oligonucleotides; signal transduction inhibitors; sizofiran; sobuzoxane; sodium borocaptate; sodium phenylacetate; solverol; somatomedin binding protein; sonermin; sparfosic acid; spicamycin D; spiromustine; splenopentin; spongistatin

16

1; squalamine; stipiamide; stromelysin inhibitors; sulfinosine; superactive vasoactive intestinal peptide antagonist; suradista; suramin; swainsonine; tallimustine; tamoxifen methiodide; tauromustine; tazarotene; tecogalan sodium; tegafur; tellurapyrylium; telomerase inhibitors; temoporfin; teniposide; tetrachlorodecaoxide; tetrazomine; thaliblastine; thiocoraline; thrombopoietin; thrombopoietin mimetic; thymalfasin; thymopoietin receptor agonist; thymotrnan; thyroid stimulating hormone; tin ethyl etiopurpurin; tirapazamine; titanocene bichloride; topsentin; toremifene; translation inhibitors; tretinoin; triacetyluridine; triciribine; trimetrexate; triptorelin; tropisetron; turosteride; tyrosine kinase inhibitors; tyrphostins; UBC inhibitors; ubenimex; urogenital sinus-derived growth inhibitory factor; urokinase receptor antagonists; vapreotide; variolin B; velaresol; veramine; verdins; verteporfin; vinorelbine; vinoxaltine; vitaxin; vorozole; zanoterone; zeniplatin; zilascorb; and zinostatin stimalamer.

Specific second active agents include, but are not limited to, oblimersen (Genasense®), remicade, docetaxel, celecoxib, melphalan, dexamethasone (Decadron®), steroids, gemcitabine, cisplatin, temozolomide, etoposide, cyclophosphamide, temodar, carboplatin, procarbazine, gliadel, tamoxifen, topotecan, methotrexate, Arisa®, taxol, taxotere, fluorouracil, leucovorin, irinotecan, xeloda, CPT-11, interferon alpha, pegylated interferon alpha (e.g., PEG INTRON-A), capecitabine, cisplatin, thiotepa, fludarabine, carboplatin, liposomal daunorubicin, cytarabine, doxorubicin, paclitaxel, vinblastine, IL-2, GM-CSF, dacarbazine, vinorelbine, zoledronic acid, palmitronate, biaxin, busulphan, prednisone, bisphosphonate, arsenic trioxide, vincristine, doxorubicin (Doxil®), paclitaxel, ganciclovir, adriamycin, estramustine sodium phosphate (Emcyt®), sulindac, and etoposide.

5.3 Methods of Treatments and Prevention

Methods of this invention encompass methods of treating, preventing and/or managing various types of cancer and diseases and disorders associated with, or characterized by, undesired angiogenesis. As used herein, unless otherwise specified, the term “treating” refers to the administration of a compound of the invention or other additional active agent after the onset of symptoms of the particular disease or disorder. As used herein, unless otherwise specified, the term “preventing” refers to the administration prior to the onset of symptoms, particularly to patients at risk of cancer, and other diseases and disorders associated with, or characterized by, undesired angiogenesis. The term “prevention” includes the inhibition of a symptom of the particular disease or disorder. Patients with familial history of cancer and diseases and disorders associated with, or characterized by, undesired angiogenesis are preferred candidates for preventive regimens. As used herein and unless otherwise indicated, the term “managing” encompasses preventing the recurrence of the particular disease or disorder in a patient who had suffered from it, and/or lengthening the time a patient who had suffered from the disease or disorder remains in remission.

As used herein, the term “cancer” includes, but is not limited to, solid tumors and blood born tumors. The term “cancer” refers to disease of skin tissues, organs, blood, and vessels, including, but not limited to, cancers of the bladder, bone or blood, brain, breast, cervix, chest, colon, endometrium, esophagus, eye, head, kidney, liver, lymph nodes, lung, mouth, neck, ovaries, pancreas, prostate, rectum, stomach, testis, throat, and uterus. Specific cancers include, but are not limited to, advanced malignancy, amyloidosis, neuroblastoma, meningioma, hemangiopericytoma, multiple brain metastase, glioblastoma multiforms, glioblastoma, brain stem glioma, poor prognosis malignant brain tumor,

US 9,101,622 B2

17

malignant glioma, recurrent malignant glioma, anaplastic astrocytoma, anaplastic oligodendroglioma, neuroendocrine tumor, rectal adenocarcinoma, Dukes C & D colorectal cancer, unresectable colorectal carcinoma, metastatic hepatocellular carcinoma, Kaposi's sarcoma, karotype acute myeloblastic leukemia, Hodgkin's lymphoma, non-Hodgkin's lymphoma, cutaneous T-Cell lymphoma, cutaneous B-Cell lymphoma, diffuse large B-Cell lymphoma, low grade follicular lymphoma, malignant melanoma, malignant mesothelioma, malignant pleural effusion mesothelioma syndrome, peritoneal carcinoma, papillary serous carcinoma, gynecologic sarcoma, soft tissue sarcoma, scleroderma, cutaneous vasculitis, Langerhans cell histiocytosis, leiomyosarcoma, fibrodysplasia ossificans progressive, hormone refractory prostate cancer, resected high-risk soft tissue sarcoma, unresectable hepatocellular carcinoma, Waldenstrom's macroglobulinemia, smoldering myeloma, indolent myeloma, fallopian tube cancer, androgen independent prostate cancer, androgen dependent stage IV non-metastatic prostate cancer, hormone-insensitive prostate cancer, chemotherapy-insensitive prostate cancer, papillary thyroid carcinoma, follicular thyroid carcinoma, medullary thyroid carcinoma, and leiomyoma. In a specific embodiment, the cancer is metastatic. In another embodiment, the cancer is refractory or resistance to chemotherapy or radiation; in particular, refractory to thalidomide.

As used herein to refer to diseases and conditions other than cancer, the terms "diseases or disorders associated with, or characterized by, undesired angiogenesis," "diseases or disorders associated with undesired angiogenesis," and "diseases or disorders characterized by undesired angiogenesis" refer to diseases, disorders and conditions that are caused, mediated or attended by undesired, unwanted or uncontrolled angiogenesis, including, but not limited to, inflammatory diseases, autoimmune diseases, genetic diseases, allergic diseases, bacterial diseases, ocular neovascular diseases, choroidal neovascular diseases, and retina neovascular diseases.

Examples of such diseases or disorders associated with undesired angiogenesis include, but are not limited to, diabetic retinopathy, retinopathy of prematurity, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, proliferative vitreoretinopathy, trachoma, myopia, optic pits, epidemic keratoconjunctivitis, atopic keratitis, superior limbic keratitis, pterygium keratitis sicca, sjogrens, acne rosacea, phlyctenulosis, syphilis, lipid degeneration, bacterial ulcer, fungal ulcer, Herpes simplex infection, Herpes zoster infection, protozoan infection, Kaposi sarcoma, Mooren ulcer, Terrien's marginal degeneration, marginal keratolysis, rheumatoid arthritis, systemic lupus, polyarteritis, trauma, Wegeners sarcoidosis, Scleritis, Steven's Johnson disease, peripheral radial keratotomy, sickle cell anemia, sarcoid, pseudoxanthoma elasticum, Pagets disease, vein occlusion, artery occlusion, carotid obstructive disease, chronic uveitis, chronic vitritis, Lyme's disease, Eales disease, Beheet's disease, retinitis, choroiditis, presumed ocular histoplasmosis, Bests disease, Stargarts disease, pars planitis, chronic retinal detachment, hyperviscosity syndromes, toxoplasmosis, rubeosis, sarcoidosis, sclerositis, soriatis, psoriasis, primary sclerosing cholangitis, proctitis, primary biliary srosis, idiopathic pulmonary fibrosis, and alcoholic hepatitis.

In specific embodiments of the invention, diseases or disorders associated with undesired angiogenesis do not include congestive heart failure, cardiomyopathy, pulmonary edema, endotoxin-mediated septic shock, acute viral myocarditis, cardiac allograft rejection, myocardial infarction, HIV, hepatitis, adult respiratory distress syndrome, bone-resorption disease, chronic obstructive pulmonary diseases, chronic pul-

18

monary inflammatory disease, dermatitis, cystic fibrosis, septic shock, sepsis, endotoxic shock, hemodynamic shock, sepsis syndrome, post ischemic reperfusion injury, meningitis, psoriasis, fibrotic disease, cachexia, graft rejection, rheumatoid spondylitis, osteoporosis, Crohn's disease, ulcerative colitis, inflammatory-bowel disease, multiple sclerosis, systemic lupus erythematosus, erythema nodosum leprosum in leprosy, radiation damage, asthma, hyperoxic alveolar injury, malaria, mycobacterial infection, and opportunistic infections resulting from HIV.

This invention encompasses methods of treating patients who have been previously treated for cancer or diseases or disorders associated with, or characterized by, undesired angiogenesis, but are non-responsive to standard therapies, as well as those who have not previously been treated. The invention also encompasses methods of treating patients regardless of patient's age, although some diseases or disorders are more common in certain age groups. The invention further encompasses methods of treating patients who have undergone surgery in an attempt to treat the disease or condition at issue, as well as those who have not. Because patients with cancer and diseases and disorders characterized by undesired angiogenesis have heterogenous clinical manifestations and varying clinical outcomes, the treatment given to a patient may vary, depending on his/her prognosis. The skilled clinician will be able to readily determine without undue experimentation specific secondary agents, types of surgery, and types of non-drug based standard therapy that can be effectively used to treat an individual patient with cancer and other diseases or disorders.

Methods encompassed by this invention comprise administering one or more immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, to a patient (e.g., a human) suffering, or likely to suffer, from cancer or a disease or disorder mediated by undesired angiogenesis.

In one embodiment of the invention, an immunomodulatory compound of the invention can be administered orally and in single or divided daily doses in an amount of from about 0.10 to about 150 mg/day. In a particular embodiment, 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™) may be administered in an amount of from about 0.1 to about 1 mg per day, or alternatively from about 0.1 to about 5 mg every other day. In a preferred embodiment, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) may be administered in an amount of from about 5 to 25 mg per day, or alternatively from about 10 to about 50 mg every other day.

In a specific embodiment, 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™) may be administered in an amount of about 1, 2, or 5 mg per day to patients with relapsed multiple myeloma. In a particular embodiment, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) may be administered initially in an amount of 5 mg/day and the dose can be escalated every week to 10, 20, 25, 30 and 50 mg/day. In a specific embodiment, Revimid™ can be administered in an amount of up to about 30 mg/day to patients with solid tumor. In a particular embodiment, Revimid™ can be administered in an amount of up to about 40 mg/day to patients with glioma.

5.3.1 Combination Therapy with a Second Active Agent

Specific methods of the invention comprise administering an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, in combination with one or more second active agents, and/or in combination with radia-

US 9,101,622 B2

19

tion therapy, blood transfusions, or surgery. Examples of immunomodulatory compounds of the invention are disclosed herein (see, e.g., section 5.1). Examples of second active agents are also disclosed herein (see, e.g., section 5.2).

Administration of the immunomodulatory compounds and the second active agents to a patient can occur simultaneously or sequentially by the same or different routes of administration. The suitability of a particular route of administration employed for a particular active agent will depend on the active agent itself (e.g., whether it can be administered orally without decomposing prior to entering the blood stream) and the disease being treated. A preferred route of administration for an immunomodulatory compound of the invention is orally. Preferred routes of administration for the second active agents or ingredients of the invention are known to those of ordinary skill in the art. See, e.g., *Physicians' Desk Reference*, 1755-1760 (56th ed., 2002).

In one embodiment of the invention, the second active agent is administered intravenously or subcutaneously and once or twice daily in an amount of from about 1 to about 1000 mg, from about 5 to about 500 mg, from about 10 to about 350 mg, or from about 50 to about 200 mg. The specific amount of the second active agent will depend on the specific agent used, the type of disease being treated or managed, the severity and stage of disease, and the amount(s) of immunomodulatory compounds of the invention and any optional additional active agents concurrently administered to the patient. In a particular embodiment, the second active agent is oblimersen (Genasense®), GM-CSF, G-CSF, EPO, taxotere, irinotecan, dacarbazine, transretinoic acid, topotecan, pentoxifylline, ciprofloxacin, dexamethasone, vincristine, doxorubicin, COX-2 inhibitor, IL2, IL8, IL18, IFN, Ara-C, vinorelbine, or a combination thereof.

In a particular embodiment, GM-CSF, G-CSF or EPO is administered subcutaneously during about five days in a four or six week cycle in an amount of from about 1 to about 750 mg/m²/day, preferably in an amount of from about 25 to about 500 mg/m²/day, more preferably in an amount of from about 50 to about 250 mg/m²/day, and most preferably in an amount of from about 50 to about 200 mg/m²/day. In a certain embodiment, GM-CSF may be administered in an amount of from about 60 to about 500 mcg/m² intravenously over 2 hours, or from about 5 to about 12 mcg/m²/day subcutaneously. In a specific embodiment, G-CSF may be administered subcutaneously in an amount of about 1 mcg/kg/day initially and can be adjusted depending on rise of total granulocyte counts. The maintenance dose of G-CSF may be administered in an amount of about 300 (in smaller patients) or 480 mcg subcutaneously. In a certain embodiment, EPO may be administered subcutaneously in an amount of 10,000 Unit 3 times per week.

In another embodiment, Revimid™ in an amount of about 25 mg/d and dacarbazine in an amount of about from 200 to 1,000 mg/m²/d are administered to patients with metastatic malignant melanoma. In a specific embodiment, Revimid™ is administered in an amount of from about 5 to about 25 mg/d to patients with metastatic malignant melanoma whose disease has progressed on treatment with dacarbazine, IL-2 or IFN. In a specific embodiment, Revimid™ is administered to patients with relapsed or refractory multiple myeloma in an amount of about 15 mg/d twice a day or about 30 mg/d four times a day in a combination with dexamethasone.

In another embodiment, an immunomodulatory compound is administered with melphalan and dexamethasone to patients with amyloidosis. In a specific embodiment, an immunomodulatory compound of the invention and steroids can be administered to patients with amyloidosis.

20

In another embodiment, an immunomodulatory compound is administered with gemcitabine and cisplatin to patients with locally advanced or metastatic transitional cell bladder cancer.

In another embodiment, an immunomodulatory compound is administered in combination with a second active ingredient as follows: temozolomide to pediatric patients with relapsed or progressive brain tumors or recurrent neuroblastoma; celecoxib, etoposide and cyclophosphamide for relapsed or progressive CNS cancer; temodar to patients with recurrent or progressive meningioma, malignant meningioma, hemangiopericytoma, multiple brain metastases, relapsed brain tumors, or newly diagnosed glioblastoma multiformis; irinotecan to patients with recurrent glioblastoma; carboplatin to pediatric patients with brain stem glioma; procarbazine to pediatric patients with progressive malignant gliomas; cyclophosphamide to patients with poor prognosis malignant brain tumors, newly diagnosed or recurrent glioblastoma multiformis; Gliadel® for high grade recurrent malignant gliomas; temozolomide and tamoxifen for anaplastic astrocytoma; or topotecan for gliomas, glioblastoma, anaplastic astrocytoma or anaplastic oligodendroglioma.

In another embodiment, an immunomodulatory compound is administered with methotrexate and cyclophosphamide to patients with metastatic breast cancer.

In another embodiment, an immunomodulatory compound is administered with temozolomide to patients with neuroendocrine tumors.

In another embodiment, an immunomodulatory compound is administered with gemcitabine to patients with recurrent or metastatic head or neck cancer. In another embodiment, an immunomodulatory compound is administered with gemcitabine to patients with pancreatic cancer.

In another embodiment, an immunomodulatory compound is administered to patients with colon cancer in combination with Arisa®, taxol and/or taxotere.

In another embodiment, an immunomodulatory compound is administered with capecitabine to patients with refractory colorectal cancer or patients who fail first line therapy or have poor performance in colon or rectal adenocarcinoma.

In another embodiment, an immunomodulatory compound is administered in combination with fluorouracil, leucovorin, and irinotecan to patients with Dukes C & D colorectal cancer or to patients who have been previously treated for metastatic colorectal cancer.

In another embodiment, an immunomodulatory compound is administered to patients with refractory colorectal cancer in combination with capecitabine, xeloda, and/or CPT-11.

In another embodiment, an immunomodulatory compound of the invention is administered with capecitabine and irinotecan to patients with refractory colorectal cancer or to patients with unresectable or metastatic colorectal carcinoma.

In another embodiment, an immunomodulatory compound is administered alone or in combination with interferon alpha or capecitabine to patients with unresectable or metastatic hepatocellular carcinoma; or with cisplatin and thiotepa to patients with primary or metastatic liver cancer.

In another embodiment, an immunomodulatory compound is administered in combination with pegylated interferon alpha to patients with Kaposi's sarcoma.

In another embodiment, an immunomodulatory compound is administered in combination with fludarabine, carboplatin, and/or topotecan to patients with refractory or relapsed or high-risk acuted myelogenous leukemia.

In another embodiment, an immunomodulatory compound is administered in combination with liposomal daunorubicin,

US 9,101,622 B2

21

topotecan and/or cytarabine to patients with unfavorable karyotype acute myeloblastic leukemia.

In another embodiment, an immunomodulatory compound is administered in combination with gemcitabine and irinotecan to patients with non-small cell lung cancer. In one embodiment, an immunomodulatory compound is administered in combination with carboplatin and irinotecan to patients with non-small cell lung cancer. In one embodiment, an immunomodulatory compound is administered with doctaxol to patients with non-small cell lung cancer who have been previously treated with carbo/VP 16 and radiotherapy.

In another embodiment, an immunomodulatory compound is administered in combination with carboplatin and/or taxotere, or in combination with carboplatin, paclitaxel and/or thoracic radiotherapy to patients with non-small cell lung cancer. In a specific embodiment, an immunomodulatory compound is administered in combination with taxotere to patients with stage IIIB or TV non-small cell lung cancer.

In another embodiment, an immunomodulatory compound of the invention is administered in combination with oblimersen (Genasense®) to patients with small cell lung cancer.

In another embodiment, an immunomodulatory compound is administered alone or in combination with a second active ingredient such as vinblastine or fludarabine to patients with various types of lymphoma, including, but not limited to, Hodgkin's lymphoma, non-Hodgkin's lymphoma, cutaneous T-Cell lymphoma, cutaneous B-Cell lymphoma, diffuse large B-Cell lymphoma or relapsed or refractory low grade follicular lymphoma.

In another embodiment, an immunomodulatory compound is administered in combination with taxotere, IL-2, IFN, GM-CSF, and/or dacarbazine to patients with various types or stages of melanoma.

In another embodiment, an immunomodulatory compound is administered alone or in combination with vinorelbine to patients with malignant mesothelioma, or stage IIIB non-small cell lung cancer with pleural implants or malignant pleural effusion mesothelioma syndrome.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of multiple myeloma in combination with dexamethasone, zoledronic acid, palmitronate, GM-CSF, biacin, vinblastine, melphalan, busulphan, cyclophosphamide, IFN, palmidronate, prednisone, bisphosphonate, celecoxib, arsenic trioxide, PEG INTRON-A, vincristine, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with relapsed or refractory multiple myeloma in combination with doxorubicin (Doxil®), vincristine and/or dexamethasone (Decadron®).

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of ovarian cancer such as peritoneal carcinoma, papillary serous carcinoma, refractory ovarian cancer or recurrent ovarian cancer, in combination with taxol, carboplatin, doxorubicin, gemcitabine, cisplatin, xeloda, paclitaxel, dexamethasone, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of prostate cancer, in combination with xeloda, 5 FU/LV, gemcitabine, irinotecan plus gemcitabine, cyclophosphamide, vincristine, dexamethasone, GM-CSF, celecoxib, taxotere, ganciclovir, paclitaxel, adriamycin, docetaxel, estramustine, Emcyt, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of

22

renal cell cancer, in combination with capecitabine, IFN, tamoxifen, IL-2, GM-CSF, Celebrex®, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of gynecologic, uterus or soft tissue sarcoma cancer in combination with IFN, a COX-2 inhibitor such as Celebrex®, and/or sulindac.

In another embodiment, an immunomodulatory compound is administered to patients with various types or stages of solid tumors in combination with celebrex, etoposide, cyclophosphamide, docetaxel, apecitabine, IFN, tamoxifen, IL-2, GM-CSF, or a combination thereof.

In another embodiment, an immunomodulatory compound is administered to patients with scleroderma or cutaneous vasculitis in combination with celebrex, etoposide, cyclophosphamide, docetaxel, apecitabine, IFN, tamoxifen, IL-2, GM-CSF, or a combination thereof.

This invention also encompasses a method of increasing the dosage of an anti-cancer drug or agent that can be safely and effectively administered to a patient, which comprises administering to a patient (e.g., a human) an immunomodulatory compound of the invention, or a pharmaceutically acceptable derivative, salt, solvate, clathrate, hydrate, or prodrug thereof. Patients that can benefit by this method are those likely to suffer from an adverse effect associated with anti-cancer drugs for treating a specific cancer of the skin, subcutaneous tissue, lymph nodes, brain, lung, liver, bone, intestine, colon, heart, pancreas, adrenal, kidney, prostate, breast, colorectal, or combinations thereof. The administration of an immunomodulatory compound of the invention alleviates or reduces adverse effects which are of such severity that it would otherwise limit the amount of anti-cancer drug.

In one embodiment, an immunomodulatory compound of the invention can be administered orally and daily in an amount of from about 0.1 to about 150 mg, and preferably from about 1 to about 50 mg, more preferably from about 2 to about 25 mg prior to, during, or after the occurrence of the adverse effect associated with the administration of an anti-cancer drug to a patient. In a particular embodiment, an immunomodulatory compound of the invention is administered in combination with specific agents such as heparin, aspirin, coumadin, or G-CSF to avoid adverse effects that are associated with anti-cancer drugs such as but not limited to neutropenia or thrombocytopenia.

In one embodiment, an immunomodulatory compound of the invention can be administered to patients with diseases and disorders associated with, or characterized by, undesired angiogenesis in combination with additional active ingredients including but not limited to anti-cancer drugs, anti-inflammatories, antihistamines, antibiotics, and steroids.

In another embodiment, this invention encompasses a method of treating, preventing and/or managing cancer, which comprises administering an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, in conjunction with (e.g. before, during, or after) conventional therapy including, but not limited to, surgery, immunotherapy, biological therapy, radiation therapy, or other non-drug based therapy presently used to treat, prevent or manage cancer. The combined use of the immunomodulatory compounds of the invention and conventional therapy may provide a unique treatment regimen that is unexpectedly effective in certain patients. Without being limited by theory, it is believed that immunomodulatory compounds of the invention may provide additive or synergistic effects when given concurrently with conventional therapy.

US 9,101,622 B2

23

As discussed elsewhere herein, the invention encompasses a method of reducing, treating and/or preventing adverse or undesired effects associated with conventional therapy including, but not limited to, surgery, chemotherapy, radiation therapy, hormonal therapy, biological therapy and immunotherapy. One or more immunomodulatory compounds of the invention and other active ingredient can be administered to a patient prior to, during, or after the occurrence of the adverse effect associated with conventional therapy.

In one embodiment, an immunomodulatory compound of the invention can be administered in an amount of from about 0.1 to about 150 mg, and preferably from about 1 to about 25 mg, more preferably from about 2 to about 10 mg orally and daily alone, or in combination with a second active agent disclosed herein (see, e.g., section 5.2), prior to, during, or after the use of conventional therapy.

In a specific embodiment of this method, an immunomodulatory compound of the invention and doxorubicin are administered to patients with non-small cell lung cancer who were previously treated with carbo/VP 16 and radiotherapy.

5.3.2 Use with Transplantation Therapy

Compounds of the invention can be used to reduce the risk of Graft Versus Host Disease (GVHD). Therefore, the invention encompasses a method of treating, preventing and/or managing cancer, which comprises administering the immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, in conjunction with transplantation therapy.

As those of ordinary skill in the art are aware, the treatment of cancer is often based on the stages and mechanism of the disease. For example, as inevitable leukemic transformation develops in certain stages of cancer, transplantation of peripheral blood stem cells, hematopoietic stem cell preparation or bone marrow may be necessary. The combined use of the immunomodulatory compound of the invention and transplantation therapy provides a unique and unexpected synergism. In particular, an immunomodulatory compound of the invention exhibits immunomodulatory activity that may provide additive or synergistic effects when given concurrently with transplantation therapy in patients with cancer.

An immunomodulatory compound of the invention can work in combination with transplantation therapy reducing complications associated with the invasive procedure of transplantation and risk of GVHD. This invention encompasses a method of treating, preventing and/or managing cancer which comprises administering to a patient (e.g., a human) an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof, before, during, or after the transplantation of umbilical cord blood, placental blood, peripheral blood stem cell, hematopoietic stem cell preparation or bone marrow. Examples of stem cells suitable for use in the methods of the invention are disclosed in U.S. provisional patent application No. 60/372,348, filed Apr. 12, 2002 by R. Hariri et al., the entirety of which is incorporated herein by reference.

In one embodiment of this method, an immunomodulatory compound of the invention is administered to patients with multiple myeloma before, during, or after the transplantation of autologous peripheral blood progenitor cell.

In another embodiment, an immunomodulatory compound is administered to patients with relapsing multiple myeloma after the stem cell transplantation.

24

In another embodiment, an immunomodulatory compound and prednisone are administered as maintenance therapy to patients with multiple myeloma following the transplantation of autologous stem cell.

In another embodiment, an immunomodulatory compound and dexamethasone are administered as salvage therapy for low risk post transplantation to patients with multiple myeloma.

In another embodiment, an immunomodulatory compound and dexamethasone are administered as maintenance therapy to patients with multiple myeloma following the transplantation of autologous bone marrow.

In another embodiment, an immunomodulatory compound is administered following the administration of high dose of melphalan and the transplantation of autologous stem cell to patients with chemotherapy responsive multiple myeloma.

In another embodiment, an immunomodulatory compound and PEG INTRO-A are administered as maintenance therapy to patients with multiple myeloma following the transplantation of autologous CD34-selected peripheral stem cell.

In another embodiment, an immunomodulatory compound is administered with post transplant consolidation chemotherapy to patients with newly diagnosed multiple myeloma to evaluate anti-angiogenesis.

In another embodiment, an immunomodulatory compound and dexamethasone are administered as maintenance therapy after DCEP consolidation, following the treatment with high dose of melphalan and the transplantation of peripheral blood stem cell to 65 years of age or older patients with multiple myeloma.

5.3.3 Cycling Therapy

In certain embodiments, the prophylactic or therapeutic agents of the invention are cyclically administered to a patient. Cycling therapy involves the administration of an active agent for a period of time, followed by a rest for a period of time, and repeating this sequential administration. Cycling therapy can reduce the development of resistance to one or more of the therapies, avoid or reduce the side effects of one of the therapies, and/or improves the efficacy of the treatment.

Consequently, in one specific embodiment of the invention, an immunomodulatory compound of the invention is administered daily in a single or divided doses in a four to six week cycle with a rest period of about a week or two weeks. The invention further allows the frequency, number, and length of dosing cycles to be increased. Thus, another specific embodiment of the invention encompasses the administration of an immunomodulatory compound of the invention for more cycles than are typical when it is administered alone. In yet another specific embodiment of the invention, an immunomodulatory compound of the invention is administered for a greater number of cycles that would typically cause dose-limiting toxicity in a patient to whom a second active ingredient is not also being administered.

In one embodiment, an immunomodulatory compound of the invention is administered daily and continuously for three or four weeks at a dose of from about 0.1 to about 150 mg/d followed by a break of one or two weeks. Actimid™ is preferably administered daily and continuously at an initial dose of 0.1 to 5 mg/d with dose escalation (every week) by 1 to 10 mg/d to a maximum dose of 50 mg/d for as long as therapy is tolerated. In a particular embodiment, Revimid™ is administered in an amount of about 5, 10, or 25 mg/day, preferably in an amount of about 10 mg/day for three to four weeks, followed by one week or two weeks of rest in a four or six week cycle.

In one embodiment of the invention, an immunomodulatory compound of the invention and a second active ingredient are administered orally, with administration of an immunomodulatory compound of the invention occurring 30 to 60 minutes prior to a second active ingredient, during a cycle of four to six weeks. In another embodiment of the invention, the combination of an immunomodulatory compound of the invention and a second active ingredient is administered by intravenous infusion over about 90 minutes every cycle. In a specific embodiment, one cycle comprises the administration of from about 10 to about 25 mg/day of Revimid™ and from about 50 to about 200 mg/m²/day of a second active ingredient daily for three to four weeks and then one or two weeks of rest. In another specific embodiment, each cycle comprises the administration of from about 5 to about 10 mg/day of Actimid™ and from about 50 to about 200 mg/m²/day of a second active ingredient for 3 to 4 weeks followed by one or two weeks of rest. Typically, the number of cycles during which the combinatorial treatment is administered to a patient will be from about one to about 24 cycles, more typically from about two to about 16 cycles, and even more typically from about four to about three cycles.

5.4 Pharmaceutical Compositions and Dosage Forms

Pharmaceutical compositions can be used in the preparation of individual, single unit dosage forms. Pharmaceutical compositions and dosage forms of the invention comprise an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof. Pharmaceutical compositions and dosage forms of the invention can further comprise one or more excipients.

Pharmaceutical compositions and dosage forms of the invention can also comprise one or more additional active ingredients. Consequently, pharmaceutical compositions and dosage forms of the invention comprise the active ingredients disclosed herein (e.g., an immunomodulatory compound and a second active agent). Examples of optional second, or additional, active ingredients are disclosed herein (see, e.g., section 5.2).

Single unit dosage forms of the invention are suitable for oral, mucosal (e.g., nasal, sublingual, vaginal, buccal, or rectal), parenteral (e.g., subcutaneous, intravenous, bolus injection, intramuscular, or intraarterial), topical (e.g., eye drops or other ophthalmic preparations), transdermal or transcutaneous administration to a patient. Examples of dosage forms include, but are not limited to: tablets; caplets; capsules, such as soft elastic gelatin capsules; cachets; troches; lozenges; dispersions; suppositories; powders; aerosols (e.g., nasal sprays or inhalers); gels; liquid dosage forms suitable for oral or mucosal administration to a patient, including suspensions (e.g., aqueous or non-aqueous liquid suspensions, oil-in-water emulsions, or a water-in-oil liquid emulsions), solutions, and elixirs; liquid dosage forms suitable for parenteral administration to a patient; eye drops or other ophthalmic preparations suitable for topical administration; and sterile solids (e.g., crystalline or amorphous solids) that can be reconstituted to provide liquid dosage forms suitable for parenteral administration to a patient.

The composition, shape, and type of dosage forms of the invention will typically vary depending on their use. For example, a dosage form used in the acute treatment of a disease may contain larger amounts of one or more of the active ingredients it comprises than a dosage form used in the chronic treatment of the same disease. Similarly, a parenteral dosage form may contain smaller amounts of one or more of the active ingredients it comprises than an oral dosage form used to treat the same disease. These and other ways in which

specific dosage forms encompassed by this invention will vary from one another will be readily apparent to those skilled in the art. See, e.g., *Remington's Pharmaceutical Sciences*, 18th ed., Mack Publishing, Easton, Pa. (1990).

Typical pharmaceutical compositions and dosage forms comprise one or more excipients. Suitable excipients are well known to those skilled in the art of pharmacy, and non-limiting examples of suitable excipients are provided herein. Whether a particular excipient is suitable for incorporation into a pharmaceutical composition or dosage form depends on a variety of factors well known in the art including, but not limited to, the way in which the dosage form will be administered to a patient. For example, oral dosage forms such as tablets may contain excipients not suited for use in parenteral dosage forms. The suitability of a particular excipient may also depend on the specific active ingredients in the dosage form. For example, the decomposition of some active ingredients may be accelerated by some excipients such as lactose, or when exposed to water. Active ingredients that comprise primary or secondary amines are particularly susceptible to such accelerated decomposition. Consequently, this invention encompasses pharmaceutical compositions and dosage forms that contain little, if any, lactose other mono- or disaccharides. As used herein, the term "lactose-free" means that the amount of lactose present, if any, is insufficient to substantially increase the degradation rate of an active ingredient.

Lactose-free compositions of the invention can comprise excipients that are well known in the art and are listed, for example, in the *U.S. Pharmacopeia* (USP) 25-NF20 (2002). In general, lactose-free compositions comprise active ingredients, a binder/filler, and a lubricant in pharmaceutically compatible and pharmaceutically acceptable amounts. Preferred lactose-free dosage forms comprise active ingredients, microcrystalline cellulose, pre-gelatinized starch, and magnesium stearate.

This invention further encompasses anhydrous pharmaceutical compositions and dosage forms comprising active ingredients, since water can facilitate the degradation of some compounds. For example, the addition of water (e.g., 5%) is widely accepted in the pharmaceutical arts as a means of simulating long-term storage in order to determine characteristics such as shelf-life or the stability of formulations over time. See, e.g., Jens T. Carstensen, *Drug Stability: Principles & Practice*, 2d. Ed., Marcel Dekker, NY, N.Y., 1995, pp. 379-80. In effect, water and heat accelerate the decomposition of some compounds. Thus, the effect of water on a formulation can be of great significance since moisture and/or humidity are commonly encountered during manufacture, handling, packaging, storage, shipment, and use of formulations.

Anhydrous pharmaceutical compositions and dosage forms of the invention can be prepared using anhydrous or low moisture containing ingredients and low moisture or low humidity conditions. Pharmaceutical compositions and dosage forms that comprise lactose and at least one active ingredient that comprises a primary or secondary amine are preferably anhydrous if substantial contact with moisture and/or humidity during manufacturing, packaging, and/or storage is expected.

An anhydrous pharmaceutical composition should be prepared and stored such that its anhydrous nature is maintained. Accordingly, anhydrous compositions are preferably packaged using materials known to prevent exposure to water such that they can be included in suitable formulary kits. Examples

27

of suitable packaging include, but are not limited to, hermetically sealed foils, plastics, unit dose containers (e.g., vials), blister packs, and strip packs.

The invention further encompasses pharmaceutical compositions and dosage forms that comprise one or more compounds that reduce the rate by which an active ingredient will decompose. Such compounds, which are referred to herein as “stabilizers,” include, but are not limited to, antioxidants such as ascorbic acid, pH buffers, or salt buffers.

Like the amounts and types of excipients, the amounts and specific types of active ingredients in a dosage form may differ depending on factors such as, but not limited to, the route by which it is to be administered to patients. However, typical dosage forms of the invention comprise an immunomodulatory compound of the invention or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof in an amount of from about 0.10 to about 150 mg. Typical dosage forms comprise an immunomodulatory compound of the invention or a pharmaceutically acceptable salt, solvate, hydrate, stereoisomer, clathrate, or prodrug thereof in an amount of about 0.1, 1, 2, 5, 7.5, 10, 12.5, 15, 17.5, 20, 25, 50, 100, 150 or 200 mg. In a particular embodiment, a preferred dosage form comprises 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™) in an amount of about 1, 2, 5, 10, 25 or 50 mg. In a specific embodiment, a preferred dosage form comprises 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) in an amount of about 5, 10, 25 or 50 mg. Typical dosage forms comprise the second active ingredient in an amount of 1 to about 1000 mg, from about 5 to about 500 mg, from about 10 to about 350 mg, or from about 50 to about 200 mg. Of course, the specific amount of the anti-cancer drug will depend on the specific agent used, the type of cancer being treated or managed, and the amount(s) of an immunomodulatory compound of the invention and any optional additional active agents concurrently administered to the patient.

5.4.1 Oral Dosage Forms

Pharmaceutical compositions of the invention that are suitable for oral administration can be presented as discrete dosage forms, such as, but are not limited to, tablets (e.g., chewable tablets), caplets, capsules, and liquids (e.g., flavored syrups). Such dosage forms contain predetermined amounts of active ingredients, and may be prepared by methods of pharmacy well known to those skilled in the art. See generally, *Remington's Pharmaceutical Sciences*, 18th ed., Mack Publishing, Easton, Pa. (1990).

Typical oral dosage forms of the invention are prepared by combining the active ingredients in an intimate admixture with at least one excipient according to conventional pharmaceutical compounding techniques. Excipients can take a wide variety of forms depending on the form of preparation desired for administration. For example, excipients suitable for use in oral liquid or aerosol dosage forms include, but are not limited to, water, glycols, oils, alcohols, flavoring agents, preservatives, and coloring agents. Examples of excipients suitable for use in solid oral dosage forms (e.g., powders, tablets, capsules, and caplets) include, but are not limited to, starches, sugars, micro-crystalline cellulose, diluents, granulating agents, lubricants, binders, and disintegrating agents.

Because of their ease of administration, tablets and capsules represent the most advantageous oral dosage unit forms, in which case solid excipients are employed. If desired, tablets can be coated by standard aqueous or nonaqueous techniques. Such dosage forms can be prepared by any of the methods of pharmacy. In general, pharmaceutical compositions and dosage forms are prepared by uniformly and inti-

28

ately admixing the active ingredients with liquid carriers, finely divided solid carriers, or both, and then shaping the product into the desired presentation if necessary.

For example, a tablet can be prepared by compression or molding. Compressed tablets can be prepared by compressing in a suitable machine the active ingredients in a free-flowing form such as powder or granules, optionally mixed with an excipient. Molded tablets can be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

Examples of excipients that can be used in oral dosage forms of the invention include, but are not limited to, binders, fillers, disintegrants, and lubricants. Binders suitable for use in pharmaceutical compositions and dosage forms include, but are not limited to, corn starch, potato starch, or other starches, gelatin, natural and synthetic gums such as acacia, sodium alginate, alginic acid, other alginates, powdered tragacanth, guar gum, cellulose and its derivatives (e.g., ethyl cellulose, cellulose acetate, carboxymethyl cellulose calcium, sodium carboxymethyl cellulose), polyvinyl pyrrolidone, methyl cellulose, pre-gelatinized starch, hydroxypropyl methyl cellulose, (e.g., Nos. 2208, 2906, 2910), microcrystalline cellulose, and mixtures thereof.

Suitable forms of microcrystalline cellulose include, but are not limited to, the materials sold as AVICEL-PH-101, AVICEL-PH-103 AVICEL RC-581, AVICEL-PH-105 (available from FMC Corporation, American Viscose Division, Avicel Sales, Marcus Hook, Pa.), and mixtures thereof. A specific binder is a mixture of microcrystalline cellulose and sodium carboxymethyl cellulose sold as AVICEL RC-581. Suitable anhydrous or low moisture excipients or additives include AVICEL-PH-103™ and Starch 1500 LM.

Examples of fillers suitable for use in the pharmaceutical compositions and dosage forms disclosed herein include, but are not limited to, talc, calcium carbonate (e.g., granules or powder), microcrystalline cellulose, powdered cellulose, dextrates, kaolin, mannitol, silicic acid, sorbitol, starch, pre-gelatinized starch, and mixtures thereof. The binder or filler in pharmaceutical compositions of the invention is typically present in from about 50 to about 99 weight percent of the pharmaceutical composition or dosage form.

Disintegrants are used in the compositions of the invention to provide tablets that disintegrate when exposed to an aqueous environment. Tablets that contain too much disintegrant may disintegrate in storage, while those that contain too little may not disintegrate at a desired rate or under the desired conditions. Thus, a sufficient amount of disintegrant that is neither too much nor too little to detrimentally alter the release of the active ingredients should be used to form solid oral dosage forms of the invention. The amount of disintegrant used varies based upon the type of formulation, and is readily discernible to those of ordinary skill in the art. Typical pharmaceutical compositions comprise from about 0.5 to about 15 weight percent of disintegrant, preferably from about 1 to about 5 weight percent of disintegrant.

Disintegrants that can be used in pharmaceutical compositions and dosage forms of the invention include, but are not limited to, agar-agar, alginic acid, calcium carbonate, microcrystalline cellulose, croscarmellose sodium, crospovidone, polacrillin potassium, sodium starch glycolate, potato or tapioca starch, other starches, pre-gelatinized starch, other starches, clays, other alginates, other celluloses, gums, and mixtures thereof.

Lubricants that can be used in pharmaceutical compositions and dosage forms of the invention include, but are not limited to, calcium stearate, magnesium stearate, mineral oil, light mineral oil, glycerin, sorbitol, mannitol, polyethylene

glycol, other glycols, stearic acid, sodium lauryl sulfate, talc, hydrogenated vegetable oil (e.g., peanut oil, cottonseed oil, sunflower oil, sesame oil, olive oil, corn oil, and soybean oil), zinc stearate, ethyl oleate, ethyl laureate, agar, and mixtures thereof. Additional lubricants include, for example, a syloid silica gel (AEROSIL200, manufactured by W. R. Grace Co. of Baltimore, Md.), a coagulated aerosol of synthetic silica (marketed by Degussa Co. of Plano, Tex.), CAB-O-SIL (a pyrogenic silicon dioxide product sold by Cabot Co. of Boston, Mass.), and mixtures thereof. If used at all, lubricants are typically used in an amount of less than about 1 weight percent of the pharmaceutical compositions or dosage forms into which they are incorporated.

A preferred solid oral dosage form of the invention comprises an immunomodulatory compound of the invention, anhydrous lactose, microcrystalline cellulose, polyvinylpyrrolidone, stearic acid, colloidal anhydrous silica, and gelatin.

5.4.2 Delayed Release Dosage Forms

Active ingredients of the invention can be administered by controlled release means or by delivery devices that are well known to those of ordinary skill in the art. Examples include, but are not limited to, those described in U.S. Pat. Nos. 3,845,770; 3,916,899; 3,536,809; 3,598,123; and 4,008,719, 5,674,533, 5,059,595, 5,591,767, 5,120,548, 5,073,543, 5,639,476, 5,354,556, and 5,733,566, each of which is incorporated herein by reference. Such dosage forms can be used to provide slow or controlled-release of one or more active ingredients using, for example, hydropropylmethyl cellulose, other polymer matrices, gels, permeable membranes, osmotic systems, multilayer coatings, microparticles, liposomes, microspheres, or a combination thereof to provide the desired release profile in varying proportions. Suitable controlled-release formulations known to those of ordinary skill in the art, including those described herein, can be readily selected for use with the active ingredients of the invention. The invention thus encompasses single unit dosage forms suitable for oral administration such as, but not limited to, tablets, capsules, gelcaps, and caplets that are adapted for controlled-release.

All controlled-release pharmaceutical products have a common goal of improving drug therapy over that achieved by their non-controlled counterparts. Ideally, the use of an optimally designed controlled-release preparation in medical treatment is characterized by a minimum of drug substance being employed to cure or control the condition in a minimum amount of time. Advantages of controlled-release formulations include extended activity of the drug, reduced dosage frequency, and increased patient compliance. In addition, controlled-release formulations can be used to affect the time of onset of action or other characteristics, such as blood levels of the drug, and can thus affect the occurrence of side (e.g., adverse) effects.

Most controlled-release formulations are designed to initially release an amount of drug (active ingredient) that promptly produces the desired therapeutic effect, and gradually and continually release of other amounts of drug to maintain this level of therapeutic or prophylactic effect over an extended period of time. In order to maintain this constant level of drug in the body, the drug must be released from the dosage form at a rate that will replace the amount of drug being metabolized and excreted from the body. Controlled-release of an active ingredient can be stimulated by various conditions including, but not limited to, pH, temperature, enzymes, water, or other physiological conditions or compounds.

5.4.3 Parenteral Dosage Forms

Parenteral dosage forms can be administered to patients by various routes including, but not limited to, subcutaneous, intravenous (including bolus injection), intramuscular, and intraarterial. Because their administration typically bypasses patients' natural defenses against contaminants, parenteral dosage forms are preferably sterile or capable of being sterilized prior to administration to a patient. Examples of parenteral dosage forms include, but are not limited to, solutions ready for injection, dry products ready to be dissolved or suspended in a pharmaceutically acceptable vehicle for injection, suspensions ready for injection, and emulsions.

Suitable vehicles that can be used to provide parenteral dosage forms of the invention are well known to those skilled in the art. Examples include, but are not limited to: Water for Injection USP; aqueous vehicles such as, but not limited to, Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, and Lactated Ringer's Injection; water-miscible vehicles such as, but not limited to, ethyl alcohol, polyethylene glycol, and polypropylene glycol; and non-aqueous vehicles such as, but not limited to, corn oil, cottonseed oil, peanut oil, sesame oil, ethyl oleate, isopropyl myristate, and benzyl benzoate.

Compounds that increase the solubility of one or more of the active ingredients disclosed herein can also be incorporated into the parenteral dosage forms of the invention. For example, cyclodextrin and its derivatives can be used to increase the solubility of an immunomodulatory compound of the invention and its derivatives. See, e.g., U.S. Pat. No. 5,134,127, which is incorporated herein by reference.

5.4.4 Topical and Mucosal Dosage Forms

Topical and mucosal dosage forms of the invention include, but are not limited to, sprays, aerosols, solutions, emulsions, suspensions, eye drops or other ophthalmic preparations, or other forms known to one of skill in the art. See, e.g., *Remington's Pharmaceutical Sciences*, 16th and 18th eds., Mack Publishing, Easton, Pa. (1980 & 1990); and *Introduction to Pharmaceutical Dosage Forms*, 4th ed., Lea & Febiger, Philadelphia (1985). Dosage forms suitable for treating mucosal tissues within the oral cavity can be formulated as mouthwashes or as oral gels.

Suitable excipients (e.g., carriers and diluents) and other materials that can be used to provide topical and mucosal dosage forms encompassed by this invention are well known to those skilled in the pharmaceutical arts, and depend on the particular tissue to which a given pharmaceutical composition or dosage form will be applied. With that fact in mind, typical excipients include, but are not limited to, water, acetone, ethanol, ethylene glycol, propylene glycol, butane-1,3-diol, isopropyl myristate, isopropyl palmitate, mineral oil, and mixtures thereof to form solutions, emulsions or gels, which are non-toxic and pharmaceutically acceptable. Moisturizers or humectants can also be added to pharmaceutical compositions and dosage forms if desired. Examples of such additional ingredients are well known in the art. See, e.g., *Remington's Pharmaceutical Sciences*, 16th and 18th eds., Mack Publishing, Easton, Pa. (1980 & 1990).

The pH of a pharmaceutical composition or dosage form may also be adjusted to improve delivery of one or more active ingredients. Similarly, the polarity of a solvent carrier, its ionic strength, or tonicity can be adjusted to improve delivery. Compounds such as stearates can also be added to pharmaceutical compositions or dosage forms to advantageously alter the hydrophilicity or lipophilicity of one or more active ingredients so as to improve delivery. In this regard, stearates can serve as a lipid vehicle for the formulation, as an emulsifying agent or surfactant, and as a delivery-

enhancing or penetration-enhancing agent. Different salts, hydrates or solvates of the active ingredients can be used to further adjust the properties of the resulting composition.

5.4.5 Kits

Typically, active ingredients of the invention are preferably not administered to a patient at the same time or by the same route of administration. This invention therefore encompasses kits which, when used by the medical practitioner, can simplify the administration of appropriate amounts of active ingredients to a patient.

A typical kit of the invention comprises a dosage form of an immunomodulatory compound of the invention, or a pharmaceutically acceptable salt salt, solvate, hydrate, stereoisomer, prodrug, or clathrate thereof. Kits encompassed by this invention can further comprise additional active ingredients such as oblimersen (Genasense®), melphalan, G-CSF, GM-CSF, EPO, topotecan, dacarbazine, irinotecan, taxotere, IFN, COX-2 inhibitor, pentoxifylline, ciprofloxacin, dexamethasone, IL2, IL8, IL18, Ara-C, vinorelbine, isotretinoin, 13 cis-retinoic acid, or a pharmacologically active mutant or derivative thereof, or a combination thereof. Examples of the additional active ingredients include, but are not limited to, those disclosed herein (see, e.g., section 5.2).

Kits of the invention can further comprise devices that are used to administer the active ingredients. Examples of such devices include, but are not limited to, syringes, drip bags, patches, and inhalers.

Kits of the invention can further comprise cells or blood for transplantation as well as pharmaceutically acceptable vehicles that can be used to administer one or more active ingredients. For example, if an active ingredient is provided in a solid form that must be reconstituted for parenteral administration, the kit can comprise a sealed container of a suitable vehicle in which the active ingredient can be dissolved to form a particulate-free sterile solution that is suitable for parenteral administration. Examples of pharmaceutically acceptable vehicles include, but are not limited to: Water for Injection USP; aqueous vehicles such as, but not limited to, Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, and Lactated Ringer's Injection; water-miscible vehicles such as, but not limited to, ethyl alcohol, polyethylene glycol, and polypropylene glycol; and non-aqueous vehicles such as, but not limited to, corn oil, cottonseed oil, peanut oil, sesame oil, ethyl oleate, isopropyl myristate, and benzyl benzoate.

6. EXAMPLES

Certain embodiments of the invention are illustrated by the following non-limiting examples.

6.1 Modulation of Cytokine Production

A series of non-clinical pharmacology and toxicology studies have been performed to support the clinical evaluation of an immunomodulatory compound of the invention in human subjects. These studies were performed in accordance with internationally recognized guidelines for study design and in compliance with the requirements of Good Laboratory Practice (GLP), unless otherwise noted.

Inhibition of TNF- α production following LPS-stimulation of human PBMC and human whole blood by 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™), 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione and thalidomide (Revimid™) was investigated in vitro (Muller et al., *Bioorg. Med. Chem. Lett.* 9:1625-1630, 1999). The IC₅₀'s of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione for inhibiting production of TNF- α following LPS-stimulation of PBMC and human whole blood were ~24 nM (6.55 ng/mL) and ~25 nM (6.83 ng/mL), respectively. In vitro studies suggest a pharmacological activity profile for 3-(4-amino-1-oxo-1,3-dihydro-

isoindol-2-yl)-piperidine-2,6-dione that is similar to, but at least 200 times more potent than, thalidomide. In vitro studies have also demonstrated that concentrations of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione of 2.73 to 27.3 ng/mL (0.01 to 0.1 μ M) achieved 50% inhibition of the proliferation of MM.1S and Hs Sultan cells.

The IC₅₀'s of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione for inhibiting production of TNF- α following LPS-stimulation of PBMC and human whole blood were ~100 nM (25.9 ng/mL) and ~480 nM (103.6 ng/mL), respectively. Thalidomide, in contrast, had an IC₅₀ of ~194 μ M (50.2 μ g/mL) for inhibiting production of TNF- α following LPS-stimulation of PBMC. In vitro studies suggest a pharmacological activity profile for 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione that is similar to, but 50 to 2000 times more potent than, thalidomide. It has been shown that the compound is approximately 50-100 times more potent than thalidomide in stimulating the proliferation of T-cells following primary induction by T-cell receptor (TCR) activation. 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione is also approximately 50 to 100 times more potent than thalidomide in augmenting the production of IL-2 and IFN- γ following TCR activation of PBMC (IL-2) or T-cells (IFN- γ). In addition, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione exhibited dose-dependent inhibition of LPS-stimulated production of the pro-inflammatory cytokines TNF- α , IL-1 β , and IL-6 by PBMC while it increased production of the anti-inflammatory cytokine IL-10.

6.2 Inhibition of MM Cell Proliferation

The ability of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) and thalidomide for comparison to effect the proliferation of MM cell lines has been investigated in an in vitro study. Uptake [³H]-thymidine by different MM cell lines MM.1S, Hs Sultan, U266 and RPMI-8226) was measured as an indicator of cell proliferation. Cells were incubated in the presence of compounds for 48 hours; [³H]-thymidine was included for the last 8 hours of the incubation period. Addition of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione to MM.1S and Hs Sultan cells resulted in 50% inhibition of cell proliferation at concentrations of 0.4 μ M and 1 μ M, respectively. In contrast, addition of thalidomide at concentrations up to 100 μ M resulted in only 15% and 20% inhibition of cell proliferation in MM.1S and Hs Sultan cells, respectively. These data are summarized in FIG. 1.

6.3 Toxicology Studies

The effects of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) on cardiovascular and respiratory function are investigated in anesthetized dogs. Two groups of Beagle dogs (2/sex/group) are used. One group receives three doses of vehicle only and the other receives three ascending doses of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (2, 10, and 20 mg/kg). In all cases, doses of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione or vehicle are successively administered via infusion through the jugular vein separated by intervals of at least 30 minutes.

The cardiovascular and respiratory changes induced by 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione are minimal at all doses when compared to the vehicle control group. The only statistically significant difference between the vehicle and treatment groups is a small increase in arterial blood pressure (from 94 mmHg to 101 mmHg) following administration of the low dose of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. This effect lasts approximately 15 minutes and is not seen at higher doses. Deviations in femoral blood flow, res-

piratory parameters, and Qtc interval are common to both the control and treated groups and are not considered treatment-related.

6.4 Cycling Therapy in Patients

In a specific embodiment, an immunomodulatory compound of the invention are cyclically administered to patients with cancer. Cycling therapy involves the administration of a first agent for a period of time, followed by a rest for a period of time and repeating this sequential administration. Cycling therapy can reduce the development of resistance to one or more of the therapies, avoid or reduce the side effects of one of the therapies, and/or improves the efficacy of the treatment.

In a specific embodiment, prophylactic or therapeutic agents are administered in a cycle of about 4 to 6 weeks, about once or twice every day. One cycle can comprise the administration of a therapeutic or prophylactic agent for three to four weeks and at least a week or two weeks of rest. The number of cycles administered is from about one to about 24 cycles, more typically from about two to about 16 cycles, and more typically from about four to about eight cycles.

For example, in a cycle of four weeks, on day 1, the administration of 25 mg/d of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione is started. On day 22, the administration of the compound is stopped for a week of rest. On day 29, the administration of 25 mg/d 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidin-2,6-dione is begun.

6.5 Clinical Studies in Patients

6.5.1 Treatment of Relapsed Multiple Myeloma

4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (Actimid™) was administered to patients with relapsed/refractory multiple myeloma. The study was conducted in compliance with Good Clinical Practices. Patients were at least 18 years old, had been diagnosed with multiple myeloma (with paraprotein in serum and/or urine), and were considered refractory to treatment after at least two cycles of treatment, or have relapsed after two cycles of treatment.

Patients who have progressive disease, according to the Southwest Oncology Group (SWOG) criteria, on their prior regimen are considered treatment refractory. Relapse following remission is defined as >25% increase in M component from baseline levels; reappearance of the M paraprotein that had previously disappeared; or a definite increase in the size and number of lytic bone lesions recognized on radiographs. Patients may have had prior therapy with thalidomide, provided they were able to tolerate the treatment. A Zubrod performance status of 0 to 2 is required for all patients.

4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione is administered to patients at doses of 1, 2, 5, or 10 mg/day for up to four weeks; at each dose level, three patients are initially enrolled. Dosing occurs at approximately the same time each morning; all doses are administered in the fasted state (no eating for at least two hours prior to dosing and two hours after dosing). 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione doses are administered in an ascending fashion such that patients in the first cohort receive the lowest

dose of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione (1 mg/day) and escalation to the next higher dose level occurs only following the establishment of safety and tolerability at the current dose. If one out of three patients at any dose level experience dose limiting toxicity (DLT), three additional patients are enrolled at that dose. If none of the three additional patients experience DLT, escalation to the next dose level occurs; dose escalations continue in a similar fashion until the MTD is established or the maximum daily dose (10 mg/day) is attained. However, if one of the three additional patients enrolled experiences DLT, the MTD has been reached. If two or more of the three additional patients enrolled experience DLT, the MTD is judged to have been exceeded and three additional patients are enrolled at the preceding dose level to confirm the MTD. Once the MTD has been identified, four additional patients are enrolled at that dose level so that a total of 10 patients is treated at the MTD.

Blood sampling for analysis of pharmacokinetic parameters is performed on Days 1 and 28 according to the following sampling schedule: pre-dose, 0.25, 0.5, 0.75, 1, 1.5, 2, 2.5, 3, 4, 6, 8, 10, 12, 18, and 24 hours post-dose. An additional blood sample is collected at each weekly visit for the determination of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione levels. Total urine collections are also made with urine pooled according to the following time intervals post-dose: 0 to 4, 4 to 8, 8 to 12, and 12 to 24 hours. Safety assessments are made by monitoring adverse events, vital signs, ECGs, clinical laboratory evaluations (blood chemistry, hematology, lymphocyte phenotyping, and urinalysis), and physical examination at specific times during the study.

Results of interim pharmacokinetic analyses obtained following single- and multiple-dose administration of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione to multiple myeloma patients are presented below in Tables 1 and 2. These data show that 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione was steadily absorbed at all dose levels in relapsed multiple myeloma patients. Maximum plasma concentrations occurred at a median T_{max} of between 2.5 and 2.8 hours post-dose at Day 1 and between 3 and 4 hours post-dose at Week 4. At all doses, plasma concentrations declined in a monophasic manner after reaching C_{max} . The start of the elimination phase occurred between 3 and 10 hours post-dose at Day 1 and Week 4, respectively.

These data also showed that after 4 weeks of dosing, 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione accumulated to a small extent (mean accumulation ratios ~1.02 to 1.52 and ~0.94 to 1.62 for C_{max} and $AUC_{(0-\tau)}$, respectively). There was almost a dose proportional increase in $AUC_{(0-\tau)}$ and C_{max} values with increasing dose. A five-fold higher dose of 4-(amino)-2-(2,6-dioxo(3-piperidyl))-isoindoline-1,3-dione produced a 3.2- and 2.2-fold increase in C_{max} at Day 1 and Week 4, respectively. Similarly, a 5-fold increase in dose resulted in a 3.6- and 2.3-fold increase in $AUC_{(0-\tau)}$, at Day 1 and Week 4, respectively.

TABLE 1

| Pharmacokinetic parameters of Actimid™ in relapsed multiple myeloma patients | | | | |
|--|-----------|----------------|----------------|-----------------|
| Parameter | | 1 mg | 2 mg | 5 mg |
| | | (N = 6) | (N = 2) | (N = 3) |
| Day 1 | | | | |
| C_{max} | ng/mL | 15.03 (4.04) | 24.4* (12.1) | 48.56 (14.03) |
| t_{max} | h | 3.3 (2.6) | 2.7* (0.3) | 2.3 (0.3) |
| $AUC_{(0-\infty)}$ | ng · h/mL | 152.90 (36.62) | 279.18 (51.10) | 593.10 (335.23) |

US 9,101,622 B2

35

TABLE 1-continued

| Pharmacokinetic parameters of Actimid™ in relapsed multiple myeloma patients | | | | |
|--|--------|-----------------|-----------------|-----------------|
| Parameter | | 1 mg (N = 6) | 2 mg (N = 2) | 5 mg (N = 3) |
| AUC _(0-τ) | | 134.21 (27.14) | 249.57 (29.26) | 520.94 (267.32) |
| t _{1/2} | h | 7.3 (3.4) | 6.3 (1.4) | 6.5 (2.2) |
| CL/F | mL/min | 114.75 (29.20) | 121.43 (22.22) | 182.31 (117.06) |
| Vz/f | L | 69.55 (44.97) | 65.31 (2.80) | 87.24 (22.61) |

τ = 24 hours

N/A = not available

36

TABLE 2

| Pharmacokinetic parameters of Actimid™ following multiple oral doses (1, 2, and 5 mg/day) in relapsed multiple myeloma patients | | | | |
|---|-----------|-----------------|-----------------|-----------------|
| Parameter | | 1 mg (N = 5) | 2 mg (N = 2) | 5 mg (N = 3) |
| Week 4 | | | | |
| C _{max} | ng/mL | 23.20 (7.48) | 30.05* (15.64) | 58.07 (38.08) |
| t _{max} | h | 3.6 (1.5) | 2.8* (0.3) | 5.0 (2.6) |
| AUC _(0-∞) | ng · h/mL | N/A | N/A | N/A |
| AUC _(0-τ) | | 239.31 (122.59) | 269.36 (186.34) | 597.24 (354.23) |
| t _{1/2} | h | 6.2* (0.6) | 7.7 (2.8) | 7.8 (4.0) |
| CL/F | mL/min | 87.85 (48.48) | 162.68 (112.54) | 207.50 (175.41) |
| Vz/f | L | 41.35* (8.84) | 95.04 (35.39) | 103.95 (27.25) |

τ = 24 hours

N/A = not available

*N = 3 patients

6.5.2 Treatment of Relapsed Multiple Myeloma

Two Phase 1 clinical studies of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) have been conducted to identify the maximum tolerated dose (MTD) in patients with refractory or relapsed multiple myeloma. These studies have also characterized the safety profile of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione when ascending doses of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione were given orally for up to 4 weeks. Patients started 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione treatment at 5 mg/day with subsequent escalation to 10, 25, and 50 mg/day. Patients were enrolled for 28 days at their assigned dose, with the option of extended treatment for those who did not exhibit disease progression or experience dose limiting toxicity (DLT). Patients were evaluated for adverse events at each visit and the severity of these events was graded according to the National Cancer Institute (NCI) Common Toxicity Criteria. Patients were discontinued if they experienced DLT (Grade 3 or greater non-hematological, or Grade 4 hematological toxicity).

In this study, 27 patients were enrolled. All patients had relapsed multiple myeloma and 18 (72%) were refractory to salvage therapy. Among these patients, 15 had undergone prior autologous stem cell transplantation and 16 patients had received prior thalidomide treatment. The median number of prior regimens was 3 (range 2 to 6).

Blood and urine samples were collected for analysis of pharmacokinetic parameters on Days 1 and 28. Blood samples were collected according to the following sampling schedule: pre-dose, 0.25, 0.5, 0.75, 1, 1.5, 2, 2.5, 3, 4, 6, 8, 10, 12, 18, and 24 hours post-dose. In addition, a blood sample was collected at each weekly clinic visit for 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione determination. Total urine was collected and pooled according to

the following time intervals post-dose: 0 to 4, 4 to 8, 8 to 12, and 12 to 24 hours. Response to treatment was assessed by M-protein quantification (by immunoelectrophoresis) from serum and a 24-hour urine collection, with creatinine clearance and 24-hour protein calculations undertaken at screening, baseline, Weeks 2 and 4, and monthly thereafter (or upon early termination). Bone marrow aspirations and/or tissue biopsy are also performed at Months 3, 6 and 12 if a patient's paraprotein serum concentration or 24-hour urine protein excretion declined to the next lower level, based on best response criteria. Preliminary results for the 28-day treatment period are summarized below.

Preliminary pharmacokinetic analyses based on these two studies indicated that AUC and C_{max} values increase proportionally with dose following single and multiple doses in multiple myeloma patients (as was seen in healthy volunteers). Further, there was no evidence of accumulation with multiple dosing as single dose AUC_(0-τ) was comparable to multiple dose AUC_(0-τ) following the same dose of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. Similar to healthy volunteer studies, double peaks were observed. Exposure in multiple myeloma patients appeared to be slightly higher based on C_{max} and AUC values as compared to healthy male volunteers while clearance in multiple myeloma patients was lower than it was in healthy volunteers, consistent with their poorer renal function (both as a consequence of their age and their disease). Finally, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione half-life in patients was shorter than in healthy volunteers (mean 8 hours, ranging up to 17 hours).

In this study, the first cohort of 3 patients was treated for 28 days at 5 mg/day without any dose limiting toxicity (DLT). The second cohort of 3 patients subsequently commenced therapy at 10 mg/day. Patients in the second 10 mg/day of

US 9,101,622 B2

37

3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione cohort tolerated treatment well.

6.5.3 Treatment of Solid Tumors

Study with 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) was conducted in patients with varying types of solid tumors, including malignant melanoma (13), carcinoma of the pancreas (2), carcinoid-unknown primary (1), renal carcinoma (1), breast carcinoma (1) and NSCLC (2). Patients received 5 mg/day 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione for seven days and are subsequently escalated every seven days to 10 mg/day, 25 mg/day, and 50 mg/day for a total of 4 weeks of treatment. Patients who, experienced clinical benefit were permitted to continue on treatment as Named Patients.

The study initially enrolled 20 patients and was subsequently amended to enroll 16 additional patients (adrenal carcinoma, NSCLC, malignant mesothelioma, breast cancer, malignant melanoma (8), renal cell cancer (4)) at a higher dose. The 16 additional patients were given weekly escalating doses of 25 mg/day, 50 mg/day, 75 mg/day, 100 mg/day, 125 mg/day, and 150 mg/day over a 6-week period with continuing treatment for an additional six weeks.

The study of Phase 1 study was designed to determine a maximum tolerated dose (MTD) of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione in patients with refractory solid tumors and/or lymphoma, as well as to characterize the pharmacokinetic and side effect profiles of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione in this patient population. The study design dictates that at least 3 patients must be enrolled at a dose level and have completed 28 days of treatment prior to enrollment of patients at the next higher dose level. Patients in the first cohort began dosing at 5 mg/day of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione. Patients will be escalated to 10, 20, 25, and 30 mg/day provided there is no toxicity.

In this study, the MTD is defined as the highest dose level in which fewer than two of six patients treated did not experience Grade 3 or greater non-hematological toxicity or Grade 4 or greater hematological toxicity. If, at any given dose level in either study, one out of three patients experiences toxicity, three additional patients must be treated at that particular dose. If, however, two out of six patients experience DLT, the MTD is judged to have been exceeded. No further dose escalations are to occur and additional patients are to be enrolled at the previous dose level. The dose of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione administered is escalated until the MTD is achieved or the maximum daily dose of is reached.

No DLTs were reported in the initial group of 20 patients enrolled in the study. Thirteen of the original 20 trial patients, along with 2 non-trial patients, continued on treatment as named patients at doses up to 150 mg/day.

6.5.4 Treatment of Gliomas

This study was performed to find toxicity in patients with recurrent, high-grade gliomas. The study is designed such that patients are given increasingly higher doses of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione until a maximum tolerated dose (MTD) is established. The study also seeks to obtain preliminary toxicity information and pharmacokinetic data on 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione, as well as to develop exploratory data concerning surrogate end points of angiogenic activity in vivo using functional neuro-imaging studies, and in vitro assays of serum angiogenic peptides.

Patients enrolled in the first cohort receive 2.5 mg/m²/day for a 4-week cycle. During each 4-week cycle of therapy, 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,

38

6-dione administered once daily for 3 weeks followed by a week of rest. Patients who complete a treatment cycle may receive another cycle of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione treatment if two criteria are met. First, the patient must have stable disease or have experienced a partial response or complete response, or the patient is benefiting from the therapy with 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione as evidenced by a decrease in tumor-related symptoms such as neurological deficits. Second, the patient must have recovered from toxicity related to 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione which occurred in the prior cycle by Day 42 or sooner (28-day cycle plus limit of 2 weeks to recover) as evidenced by a return to Grade ≤ 1 toxicity level. Patients who experience DLT in the previous cycle should have their dose modified. DLT is defined as a non-hematological event Grade ≥ 3 toxicity or hematological event of Grade 4 toxicity thought to be related to the study medication. Patients who experience DLT in the first cycle and have no response to therapy are removed from the study.

3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione doses are subsequently escalated to 5, 8, 11, 15, and 20 mg/m²/day to a maximum total daily dose of 40 mg. Patients continue to receive 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione on a 4-week cycle per dose level until one of the off-study criteria are met.

Three patients are enrolled in each cohort. If at least one DLT occurs, three additional patients are added to the cohort at that particular dose level. If two DLTs occur, the MTD, defined as the dose at which fewer than one-third of patients at each dose level experiences DLT has been exceeded and four more patients are treated at the previous dose.

Patients who experience DLT during the first 4-week cycle are removed from the study, except if they have a response to therapy. For patients who have completed their first 4-week cycle of without DLT, but who subsequently experience Grade 3 or 4 hematological and/or nonhematological toxicity, treatment is suspended for a minimum of a week. If the toxicity resolves to <Grade 2 within three weeks, the patient is treated at two dose levels lower than the dose that caused the toxicity (or a 50% reduction if the patient was treated at the first or second dose level). Patients in whom Grade 3 or 4 toxicity does not resolve to <Grade 1 within three weeks, or those who have another Grade 3 toxicity at the reduced dose are removed from the study.

Pharmacokinetic sampling is performed prior the first dose of 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Day 1) and 0.5, 1, 2, 4, 6, 8, 24, and 48 hours thereafter. Sampling is also conducted pre-dose on Days 7 and 21 and 0.5, 1, 2, 4, 6, 8, and 24 post-dose on Day 21 to evaluate steady-state 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione levels.

6.5.5 Treatment of Metastatic Melanoma

Patients with metastatic melanoma were started on 3-(4-amino-1-oxo-1,3-dihydro-isoindol-2-yl)-piperidine-2,6-dione (Revimid™) at 5 mg/day for seven days. The dose was then increased every seven days to 10 mg/day, 25 mg/day, and 50 mg/day, respectively, for a total of four weeks on therapy. Five of the 13 melanoma patients who were treated under this regimen either showed disease stabilization or a partial response in the first four weeks of treatment. Tumor response was seen in cutaneous and subcutaneous lesions (five patients), lymph nodes (two patients), and liver (one patient). The duration of response was approximately six months. The result suggests that the compound appears is a promising new anti-cancer agent and has both antiangiogenic and immunomodulatory properties.

US 9,101,622 B2

39


6.5.6 Treatment of Relapsed or Refractory Multiple Myeloma

Patients with relapsed and refractory Dune-Salmon stage III multiple myeloma, who have either failed at least three previous regimens or presented with poor performance status, neutropenia or thrombocytopenia, are treated with up to four cycles of combination of melphalan (50 mg intravenously), an immunomodulatory compound of the invention (about 1 to 150 mg orally daily), and dexamethasone (40 mg/day orally on days 1 to 4) every four to six weeks. Maintenance treatment consisting of daily an immunomodulatory compound of the invention and monthly dexamethasone are continued until the disease progression. The therapy using an immunomodulatory compound of the invention in combination with melphalan and dexamethasone is highly active and generally tolerated in heavily pretreated multiple myeloma patients whose prognosis is otherwise poor.

The embodiments of the invention described above are intended to be merely exemplary, and those skilled in the art will recognize, or will be able to ascertain using no more than routine experimentation, numerous equivalents of specific compounds, materials, and procedures. All such equivalents are considered to be within the scope of the invention and are encompassed by the appended claims.


What is claimed is:

1. A method of treating multiple myeloma, which comprises administering in a 28 day cycle to a patient having multiple myeloma: (a) about 1 to about 50 mg per day of a compound having the formula:



or a pharmaceutically acceptable salt, solvate or stereoisomer thereof for 21 consecutive days followed by seven consecutive days of rest from administration of said compound, and (b) a therapeutically effective amount of dexamethasone during such 28 day cycle, wherein the patient has not received previous treatment for multiple myeloma.

2. The method of claim 1, wherein the compound is



40

and is not a pharmaceutically acceptable salt, solvate or stereoisomer thereof.

3. The method of claim 1, wherein the compound is a pharmaceutically acceptable salt.

4. The method of claim 1, wherein the compound is a pharmaceutically acceptable solvate.

5. The method of claim 1, wherein the compound is a pharmaceutically acceptable stereoisomer.

6. The method of claim 1, wherein the patient is ineligible for stem cell transplantation.

7. The method of claim 1, wherein the patient is 65 years of age or older.

8. The method of claim 1, wherein the patient is younger than 65 years of age.

9. The method of claim 6, wherein the patient is younger than 65 years of age.

10. The method of claim 1, wherein the compound is administered in an amount of about 25 mg per day.

11. The method of claim 1, wherein the compound is administered in an amount of about 20 mg per day.

12. The method of claim 1, wherein the compound is administered in an amount of about 15 mg per day.

13. The method of claim 1, wherein the compound is administered in an amount of about 10 mg per day.

14. The method of claim 1, wherein the compound is administered in an amount of about 5 mg per day.

15. The method of claim 1, wherein the compound is administered in an amount of about 2.5 mg per day.

16. The method of claim 1, wherein the compound is administered orally.

17. The method of claim 16, wherein the compound is administered in the form of a capsule or tablet.

18. The method of claim 17, wherein the compound is administered in a capsule in an amount from about 1 mg to about 50 mg.

19. The method of claim 18, wherein the compound is administered in a capsule in an amount of about 2.5 mg, about 5 mg, about 10 mg, about 15 mg, about 20 mg or about 25 mg.

20. The method of claim 17, wherein the capsule comprises the compound, lactose anhydrous, microcrystalline cellulose, croscarmellose sodium and magnesium stearate.

21. The method of claim 1, wherein the dexamethasone is administered in an amount of 40 mg per day on days 1-4 of each 28 day cycle.

22. The method of claim 1, wherein the dexamethasone is administered in an amount of 40 mg per day on days 1, 8, 15 and 22 of each 28 day cycle.

23. The method of claim 1, which further comprises administering a therapeutically effective amount of an additional active agent.

24. The method of claim 23, wherein the additional active agent is melphalan, doxorubicin, vincristine, prednisone, cyclophosphamide, biacin, a proteasome inhibitor, or a combination thereof.

25. The method of claim 1, wherein the solvate is hydrate.

* * * * *