



US 20130040906A1

(19) **United States**(12) **Patent Application Publication**
Kuntz et al.(10) **Pub. No.: US 2013/0040906 A1**(43) **Pub. Date: Feb. 14, 2013**(54) **INHIBITORS OF HUMAN EZH2, AND
METHODS OF USE THEREOF***G01N 33/53* (2006.01)*C12Q 1/68* (2006.01)*A61P 35/00* (2006.01)*A61P 35/02* (2006.01)(76) Inventors: **Kevin W. Kuntz**, Woburn, MA (US);
Sarah K. Knutson, Cambridge, MA
(US); **Timothy James Nelson Wigle**,
Waltham, MA (US)(52) **U.S. Cl. 514/46; 436/501; 435/7.92; 435/6.12;
436/94**(21) Appl. No.: **13/418,242**(57) **ABSTRACT**(22) Filed: **Mar. 12, 2012****Related U.S. Application Data**(63) Continuation-in-part of application No. 13/230,703,
filed on Sep. 12, 2011.(60) Provisional application No. 61/381,684, filed on Sep.
10, 2010.**Publication Classification**(51) **Int. Cl.***A61K 31/7076* (2006.01)*G01N 27/26* (2006.01)

The invention relates to inhibition of wild-type and certain mutant forms of human histone methyltransferase EZH2, the catalytic subunit of the PRC2 complex which catalyzes the mono- through tri-methylation of lysine 27 on histone H3 (H3-K27). In one embodiment the inhibition is selective for the mutant form of the EZH2, such that trimethylation of H3-K27, which is associated with certain cancers, is inhibited. The methods can be used to treat cancers including follicular lymphoma and diffuse large B-cell lymphoma (DLBCL). Also provided are methods for identifying small molecule selective inhibitors of the mutant forms of EZH2 and also methods for determining responsiveness to an EZH2 inhibitor in a subject.

FIG. 1




FIG. 2




FIG. 3




FIG. 4





a *EZH2* Mutation**c** *KDM6A/UTX* mutation**b** *EZH2* overexpression**d** *PHF19/PCL3* overexpression

FIG. 5

LEGEND

- 1 = marker
- 2 = EZH2: WT
- 3 = EZH2: Y641F
- 4 = EZH2: Y641H
- 5 = EZH2: Y641N
- 6 = EZH2: Y641S




FIG. 6




FIG. 7




FIG. 8

Compound 75 potency against EZH2 WT and Y641 mutants




FIG. 9




FIG. 10




FIG. 11




FIG. 12





FIG. 13

Inhibitor A



Inhibitor B




FIG. 14





FIG. 15



INHIBITORS OF HUMAN EZH2, AND METHODS OF USE THEREOF

RELATED REFERENCE

[0001] This application is a continuation-in-part of the U.S. patent application Ser. No. 13/230,703, filed Sep. 12, 2011, which claims priority to, and the benefit of, U.S. Ser. No. 61/381,684, filed Sep. 10, 2010, each of which is incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

[0002] This invention relates to inhibition of wild-type and certain mutant forms of human histone methyltransferase EZH2, the catalytic subunit of the PRC2 complex which catalyzes the mono- through tri-methylation of lysine 27 on histone H3 (H3-K27), methods for treating cancers including follicular lymphoma and diffuse large B-cell lymphoma (DLBCL) and methods for determining responsiveness to an EZH2 inhibitor in a subject.

BACKGROUND

[0003] In eukaryotic cells DNA is packaged with histones to form chromatin. Approximately 150 base pairs of DNA are wrapped twice around an octamer of histones (two each of histones 2A, 2B, 3 and 4) to form a nucleosome, the basic unit of chromatin. Changes in the ordered structure of chromatin can lead to alterations in transcription of associated genes. This process is highly controlled because changes in gene expression patterns can profoundly affect fundamental cellular processes, such as differentiation, proliferation and apoptosis. Control of changes in chromatin structure (and hence of transcription) is mediated by covalent modifications to histones, most notably of their N-terminal tails. These modifications are often referred to as epigenetic because they can lead to heritable changes in gene expression, but they do not affect the sequence of the DNA itself. Covalent modifications (for example, methylation, acetylation, phosphorylation and ubiquitination) of the side chains of amino acids are enzymatically mediated.

[0004] The selective addition of methyl groups to specific amino acid sites on histones is controlled by the action of a unique family of enzymes known as histone methyltransferases (HMTs). The level of expression of a particular gene is influenced by the presence or absence of one or more methyl groups at a relevant histone site. The specific effect of a methyl group at a particular histone site persists until the methyl group is removed by a histone demethylase, or until the modified histone is replaced through nucleosome turnover. In a like manner, other enzyme classes can decorate DNA and histones with other chemical species, and still other enzymes can remove these species to provide control of gene expression.

[0005] The orchestrated collection of biochemical systems behind transcriptional regulation must be tightly controlled in order for cell growth and differentiation to proceed optimally. Disease states result when these controls are disrupted by aberrant expression and/or activity of the enzymes responsible for DNA and histone modification. In human cancers, for example, there is a growing body of evidence to suggest that dysregulated epigenetic enzyme activity contributes to the uncontrolled cell proliferation associated with cancer as well as other cancer-relevant phenotypes such as enhanced cell migration and invasion. Beyond cancer, there is growing

evidence for a role of epigenetic enzymes in a number of other human diseases, including metabolic diseases (such as diabetes), inflammatory diseases (such as Crohn's disease), neurodegenerative diseases (such as Alzheimer's disease), and cardiovascular diseases. Therefore, selectively modulating the aberrant action of epigenetic enzymes holds great promise for the treatment of a range of diseases.

Histone Methyltransferase EZH2

[0006] Polycomb group (PcG) and trithorax group (trxG) proteins are known to be part of the cellular memory system. Francis et al. (2001) *Nat Rev Mol Cell Biol* 2:409-21; Simon et al. (2002) *Curr Opin Genet Dev* 12:210-8. Both groups of proteins are involved in maintaining the spatial patterns of homeotic box (Hox) gene expression, which are established early in embryonic development by transiently expressed segmentation genes. In general, PcG proteins are transcriptional repressors that maintain the "off state," and trxG proteins are transcriptional activators that maintain the "on state." Because members of PcG and trxG proteins contain intrinsic histone methyltransferase (HMTase) activity, PcG and trxG proteins may participate in cellular memory through methylation of core histones. Beisel et al. (2002) *Nature* 419:857-62; Cao et al. (2002) *Science* 298:1039-43; Czermin et al. (2002) *Cell* 111:185-96; Kuzmichev et al. (2002) *Genes Dev* 16:2893-905; Milne et al. (2002) *Mol Cell* 10:1107-17; Muller et al. (2002) *Cell* 111:197-208; Nakamura et al. (2002) *Mol Cell* 10:1119-28.

[0007] Biochemical and genetic studies have provided evidence that *Drosophila* PcG proteins function in at least two distinct protein complexes, the Polycomb repressive complex 1 (PRC1) and the ESC-E(Z) complex (also known as Polycomb repressive complex 2 (PRC2)), although the compositions of the complexes may be dynamic. Otte et al. (2003) *Curr Opin Genet Dev* 13:448-54. Studies in *Drosophila* (Czermin et al. (supra); Muller et al. (supra)) and mammalian cells (Cao et al. (supra); Kuzmichev et al. (supra)) have demonstrated that the ESC-E(Z)/EED-EZH2 (i.e., PRC2) complexes have intrinsic histone methyltransferase activity. Although the compositions of the complexes isolated by different groups are slightly different, they generally contain EED, EZH2, SUZ12, and RbAp48 or *Drosophila* homologs thereof. However, a reconstituted complex comprising only EED, EZH2, and SUZ12 retains histone methyltransferase activity for lysine 27 of histone H3. U.S. Pat. No. 7,563,589 (incorporated by reference).

[0008] Of the various proteins making up PRC2 complexes, EZH2 (Enhancer of Zeste Homolog 2) is the catalytic subunit. The catalytic site of EZH2 in turn is present within a SET domain, a highly conserved sequence motif (named after Su(var)3-9, Enhancer of Zeste, Trithorax) that is found in several chromatin-associated proteins, including members of both the Trithorax group and Polycomb group. SET domain is characteristic of all known histone lysine methyltransferases except the H3-K79 methyltransferase DOT1.

[0009] In addition to Hox gene silencing, PRC2-mediated histone H3-K27 methylation has been shown to participate in X-inactivation. Plath et al. (2003) *Science* 300:131-5; Silva et al. (2003) *Dev Cell* 4:481-95. Recruitment of the PRC2 complex to Xi and subsequent trimethylation on histone H3-K27 occurs during the initiation stage of X-inactivation and is dependent on Xist RNA. Furthermore, EZH2 and its associated histone H3-K27 methyltransferase activity was found to

mark differentially the pluripotent epiblast cells and the differentiated trophectoderm. Erhardt et al. (2003) *Development* 130:4235-48).

[0010] Consistent with a role of EZH2 in maintaining the epigenetic modification patterns of pluripotent epiblast cells, Cre-mediated deletion of EZH2 results in loss of histone H3-K27 methylation in the cells. Erhardt et al. (supra). Further, studies in prostate and breast cancer cell lines and tissues have revealed a strong correlation between the levels of EZH2 and SUZ12 and the invasiveness of these cancers (Bracken et al. (2003) *EMBO J* 22:5323-35; Kirmizis et al. (2003) *Mol Cancer Ther* 2:113-21; Kleer et al. (2003) *Proc Natl Acad Sci USA* 100:11606-11; Varambally et al. (2002) *Nature* 419: 624-9), indicating that dysfunction of the PRC2 complex may contribute to cancer.

[0011] Recently, somatic mutations of EZH2 were reported to be associated with follicular lymphoma (FL) and the germinal center B cell-like (GCB) subtype of diffuse large B-cell lymphoma (DLBCL). Morin et al. (2010) *Nat Genet* 42:181-5. In all cases, occurrence of the mutant EZH2 gene was found to be heterozygous, and expression of both wild-type and mutant alleles was detected in the mutant samples profiled by transcriptome sequencing. Currently, the standard of care for the treatment of most cases of DLBCL is the R—CHOP regimen. However, the outcome of this regimen is far from satisfactory. Therefore, there is a great medical need to identify novel and effective therapies, optionally based on the genetic profiles of the subject.

SUMMARY OF THE INVENTION

[0012] The invention is based upon the discovery that cells expressing certain EZH2 mutants are more responsive to EZH2 inhibitors than cells expressing wild type EZH2.

[0013] The invention features a method for treating or alleviating a symptom of cancer or precancerous condition in a subject administering to a subject expressing a mutant EZH2 comprising a mutation in the substrate pocket domain as defined in SEQ ID NO: 6 a therapeutically effective amount of an EZH2 inhibitor.

[0014] The invention also features a method of determining a responsiveness of a subject having a cancer or a precancerous condition to an EZH2 inhibitor by providing a sample from the subject; and detecting a mutation in the EZH2 substrate pocket domain as defined in SEQ ID NO: 6; and the presence of said mutation indicates the subject is responsive to the EZH2 inhibitor.

[0015] The mutant EZH2 of the present invention is a mutant EZH2 polypeptide or a nucleic acid sequence encoding a mutant EZH2 polypeptide. Preferably, the mutant EZH2 comprises a mutation at amino acid position 677, 687, 674, 685, or 641 of SEQ ID NO: 1. More preferably, mutation is selected from the group consisting of a substitution of glycine (G) for the wild type residue alanine (A) at amino acid position 677 of SEQ ID NO: 1 (A677G); a substitution of valine (V) for the wild type residue alanine (A) at amino acid position 687 of SEQ ID NO: 1 (A687V); a substitution of methionine (M) for the wild type residue valine (V) at amino acid position 674 of SEQ ID NO: 1 (V674M); a substitution of histidine (H) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685H); a substitution of cysteine (C) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685C); a substitution of phenylalanine (F) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641F); a sub-

stitution of histidine (H) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641H); a substitution of asparagine (N) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641N); a substitution of serine (S) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641S); and a substitution of cysteine (C) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641C).

[0016] The subject of the present invention includes any human subject who has been diagnosed with, has symptoms of, or is at risk of developing a cancer or a precancerous condition. For example, the cancer is lymphoma, leukemia or melanoma. Preferably, the lymphoma is non-Hodgkin lymphoma, follicular lymphoma or diffuse large B-cell lymphoma. Alternatively, the leukemia is chronic myelogenous leukemia (CML). The precancerous condition is myelodysplastic syndromes (MDS, formerly known as preleukemia).

[0017] Preferred EZH2 inhibitor for the methods of the present invention is selected from compounds listed in Table 1.

[0018] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. In the specification, the singular forms also include the plural unless the context clearly dictates otherwise. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. All publications, patent applications, patents and other references mentioned herein are incorporated by reference. The references cited herein are not admitted to be prior art to the claimed invention. In the case of conflict, the present specification, including definitions, will control. In addition, the materials, methods and examples are illustrative only and are not intended to be limiting.

[0019] Other features and advantages of the invention will be apparent from the following detailed description and claims.

BRIEF DESCRIPTION OF THE FIGURES

[0020] FIG. 1 is two graphs establishing that B-cell lymphoma-associated mutants of EZH2 are active histone methyltransferases. In vitro methyltransferase activity of PRC2 complexes containing wild-type and various Y641 mutants of EZH2 was measured as (A) methyl transfer reactions using a peptide (H3 21-44) as substrate, and (B) methyl transfer reactions using avian nucleosomes as substrate. Symbols: wild-type (●), Y641F (○), Y641H (□), Y641N (■), and Y641S (▲). CPM is counts per minute, referring to scintillation counting as a result of ³H radiation.

[0021] FIG. 2 is four graphs establishing that PRC2 complexes containing mutant EZH2 preferentially catalyze di- and tri-methylation of histone H3-K27. (A) Methyltransferase activity of mutant and wild-type (WT) complexes on unmethylated peptide (open bars), monomethylated peptide (hashed bars), and dimethylated peptide (closed bars). (B) Affinity for peptide substrates as judged by $K_{1/2}$ is similar across all peptide methylation states for PRC2 complexes containing wild-type (○), Y641F (●), Y641H (□), Y641N (■), and Y641S (▲) EZH2. Note that the variation in $K_{1/2}$ values across all substrates and all enzyme forms is less than 3.5-fold. For any particular methylation state of substrate the variation in $K_{1/2}$ value is less than 2-fold. (C) Enzyme turn-

over number (k_{cat}) varies with substrate methylation status in opposing ways for WT and Y641 mutants of EZH2. The k_{cat} decreases with increasing K27 methylation states for wild-type (\circ), but increases for Y641F (\bullet), Y641H (\square), Y641N (\blacksquare), and Y641S (\blacktriangle) mutants of EZH2. (D) Catalytic efficiency ($k_{cat}/K_{1/2}$) decreases with increasing K27 methylation states for wild-type (\circ), but increases for Y641F (\bullet), Y641H (\square), Y641N (\blacksquare), and Y641S (\blacktriangle) mutants of EZH2. In panels B-D, the lines drawn to connect the data points are not intended to imply any mathematical relationship; rather, they are merely intended to serve as visual aides.

[0022] FIG. 3A is a trio of graphs depicting predicted relative levels of H3-K27me3 (top panel), H3-K27me2 (middle panel), and H3-K27me1 (bottom panel) for cells containing different EZH2 mutants. Simulations were performed using a coupled enzyme steady state velocity equation and the steady state kinetic parameters shown in Table 2. All values are relative to the homozygous WT EZH2-containing cells and assume saturating concentrations of intracellular SAM, relative to K_m and intracellular nucleosome concentrations similar to K_m .

[0023] FIG. 3B is a series of Western blot analyses of relative patterns of H3-K27 methylation status for lymphoma cell lines homozygous for WT EZH2, or heterozygous for the indicated EZH2 Y641 mutation. Panels from top to bottom depict the results of probing with antibodies specific for the following: total EZH2; H3-K27me3; H3-K27me2; H3-K27me1; and total histone H3 as loading control.

[0024] FIG. 4 depicts selected proposed mechanisms leading to aberrantly high levels of trimethylation on histone H3-K27 in cancer. These include: a) mutation of Y641 in EZH2 resulting in a change in substrate preference from the nonmethylated to the mono- and di-methylated histone H3-K27; b) overexpression of EZH2; c) mutations in UTX that inactivate enzyme function, causing a decrease in demethylation of H3-K27me3; and d) overexpression of the PRC2 complex subunit PHF19/PCL3 that leads to increases in recruitment of the PRC2 complex to specific genes and an increase in histone H3-K27 trimethylation. In all four models the alteration leads to aberrant histone H3-K27 trimethylation in the proximal promoter regions of genes resulting in transcriptional repression of key genes in cancer.

[0025] FIG. 5 depicts a SDS-PAGE gel showing that the expression levels of each of the five-component PRC2 complexes are similar with mutant and wild-type EZH2.

[0026] FIG. 6 is a pair of tables showing that mutant and wild-type (WT) PRC2 complexes display strong substrate preference for H3-K27-containing peptides. Each enzyme was tested against a panel of overlapping 15-mer peptides covering all of H3 and H4. Activity was measured as velocity (CPM per minute), and the reported value represents the mean of two independent determinations for each reaction. For all the complexes the most favored peptide was H3:16-30. WT complex had greater than 6-fold more activity against this peptide than any of the mutant complexes.

[0027] FIG. 7 is a graph depicting inhibitory potency of S-adenosyl-L-homocysteine (SAH) against EZH2 WT and Y641 mutants of EZH2. The X axis shows log concentration of SAH; the Y axis shows percent inhibition.

[0028] FIG. 8 is a graph depicting inhibitory potency of Compound 75 against EZH2 WT and Y641 mutants of EZH2. The X axis shows log concentration of Compound 75; the Y axis shows percent inhibition.

[0029] FIG. 9 depicts a Western Blot analysis of relative levels of H3-K27me1, me2 and me3 in a cell line pane, including multiple DLBCL lines expressing WT or Y641 mutant EZH2. Histones were extracted from the cell lines shown, fractionated by SDS-PAGE on a 4-20% gel, transferred to nitrocellulose membranes, and probed with antibodies to Histone H3, H3-K27me1, me2, or me3.

[0030] FIG. 10 depicts an immunocytochemistry analysis of H3 and H3-K27me3 levels in a panel of WT and Y641 mutant lymphoma cell lines. Cell pellets from the indicated cell lines were fixed and embedded in paraffin. Slides were prepared and levels of H3 and H3-K27me3 were evaluated by immunocytochemistry using antibodies to histone H3, or H3-K27me3.

[0031] FIG. 11 depicts an immunocytochemistry analysis of H3 and H3-K27me2 levels in a panel of WT and Y641 mutant lymphoma cell lines. Cell pellets from the indicated cell lines were fixed and embedded in paraffin. Slides were prepared and levels of H3 and H3-K27me2 were evaluated by immunocytochemistry using antibodies to histone H3, or H3-K27me2.

[0032] FIG. 12 is a graph depicting the inhibition of global H3-K27me3 levels by EZH2 inhibitor treatment in Y641 mutant WSU-DLCL2 cells. WSU-DLCL2 cells were treated for 4 days with the indicated concentrations of EZH2 inhibitor A or B. Following compound treatment, histones were extracted, fractionated by SDS-PAGE on a 4-20% gel, transferred to nitrocellulose membranes, and probed with antibodies to Histone H3, or H3-K27me3.

[0033] FIG. 13 is a graph showing that the EZH2 inhibitors can block proliferation of a Y641 mutant WSU-DLCL2 cells, but has little effect on non Y641 mutant OCI-LY19 cells. Cells were incubated in the presence of increasing concentrations of EZH2 inhibitor A or B for eleven days. Vehicle treated (DMSO) cells were included as controls. Cell number and viability was determined using the Guava Viacount assay in a Guava EasyCyte Plus instrument. Cells were split and media and compound was replenished every 3-4 days.

[0034] FIG. 14 is a graph showing the presence of an EZH2 (Y641) mutation and/or high H3-K27me3 and low H3-K27me2 levels predict sensitivity to EZH2 inhibitors. Cell lines were maintained in the presence of increasing concentrations of one EZH2 inhibitor up to 25 μ M. Viable cells counts were used to derive IC_{90} values after 11 days of treatment. Results are plotted with cell lines segregated according to EZH2 mutational status (A), or segregated according to H3-K27me2 and H3-K27me3 levels (B). In both plots, the line shows the average IC_{90} values from the indicated cell line group.

[0035] FIG. 15 is a panel of graphs showing velocity vs. enzyme concentration measured for wild-type, A677G or A687V EZH2. Biotinylated peptides representing histone H3 residues 21-44 containing un-, mono-, di- or trimethyl lysine 27 (H3K27me0 through H3K27me3) were assayed at a fixed concentration with a dilution series of the indicated EZH2 enzyme. Timepoints were sampled over the course of 90 minutes and 3H-SAM incorporation at lysine 27 of the H3 peptide was measured by capturing the peptide in a Flashplate and reading the counts per minute (CPM). Linear regression of the timecourse yielded enzyme velocity in CPM per minute (CPM/min) which was plotted as a function of enzyme concentration.

DETAILED DESCRIPTION

[0036] Chromatin structure is important in gene regulation and epigenetic inheritance. Post-translational modifications of histones, such as methylation are involved in the establishment and maintenance of higher-order chromatin structure.

EZH2 Mutants

[0037] EZH2 is a histone methyltransferase that is the catalytic subunit of the PRC2 complex which catalyzes the mono-through tri-methylation of lysine 27 on histone H3 (H3-K27).

[0038] Point mutations of the EZH2 gene at a single amino acid residue (e.g., Tyr641, herein referred to as Y641) of EZH2 have been reported to be linked to subsets of human B-cell lymphoma. Morin et al. (2010) *Nat Genet* 42(2):181-5. In particular, Morin et al. reported that somatic mutations of tyrosine 641 (Y641F, Y641H, Y641N, and Y641S) of EZH2 were associated with follicular lymphoma (FL) and the germinal center B cell-like (GCB) subtype of diffuse large B-cell lymphoma (DLBCL). The mutant allele is always found associated with a wild-type allele (heterozygous) in disease cells, and the mutations were reported to ablate the enzymatic activity of the PRC2 complex for methylating an unmodified peptide substrate.

[0039] The present invention is based in part upon the surprising discovery that cells expressing an EZH2 mutant are more sensitive to EZH2 inhibitors of the instant invention than cells expressing wild type EZH2. Accordingly, an aspect of the present invention relates to methods for treating or alleviating a symptom of cancer or precancerous condition in a subject by administering to a subject expressing a mutant EZH2 a therapeutically effective amount of an EZH2 inhibitor. The mutant EZH2 of the present invention refers to a mutant EZH2 polypeptide or a nucleic acid sequence encoding a mutant EZH2 polypeptide. Preferably the mutant EZH2 comprises one or more mutations in its substrate pocket domain as defined in SEQ ID NO: 6. For example, the mutation may be a substitution, a point mutation, a nonsense mutation, a miss sense mutation, a deletion, or an insertion. Exemplary substitution amino acid mutation includes a substitution at amino acid position 677, 687, 674, 685, or 641 of SEQ ID NO: 1, such as, but is not limited to a substitution of glycine (G) for the wild type residue alanine (A) at amino acid position 677 of SEQ ID NO: 1 (A677G); a substitution of valine (V) for the wild type residue alanine (A) at amino acid position 687 of SEQ ID NO: 1 (A687V); a substitution of methionine (M) for the wild type residue valine (V) at amino acid position 674 of SEQ ID NO: 1 (V674M); a substitution of histidine (H) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685H); a substitution of cysteine (C) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685C); a substitution of phenylalanine (F) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641F); a substitution of histidine (H) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641H); a substitution of asparagine (N) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641N); a substitution of serine (S) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641S); or a substitution of cysteine (C) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641C).

[0040] The mutation of the present invention may also include a substitution of serine (S) for the wild type residue asparagine (N) at amino acid position 322 of SEQ ID NO: 3 (N322S), a substitution of glutamine (Q) for the wild type residue arginine (R) at amino acid position 288 of SEQ ID NO: 3 (R288Q), a substitution of isoleucine (I) for the wild type residue threonine (T) at amino acid position 573 of SEQ ID NO: 3 (T573I), a substitution of glutamic acid (E) for the wild type residue aspartic acid (D) at amino acid position 664 of SEQ ID NO: 3 (D664E), a substitution of glutamine (Q) for the wild type residue arginine (R) at amino acid position 458 of SEQ ID NO: 5 (R458Q), a substitution of lysine (K) for the wild type residue glutamic acid (E) at amino acid position 249 of SEQ ID NO: 3 (E249K), a substitution of cysteine (C) for the wild type residue arginine (R) at amino acid position 684 of SEQ ID NO: 3 (R684C), a substitution of histidine (H) for the wild type residue arginine (R) at amino acid position 628 of SEQ ID NO: 21 (R628H), a substitution of histidine (H) for the wild type residue glutamine (Q) at amino acid position 501 of SEQ ID NO: 5 (Q501H), a substitution of asparagine (N) for the wild type residue aspartic acid (D) at amino acid position 192 of SEQ ID NO: 3 (D192N), a substitution of valine (V) for the wild type residue aspartic acid (D) at amino acid position 664 of SEQ ID NO: 3 (D664V), a substitution of leucine (L) for the wild type residue valine (V) at amino acid position 704 of SEQ ID NO: 3 (V704L), a substitution of serine (S) for the wild type residue proline (P) at amino acid position 132 of SEQ ID NO: 3 (P132S), a substitution of lysine (K) for the wild type residue glutamic acid (E) at amino acid position 669 of SEQ ID NO: 21 (E669K), a substitution of threonine (T) for the wild type residue alanine (A) at amino acid position 255 of SEQ ID NO: 3 (A255T), a substitution of valine (V) for the wild type residue glutamic acid (E) at amino acid position 726 of SEQ ID NO: 3 (E726V), a substitution of tyrosine (Y) for the wild type residue cysteine (C) at amino acid position 571 of SEQ ID NO: 3 (C571Y), a substitution of cysteine (C) for the wild type residue phenylalanine (F) at amino acid position 145 of SEQ ID NO: 3 (F145C), a substitution of threonine (T) for the wild type residue asparagine (N) at amino acid position 693 of SEQ ID NO: 3 (N693T), a substitution of serine (S) for the wild type residue phenylalanine (F) at amino acid position 145 of SEQ ID NO: 3 (F145S), a substitution of histidine (H) for the wild type residue glutamine (Q) at amino acid position 109 of SEQ ID NO: 21 (Q109H), a substitution of cysteine (C) for the wild type residue phenylalanine (F) at amino acid position 622 of SEQ ID NO: 21 (F622C), a substitution of arginine (R) for the wild type residue glycine (G) at amino acid position 135 of SEQ ID NO: 3 (G135R), a substitution of glutamine (Q) for the wild type residue arginine (R) at amino acid position 168 of SEQ ID NO: 5 (R168Q), a substitution of arginine (R) for the wild type residue glycine (G) at amino acid position 159 of SEQ ID NO: 3 (G159R), a substitution of cysteine (C) for the wild type residue arginine (R) at amino acid position 310 of SEQ ID NO: 5 (R310C), a substitution of histidine (H) for the wild type residue arginine (R) at amino acid position 561 of SEQ ID NO: 3 (R561H), a substitution of histidine (H) for the wild type residue arginine (R) at amino acid position 634 of SEQ ID NO: 21 (R634H), a substitution of arginine (R) for the wild type residue glycine (G) at amino acid position 660 of SEQ ID NO: 3 (G660R), a substitution of cysteine (C) for the wild type residue tyrosine (Y) at amino acid position 181 of SEQ ID NO: 3 (Y181C), a substitution of arginine (R) for the wild type residue histidine (H) at amino acid position 297 of SEQ

ID NO: 3 (H297R), a substitution of serine (S) for the wild type residue cysteine (C) at amino acid position 612 of SEQ ID NO: 21 (C612S), a substitution of tyrosine (Y) for the wild type residue histidine (H) at amino acid position 694 of SEQ ID NO: 3 (H694Y), a substitution of alanine (A) for the wild type residue aspartic acid (D) at amino acid position 664 of SEQ ID NO: 3 (D664A), a substitution of threonine (T) for the wild type residue isoleucine (I) at amino acid position 150 of SEQ ID NO: 3 (I150T), a substitution of arginine (R) for the wild type residue isoleucine (I) at amino acid position 264 of SEQ ID NO: 3 (I264R), a substitution of leucine (L) for the wild type residue proline (P) at amino acid position 636 of SEQ ID NO: 3 (P636L), a substitution of threonine (T) for the wild type residue isoleucine (I) at amino acid position 713 of SEQ ID NO: 3 (I713T), a substitution of proline (P) for the wild type residue glutamine (Q) at amino acid position 501 of SEQ ID NO: 5 (Q501P), a substitution of glutamine (Q) for the wild type residue lysine (K) at amino acid position 243 of SEQ ID NO: 3 (K243Q), a substitution of aspartic acid (D) for the wild type residue glutamic acid (E) at amino acid position 130 of SEQ ID NO: 5 (E130D), a substitution of glycine (G) for the wild type residue arginine (R) at amino acid position 509 of SEQ ID NO: 3 (R509G), a substitution of histidine (H) for the wild type residue arginine (R) at amino acid position 566 of SEQ ID NO: 3 (R566H), a substitution of histidine (H) for the wild type residue aspartic acid (D) at amino acid position 677 of SEQ ID NO: 3 (D677H), a substitution of asparagine (N) for the wild type residue lysine (K) at amino acid position 466 of SEQ ID NO: 5 (K466N), a substitution of histidine (H) for the wild type residue arginine (R) at amino acid position 78 of SEQ ID NO: 3 (R78H), a substitution of methionine (M) for the wild type residue lysine (K) at amino acid position 1 of SEQ ID NO: 6 (K6M), a substitution of leucine (L) for the wild type residue serine (S) at amino acid position 538 of SEQ ID NO: 3 (S538L), a substitution of glutamine (Q) for the wild type residue leucine (L) at amino acid position 149 of SEQ ID NO: 3 (L149Q), a substitution of valine (V) for the wild type residue leucine (L) at amino acid position 252 of SEQ ID NO: 3 (L252V), a substitution of valine (V) for the wild type residue leucine (L) at amino acid position 674 of SEQ ID NO: 3 (L674V), a substitution of valine (V) for the wild type residue alanine (A) at amino acid position 656 of SEQ ID NO: 3 (A656V), a substitution of aspartic acid (D) for the wild type residue alanine (A) at amino acid position 731 of SEQ ID NO: 3 (Y731D), a substitution of threonine (T) for the wild type residue alanine (A) at amino acid position 345 of SEQ ID NO: 3 (A345T), a substitution of aspartic acid (D) for the wild type residue alanine (A) at amino acid position 244 of SEQ ID NO: 3 (Y244D), a substitution of tryptophan (W) for the wild type residue cysteine (C) at amino acid position 576 of SEQ ID NO: 3 (C576W), a substitution of lysine (K) for the wild type residue asparagine (N) at amino acid position 640 of SEQ ID NO: 3 (N640K), a substitution of lysine (K) for the wild type

residue asparagine (N) at amino acid position 675 of SEQ ID NO: 3 (N675K), a substitution of tyrosine (Y) for the wild type residue aspartic acid (D) at amino acid position 579 of SEQ ID NO: 21 (D579Y), a substitution of isoleucine (I) for the wild type residue asparagine (N) at amino acid position 693 of SEQ ID NO: 3 (N693I), and a substitution of lysine (K) for the wild type residue asparagine (N) at amino acid position 693 of SEQ ID NO: 3 (N693K).

[0041] The mutation of the present invention may be a frameshift at amino acid position 730, 391, 461, 441, 235, 254, 564, 662, 715, 405, 685, 64, 73, 656, 718, 374, 592, 505, 730, or 363 of SEQ ID NO: 3, 5 or 21 or the corresponding nucleotide position of the nucleic acid sequence encoding SEQ ID NO: 3, 5, or 21. The mutation of the EZH2 may also be an insertion of a glutamic acid (E) between amino acid positions 148 and 149 of SEQ ID NO: 3, 5 or 21. Another example of EZH2 mutation is a deletion of glutamic acid (E) and leucine (L) at amino acid positions 148 and 149 of SEQ ID NO: 3, 5 or 21. The mutant EZH2 may further comprise a nonsense mutation at amino acid position 733, 25, 317, 62, 553, 328, 58, 207, 123, 63, 137, or 60 of SEQ ID NO: 3, 5 or 21.

[0042] It has also been unexpectedly discovered that the wild-type (WT) EZH2 enzyme displays greatest catalytic efficiency (kcat/K) for the zero- to mono-methylation reaction of H3-K27 and lesser efficiency for subsequent (mono- to di- and di- to tri-methylation) reactions; whereas, in stark contrast, the disease-associated Y641 mutations display very limited ability to perform the first methylation reaction but have enhanced catalytic efficiency for the subsequent reactions relative to wild-type enzyme. These results imply that the malignant phenotype of disease exploits the combined activities of a H3-K27 mono-methylating enzyme (PRC2 containing WT EZH2 or EZH1) together with PRC2 containing mutant EZH2 for augmented conversion of H3-K27 to the tri-methylated form (H3-K27me3). Therefore, an aspect of the present invention relates to inhibiting the activity of EZH2, including certain mutant forms of EZH2. In one embodiment the present invention relates to inhibiting selectively the activity of certain mutant forms of EZH2.

[0043] While not intending to be bound by any one theory, it is hypothesized that the mutation of EZH2 in its substrate pocket domain may facilitate multiple rounds of H3-K27 methylation by impacting the H-bonding pattern and/or steric crowding in the active site of the enzyme-bisubstrate ternary complex, affecting the formation of a proper water channel for deprotonation of the reacting lysine.

[0044] Human EZH2 nucleic acids and polypeptides have previously been described. See, e.g., Chen et al. (1996) *Genomics* 38:30-7 [746 amino acids]; Swiss-Prot Accession No. Q15910 [746 amino acids]; GenBank Accession Nos. NM_004456 and NP_004447 (isoform a [751 amino acids]); and GenBank Accession Nos. NM_152998 and NP_694543 (isoform b [707 amino acids]), each of which is incorporated herein by reference in its entirety.

Amino acid sequence of human EZH2 (Swiss-Prot Accession No. Q15910)
(SEQ ID NO: 1)
MGQTGKKSEKGPVCRKRVKSEYMRLRQLKRFRADEVKSMFSSNRQKILERTEILNQEW
KQRRIQPVHILTSVSSLRGTRECSVTSDDLDFPTQVPLKTLNAVASVPIMYSWSPLQQNF
MVEDETVLHNIPIYMGDEVLDQDGTFFIEELIKNYDGKVHGDRECGFINDEIFVELVNALGQ
YNDDDDDDDGDDPEEREKQKDLDRDDKESRPPKFPSPDKIFEAISSMFPDKGTAEEL

- continued

KEKYKELTEQQLPGALPPECTPNIDGPNKSVQREQLHSFHTLFCRRCFKYDCFLHPFH
ATPNTYKRKNTETALDNKPCGPQCYQHLEGAKEFAAALTAERIKTPPKRPGRRRGRRLPN
NSSRPSPTPTINVLESKDTSDREAGTETGGENNDKEEEKKDETSSSSEANSRCQTPIKM
KPNIEPPENVEWGSAGEASMFVRLIGTYDNFCAIARLIGTKTCRQVYEFVKESSIIAPA
PAEDVDTPPRKKRKHRLWAAHCRKIQLKKDGSSNHVYNYQPCDHPRQPCDSSCPCVIAQ
NFCEKFCQCSSECQNRFPGCRCKAQCNKQPCCYLAVRECDPDLCLTCGAADHWDKKNVS
CKNCISIQRSKKHLLLLAPSDVAGWGIFIKDPVQKNEFISEYCGEIIISQDEADRRGKVYDK
YMCSFLFNLNDFVVDATRKGNKIRFANHSVNPNCYAKVMMVNGDHRIGIFAKRAIQTGE
ELFFDYRYSQADALKYVGIEREMEIP

mRNA sequence of human EZH2, transcript variant 1 (GenBank
Accession No. NM_004456)

(SEQ ID NO: 2)

ggcgcgcttgattgggctgggggggccaataaaaagcgatggcgattgggctgccgct
ttggcgctcggtccggtcgctccgacaccgggtgggactcagaaggcagtgaggcccg
gcggcgcgcgcgcgcgcgcgggggcgacgcgcgggaacaacgcgagtcggcgcgcggg
acgaagaataatcatgggccagactgggaagaaatctgagaaggaccagtttgttgcg
gaagcgtgtaaaatcagagtacatgcgactgagacagctcaagaggttcagacgagctga
tgaagtaaaagagtatgttttagttccaatcgtcagaaaattttggaagaacggaatctt
aaaccaagaatggaaacagcgaaggatacacgcctgtgcacatcctgacttctgtgagctc
attgcgcgggactaggagtggtcggtgaccagtgacttgattttccaacacaagtcatt
ccattaaagactctgaatgcagttgcttcagtaaccataatgtattcttggtctccct
acagcagaattttatggtggaagatgaaactgtttacataacattccttatatgggaga
tgaagtttttagatcaggatggtagctttcattgaagaactaataaaaaattatgatgggaa
agtacacgggtagagaaatgtgggtttataaatgatgaaatttttgtggagttggtgaa
tgcccttggtcaatataatgatgatgacgatgatgatggagacgatcctgaagaaag
agaagaaaagcagaaagatctggaggatcacccagatgataaagaaagcccccacctcg
gaaatttccttctgataaaatttttgaagccatttcctcaatgttttcagataagggcac
agcagaagaactaaaggaaaaatataaagaactcacgaacagcagctcccaggcgact
tcctcctgaatgtacccccacatagatggaccaaagtctaatctgttcagagagagca
aagcttacctcctttcatacgctttttctgtaggcgatgttttaaatatgactgcttcct
acatcgtaagtgaattattctttcatgcaacaccaacacttataagcggaagaacac
agaaacagctctagacaacaaacctgtggaccacagtggtaccagcatttgaggaggagc
aaaggagtttgctgctgctctcaccgctgagcggataaagacccacaaacgtccagg
aggcgcgagaagaggacggcttccaataacagtagcaggccagcaccaccaccattaa
tgtgctggaatcaaaggatacagacagtgatagggaagcagggactgaaacggggggaga
gaacaatgataaagaagaagaagaagaagatgaaacttcgagctcctctgaagcaaa
ttctcgggtgtcaaacaccaataaagatgaagccaaatattgaacctcctgagaatgtgga
gtggagtggtgctgaagcctcaatgttttagagtccctcattggcacttactatgacaattt
ctgtgccattgctagggttaattgggacaaaacatgtagacaggtgtatgagtttagagt
caagaatctagcatcatagctccagctcccgctgaggatgtggatactcctccaaggaa

- continued

aaagaagaggaaacacccggttggtgggtgcacactgcagaaagatacagctgaaaaagga
cggtcctcctaaccatggtttacaactatcaaccctgtgatcatccacggcagccttgatga
cagttcgtgcccctgtgtgatagcacaaaattttgtgaaaagttttgtcaatgtagttc
agagtgtcaaaacccgctttccgggatgccgtgcaaagcacagtgcacaccaagcagtg
cccgtgtacctgggtgtccgagagtggtgacctgacctgtgtcttacttgtggagccgc
tgaccattgggacagtaaaaaatgtgtcctgcaagaactgcagattcagcggggctccaa
aaagcatctattgctggcaccatctgacgtggcaggctgggggattttatcaaagatcc
tgtgcagaaaaatgaattcatctcagaatactgtggagagattttctcaagatgaagc
tgacagaagagggaaagtgtatgataaatacatgtgcagctttctgttcaacttgaacaa
tgattttgtgggtggatgcaacccgcaagggttaaaaaattcgttttgcaaatcattcggt
aaatccaaactgctatgcaaaagtattgatggtaacgggtgatcacaggataggtatttt
tgccaagagagccatccagactggcgaagagctgtttttgattacagatacagccaggc
tgatgccctgaagtatgtcggcatcgaaagagaaatggaaatcccttgacatctgctacc
tcctccccctcctctgaaacagctgccttagcttcaggaacctcgagtactgtgggcaa
tttagaaaaagaacatgcagtttgaaattctgaatttgcaaagtactgaagaataattt
atagtaatgagtttaaaaatcaactttttattgccttctcaccagctgcaaaagtgtttg
taccagtgaatttttgcaataatgcagtatggtacatttttcaactttgaataaagaata
cttgaacttgctctgttgaatc

Full amino acid of EZH2, isoform a (GenBank Accession No.
NP_004447)

(SEQ ID NO: 3)

MGQTGKKSEKGPVCRWRKRVKSEYMLRLQLKRFRRADEVKSMFSSNRQKILERTEILNQEW
KQRRIQPVHILTSVSSLRGTRECSVTSDLDFPTQVIPLKTLNAVASVPIMYSWSPLQQNF
MVEDETVLHNIPIYMGDEVLDQDGTFIGEELIKNYDGKVHGDRECGFINDEIFVELVNALGQ
YNDDDDDDDGDDPEEREKQKDLERDDKESRPPKFPSPDKIFEAISSMFPDKGTAEEL
KEYKELTEQQLPGALPPECTPNIDGPNKSVQREQSLHSFHTLFCRRCFKYDCFLHRKC
NYSFHATPNTYKRIKNTETALDNKPCGPQCYQHLEGAKEFAAALTAERIKTPPKRPGGRRR
GRLPNNSSRPSTPTINVLESKDTSDREAGTETGGENNDKEEEKKDETSSSSSEANSRCQ
TPIKMKPNIEPPENVEWSGAEASMPRVLIGTYDNCFAIARLIGTKTCRQVYEFVKES
IIAPAPAEDVDTPPRKKRKHRLWAAHCRKIQLKKDGSNHVYNYQPCDHRQPCDSSCP
CVIAQNFCEKFCQCSSECQNRFPGRCKAQCNKQCPCYLAVRECDPDLCLTCGAADHWD
SKNVSCNKCSIQRGSKKHLALLAPSDVAGWGIFIKDPVQKNEFISEYCGEIIISQDEADRRG
KVYDKYMCSEFLNLNDFVVDATRKGNKIRFANHSVNPNCYAKVMVMNGDHRIGIFAKRA
IQTGEELFFDYRYSQADALKYVGIEREMEIP

mRNA sequence of human EZH2, transcript variant 2 (GenBank
Accession No. NM_152998)

(SEQ ID NO: 4)

ggcggcgcttgattgggctgggggggccaataaaagcgatggcgattgggctgccgcgt
ttggcgctcgggtccgggtcgcgtccgacacccgggtgggactcagaaggcagtgaggcccg
gcggcgggcgggcgggcgcgggggcgacgcggggaacaacgcgagtcggcgcgcggg
acgaagaataatcatgggccagactgggaagaaatctgagaaggaccagtttgttgcg
gaagcgtgtaaaatcagagtacatgcgactgagacagctcaagaggttcagacgagctga

- continued

tgaagtaaagagtatgtttagttccaatcgtcagaaaaatttggaaagaacggaaatctt
aaaccaagaatggaacagcgaaggatacagcctgtgcacatcctgacttctgtgagctc
attgcgcgaggactaggaggtggaagatgaaactgtttacataacattccttatatggg
agatgaagttttagatcaggatggtactttcattgaagaactaataaaaattatgatgg
gaaagtacacgggtagagaatgtgggtttataaatgatgaaatttttgtggagtgggt
gaatgcccttggcfaatataatgatgatgacgatgatgatgatggagacgatcctgaaga
aagagaagaaaagcagaagatctggaggatcacggagatgataaagaaagccgccacc
tcggaaatttcttctgataaaattttgaagccatttctcfaatgtttccagataaggg
cacagcagaagaactaaaggaaaaatataagaactcacgaacagcagctccaggcgc
acttctctctgaatgtaccccaacatagatggacaaatgctaaatctgttcagagaga
gcaaagcttacactcctttcatacgttttctgtaggcgatgttttaatatgactgctt
cctacatccttttcatgcaacacccaacacttataagcgaagaacacagaacagctct
agacaacaaaccttgtggaccacagtgttaccagcatttggaggaggcaaaggagtttgc
tgctgctctcaccgctgagcggataaagacccccacaaaacgtccaggaggccgcagaag
aggacggcttcccaataacagtagcaggccagcaccaccatfaatgtgctggaatc
aaaggatacagacagtgataggaagcagggaactgaaacggggggagagaacaatgataa
agaagaagaagagaagaaagatgaaacttcgagctcctctgaagcaaatctcgggtgtca
aacccaataaagatgaagccaaatattgaacctcctgagaatgtggagtggagtgggtgc
tgaagcctcaatgtttagagtctcattggcacttactatgacaatttctgtgccattgc
taggttaattgggacaaaacatgtagacaggtgtatgagtttagagtcaaagaatctag
catcatagctccagctcccgtgaggatgtggatactcctccaaggaaaaagaaggaa
acaccgggtgtgggctgcacactgcagaaagatacagctgaaaaaggacggctcctctaa
ccatgtttacaactatcaacctgtgatcatccacggcagcCttgtgacagttcgtgccc
ttgtgtgatagcacaaaaattttgtgaaaagtgttgcfaatgttagttcagagtgtcaaaa
ccgcttccggggtgcccgtgcaaagcacagtgcacaccaagcagtgcccgtgctacct
ggctgtccgagagtgtgacctgacctctgtcttacttgtggagccgctgaccattggga
cagtaaaaatgtgtcctgcaagaactgcagtattcagcggggctccaaaagcatctatt
gctggcaccatctgacgtggcaggctgggggattttatcaaagatcctgtgcagaaaaa
tgaattcatctcagaatactgtggagagattatttctcaagatgaagctgacagaagagg
gaaagtgtatgataaatacatgtgcagcttctgttcaacttgaacaatgattttgtgggt
ggatgcaaccgcgaagggtacaaaattcgttttgcaaatcattcggtaaatccaaactg
ctatgcaaaagtattgatggttaacggtgatcacaggataggtatttttgccaagagagc
catccagactggcgaagagctgtttttgattacagatacagccaggctgatgccctgaa
gtatgtcggcatcgaaagagaaatggaaatcccttgacatctgctacctcctccccctc
ctctgaaacagctgccttagcttcaggaacctcgagtactgtgggcaatttagaaaaaga
acatgcagtttgaattctgaatttgcaaagtactgtaagaataatttatagtaatgagt
ttaaaaatcaactttttatgccttctcaccagctgcaaagtgttttgtaccagtgaatt
tttgcataaatgcagtatggtacatttttcaacttgaataaagaataacttgaacttgtc
cttgttgaatc

-continued

Full amino acid of EZH2, isoform b (GenBank Accession No. NP_694543)

(SEQ ID NO: 5)

MGQTGKKSEKGPVCWRKRVKSEYMRLRQLKRFRRADEVKSMFSSNRQKIL
 ERTEILNQEWKQRRIQPVHILTSVSSLRGTREREDEVTLHNIPLYMGDEVL
 DQDGTFFIEELIKNYDGKVHGDRECGFINDEIFVELVNALGQYNDDDDDD
 GDDPEEREKQKDLHRDDKESRPPRKFP SDKIFEAISSMFPDKGTAE
 LKEYKELTEQQLPGALPPECTPNIDGPNKSVQREQSLHSFHTLFCRR
 FKYDCFLHPFHATPNTYKRKNTETALDNKPCGPQCYQHLEGAKEFAAALT
 AERIKTPPKRPGGRRRGRLPNNSSRPSTPTINVLESKDTSDREAGTETG
 GENNDKEEEKKDETSSSSSEANSRCQTPIKMKPNIEPPENVEWSGAEAS
 FRVLIGTYDYNFCAIARLIGTKTCRQVYEFVRKESSIIAPAPAEDVDTP
 RKKKRKHRLWAAHCRKIQLKKGSSNHVYNYQPCDHPRQPCDSSPCVIA
 QNFCEKFCQCSSECQNRFPGCRCKAQCNKQCPCYLAVRECDPDLCLTCG
 AADHWDSKNVSKNCSIQRGSKHLLAPSDVAGWGIKDPVQKNEFIS
 EYCGEIIISQDEADRRGKVYDKYMCFLFNLNDFVVDATRKGNKIRFANH
 SVNPNYAKVMMVNGDHRIGIFAKRAIQTGEELFFDYRYSQADALKYVGI
 EREMEIP

Full amino acid of EZH2, isoform e (GenBank Accession No. NP_001190178.1)

(SEQ ID NO: 21)

MGQTGKKSEKGPVCWRKRVKSEYMRLRQLKRFRRADEVKSMFSSNRQKILERTEILNQEWKQRRIQPVHI
 LTSCSVTSDLDFPTQVIPLKTLNAVASVPIMYSWSPLQQNFMVEDEVTLHNIPLYMGDEVLDQDGTFFIEEL
 IKNYDGKVHGDRECGFINDEIFVELVNALGQYNDDDDDDGDDPEEREKQKDLHRDDKESRPPRKFP
 SDKIFEAISSMFPDKGTAEELKEYKELTEQQLPGALPPECTPNIDGPNKSVQREQSLHSFHTLFCRR
 FKYDCFLHPFHATPNTYKRKNTETALDNKPCGPQCYQHLEGAKEFAAALTAERIKTPPKRPGGRRRGRLP
 NNSRPSTPTINVLESKDTSDREAGTETGGENNDKEEEKKDETSSSSEANSRCQTPIKMKPNIEPPEN
 VEWSGAEASMFRLIGTYDYNFCAIARLIGTKTCRQVYEFVRKESSIIAPAPAEDVDTPPKKKRKHRLW
 AAHCRKIQLKKGQNRFPGCRCKAQCNKQCPCYLAVRECDPDLCLTCGAADHWDSKNVSKNCSIQRGSK
 KHLAPSDVAGWGIKDPVQKNEFISEYCGEIIISQDEADRRGKVYDKYMCFLFNLNDFVVDATRK
 GNKIRFANHSVNPNYAKVMMVNGDHRIGIFAKRAIQTGEELFFDYRYSQADALKYVGI EREMEIP

Homo sapiens enhancer of zeste homolog 2 (*Drosophila*) (EZH2), transcript variant 5, mRNA (GenBank Accession No. NM_001203249.1)


(SEQ ID NO: 22)

GACGACGTTCCGCGCGGGGAACGAGTAGCTTCGCCTCTGACGTTTCCCCACGACGACCCCGAAATC
 CCCCTGAGCTCCGCGGCTCGCGGCTGCCCTCGCCGCTGCTGGCTTTATGCTAAGTTTGAGGAAGA
 GTCGAGCTGCTCTGCTCTCTATTGATTGTGTTTCTGGAGGCGCTCTGTTGAATCCCACTTCATTGTGT
 ACATCCCTTCCGTTCCCCCCAAAACTGTGCCACAGGGTTACTTTTGAAGCGGAGGAATCGAGAA
 GCACGATCTTTTGAAAACCTTGGTGAACGCCTAAATAATCATGGCCAGACTGGGAAGAAATCTGAGAAG
 GGACCAGTTTGTGGCGGAAGCGTGTAATAATCAGAGTACATGCGACTGAGACAGCTCAAGAGGTTGAGAC
 GAGCTGATGAAGTAAAGAGTATGTTTAGTTCCAATCGTCAGAAAATTTTGAAAGAACGGAAATCTTAA
 CCAAGATGGAACAGCGAAGGATACAGCCTGTGCACATCCTGACTTCTTGTTCGGTGACCAGTGACTTG
 GATTTTCCAACACAAGTCATCCATTAAAGACTCTGAATGCAGTTGCTTCAGTACCCATAATGTATTCTT
 GGTCTCCCTACAGCAGAAATTTATGGTGAAGATGAACTGTTTTACATAACATTCTTATATGGGAGA

- continued

TGAAGTTT TAGATCAGGATGGTACTTT CATTGAAGAACTAATAAAAAATTATGATGGGAAAGTACACGGG
GATAGAGAATGTGGGTTTATAAATGATGAAATTTTGTGGAGTTGGTGAATGCCCTTGGTCAATATAATG
ATGATGACGATGATGATGATGGAGACGATCCTGAAGAAAGAGAAGAAAAGCAGAAAGATCTGGAGGATCA
CCGAGATGATAAAGAAAGCCGCCACCTCGGAAATTTCTTCTGATAAAATTTTGAAGCCATTTCTCA
ATGTTTCCAGATAAGGGCACAGCAGAGAAGAACTAAAGGAAAAATATAAGAAGCTCACCGAACAGCAGCTCC
CAGGCGCACTTCTCTGAATGTACCCCAACATAGATGGACCAATGCTAAATCTGTTCAGAGAGAGCA
AAGCTTACACTCCTTTTATACGCTTTTCTGTAGGCGATGTTTTAAATATGACTGCTTCCTACATCCTTTT
CATGCAACACCCAACTTTATAAGCGGAAGAACACAGAAACAGCTCTAGACAACAACTTGTGGACCAC
AGTGTACCAGCATTTGGAGGGAGCAAGGAGTTTGCTGCTGCTCTCACCGTGAGCGGATAAAGACCCC
ACCAAAACGTCAGGAGGCGCAGAAAGGACGGCTTCCAATAACAGTAGCAGGCCAGCACCCCCACC
ATTAATGTCTGGAATCAAAGGATACAGACAGTGATAGGGAAGCAGGACTGAAACGGGGGAGAGAACA
ATGATAAAGAAGAAGAAGAGAAGAAAGATGAACTTCGAGCTCCTCTGAAGCAAATTCGCGTGTCAAAC
ACCAATAAAGATGAAGCCAATATTGAACCTCCTGAGAATGTGGAGTGGAGTGGTGAAGCCTCAATG
TTTAGAGTCCTCATTGGCACTTACTATGACAATTTCTGTGCCATTGCTAGGTAAATTGGGACAAAACAT
GTAGACAGGTGTATGAGTTTAGAGTCAAAGAACTAGCATCATAGCTCCAGCTCCCGCTGAGGATGTGGA
TACTCTCCAAAGGAAAAAGAGAGAAACACCGGTTGTGGGCTGCACACTGCAGAAAGATACAGCTGAAA
AAGGGTCAAAACCGCTTTCCGGGATGCCGCTGCAAAGCACAGTGCAACACCAAGCAGTGCCCGTGCTACC
TGGCTGTCCGAGAGTGTGACCTGACCTCTGTCTTACTTGTGGAGCCGCTGACCATTGGGACAGTAAAA
TGTGTCTGCAAGAACTGCAGTATTACGCGGGCTCCAAAAAGCATCTATTGCTGGCACCATCTGACGTG
GCAGGCTGGGGGATTTTATCAAAGATCCTGTGCAGAAAAATGAATTCATCTCAGAATACTGTGGAGAGA
TTATTTCTCAAGATGAAGCTGACAGAAGAGGAAAGTGTATGATAAATACATGTGCAGCTTTCTGTTCAA
CTTGAAACATGATTTTGTGGTGGATGCAACCCGCAAGGGTAACAAAATTCGTTTGCAAATCATTTCGGTA
AATCCAACTGCTATGCAAAAGTTATGATGGTTAACGGTGATCACAGGATAGGTATTTTGCCAAGAGAG
CCATCCAGACTGGCGAAGAGCTGTTTTTGATTACAGATACAGCCAGGCTGATGCCCTGAAGTATGTCGG
CATCGAAAGAGAAATGGAAATCCCTTGACATCTGCTACCTCCTCCCCCTCCTCTGAAACAGCTGCCTTA
GCTTCAGGAACCTCGAGTACTGTGGGCAATTTAGAAAAAGAACATGCAGTTTGAAATTCGAATTTGCAA
AGTACTGTAAAGAATAATTTATAGTAATGAGTTTAAAAATCAACTTTTTATTGCCCTTCTACCAGCTGCAA
AGTGTTTTGTACCAGTGAATTTTGAATAATGCAGTATGGTACATTTTCAACTTTGAATAAAGAATAC
TTGAACCTTGTCCTTGTGTAATC

[0045] A structure model of partial EZH2 protein based on the A chain of nuclear receptor binding SET domain protein 1 (NSD1) is provided below. This model corresponds to amino acid residues 533-732 of EZH2 sequence of SEQ ID NO: 1.



[0046] The corresponding amino acid sequence of this structure model is provided below. The residues in the substrate pocket domain are underlined. The residues in the SET domain are shown italic.

(SEQ ID NO: 6)
 SPCPVIAQNFCCKFCQCSSECQNRFPGRCKAQCNKQPCYLAVRECDPDLCLTCGAA
 DHWDSKNVSCNKNCISIQRGSKK HILLAPSDVAGWGIFIKDPVQKNEFISEY⁶⁴¹ *CGEIISQDEA*
*DRRGKVYDKYMCSEFLNNDVY*⁶⁷⁴ *YDA*⁶⁷⁷ *TRKGNKIK*⁶⁸⁵ *FA*⁶⁸⁷ *NHSVNPNCYAKVMMVNGD*
HRIGIFAKRAIQTGEELFFDYRSQAD

[0047] The catalytic site of EZH2 is believed to reside in a conserved domain of the protein known as the SET domain. The amino acid sequence of the SET domain of EZH2 is provided by the following partial sequence spanning amino acid residues 613-726 of Swiss-Prot Accession No. Q15910 (SEQ ID NO: 1):

(SEQ ID NO: 7)
 HLLLAPSDVAGWGIFIKDPVQKNEFISEYCGEIISQDEADRRGKVYDKYMCSEFLNNDVVD
 ATRKGNKIRFANHSVNPNCYAKVMMVNGDHRIGIFAKRAIQTGEELFFDY.

[0048] The tyrosine (Y) residue shown underlined in SEQ ID NO: 7 is Tyr641 (Y641) in Swiss-Prot Accession No. Q15910 (SEQ ID NO: 1).

[0049] The SET domain of GenBank Accession No. NP_004447 (SEQ ID NO: 3) spans amino acid residues 618-731 and is identical to SEQ ID NO:6. The tyrosine residue corresponding to Y641 in Swiss-Prot Accession No. Q15910 shown underlined in SEQ ID NO: 7 is Tyr646 (Y646) in GenBank Accession No. NP_004447 (SEQ ID NO: 3).

[0050] The SET domain of GenBank Accession No. NP_694543 (SEQ ID NO: 5) spans amino acid residues 574-687 and is identical to SEQ ID NO: 7. The tyrosine residue corresponding to Y641 in Swiss-Prot Accession No. Q15910 shown underlined in SEQ ID NO: 7 is Tyr602 (Y602) in GenBank Accession No. NP_694543 (SEQ ID NO: 5).

[0051] The nucleotide sequence encoding the SET domain of GenBank Accession No. NP_004447 is

(SEQ ID NO: 8)
 catctattgctggcaccatctgacgtggcaggctgggggatttttatcaaagatcctgtgcaga
 aaaatgaattcatctcagaatactgtggagagattatttctcaagatgaagctgacagaagagg
 gaaagtgtatgataaaatcatgtgcagctttctgttcaactgaacaatgattttgtggtggat
 gcaacccgcaagggttaacaaattcgttttgcaaatcattcggttaaatccaaactgctatgcaa
 aagttatgatggttaacgggtgatcacaggataggtatttttccaagagagccatccagactgg
 cgaagagctgttttttgattac,

where the codon encoding Y641 is shown underlined.

[0052] For purposes of this application, amino acid residue Y641 of human EZH2 is to be understood to refer to the tyrosine residue that is or corresponds to Y641 in Swiss-Prot Accession No. Q15910.

Full amino acid sequence of Y641 mutant EZH2

(SEQ ID NO: 9)
 MGQTGKKSEKGPVCWRKRVKSEYMRLRQLKRFRRADEVKSMFSSNRQKILERTILNQEW
 KQRRIQPVHILTSVSSLRGTRECSVTSDDLDFPTQVIPLKTLNAVASVPIMYSWSPLQQNF
 MVEDETVLHNPYMGDEVLDQDGTFFIEELIKNYDGKVGHDRECGFINDEIFVELVNALGQ
 YNDDDDDDGDDPEEREKQKDLEDHRDDKESRPPRKFPSPDKIFEAISSMFPDKGTAEEL
 KEKYKELTEQQLPGALPPECTPNIDGPNKSVQREQSLHSFHTLFCRRCFKYDCFLHPFH

-continued

ATPNTYKRKNTETALDNKPCGPQCYQHLEGAKEFAAALTAERIKTPPKRPGRRRGRLPN
 NSSRPSTPTINVLESKDTSREAGTETGGENNDKEEEKKDETSSSEANSRCQTPIKM
 KPNIIPPENVEWSGAEASMPRVLIGTYDNFCAIARLIGTKTCRQVYEFVKESSIIAPA
 PAEDVDTPPRKKRKHRLWAAHCRKIQLKKDGSSNHVYNYQPCDHPRQPCDSSCPCVIAQ
 NFCEKFCQCSSECQNRFPGCRCKAQCNTKQPCYLAVERCDPDLCLTCGAADHWDKSNVS
 CKNCISIQRGSKKHLALLAPSDVAGWGIFIKDPVQKNEFISEXCGEIIISQDEADRRGKVYDK
 YMCSTFLNNDNFVVDATRKGNKIRFANHNVNPNKYAKVMMVNGDHRIGIFAKRAIQTGE
 ELFFDYRYSQADALKYVGIEREMEIP
 Wherein x can be any amino acid residue other than tyrosine (Y)

[0053] Also for purposes of this application, a Y641 mutant of human EZH2, and, equivalently, a Y641 mutant of EZH2, is to be understood to refer to a human EZH2 in which the amino acid residue corresponding to Y641 of wild-type human EZH2 is substituted by an amino acid residue other than tyrosine.

[0054] In one embodiment the amino acid sequence of a Y641 mutant of EZH2 differs from the amino acid sequence of wild-type human EZH2 only by substitution of a single

amino acid residue corresponding to Y641 of wild-type human. EZH2 by an amino acid residue other than tyrosine.

[0055] In one embodiment the amino acid sequence of a Y641 mutant of EZH2 differs from the amino acid sequence of wild-type human EZH2 only by substitution of phenylalanine (F) for the single amino acid residue corresponding to Y641 of wild-type human EZH2. The Y641 mutant of EZH2 according to this embodiment is referred to herein as a Y641F mutant or, equivalently, Y641F.

Y641F

(SEQ ID NO: 10)

MGQTGKKSEKGPVCRKRVKSEYMRLRQLKRFRRADEVKSMFSSNRQKILERTTEILNQEW
 KQRRIQPVHILTSVSSSLRGTRRECSVTSLDFTQVPLKTLNAVASVPIMYSWSPLQQNF
 MVEDETVLHNIPYMGDEVLDQDGTFFIEELIKNYDGKVHGDRECGFINDEIFVELVNALGQ
 YNDDDDDDGDDPEEREKQKDLDRDDKESRPPRKFPSPDKIFEAISSMFPDKGTADEL
 KEKYKELTEQQLPGALPPECTPNIDGPNKSVQREQSLHSFHTLFCRRCKFYDCFLHPFH
 ATPNTYKRKNTETALDNKPCGPQCYQHLEGAKEFAAALTAERIKTPPKRPGRRRGRLPN
 NSSRPSTPTINVLESKDTSREAGTETGGENNDKEEEKKDETSSSEANSRCQTPIKM
 KPNIIPPENVEWSGAEASMPRVLIGTYDNFCAIARLIGTKTCRQVYEFVKESSIIAPA
 PAEDVDTPPRKKRKHRLWAAHCRKIQLKKDGSSNHVYNYQPCDHPRQPCDSSCPCVIAQ
 NFCEKFCQCSSECQNRFPGCRCKAQCNTKQPCYLAVERCDPDLCLTCGAADHWDKSNVS
 CKNCISIQRGSKKHLALLAPSDVAGWGIFIKDPVQKNEFISEFCGEIIISQDEADRRGKVYDK
 YMCSTFLNNDNFVVDATRKGNKIRFANHNVNPNKYAKVMMVNGDHRIGIFAKRAIQTGE
 ELFFDYRYSQADALKYVGIEREMEIP

[0056] In one embodiment the amino acid sequence of a Y641 mutant of EZH2 differs from the amino acid sequence of wild-type human EZH2 only by substitution of histidine (H) for the single amino acid residue corresponding to Y641 of wild-type human EZH2. The Y641 mutant of EZH2 according to this embodiment is referred to herein as a Y641H mutant or, equivalently, Y641H.

Y641H

(SEQ ID NO: 11)

MGQTGKKSEKGPVCRKRVKSEYMRLRQLKRFRRADEVKSMFSSNRQKILERTTEILNQEW
 KQRRIQPVHILTSVSSSLRGTRRECSVTSLDFTQVPLKTLNAVASVPIMYSWSPLQQNF
 MVEDETVLHNIPYMGDEVLDQDGTFFIEELIKNYDGKVHGDRECGFINDEIFVELVNALGQ

-continued

YNDDDDDDGDDPEEREKQKDLLEDHRDDKESRPPRFPSDKIFEAISSMFPDKGTAEEL
 KEKYKELTEQQLPGALPPECTPNIDGPNASVQREQSLHSFHTLFCRRCFKYDCFLHPFH
 ATPNTYKRKNTETALDNKPCGPQCYQHLEGAKEFAAALTAERI KTPPKRPGRRRGRLPN
 NSSRPSTPTINVLESKD TSDREAGTETGGENNDKEEEKKDETSSSEANSRCQTPIKM
 KPNI EPPENVEWSGAEASMFVRLIGTYDNFCAIARLIGTKTCRQVVEFRVKESSIAPA
 PAEDVDTPPRKKRKHRLWAAHCRKIQLKKDGSSNHVYNYQPCDHPRQPCDSSCPCVIAQ
 NFCEKFCQCSSECQNRFPGCRCKAQCN TKQPCYLA VRECDPDLCLTCGAADHWDSKNVS
 CKNC SIQRGSKKHL L LAPS DVAGW G I F I K D P V Q K N E F I S E H C G E I I S Q D E A D R R G K V Y D K
 YMCSFLFNLNND FVVDATRKGNKIRFANH SVNPNCYAKVMVNGDHRIGIFAKRAIQTGE
 ELFFDYRYSQADALKYVGIEREMEIP

[0057] In one embodiment the amino acid sequence of a Y641 mutant of EZH2 differs from the amino acid sequence of wild-type human EZH2 only by substitution of asparagine (N) for the single amino acid residue corresponding to Y641 of wild-type human EZH2. The Y641 mutant of EZH2 according to this embodiment is referred to herein as a Y641N mutant or, equivalently, Y641N.

Y641N
 (SEQ ID NO: 12)
 MGQTGKKSEKGPVCWRKRVKSEYMRLRQLKRFRRADEVKSMFSSNRQKILERTEILNQEW
 KQRRIQPVHILTSVSSLRGTRECSVTSDLDFPTQVIPLKTLNAVASVPIMYSWSPLQQNF
 MVEDETVLHNIPYMGDEVLDQDGT FIEELIKNYDGKVHGDRECGFINDEIFVELVNALGQ
 YNDDDDDDGDDPEEREKQKDLLEDHRDDKESRPPRFPSDKIFEAISSMFPDKGTAEEL
 KEKYKELTEQQLPGALPPECTPNIDGPNASVQREQSLHSFHTLFCRRCFKYDCFLHPFH
 ATPNTYKRKNTETALDNKPCGPQCYQHLEGAKEFAAALTAERI KTPPKRPGRRRGRLPN
 NSSRPSTPTINVLESKD TSDREAGTETGGENNDKEEEKKDETSSSEANSRCQTPIKM
 KPNI EPPENVEWSGAEASMFVRLIGTYDNFCAIARLIGTKTCRQVVEFRVKESSIAPA
 PAEDVDTPPRKKRKHRLWAAHCRKIQLKKDGSSNHVYNYQPCDHPRQPCDSSCPCVIAQ
 NFCEKFCQCSSECQNRFPGCRCKAQCN TKQPCYLA VRECDPDLCLTCGAADHWDSKNVS
 CKNC SIQRGSKKHL L LAPS DVAGW G I F I K D P V Q K N E F I S E N C G E I I S Q D E A D R R G K V Y D K
 YMCSFLFNLNND FVVDATRKGNKIRFANH SVNPNCYAKVMVNGDHRIGIFAKRAIQTGE
 ELFFDYRYSQADALKYVGIEREMEIP

[0058] In one embodiment the amino acid sequence of a Y641 mutant of EZH2 differs from the amino acid sequence of wild-type human EZH2 only by substitution of serine (S) for the single amino acid residue corresponding to Y641 of wild-type human EZH2. The Y641 mutant of EZH2 according to this embodiment is referred to herein as a Y641S mutant or, equivalently, Y641S.

Y641S
 (SEQ ID NO: 13)
 MGQTGKKSEKGPVCWRKRVKSEYMRLRQLKRFRRADEVKSMFSSNRQKILERTEILNQEW
 KQRRIQPVHILTSVSSLRGTRECSVTSDLDFPTQVIPLKTLNAVASVPIMYSWSPLQQNF
 MVEDETVLHNIPYMGDEVLDQDGT FIEELIKNYDGKVHGDRECGFINDEIFVELVNALGQ

-continued

YNDDDDDDGDDPEEREKQKDLEDHRDDKESRPPRKFPSPDKIFEAISSMFPDGTAEEL
 KEKYKELTEQQLPGALPPECTPNIDGPNASVQREQSLHSFHTLFCRRCFKYDCFLHPFH
 ATPNTYKRKNTETALDNKPCGPQCYQHLEGAKEFAAALTAERIKTPPKRPGRRRRLPN
 NSSRPSTPTINVLESKDTSDREAGTETGGENNDKEEEKKDETSSSEANSRCQTPIKM
 KPNIEPPENVEWSGAEASMFVRLIGTYDYNFCAIARLIGTKTCRQVYEFVKESSIIAPA
 PAEDVDTPPRKKRKHRLWAAHCRKIQKKDGSSNHVINYQPCDHPQPCDSSCPCVIAQ
 NFCEKFCQCSSECQNRFPQCRCKAQCNKQPCPCYLAVRECDPDLCLTCGAADHWDSKNVS
 CKNCISIQRSGSKHLLAPSDVAGWIGIFIKDPVQKNEFISESCGEIISQDEADRRGKVYDK
 YMCSEFLNLDNFVVDATRGKNGKIRFANHNSVNPNCYAKVMMVNGDHRIGIFAKRAIQTGE
 ELFFDYRYSQADALKYVGIEREMEIP

[0059] In one embodiment the amino acid sequence of a Y641 mutant of EZH2 differs from the amino acid sequence of wild-type human EZH2 only by substitution of cysteine (C) for the single amino acid residue corresponding to Y641 of wild-type human EZH2. The Y641 mutant of EZH2 according to this embodiment is referred to herein as a Y641C mutant or, equivalently, Y641C.

Y641C

(SEQ ID NO: 14)

MGQTGKKSEKGPVCRKRVKSEYMRLRQLKRFRRADEVKSMFSSNRQKILERTTEILNQEW
 KQRIQPVHILTSVSSLRGTRECSVTSDLDFTQVIPLKTLNAVASVPIMYSWSPLQQNF
 MVEDETVLHNIPYMGDEVLDQDGTFFIEELIKNYDGKVHGDRECGFINDEIFVELVNALGQ
 YNDDDDDDGDDPEEREKQKDLEDHRDDKESRPPRKFPSPDKIFEAISSMFPDGTAEEL
 KEKYKELTEQQLPGALPPECTPNIDGPNASVQREQSLHSFHTLFCRRCFKYDCFLHPFH
 ATPNTYKRKNTETALDNKPCGPQCYQHLEGAKEFAAALTAERIKTPPKRPGRRRRLPN
 NSSRPSTPTINVLESKDTSDREAGTETGGENNDKEEEKKDETSSSEANSRCQTPIKM
 KPNIEPPENVEWSGAEASMFVRLIGTYDYNFCAIARLIGTKTCRQVYEFVKESSIIAPA
 PAEDVDTPPRKKRKHRLWAAHCRKIQKKDGSSNHVINYQPCDHPQPCDSSCPCVIAQ
 NFCEKFCQCSSECQNRFPQCRCKAQCNKQPCPCYLAVRECDPDLCLTCGAADHWDSKNVS
 CKNCISIQRSGSKHLLAPSDVAGWIGIFIKDPVQKNEFISECGEIIISQDEADRRGKVYDK
 YMCSEFLNLDNFVVDATRGKNGKIRFANHNSVNPNCYAKVMMVNGDHRIGIFAKRAIQTGE
 ELFFDYRYSQADALKYVGIEREMEIP

[0060] In one embodiment the amino acid sequence of a A677 mutant of EZH2 differs from the amino acid sequence of wild-type human EZH2 only by substitution of a non-alanine amino acid, preferably glycine (G) for the single

amino acid residue corresponding to A677 of wild-type human EZH2. The A677 mutant of EZH2 according to this embodiment is referred to herein as an A677 mutant, and preferably an A677G mutant or, equivalently, A677G.

A677

(SEQ ID NO: 15)

MGQTGKKSEKGPVCRKRVKSEYMRLRQLKRFRRADEVKSMFSSNRQKILERTTEILNQEW
 KQRIQPVHILTSVSSLRGTRECSVTSDLDFTQVIPLKTLNAVASVPIMYSWSPLQQNF
 MVEDETVLHNIPYMGDEVLDQDGTFFIEELIKNYDGKVHGDRECGFINDEIFVELVNALGQ
 YNDDDDDDGDDPEEREKQKDLEDHRDDKESRPPRKFPSPDKIFEAISSMFPDGTAEEL

-continued

KEKYKELTEQQLPGALPPECTPNIDGPNASVQREQLHSFHTLFCRRCFKYDCFLHPPH
 ATPNTYKRKNTETALDNKPCGPQCYQHLEGAKEFAAALTAERIKTPPKRPGRRRGRLPN
 NSSRPSTPTINVLESKDTSDREAGTETGGENNDKEEEKKDETSSSEANSRCQTPIKM
 KPNIEPPENVEWSGAEASMFVRLIGTYDNPFCIAIRLIGTKTCRQVYEFVKESSIIAPA
 PAEDVDTPPRKKRKHRLWAAHCRKIQLKKDGSSNHVYNYQPCDHPRQPCDSSCPCVIAQ
 NFCEKFCQCSSECQNRFPGRCKAQCNKQPCYLAVERCDPDLCLTCGAADHWDKSNVS
 CKNC SIQRGSKKHL L LAPS DVAGW G I F I K D P V Q K N E F I S E Y C G E I I S Q D E A D R R G K V Y D K
 Y M C S F L F N L N D F V V D X T R K G N K I R F A N H S V N P N C Y A K V M M V N G D H R I G I F A K R A I Q T G E
 ELFFDYRYSQADALKYVGIEREMEIP
 wherein X is preferably a glycine (G).

[0061] In one embodiment the amino acid sequence of a A687 mutant of EZH2 differs from the amino acid sequence of wild-type human EZH2 only by substitution of a non-alanine amino acid, preferably valine (V) for the single amino

acid residue corresponding to A687 of wild-type human EZH2. The A687 mutant of EZH2 according to this embodiment is referred to herein as an A687 mutant and preferably an A687V mutant or, equivalently, A687V.

A687

(SEQ ID NO: 16)

MGQTGKKSEKGPVCRKRVKSEYMRLRQLKRFRRADEVKSMFSSNRQKILERTTEILNQEW
 KQRRIQPVHILTSVSSLRGTRECSVTSDLDFPTQVIPLKTLNAVASVPIMYSWSPLQQNF
 MVEDETVLHNIPYMGDEVLDQDGTFFIEELIKNYDGKVHGDRECGFINDEIFVELVNALGQ
 YNDDDDDDGDDPEEREKQKDLDRDDKESRPKRKFPSPDKIFEAISSMFPDKGTAEEL
 KEKYKELTEQQLPGALPPECTPNIDGPNASVQREQLHSFHTLFCRRCFKYDCFLHPPH
 ATPNTYKRKNTETALDNKPCGPQCYQHLEGAKEFAAALTAERIKTPPKRPGRRRGRLPN
 NSSRPSTPTINVLESKDTSDREAGTETGGENNDKEEEKKDETSSSEANSRCQTPIKM
 KPNIEPPENVEWSGAEASMFVRLIGTYDNPFCIAIRLIGTKTCRQVYEFVKESSIIAPA
 PAEDVDTPPRKKRKHRLWAAHCRKIQLKKDGSSNHVYNYQPCDHPRQPCDSSCPCVIAQ
 NFCEKFCQCSSECQNRFPGRCKAQCNKQPCYLAVERCDPDLCLTCGAADHWDKSNVS
 CKNC SIQRGSKKHL L LAPS DVAGW G I F I K D P V Q K N E F I S E Y C G E I I S Q D E A D R R G K V Y D K
 Y M C S F L F N L N D F V V D A T R K G N K I R F X N H S V N P N C Y A K V M M V N G D H R I G I F A K R A I Q T G E
 ELFFDYRYSQADALKYVGIEREMEIP
 wherein X is preferably a valine (V).

[0062] In one embodiment the amino acid sequence of a R685 mutant of EZH2 differs from the amino acid sequence of wild-type human EZH2 only by substitution of a non-arginine amino acid, preferably histidine (H) or cysteine (C) for the single amino acid residue corresponding to R685 of wild-type human EZH2. The R685 mutant of EZH2 according to this embodiment is referred to herein as an R685 mutant and preferably an R685C mutant or an R685H mutant or, equivalently, R685H or R685C.

A685

(SEQ ID NO: 17)

MGQTGKKSEKGPVCRKRVKSEYMRLRQLKRFRRADEVKSMFSSNRQKILERTTEILNQEW
 KQRRIQPVHILTSVSSLRGTRECSVTSDLDFPTQVIPLKTLNAVASVPIMYSWSPLQQNF
 MVEDETVLHNIPYMGDEVLDQDGTFFIEELIKNYDGKVHGDRECGFINDEIFVELVNALGQ

-continued

YNDDDDDDGDDPEEREKQKDLHRDDKESRPPRFPSDKIFEAISSMFPDKGTAEEL
 KEKYKELTEQQLPGALPPECTPNIDGPNASVQREQLHSFHTLFCRRCFKYDCFLHPFH
 ATPNTYKRKNTETALDNKPCGPQCYQHLEGAKEFAAALTAERIKTPPKRPGRRRGRLPN
 NSSRPSTPTINVLESKDTSDREAGTETGGENNDKEEEKKDETSSSSEANSRCQTPIKM
 KPNIEPPENVEWSGAEASMPFVLIGTYDNFCAIARLIGTKTCRQVYEFVKESSIIAPA
 PAEDVDTPPRKKRKHRLWAAHCRKIQLKKDGSNNHVNYQPCDHPRQPCDSSCPCVIAQ
 NFCEKFCQCSSECQNRFPGCRCKAQCNTKQCPCYLAVRECDPDLCLTCGAADHWDSKNVS
 CKNCISIQGSKKHLALLAPSDVAGWGFIFIKDPVQKNEFISEYCGEIIISQDEADRRGKVYDK
 YMCSEFLNNDNFVVDATRKGNKIXFANHVSVPNCYAKVMMVNGDHRIGIFAKRAIQTGE
 ELFFDYRYSQADALKYVGIEREMEIP
 Wherein X is preferably a cysteine (C) or a histidine (H).

[0063] In one embodiment the amino acid sequence of a mutant of EZH2 differs from the amino acid sequence of wild-type human EZH2 in one or more amino acid residues in its substrate pocket domain as defined in SEQ ID NO: 6. The mutant of EZH2 according to this embodiment is referred to herein as an EZH2 mutant.

tein EZH1 as the catalytic subunit. Shen, X. et al. (2008) *Mol Cell* 32:491-502. Hence, catalytic coupling between the mutant EZH2 species and PRC2 complexes containing either WT-EZH2 or WT-EZH1 would suffice to augment H3-K27me2/3 formation, and thus produce the attendant malignant phenotype. The data therefore suggest that the

Mutant EZH2 comprising one or more mutations in the substrate pocket domain

(SEQ ID NO: 18)

MGQTGKKSEKGPVCWRKRVKSEYMRLRQLKRFRRADEVKSMFSSNRQKILERTTEILNQEW
 KQRIQFVHILTSVSSLRGTRECSVTSDLDFPTQVIPLKTLNAVASVPIMYSWSPLQQNF
 MVEDETVLHNIPIYMGDEVLDQDGTFIIEELIKNYDGKVHGDRECGFINDEIFVELVNALGQ
 YNDDDDDDGDDPEEREKQKDLHRDDKESRPPRFPSDKIFEAISSMFPDKGTAEEL
 KEKYKELTEQQLPGALPPECTPNIDGPNASVQREQLHSFHTLFCRRCFKYDCFLHPFH
 ATPNTYKRKNTETALDNKPCGPQCYQHLEGAKEFAAALTAERIKTPPKRPGRRRGRIPN
 NSSRPSTPTINVLESKDTSDREAGTETGGENNDKEEEKKDETSSSSEANSRCQTPIKM
 KPNIEPPENVEWSGAEASMPFVLIGTYDNFCAIARLIGTKTCRQVYEFVKESSIIAPA
 PAEDVDTPPRKKRKHRLWAAHCRKIQLKKDGSNNHVNYQPCDHPRQPCDSSCPCVIAQ
 NFCEKFCQCSSECQNRFPGCRCKAQCNTKQCPCYLAVRECDPDLCLTCGAADHWDSKNVS
 CKNCISIQGSKKHLALLAPSDVAGWGFIFIKDPVQKNEFISEXCGEIIISQDEADRRGKVYDK
 YMXXXLXNLDNFXXDXTRKGNKXXXXHVSVPNCYAKVMMVNGDHRXGIFAKRAIQTGE
 ELFXDXRYSXADALKYVGIEREMEIP
 wherein X can be any amino acid except the corresponding wild type residue.

[0064] The implications of the present results for human disease are made clear by the data summarized in Table 2 (see Example 5). Cells heterozygous for EZH2 would be expected to display a malignant phenotype due to the efficient formation of H3-K27me1 by the WT enzyme and the efficient, subsequent transition of this progenitor species to H3-K27me2, and, especially, H3-K27me3, by the mutant enzyme form(s).

[0065] It has been reported that H3-K27me1 formation is not exclusively dependent on WT-EZH2 catalysis. Knockout studies of EZH2 and of another PRC2 subunit, EED, have demonstrated H3-K27me1 formation can be catalyzed by PRC2 complexes containing either EZH2 or the related pro-

malignant phenotype of follicular lymphoma (FL) and diffuse large B-cell lymphoma (DLBCL) of the germinal center B cell (GCB) subtype, associated with expression of mutant forms of EZH2, is the result of an overall gain of function with respect to formation of the trimethylated form of H3-K27. This interpretation of the data also helps to reconcile the existence of cancer-associated overexpression of EZH2 or PRC2 associated proteins (e.g., PHF19/PCL3) and also loss-of-function genotypes for the histone H3-K27 demethylase UTX. Loss of UTX activity would be enzymatically equivalent to a gain of function for EZH2, in either situation resulting in greater steady state levels of tri-methylated H3-K27 in cancer cells (FIG. 4).

[0066] The mono-, di-, and tri-methylation states of histone H3-K27 are associated with different functions in transcriptional control. Histone H3-K27 monomethylation is associated with active transcription of genes that are poised for transcription. Cui et al. (2009) *Cell Stem Cell* 4:80-93; Barski (2007) *Cell* 129:823-37. In contrast, trimethylation of histone H3-K27 is associated with either transcriptionally repressed genes or genes that are poised for transcription when histone H3-K4 trimethylation is in cis. Cui et al. (supra); Kirmizis et al. (2007) *Genes Dev* 18:1592-1605; Bernstein et al. (2006) *Cell* 125:315-26. Taken together, alterations in the PRC2 complex activity reported in cancer, including the Y641 mutation of EZH2, are predicted to result in an increase in the trimethylated state of histone H3-K27 and thus to result in transcriptional repression.

[0067] Another discovery of the present invention is that cells expressing a mutant EZH2 comprising a mutation in the substrate pocket domain as defined in SEQ ID NO: 6 are, in general, more sensitive to small molecule EZH2 inhibitors than cells expressing wild type (WT) EZH2. Specifically, cells expressing Y641 mutant EZH2 show reduced growing, dividing or proliferation, or even undergo apoptosis or necrosis after the treatment of EZH2 inhibitors. In contrast, cells expressing WT EZH2 are not responsive to the anti-proliferative effect of the EZH2 inhibitors (FIGS. 13 and 14). Particularly, cells expressing a substitution mutation at amino acid position 677 of SEQ ID NO: 1 show greater sensitivity to EZH2 inhibitors than cells expressing other EZH2 mutants (Example 19).

[0068] EZH2 and other protein methyltransferases have been suggested to be attractive targets for drug discovery. Copeland et al. (2009) *Nat Rev Drug Discov* 8:724-32; Copeland et al. (2010) *Curr Opin Chem Biol* 14(4):505-10; Pollock et al. (2010) *Drug Discovery Today: Therapeutic Strategies* 6(1):71-9. The present data also suggest an experimental strategy for development of FL and GCB lymphoma-specific drugs. As the differences in substrate recognition between the WT and disease-associated mutants derive from transition state interactions, small molecule inhibitors that selectively mimic the transition state of the mutant EZH2 over that of the WT enzyme should prove to be effective in blocking H3-K27 methylation in mutation-bearing cells. Inhibitors of this type would be expected to display a large therapeutic index, as target-mediated toxicity would be minimal for any cells bearing only the WT enzyme. Transition state mimicry has proved to be an effective strategy for drug design in many disease areas. See, for example, Copeland, R. A. *Enzymes: A Practical Introduction to Structure, Mechanism and Data Analysis*. 2nd ed, (Wiley, 2000).

[0069] The present results point to a previously unrecognized, surprising dependency on enzymatic coupling between enzymes that perform H3-K27 mono-methylation and certain mutant forms of EZH2 for pathogenesis in follicular lymphoma and diffuse large B-cell lymphoma. While not intending to be bound by any one theory, it is believed the data constitute the first example of a human disease that is dependent on such coupling of catalytic activity between normal (WT) and disease-associated mutant (such as Y641) enzymes.

[0070] An aspect of the present invention is a method for treating or alleviating a symptom of cancer or precancerous condition in a subject by administering to a subject expressing a mutant EZH2 comprising a mutation in the substrate pocket

domain as defined in SEQ ID NO: 6 a therapeutically effective amount of an EZH2 inhibitor.

[0071] Another aspect of the invention is a method for inhibiting in a subject conversion of H3-K27 to trimethylated H3-K27. The inhibition can involve inhibiting in a subject conversion of unmethylated H3-K27 to monomethylated H3-K27, conversion of monomethylated H3-K27 to dimethylated H3-K27, conversion of dimethylated H3-K27 to trimethylated H3-K27, or any combination thereof, including, for example, conversion of monomethylated H3-K27 to dimethylated H3-K27 and conversion of dimethylated H3-K27 to trimethylated H3-K27. As used herein, unmethylated H3-K27 refers to histone H3 with no methyl group covalently linked to the amino group of lysine 27. As used herein, monomethylated H3-K27 refers to histone H3 with a single methyl group covalently linked to the amino group of lysine 27. Monomethylated H3-K27 is also referred to herein as H3-K27me1. As used herein, dimethylated H3-K27 refers to histone H3 with two methyl groups covalently linked to the amino group of lysine 27. Dimethylated H3-K27 is also referred to herein as H3-K27me2. As used herein, trimethylated H3-K27 refers to histone H3 with three methyl groups covalently linked to the amino group of lysine 27. Trimethylated H3-K27 is also referred to herein as H3-K27me3.

[0072] Histone H3 is a 136 amino acid long protein, the sequence of which is known. See, for example, GenBank Accession No. CAB02546, the content of which is incorporated herein by reference. As disclosed further herein, in addition to full-length histone H3, peptide fragments of histone H3 comprising the lysine residue corresponding to K27 of full-length histone H3 can be used as substrate for EZH2 (and likewise for mutant forms of EZH2) to assess conversion of H3-K27m1 to H3-K27m2 and conversion of H3-K27m2 to H3-K27m3. In one embodiment, such peptide fragment corresponds to amino acid residues 21-44 of histone H3. Such peptide fragment has the amino acid sequence LATKAARKSAPATGGVKKPHRYRP (SEQ ID NO: 19).

[0073] The method of the present invention involves administering to a subject expressing a mutant EZH2 a therapeutically effective amount of an inhibitor of EZH2, wherein the inhibitor inhibits histone methyltransferase activity of EZH2, thereby inhibiting conversion of H3-K27 to trimethylated H3-K27 in the subject and thus treating or alleviating a symptom of cancer or disorders associated with abnormal histone methylation levels in the subject.

[0074] A subject expressing a mutant EZH2 of the present invention refers to a subject having a detectable amount of a mutant EZH2 polypeptide. Preferably the mutant EZH2 polypeptide comprises one or more mutations in its substrate pocket domain as defined in SEQ ID NO: 6. Exemplary mutation includes a substitution at amino acid position 677, 687, 674, 685, or 641 of SEQ ID NO: 1, such as, but is not limited to a substitution of glycine (G) for the wild type residue alanine (A) at amino acid position 677 of SEQ ID NO: 1 (A677G); a substitution of valine (V) for the wild type residue alanine (A) at amino acid position 687 of SEQ ID NO: 1 (A687V); a substitution of methionine (M) for the wild type residue valine (V) at amino acid position 674 of SEQ ID NO: 1 (V674M); a substitution of histidine (H) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685H); a substitution of cysteine (C) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685C); a substitution of phenylalanine (F) for the wild type residue tyrosine (Y) at amino acid position 641 of

SEQ ID NO: 1 (Y641F); a substitution of histidine (H) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641H); a substitution of asparagine (N) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641N); a substitution of serine (S) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641S); or a substitution of cysteine (C) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641C). More preferably, the subject has a detectable amount of a mutant EZH2 polypeptide comprising a substitution of glycine (G) for the wild type residue alanine (A) at amino acid position 677 of SEQ ID NO: 1 (A677G); a substitution of valine (V) for the wild type residue alanine (A) at amino acid position 687 of SEQ ID NO: 1 (A687V); a substitution of histidine (H) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685H); or a substitution of cysteine (C) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685C).

[0075] Alternatively a subject expressing a mutant EZH2 of the present invention refers to a subject having a detectable amount of a nucleic acid sequence encoding a mutant EZH2 polypeptide. Preferably a nucleic acid sequence encoding a mutant EZH2 polypeptide comprises one or more mutations in its substrate pocket domain as defined in SEQ ID NO: 6. Exemplary mutation includes a substitution at amino acid position 677, 687, 674, 685, or 641 of SEQ ID NO: 1, such as, but is not limited to a substitution of glycine (G) for the wild type residue alanine (A) at amino acid position 677 of SEQ ID NO: 1 (A677G); a substitution of valine (V) for the wild type residue alanine (A) at amino acid position 687 of SEQ ID NO: 1 (A687V); a substitution of methionine (M) for the wild type residue valine (V) at amino acid position 674 of SEQ ID NO: 1 (V674M); a substitution of histidine (H) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685H); a substitution of cysteine (C) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685C); a substitution of phenylalanine (F) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641F); a substitution of histidine (H) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641H); a substitution of asparagine (N) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641N); a substitution of serine (S) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641S); or a substitution of cysteine (C) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641C). More preferably, the subject has a detectable amount of nucleic acid sequence encoding a mutant EZH2 polypeptide comprising a substitution of glycine (G) for the wild type residue alanine (A) at amino acid position 677 of SEQ ID NO: 1 (A677G); a substitution of valine (V) for the wild type residue alanine (A) at amino acid position 687 of SEQ ID NO: 1 (A687V); a substitution of histidine (H) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685H); or a substitution of cysteine (C) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685C).

Detection of EZH2 Mutants

[0076] A mutant EZH2 polypeptide can be detected using any suitable method. For example, a mutant EZH2 polypeptide can be detected using an antibody that binds specifically to the mutant EZH2 polypeptide or to a peptide fragment that

is characteristic of the mutant EZH2 polypeptide. A peptide fragment that is characteristic of the mutant EZH2 polypeptide may include, for example, a SET domain as provided in SEQ ID NO: 7, except for substitution of one or more residues in the substrate pocket domain as defined in SEQ ID NO: 6 by an amino acid residue other than the wild type residue. In another embodiment, a peptide fragment that is characteristic of the mutant EZH2 polypeptide may include, for example, a 10-113 amino acid fragment of the SET domain as provided in SEQ ID NO: 7, except for substitution of one or more residues in the substrate pocket domain as defined in SEQ ID NO: 6 by an amino acid residue other than the wild type residue, provided that the fragment includes the amino acid residue corresponding to a mutation of EZH2. An antibody is considered to bind specifically to the mutant EZH2 polypeptide or to a peptide fragment that is characteristic of the mutant EZH2 polypeptide if it binds to that mutant EZH2 polypeptide or peptide fragment thereof but not to the corresponding wild-type EZH2 polypeptide or peptide fragment thereof. In one embodiment, such antibody is considered to bind specifically to the mutant EZH2 polypeptide or to a peptide fragment that is characteristic of the mutant EZH2 polypeptide if it binds to that mutant EZH2 polypeptide or peptide fragment thereof with an affinity that is at least ca. 100-fold greater than for the corresponding wild-type EZH2 polypeptide or peptide fragment thereof. In one embodiment, such antibody is considered to bind specifically to the mutant EZH2 polypeptide or to a peptide fragment that is characteristic of the mutant EZH2 polypeptide if it binds to that mutant EZH2 polypeptide or peptide fragment thereof with an affinity that is at least ca. 1000-fold greater than for the corresponding wild-type EZH2 polypeptide or peptide fragment thereof. The antibody can be used, for example, in an enzyme-linked immunosorbent assay (ELISA) or Western blot assay. The antibody may be monoclonal, polyclonal, chimeric, or an antibody fragment. The step of detecting the reaction product may be carried out with any suitable immunoassay.

[0077] In one embodiment the antibody is a monoclonal antibody. A monoclonal antibody can be prepared according to conventional methods well known in the art. See, for example, Köhler and Milstein (1975) *Nature* 256 (5517):495-7.

[0078] As another example, a mutant EZH2 polypeptide can be detected using mass spectrometry (MS), e.g., electrospray ionization coupled with time-of-flight (ESI-TOF) or matrix-assisted laser desorption/ionization coupled with time-of-flight (MALDI-TOF). Such methods are well known in the art. The analysis will involve identification of one or more peptide fragments comprising the mutation of interest, for example, a peptide 12 to 24 amino acids long comprising a sequence spanning the amino acid corresponding to a mutation in wild-type EZH2.

[0079] A nucleic acid sequence encoding a mutant EZH2 polypeptide or a peptide fragment that is characteristic of the mutant EZH2 polypeptide can be detected using any suitable method. For example, a nucleic acid sequence encoding a mutant EZH2 polypeptide can be detected using whole-genome resequencing or target region resequencing (the latter also known as targeted resequencing) using suitably selected sources of DNA and polymerase chain reaction (PCR) primers in accordance with methods well known in the art. See, for example, Bentley (2006) *Curr Opin Genet Dev.* 16:545-52, and Li et al. (2009) *Genome Res* 19:1124-32. The method typically and generally entails the steps of genomic DNA

purification, PCR amplification to amplify the region of interest, cycle sequencing, sequencing reaction cleanup, capillary electrophoresis, and data analysis. High quality PCR primers to cover region of interest are designed using in silico primer design tools. Cycle sequencing is a simple method in which successive rounds of denaturation, annealing, and extension in a thermal cycler result in linear amplification of extension products. The products are typically terminated with a fluorescent tag that identifies the terminal nucleotide base as G, A, T, or C. Unincorporated dye terminators and salts that may compete for capillary electrophoretic injection are removed by washing. During capillary electrophoresis, the products of the cycle sequencing reaction migrate through capillaries filled with polymer. The negatively charged DNA fragments are separated by size as they move through the capillaries toward the positive electrode. After electrophoresis, data collection software creates a sample file of the raw data. Using downstream software applications, further data analysis is performed to translate the collected color data images into the corresponding nucleotide bases. Alternatively or in addition, the method may include the use of microarray-based targeted region genomic DNA capture and/or sequencing. Kits, reagents, and methods for selecting appropriate PCR primers and performing resequencing are commercially available, for example, from Applied Biosystems, Agilent, and NimbleGen (Roche Diagnostics GmbH). Methods such as these have been used to detect JAK2 and myeloproliferative leukemia gene (MPL) mutations and to diagnose polycythemia vera, essential thrombocythemia, and idiopathic myelofibrosis. For use in the instant invention, PCR primers may be selected so as to amplify, for example, at least a relevant portion of SEQ ID NO: 8 (above).

[0080] Alternatively or in addition, a nucleic acid sequence encoding a mutant EZH2 polypeptide may be detected using a Southern blot in accordance with methods well known in the art. In one embodiment a DNA sequence encoding a mutant EZH2 polypeptide is detected using nucleic acid hybridization performed under highly stringent conditions. A nucleic acid probe is selected such that its sequence is complementary to a target nucleic acid sequence that includes a codon for the mutant amino acid corresponding to a mutation of wild-type EZH2. Such mutation includes any mutation of EZH2, preferably one or more mutations in the substrate pocket domain as defined in SEQ ID NO: 6 of EZH2, such as, but are not limited to a substitution of glycine (G) for the wild type residue alanine (A) at amino acid position 677 of SEQ ID NO: 1 (A677G); a substitution of valine (V) for the wild type residue alanine (A) at amino acid position 687 of SEQ ID NO: 1 (A687V); a substitution of methionine (M) for the wild type residue valine (V) at amino acid position 674 of SEQ ID NO: 1 (V674M); a substitution of histidine (H) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685H); a substitution of cysteine (C) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685C); a substitution of phenylalanine (F) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641F); a substitution of histidine (H) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641H); a substitution of asparagine (N) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641N); a substitution of serine (S) for the wild type residue tyrosine (Y) at amino acid position 641 of

SEQ ID NO: 1 (Y641S); or a substitution of cysteine (C) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641C).

[0081] A sequence-specific probe is combined with a sample to be tested under highly stringent conditions. The term “highly stringent conditions” as used herein refers to parameters with which the art is familiar. Nucleic acid hybridization parameters may be found in references that compile such methods, e.g., J. Sambrook, et al., eds., *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989, or F. M. Ausubel, et al., eds., *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc., New York. More specifically, highly stringent conditions, as used herein, refers, for example, to hybridization at 65° C. in hybridization buffer (3.5×SSC, 0.02% Ficoll, 0.02% polyvinyl pyrrolidone, 0.02% bovine serum albumin (BSA), 2.5 mM NaH₂PO₄ (pH 7), 0.5% SDS, 2 mM EDTA). SSC is 0.15 M sodium chloride/0.015 M sodium citrate, pH 7; SDS is sodium dodecyl sulphate; and EDTA is ethylenediaminetetracetic acid. After hybridization, the membrane upon which the DNA is transferred is washed, for example, in 2×SSC at room temperature and then at 0.1–0.5×SSC/0.1×SDS at temperatures up to 68° C.

[0082] There are other conditions, reagents, and so forth that can be used, which result in a similar degree of stringency. The skilled artisan will be familiar with such conditions, and thus they are not given here. It will be understood, however, that the skilled artisan will be able to manipulate the conditions in a manner to permit the clear identification of EZH2-associated nucleic acids of the invention, including, in particular, nucleic acids encoding mutants of EZH2 (e.g., by using lower stringency conditions). The skilled artisan also is familiar with the methodology for screening cells and libraries for expression of such molecules, which then are routinely isolated, followed by isolation of the pertinent nucleic acid molecule and sequencing.

[0083] The subject is administered a therapeutically effective amount of an inhibitor of EZH2. As used herein, an inhibitor of EZH2 refers, generally, to a small molecule, i.e., a molecule of molecular weight less than about 1.5 kilodaltons (kDa), which is capable of interfering with the histone methyltransferase enzymatic activity of EZH2.

[0084] In one embodiment the inhibitor of EZH2 inhibits histone methyltransferase activity of wild-type EZH2. In one embodiment the inhibitor of EZH2 inhibits histone methyltransferase activity of the mutant EZH2. In one embodiment the inhibitor of EZH2 inhibits histone methyltransferase activity of wild-type EZH2 and histone methyltransferase activity of the mutant EZH2. In one embodiment the inhibitor of EZH2 selectively inhibits histone methyltransferase activity of the mutant EZH2.

[0085] As disclosed herein, certain mutants of EZH2 (such as Y641) are relatively poor catalysts for conversion of unmethylated H3-K27 to H3-K27me1 and yet unexpectedly effective catalysts for conversion of H3-K27me2 to H3-K27me3. Certain mutants of EZH2 (such as A687) prefer monomethyl H3-K27 substrate. In contrast, certain mutants of EZH2 (such as A677) show equal preference among unmethylated, monomethylated and dimethylated H3-K27. Conversely, wild-type EZH2 is a relatively effective catalyst for conversion of unmethylated H3-K27 to H3-K27me1 and yet unexpectedly ineffective catalyst for conversion of H3-K27me2 to H3-K27me3. This is important because

mono-, di- and tri-methylated states of H3-K27 exhibit different functions in transcriptional control. For example, H3-K27me1 is associated with active transcription of genes that are poised for transcription, while H3-K27me3 is associated with transcriptionally repressed genes or genes that are poised for transcription when H3-K4 trimethylation is in cis. Thus, selective inhibition of histone methyltransferase activity of the mutant of EZH2 affects selective inhibition of production of the different methylated forms of H3-K27, thereby modifying transcription associated with H3-K27 methylation levels.

[0086] An inhibitor of EZH2 “selectively inhibits” histone methyltransferase activity of the mutant EZH2 when it inhibits histone methyltransferase activity of the mutant EZH2 more effectively than it inhibits histone methyltransferase activity of wild-type EZH2. For example, in one embodiment the selective inhibitor has an IC₅₀ for the mutant EZH2 that is at least 40 percent lower than the IC₅₀ for wild-type EZH2. In one embodiment the selective inhibitor has an IC₅₀ for the mutant EZH2 that is at least 50 percent lower than the IC₅₀ for wild-type EZH2. In one embodiment the selective inhibitor has an IC₅₀ for the mutant EZH2 that is at least 60 percent lower than the IC₅₀ for wild-type EZH2. In one embodiment the selective inhibitor has an IC₅₀ for the mutant EZH2 that is at least 70 percent lower than the IC₅₀ for wild-type EZH2. In one embodiment the selective inhibitor has an IC₅₀ for the mutant EZH2 that is at least 80 percent lower than the IC₅₀ for wild-type EZH2. In one embodiment the selective inhibitor has an IC₅₀ for the mutant EZH2 that is at least 90 percent lower than the IC₅₀ for wild-type EZH2.

[0087] In one embodiment, the selective inhibitor of a mutant EZH2 exerts essentially no inhibitory effect on wild-type EZH2.

[0088] The inhibitor inhibits conversion of H3-K27me2 to H3-K27me3. In one embodiment the inhibitor is said to inhibit trimethylation of H3-K27. Since conversion of H3-K27me1 to H3-K27me2 precedes conversion of H3-K27me2 to H3-K27me3, an inhibitor of conversion of H3-K27me1 to H3-K27me2 naturally also inhibits conversion of H3-K27me2 to H3-K27me3, i.e., it inhibits trimethylation of H3-K27. It is also possible to inhibit conversion of H3-K27me2 to H3-K27me3 without inhibition of conversion of H3-K27me1 to H3-K27me2. Inhibition of this type would also result in inhibition of trimethylation of H3-K27, albeit without inhibition of dimethylation of H3-K27.


[0089] In one embodiment the inhibitor inhibits conversion of H3-K27me1 to H3-K27me2 and the conversion of H3-K27me2 to H3-K27me3. Such inhibitor may directly inhibit the conversion of H3-K27me1 to H3-K27me2 alone. Alternatively, such inhibitor may directly inhibit both the conversion of H3-K27me1 to H3-K27me2 and the conversion of H3-K27me2 to H3-K27me3.

[0090] The inhibitor inhibits histone methylase activity. Inhibition of histone methylase activity can be detected using any suitable method. The inhibition can be measured, for example, either in terms of rate of histone methylase activity or as product of histone methylase activity. Methods suitable for either of these readouts are included in the Examples below.

[0091] The inhibition is a measurable inhibition compared to a suitable negative control. In one embodiment, inhibition is at least 10 percent inhibition compared to a suitable negative control. That is, the rate of enzymatic activity or the amount of product with the inhibitor is less than or equal to 90


percent of the corresponding rate or amount made without the inhibitor. In various other embodiments, inhibition is at least 20, 25, 30, 40, 50, 60, 70, 75, 80, 90, or 95 percent inhibition compared to a suitable negative control. In one embodiment, inhibition is at least 99 percent inhibition compared to a suitable negative control. That is, the rate of enzymatic activity or the amount of product with the inhibitor is less than or equal to 1 percent of the corresponding rate or amount made without the inhibitor.

[0092] In one embodiment, the inhibitor is S-adenosyl-L-homocysteine (SAH). SAH has the structural formula



and is commercially available from a number of suppliers, including, for example, Sigma-Aldrich, St. Louis, Mo. SAH has been described as an inhibitor of transmethylation by S-adenosylmethionine-dependent methyltransferases.

[0093] In one embodiment, the inhibitor is Compound 75



or a pharmaceutically acceptable salt thereof.

[0094] In certain embodiments the invention comprises the step of performing an assay to detect a mutant of EZH2 in a sample from a subject. Assays of this type are described above. As used herein, a “sample from a subject” refers to any suitable sample containing cells or components of cells

obtained or derived from a subject. In one embodiment the sample includes cells suspected to express mutant EZH2, e.g., cancer cells. In one embodiment the sample is a blood sample. In one embodiment the sample is a biopsy sample obtained from, for example, a lymphatic tissue (e.g., lymph node) or bone marrow. In one embodiment the sample is a biopsy sample obtained from a tissue other than or in addition to a lymphatic tissue (e.g., lymph node) or bone marrow. For example, in one embodiment the sample is a biopsy from a cancer, e.g., a tumor composed of cancer cells. Cells in the sample can be isolated from other components of the sample. For example, peripheral blood mononuclear cells (PBMCs) can be isolated as a buffy coat from a blood sample that has been centrifuged in accordance with methods familiar to those of skill in the art.

[0095] When the result of the assay on a sample from a subject indicates that a mutant EZH2 is present in the sample, the subject is said to express mutant EZH2. Indeed, in one embodiment, when the result of the assay on a sample from a subject indicates that a mutant EZH2 is present in the sample, the subject is identified as a candidate for treatment with an inhibitor of EZH2, wherein the inhibitor selectively inhibits histone methyltransferase activity of the mutant EZH2.

[0096] When the result of the assay on a sample from a cancer indicates that a mutant EZH2 is present in the cancer, the cancer is said to express the mutant EZH2.

[0097] Similarly, when the result of the assay on a sample comprising cancer cells from a subject having a cancer indicates that a mutant EZH2 is present in the sample, the subject is said to express the mutant EZH2.

[0098] Detection of dimethylated H3-K27 or trimethylated H3-K27 can be accomplished using any suitable method in the art. In one embodiment, the methylation level is detected using antibodies specific for dimethylated H3-K27 or trimethylated H3-K27. For example, the isolated tissue is formalin fixed and embedded in paraffin blocks for long term preservation. The blocks can be used to prepare slides for immunohistochemical staining or fluorescent staining with antibodies against methylated H3-K27. Alternatively, whole cell lysates or histone extracts can be prepared from the isolated tissue sample and subsequently used for immunohistochemical staining, western blot analysis or fluorescent staining. In another embodiment the methylation level is detected using a polypeptide or an aptamer specific for dimethylated H3-K27 or trimethylated H3-K27. In another embodiment, the methylation level is detected using mass spectrometry (MS).

[0099] A control dimethylated H3-K27 or a control trimethylated H3-K27 can be established from a control sample, e.g., an adjacent non-tumor tissue isolated from the subject or a healthy tissue from a healthy subject. Alternatively, the control methylation level of H3-K27me2 or H3-K27me3 can be established by a pathologist with known methods in the art.

Screening Methods

[0100] An aspect of the invention is a method for identifying a test compound as an inhibitor of a mutant EZH2. In one embodiment the method includes combining an isolated mutant EZH2 with a histone substrate, a methyl group donor (such as S-adenosyl methionine (SAM)), and a test compound, wherein the histone substrate comprises a form of H3-K27 selected from the group consisting of unmethylated H3-K27, monomethylated H3-K27, dimethylated H3-K27, and any combination thereof; and performing an assay to

detect methylation of H3-K27 in the histone substrate, thereby identifying the test compound as an inhibitor of the mutant EZH2 when methylation of H3-K27 in the presence of the test compound is less than methylation of H3-K27 in the absence of the test compound. The assay to detect methylation of H3-K27 can be selected to measure the rate of methylation, the extent of methylation, or both the rate and extent of methylation.

[0101] The mutant EZH2 is isolated as a PRC2 complex or functional equivalent thereof. As used herein, the term “isolated” means substantially separated from other components with which the complex may be found as it occurs in nature. A compound can be isolated without necessarily being purified. In one embodiment the mutant of EZH2 is isolated as a complex of a mutant EZH2 together with EED and SUZ12. In another embodiment the mutant of EZH2 is isolated as a complex of a mutant EZH2 together with EED, SUZ12, and RbAp48. Under appropriate conditions, a PRC2 complex or functional equivalent thereof exhibits histone methyltransferase activity for H3-K27. In one embodiment the complex is composed of recombinantly expressed component polypeptides, e.g., EZH2, EED, SUZ12, with or without RbAp48.

[0102] The isolated mutant EZH2 is combined with a histone substrate. A histone substrate includes any suitable source of histone polypeptides or fragments thereof that can serve as substrate for EZH2. In one embodiment the histone substrate includes histones isolated from a subject. The histones can be isolated from cells of a subject using any suitable method; such methods are well known to persons skilled in the art and need not be further specified here. See, for example, Fang et al. (2004) *Methods Enzymol* 377:213-26. In accordance with the Examples below, in one embodiment the histone substrate is provided as nucleosomes. In accordance with the Examples below, in one embodiment the histone substrate is provided as avian (chicken) erythrocyte nucleosomes.

[0103] Histone substrate so provided may include an admixture of states of histone modification, including various states of H3-K27 methylation as judged by Western blotting with H3-K27 methylation state-specific antibodies. In one embodiment the histone substrate may be provided as purified full-length histone H3. Such purified full-length histone H3 may be provided as a homogeneous preparation in respect of states of H3-K27 methylation, or as an admixture of various states of H3-K27 methylation. Homogeneous preparations of isolated histone H3 in respect of states of H3-K27 methylation may be prepared in part by passage over an immunoaffinity column loaded with suitable H3-K27 methylation state-specific antibodies or by immunoprecipitation using magnetic beads coated with suitable H3-K27 methylation state-specific antibodies. Alternatively or in addition, the methylation state of H3-K27 can be characterized as part of performing the assay. For example, the starting material histone substrate might be characterized as containing 50 percent unmethylated H3-K27, 40 percent monomethylated H3-K27, 10 percent dimethylated H3-K27, and 0 percent trimethylated H3-K27.

[0104] In one embodiment the histone substrate includes a peptide library or a suitable peptide comprising one or more amino acid sequences related to histone H3, including, in particular, a sequence that encompasses H3-K27. For example, in one embodiment, the histone substrate is a peptide fragment that corresponds to amino acid residues 21-44 of histone H3. Such peptide fragment has the amino acid

sequence LATKAARKSAPATGGVKKPHRYRP (SEQ ID NO: 19). The peptide library or peptide can be prepared by peptide synthesis according to techniques well known in the art and optionally modified so as to incorporate any desired degree of methylation of lysine corresponding to H3-K27. As described in the Examples below, such peptides can also be modified to incorporate a label, such as biotin, useful in performing downstream assays. In one embodiment the label is appended to the amino (N)-terminus of the peptide(s). In one embodiment the label is appended to the carboxy (C)-terminus of the peptide(s).

[0105] H3-K27 methylation-specific antibodies are available from a variety of commercial sources, including, for example, Cell Signaling Technology (Danvers, Mass.) and Active Motif (Carlsbad, Calif.).

[0106] The isolated mutant EZH2 is combined with a test compound. As used herein, a “test compound” refers to a small organic molecule having a molecular weight of less than about 1.5 kDa. In one embodiment a test compound is a known compound. In one embodiment a test compound is a novel compound. In one embodiment, a test compound can be provided as part of a library of such compounds, wherein the library includes, for example, tens, hundreds, thousands, or even more compounds. A library of compounds may advantageously be screened in a high throughput screening assay, for example, using arrays of test compounds and robotic manipulation in accordance with general techniques well known in the art.

[0107] In certain embodiments a test compound is a compound that is a derivative of SAH or a derivative of Compound 75.

[0108] Detection of methylation of H3-K27 can be accomplished using any suitable method. In one embodiment, the source of donor methyl groups includes methyl groups that are labeled with a detectable label. The detectable label in one embodiment is an isotopic label, e.g., tritium. Other types of labels may include, for example, fluorescent labels.

[0109] Detection of formation of trimethylated H3-K27 can be accomplished using any suitable method. For example, detection of formation of trimethylated H3-K27 can be accomplished using an assay to detect incorporation of labeled methyl groups, such as described above, optionally combined with a chromatographic or other method to separate labeled products by size, e.g., polyacrylamide gel electrophoresis (PAGE), capillary electrophoresis (CE), or high pressure liquid chromatography (HPLC). Alternatively or in addition, detection of formation of trimethylated H3-K27 can be accomplished using antibodies that are specific for trimethylated H3-K27.

[0110] Detection of conversion of monomethylated H3-K27 to dimethylated H3-K27 can be accomplished using any suitable method. In one embodiment the conversion is measured using antibodies specific for monomethylated H3-K27 and dimethylated H3-K27. For example, starting amounts or concentrations of monomethylated H3-K27 and dimethylated H3-K27 may be determined using appropriate antibodies specific for monomethylated H3-K27 and dimethylated H3-K27. Following the combination of enzyme, substrate, methyl group donor, and test compound, resulting amounts or concentrations of monomethylated H3-K27 and dimethylated H3-K27 may then be determined using appropriate antibodies specific for monomethylated H3-K27 and dimethylated H3-K27. The beginning and resulting amounts or concentrations of monomethylated H3-K27 and dimethylated H3-K27 can then be compared.

Alternatively or in addition, beginning and resulting amounts or concentrations of monomethylated H3-K27 and dimethylated H3-K27 can then be compared to corresponding amounts of concentrations from a negative control. A negative control reaction, in which no test agent is included in the assay, can be run in parallel or as a historical control. Results of such control reaction can optionally be subtracted from corresponding results of the experimental reaction prior to or in conjunction with making the comparison mentioned above.

[0111] Because the dimethylated form of H3-K27 may be further methylated in the same assay, a reduction in the amount or concentration of monomethylated H3-K27 may not appear to correspond directly to an increase in dimethylated H3-K27. In this instance, it may be presumed, however, that a reduction in the amount or concentration of monomethylated H3-K27 is, by itself, reflective of conversion of monomethylated H3-K27 to dimethylated H3-K27.

[0112] Detection of conversion of dimethylated H3-K27 to trimethylated H3-K27 can be accomplished using any suitable method. In one embodiment the conversion is measured using antibodies specific for dimethylated H3-K27 and trimethylated H3-K27. For example, starting amounts or concentrations of dimethylated H3-K27 and trimethylated H3-K27 may be determined using appropriate antibodies specific for dimethylated H3-K27 and trimethylated H3-K27. Following the combination of enzyme, substrate, and test compound, resulting amounts or concentrations of dimethylated H3-K27 and trimethylated H3-K27 may then be determined using appropriate antibodies specific for dimethylated H3-K27 and trimethylated H3-K27. The beginning and resulting amounts or concentrations of dimethylated H3-K27 and trimethylated H3-K27 can then be compared. Alternatively or in addition, beginning and resulting amounts or concentrations of dimethylated H3-K27 and trimethylated H3-K27 can then be compared to corresponding amounts of concentrations from a negative control. A negative control reaction, in which no test agent is included in the assay, can be run in parallel or as a historical control. Results of such control reaction can optionally be subtracted from corresponding results of the experimental reaction prior to or in conjunction with making the comparison mentioned above.

[0113] A test agent is identified as an inhibitor of the mutant EZH2 when methylation of H3-K27 with the test compound is less than methylation of H3-K27 without the test compound. In one embodiment, a test agent is identified as an inhibitor of the mutant EZH2 when formation of trimethylated H3-K27 in the presence of the test compound is less than formation of trimethylated H3-K27 in the absence of the test compound.

[0114] An aspect of the invention is a method for identifying a selective inhibitor of a mutant EZH2. In one embodiment the method includes combining an isolated mutant EZH2 with a histone substrate, a methyl group donor (e.g., SAM), and a test compound, wherein the histone substrate comprises a form of H3-K27 selected from the group consisting of monomethylated H3-K27, dimethylated H3-K27, and a combination of monomethylated H3-K27 and dimethylated H3-K27, thereby forming a test mixture; combining an isolated wild-type EZH2 with a histone substrate, a methyl group donor (e.g., SAM), and a test compound, wherein the histone substrate comprises a form of H3-K27 selected from the group consisting of monomethylated H3-K27, dimethylated H3-K27, and a combination of monomethylated

H3-K27 and dimethylated H3-K27, thereby forming a control mixture; performing an assay to detect trimethylation of the histone substrate in each of the test mixture and the control mixture; calculating the ratio of (a) trimethylation with the mutant EZH2 and the test compound (M+) to (b) trimethylation with the mutant EZH2 without the test compound (M-); calculating the ratio of (c) trimethylation with wild-type EZH2 and the test compound (WT+) to (d) trimethylation with wild-type EZH2 without the test compound (WT-); comparing the ratio (a)/(b) with the ratio (c)/(d); and identifying the test compound as a selective inhibitor of the mutant EZH2 when the ratio (a)/(b) is less than the ratio (c)/(d). In one embodiment the method further includes taking into account a negative control without test compound for either or both of the test mixture and the control mixture.

[0115] The present invention also provides a previously unrecognized, surprising correlation of a patient's responsiveness to an EZH2 inhibitor. Accordingly, an aspect of the invention is a method for determining a responsiveness to an EZH2 inhibitor in a subject having a cancer or a precancerous condition. The method generally involves in detecting a mutation in the EZH2 substrate pocket domain as defined in SEQ ID NO: 6. For example, the mutation may be a substitution, a point mutation, a nonsense mutation, a miss sense mutation, a deletion, or an insertion described above. Preferred substitution amino acid mutation includes a substitution at amino acid position 677, 687, 674, 685, or 641 of SEQ ID NO: 1, such as, but is not limited to a substitution of glycine (G) for the wild type residue alanine (A) at amino acid position 677 of SEQ ID NO: 1 (A677G); a substitution of valine (V) for the wild type residue alanine (A) at amino acid position 687 of SEQ ID NO: 1 (A687V); a substitution of methionine (M) for the wild type residue valine (V) at amino acid position 674 of SEQ ID NO: 1 (V674M); a substitution of histidine (H) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685H); a substitution of cysteine (C) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685C); a substitution of phenylalanine (F) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641F); a substitution of histidine (H) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641H); a substitution of asparagine (N) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641N); a substitution of serine (S) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641S); or a substitution of cysteine (C) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641C).

[0116] In a preferred embodiment, the subject has cancer or a cancerous condition. For example, the cancer is lymphoma, leukemia or melanoma. Preferably, the lymphoma is non-Hodgkin lymphoma, follicular lymphoma or diffuse large B-cell lymphoma. Alternatively, the leukemia is chronic myelogenous leukemia (CML). The precancerous condition is myelodysplastic syndromes (MDS, formerly known as preleukemia).

[0117] In another preferred embodiment, the mutant EZH2 polypeptide or the nucleic acid sequence encoding the mutant EZH2 polypeptide of the present invention comprises one or more mutations in the substrate pocket domain of EZH2 as defined in SEQ ID NO: 6. More preferably, the mutant EZH2 polypeptide or the nucleic acid sequence encoding the mutant EZH2 polypeptide of the present invention comprises a sub-

stitution at amino acid position 677, 687, 674, 685, or 641 of SEQ ID NO: 1, such as, but is not limited to a substitution of glycine (G) for the wild type residue alanine (A) at amino acid position 677 of SEQ ID NO: 1 (A677G); a substitution of valine (V) for the wild type residue alanine (A) at amino acid position 687 of SEQ ID NO: 1 (A687V); a substitution of methionine (M) for the wild type residue valine (V) at amino acid position 674 of SEQ ID NO: 1 (V674M); a substitution of histidine (H) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685H); a substitution of cysteine (C) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685C); a substitution of phenylalanine (F) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641F); a substitution of histidine (H) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641H); a substitution of asparagine (N) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641N); a substitution of serine (S) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641S); or a substitution of cysteine (C) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641C).

Pharmaceutical Compositions

[0118] One or more EZH2 antagonists can be administered alone to a human patient or in pharmaceutical compositions where they are mixed with suitable carriers or excipient(s) at doses to treat or ameliorate a disease or condition as described herein. Mixtures of these EZH2 antagonists can also be administered to the patient as a simple mixture or in suitable formulated pharmaceutical compositions. For example, one aspect of the invention relates to pharmaceutical composition comprising a therapeutically effective dose of an EZH2 antagonist, or a pharmaceutically acceptable salt, hydrate, enantiomer or stereoisomer thereof; and a pharmaceutically acceptable diluent or carrier.

[0119] Techniques for formulation and administration of EZH2 antagonists may be found in references well known to one of ordinary skill in the art, such as Remington's "The Science and Practice of Pharmacy," 21st ed., Lippincott Williams & Wilkins 2005.

[0120] Suitable routes of administration may, for example, include oral, rectal, or intestinal administration; parenteral delivery, including intravenous, intramuscular, intraperitoneal, subcutaneous, or intramedullary injections, as well as intrathecal, direct intraventricular, or intraocular injections; topical delivery, including eyedrop and transdermal; and intranasal and other transmucosal delivery.

[0121] Alternatively, one may administer an EZH2 antagonist in a local rather than a systemic manner, for example, via injection of the EZH2 antagonist directly into an edematous site, often in a depot or sustained release formulation.

[0122] In one embodiment, an EZH2 antagonist is administered by direct injection into a tumor or lymph node.

[0123] Furthermore, one may administer an EZH2 antagonist in a targeted drug delivery system, for example, in a liposome coated with cancer cell-specific antibody.

[0124] The pharmaceutical compositions of the present invention may be manufactured, e.g., by conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping or lyophilizing processes.

[0125] Pharmaceutical compositions for use in accordance with the present invention thus may be formulated in a conventional manner using one or more physiologically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active EZH2 antagonists into preparations which can be used pharmaceutically. Proper formulation is dependent upon the route of administration chosen.

[0126] For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiological saline buffer. For transmucosal administration, penetrants are used in the formulation appropriate to the barrier to be permeated. Such penetrants are generally known in the art.

[0127] For oral administration, the EZH2 antagonists can be formulated readily by combining the active EZH2 antagonists with pharmaceutically acceptable carriers well known in the art. Such carriers enable the EZH2 antagonists of the invention to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated. Pharmaceutical preparations for oral use can be obtained by combining the active EZH2 antagonist with a solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients include fillers such as sugars, including lactose, sucrose, mannitol, or sorbitol; cellulose preparations such as, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropylmethyl-cellulose, sodium carboxymethylcellulose, and/or polyvinylpyrrolidone (PVP). If desired, disintegrating agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof such as sodium alginate.

[0128] Dragee cores are provided with suitable coatings. For this purpose, concentrated sugar solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for identification or to characterize different combinations of active EZH2 antagonist doses.

[0129] Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer, such as glycerol or sorbitol. The push-fit capsules can contain the active ingredients in admixture with filler such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active EZH2 antagonists may be dissolved or suspended in suitable liquids, such as fatty oils, liquid paraffin, or liquid polyethylene glycols. In addition, stabilizers may be added.

[0130] For buccal administration, the compositions may take the form of tablets or lozenges formulated in conventional manner.

[0131] For administration by inhalation, the EZH2 antagonists for use according to the present invention are conveniently delivered in the form of an aerosol spray presentation from pressurized packs or a nebuliser, with the use of a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, carbon dioxide or

other suitable gas. In the case of pressurized aerosol the dosage unit may be determined by providing a valve to deliver a metered amount. Capsules and cartridges of e.g., gelatin for use in an inhaler or insufflator may be formulated containing a powder mix of the EZH2 antagonist and a suitable powder base such as lactose or starch.

[0132] The EZH2 antagonists can be formulated for parenteral administration by injection, e.g., bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

[0133] Pharmaceutical formulations for parenteral administration include aqueous solutions of the active EZH2 antagonists in water-soluble form. Additionally, suspensions of the active EZH2 antagonists may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the EZH2 antagonists to allow for the preparation of highly concentrated solutions.

[0134] Alternatively, the active ingredient may be in powder form for reconstitution before use with a suitable vehicle, e.g., sterile pyrogen-free water.

[0135] The EZH2 antagonists may also be formulated in rectal compositions such as suppositories or retention enemas, e.g., containing conventional suppository bases, such as cocoa butter or other glycerides.

[0136] In addition to the formulations described previously, the EZH2 antagonists may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation (for example, subcutaneously or intramuscularly or by intramuscular injection). Thus, for example, the EZH2 antagonists may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives (for example, as a sparingly soluble salt).

[0137] Alternatively, other delivery systems for hydrophobic pharmaceutical EZH2 antagonists may be employed. Liposomes and emulsions are examples of delivery vehicles or carriers for hydrophobic drugs. Certain organic solvents such as dimethylsulfoxide also may be employed. Additionally, the EZH2 antagonists may be delivered using a sustained-release system, such as semi-permeable matrices of solid hydrophobic polymers containing the therapeutic agent. Various sustained-release materials have been established and are well known by those skilled in the art. Sustained-release capsules may, depending on their chemical nature, release the EZH2 antagonists for a few weeks up to over 100 days. Depending on the chemical nature and the biological stability of the therapeutic reagent, additional strategies for protein stabilization may be employed.

[0138] The pharmaceutical compositions may also comprise suitable solid or gel phase carriers or excipients. Examples of such carriers or excipients include but are not

limited to calcium carbonate, calcium phosphate, various sugars, starches, cellulose derivatives, gelatin, and polymers, such as polyethylene glycols.

Methods of Treatment

[0139] Provided herein are methods of treating, preventing or alleviating a symptom of conditions and diseases, such as cancers and precancerous conditions, the course of which can be influenced by modulating the methylation status of histones or other proteins, wherein said methylation status is mediated at least in part by the activity of EZH2. Modulation of the methylation status of histones can in turn influence the level of expression of target genes activated by methylation, and/or target genes suppressed by methylation.

[0140] For example, one aspect of the invention relates to a method for treating or alleviating a symptom of cancer or precancerous condition. The method comprises the step of administering to a subject having a cancer or a precancerous condition and expressing a mutant EZH2 a therapeutically effective amount of an inhibitor of EZH2. Preferably, a mutant EZH2 polypeptide or a nucleic acid sequence encoding a mutant EZH2 polypeptide comprises a mutation in its substrate pocket domain as defined in SEQ ID NO: 6. More preferably, a mutant EZH2 polypeptide or a nucleic acid sequence encoding a mutant EZH2 polypeptide comprises a substitution mutation at amino acid position 677, 687, 674, 685, or 641 of SEQ ID NO: 1, such as, but is not limited to a substitution of glycine (G) for the wild type residue alanine (A) at amino acid position 677 of SEQ ID NO: 1 (A677G); a substitution of valine (V) for the wild type residue alanine (A) at amino acid position 687 of SEQ ID NO: 1 (A687V); a substitution of methionine (M) for the wild type residue valine (V) at amino acid position 674 of SEQ ID NO: 1 (V674M); a substitution of histidine (H) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685H); a substitution of cysteine (C) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685C); a substitution of phenylalanine (F) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641F); a substitution of histidine (H) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641H); a substitution of asparagine (N) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641N); a substitution of serine (S) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641S); or a substitution of cysteine (C) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641C).

[0141] In one embodiment the inhibitor inhibits histone methyltransferase activity of the mutant EZH2. In one embodiment the inhibitor selectively inhibits histone methyltransferase activity of the mutant EZH2. Optionally, the cancer is a lymphoma. Preferably, the cancer is a non-Hodgkin lymphoma, a follicular lymphoma or a diffuse large B-cell lymphoma (DLBCL). Alternatively, the cancer is leukemia (such as CML) or melanoma. The precancerous condition includes, but is not limited to, myelodysplastic syndromes (MDS, formerly known as preleukemia).

[0142] Diseases such as cancers and neurological disease can be treated by administration of modulators of protein (e.g., histone) methylation, e.g., modulators of histone methyltransferase, or histone demethylase enzyme activity. Histone methylation has been reported to be involved in aberrant expression of certain genes in cancers, and in silencing of

neuronal genes in non-neuronal cells. Modulators described herein can be used to treat such diseases, i.e., to inhibit methylation of histones in affected cells.

[0143] Based at least on the fact that abnormal histone methylation has been found to be associated with certain cancers and precancerous conditions, a method for treating cancer or a precancerous condition with a mutant EZH2 in a subject comprises administering to the subject in need thereof a therapeutically effective amount of a compound that inhibits methylation or restores methylation to roughly its level in counterpart normal cells. In one embodiment a method for treating cancer or a precancerous condition in a subject comprises administering to the subject in need thereof a therapeutically effective amount of a compound that inhibits conversion of unmethylated H3-K27 to monomethylated H3-K27 (H3-K27me1). In one embodiment a method for treating cancer or a precancerous condition in a subject comprises administering to the subject in need thereof a therapeutically effective amount of a compound that inhibits conversion of monomethylated H3-K27 (H3-K27me1) to dimethylated H3-K27 (H3-K27me2). In one embodiment a method for treating cancer or a precancerous condition in a subject comprises administering to the subject in need thereof a therapeutically effective amount of a compound that inhibits conversion of H3-K27me2 to trimethylated H3-K27 (H3-K27me3). In one embodiment a method for treating cancer or a precancerous condition in a subject comprises administering to the subject in need thereof a therapeutically effective amount of a compound that inhibits both conversion of H3-K27me1 to H3-K27me2 and conversion of H3-K27me2 to H3-K27me3. It is important to note that disease-specific increase in methylation can occur at chromatin in key genomic loci in the absence of a global increase in cellular levels of histone or protein methylation. For example, it is possible for aberrant hypermethylation at key disease-relevant genes to occur against a backdrop of global histone or protein hypomethylation.

[0144] Modulators of methylation can be used for modulating cell proliferation, generally. For example, in some cases excessive proliferation may be reduced with agents that decrease methylation, whereas insufficient proliferation may be stimulated with agents that increase methylation. Accordingly, diseases that may be treated include hyperproliferative diseases, such as benign cell growth and malignant cell growth (cancer).

[0145] Exemplary cancers that may be treated include lymphomas, including non-Hodgkin lymphoma, follicular lymphoma (FL) and diffuse large B-cell lymphoma (DLBCL); melanoma; and leukemia, including CML. Exemplary precancerous condition includes myelodysplastic syndrome (MDS; formerly known as preleukemia).

[0146] Other cancers include Acute Lymphoblastic Leukemia; Acute Myeloid Leukemia; Adrenocortical Carcinoma; AIDS-Related Cancers; AIDS-Related Lymphoma; Anal Cancer; Astrocytoma, Childhood Cerebellar; Astrocytoma, Childhood Cerebral; Basal Cell Carcinoma, see Skin Cancer (non-Melanoma); Bile Duct Cancer, Extrahepatic; Bladder Cancer; Bone Cancer, osteosarcoma/Malignant Fibrous Histiocytoma; Brain Stem Glioma; Brain Tumor; Brain Tumor, Cerebellar Astrocytoma; Brain Tumor, Cerebral Astrocytoma/Malignant Glioma; Brain Tumor, Ependymoma; Brain Tumor, Medulloblastoma; Brain Tumor, Supratentorial Primitive Neuroectodermal Tumors; Brain Tumor, Visual Pathway and Hypothalamic Glioma; Breast Cancer; Bron-

chial Adenomas/Carcinoids; Burkitt's Lymphoma; Carcinoid Tumor; Carcinoid Tumor, Gastrointestinal; Carcinoma of Unknown Primary; Central Nervous System Lymphoma, Primary; Cerebellar Astrocytoma; Cervical Cancer; Childhood Cancers; Chronic Lymphocytic Leukemia; Chronic Myelogenous Leukemia; Chronic Myelogenous Leukemia, Hairy Cell; Chronic Myeloproliferative Disorders; Colon Cancer; Colorectal Cancer; Cutaneous T-Cell Lymphoma, see Mycosis Fungoides and Sezary Syndrome; Endometrial Cancer; Esophageal Cancer; Ewing's Family of Tumors; Extrahepatic Bile Duct Cancer; Eye Cancer, Intraocular Melanoma; Eye Cancer, Retinoblastoma; Gallbladder Cancer; Gastric (Stomach) Cancer; Gastrointestinal Carcinoid Tumor; Germ Cell Tumor, Extracranial; Germ Cell Tumor, Extragonadal; Germ Cell Tumor, Ovarian; Gestational Trophoblastic Tumor; Glioma; Glioma, Childhood Brain Stem; Glioma, Childhood Cerebral Astrocytoma; Glioma, Childhood Visual Pathway and Hypothalamic; Hairy Cell Leukemia; Head and Neck Cancer; Hepatocellular (Liver) Cancer, Adult (Primary); Hepatocellular (Liver) Cancer, Childhood (Primary); Hodgkin's Lymphoma; Hodgkin's Lymphoma During Pregnancy; Hypopharyngeal Cancer; Hypothalamic and Visual Pathway Glioma; Intraocular Melanoma; Islet Cell Carcinoma (Endocrine Pancreas); Kaposi's Sarcoma; Kidney (Renal Cell) Cancer; Kidney Cancer; Laryngeal Cancer; Leukemia; Lip and Oral Cavity Cancer; Liver Cancer, Adult (Primary); Liver Cancer, Childhood (Primary); Lung Cancer, Non-Small Cell; Lung Cancer, Small Cell; Lymphoma, Primary Central Nervous System; Macroglobulinemia, Waldenstrom's; Malignant Fibrous Histiocytoma of Bone/Osteosarcoma; Medulloblastoma; Melanoma; Merkel Cell Carcinoma; Mesothelioma; Mesothelioma, Adult Malignant; Metastatic Squamous Neck Cancer with Occult Primary; Multiple Endocrine Neoplasia Syndrome; Multiple Myeloma; Multiple Myeloma/Plasma Cell Neoplasm Mycosis Fungoides; Myelodysplastic Syndromes; Myelodysplastic/Myeloproliferative Diseases; Myeloid Leukemia, Adult Acute; Myeloid Leukemia, Childhood Acute; Myeloproliferative Disorders, Chronic; Nasal Cavity and Paranasal Sinus Cancer; Nasopharyngeal Cancer; Neuroblastoma; Non-Hodgkin's Lymphoma; Non-Hodgkin's Lymphoma During Pregnancy; Oral Cancer; Oral Cavity Cancer, Lip and; Oropharyngeal Cancer; Osteosarcoma/Malignant Fibrous Histiocytoma of Bone; Ovarian Cancer; Ovarian Epithelial Cancer; Ovarian Low Malignant Potential Tumor; Pancreatic Cancer; Pancreatic Cancer, Islet Cell; Paranasal Sinus and Nasal Cavity Cancer; Parathyroid Cancer; Penile Cancer; Pheochromocytoma; Pineoblastoma and Supratentorial Primitive Neuroectodermal Tumors; Pituitary Tumor; Plasma Cell Neoplasm/Multiple Myeloma; Pleuropulmonary Blastoma; Pregnancy and Breast Cancer; Prostate Cancer; Rectal Cancer; Retinoblastoma; Rhabdomyosarcoma; Salivary Gland Cancer; Sarcoma, Ewing's Family of Tumors; Sarcoma, Soft Tissue; Sarcoma, Uterine; Sezary Syndrome; Skin Cancer; Skin Cancer (non-Melanoma); Small Intestine Cancer; Soft Tissue Sarcoma; Squamous Cell Carcinoma, see Skin Cancer (non-Melanoma); Squamous Neck Cancer with Occult Primary, Metastatic; Stomach (Gastric) Cancer; Testicular Cancer; Thymoma; Thymoma and Thymic Carcinoma; Thyroid Cancer; Transitional Cell Cancer of the Renal Pelvis and Ureter; Trophoblastic Tumor, Gestational; Unknown Primary Site, Cancer of; Unusual Cancers of Childhood; Urethral Cancer; Uterine Cancer, Endometrial; Uterine Sarcoma; Vaginal Cancer; Visual Pathway and Hypo-

thalamic Glioma; Vulvar Cancer; Waldenstrom's Macroglobulinemia; Wilms' Tumor; and Women's Cancers.

[0147] Any other disease in which epigenetic methylation, which is mediated by EZH2, plays a role may be treatable or preventable using compounds and methods described herein.

[0148] For example, neurologic diseases that may be treated include epilepsy, schizophrenia, bipolar disorder or other psychological and/or psychiatric disorders, neuropathies, skeletal muscle atrophy, and neurodegenerative diseases, e.g., a neurodegenerative disease. Exemplary neurodegenerative diseases include: Alzheimer's, Amyotrophic Lateral Sclerosis (ALS), and Parkinson's disease. Another class of neurodegenerative diseases includes diseases caused at least in part by aggregation of poly-glutamine. Diseases of this class include: Huntington's Diseases, Spinalbulbar Muscular Atrophy (SBMA or Kennedy's Disease), Dentatorubropallidolusian Atrophy (DRPLA), Spinocerebellar Ataxia 1 (SCA1), Spinocerebellar Ataxia 2 (SCA2), Machado-Joseph Disease (MJD; SCA3), Spinocerebellar Ataxia 6 (SCA6), Spinocerebellar Ataxia 7 (SCAT), and Spinocerebellar Ataxia 12 (SCA12).

Combination Therapy

[0149] In one aspect of the invention, an EZH2 antagonist, or a pharmaceutically acceptable salt thereof, can be used in combination with another therapeutic agent to treat diseases such as cancer and/or neurological disorders. For example, the additional agent can be a therapeutic agent that is art-recognized as being useful to treat the disease or condition being treated by the compound of the present invention. The additional agent also can be an agent that imparts a beneficial attribute to the therapeutic composition (e.g., an agent that affects the viscosity of the composition).

[0150] The combination therapy contemplated by the invention includes, for example, administration of a compound of the invention, or a pharmaceutically acceptable salt thereof, and additional agent(s) in a single pharmaceutical formulation as well as administration of a compound of the invention, or a pharmaceutically acceptable salt thereof, and additional agent(s) in separate pharmaceutical formulations. In other words, co-administration shall mean the administration of at least two agents to a subject so as to provide the beneficial effects of the combination of both agents. For example, the agents may be administered simultaneously or sequentially over a period of time.

[0151] The agents set forth below are for illustrative purposes and not intended to be limiting. The combinations, which are part of this invention, can be the compounds of the present invention and at least one additional agent selected from the lists below. The combination can also include more than one additional agent, e.g., two or three additional agents if the combination is such that the formed composition can perform its intended function.

[0152] For example, one aspect of the invention relates to the use of an EZH2 antagonist in combination with another agent for the treatment of cancer and/or a neurological disorder. In one embodiment, an additional agent is an anticancer agent that is a compound that affects histone modifications, such as an HDAC inhibitor. In certain embodiments, an additional anticancer agent is selected from the group consisting of chemotherapeutics (such as 2CdA, 5-FU, 6-Mercaptopurine, 6-TG, Abraxane™, Accutane®, Actinomycin-D, Adriamycin®, Alimta®, all-trans retinoic acid, amethopterin, Ara-C, Azacitadine, BCNU, Blenoxane®, Camptosar®,

CeeNU®, Clofarabine, Clolar™, Cytosan®, daunorubicin hydrochloride, DaunoXome®, Dacogen®, DIC, Doxil®, Ellence®, Eloxatin®, Emcyt®, etoposide phosphate, Fludara®, FUDR®, Gemzar®, Gleevec®, hexamethylmelamine, Hycamtin®, Hydrea®, Idamycin®, Ifex®, ixabepilone, Ixempra®, L-asparaginase, Leukeran®, liposomal Ara-C, L-PAM, Lysodren, Matulane®, mithracin, Mitomycin-C, Myleran®, Navelbine®, Neutrexin®, nilotinib, Nipent®, Nitrogen Mustard, Novantrone®, Oncaspar®, Panretin®, Paraplatin®, Platinol®, prolifeprospan 20 with carmustine implant, Sandostatin®, Targretin®, Tasigna®, Taxotere®, Temodar®, TESPAs, Trisenox®, Valstar®, Velban®, Vidaza™, vincristine sulfate, VM 26, Xeloda® and Zanosar®); biologics (such as Alpha Interferon, *Bacillus Calmette-Guerin*, Bexxar®, Campath®, Ergamisol®, Erlotinib, Herceptin®, Interleukin-2, Iressa®, lenalidomide, Mylotarg®, Ontak®, Pegasys®, Revlimid®, Rituxan®, Tarceva™, Thalomid®, Velcade® and Zevalin™); small molecules (such as Tykerb®); corticosteroids (such as dexamethasone sodium phosphate, DeltaSone® and DeltaCortef®); hormonal therapies (such as Arimidex®, Aromasin®, Casodex®, Cytadren®, Eligard®, Eulexin®, Evista®, Faslodex®, Femara®, Halotestin®, Megace®, Nilandron®, Nolvadex®, Plenaxis™ and Zoladex®); and radiopharmaceuticals (such as Iodotope®, Metastron®, Phosphocol® and Samarium SM-153).

Dosage

[0153] As used herein, a “therapeutically effective amount” or “therapeutically effective dose” is an amount of an EZH2 antagonist or a combination of two or more such compounds, which inhibits, totally or partially, the progression of the condition or alleviates, at least partially, one or more symptoms of the condition. A therapeutically effective amount can also be an amount which is prophylactically effective. The amount which is therapeutically effective will depend upon the patient’s size and gender, the condition to be treated, the severity of the condition and the result sought. In one embodiment, a therapeutically effective dose refers to that amount of the EZH2 antagonists that result in amelioration of symptoms in a patient. For a given patient, a therapeutically effective amount may be determined by methods known to those of skill in the art.

[0154] Toxicity and therapeutic efficacy of EZH2 antagonists can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the maximum tolerated dose (MTD) and the ED₅₀ (effective dose for 50% maximal response). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio between MTD and ED₅₀. The data obtained from these cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. Dosage may also be guided by monitoring the EZH2 antagonist’s effect on pharmacodynamic markers of enzyme inhibition (e.g., histone methylation or target gene expression) in diseased or surrogate tissue. Cell culture or animal experiments can be used to determine the relationship between doses required for changes in pharmacodynamic markers and doses required for therapeutic efficacy can be determined in cell culture or animal experiments or early stage clinical trials. The dosage of such EZH2 antagonists lies preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. The exact formulation, route of administration and dosage can be cho-

sen by the individual physician in view of the patient’s condition. In the treatment of crises, the administration of an acute bolus or an infusion approaching the MTD may be required to obtain a rapid response.

[0155] Dosage amount and interval may be adjusted individually to provide plasma levels of the active moiety which are sufficient to maintain the methyltransferase modulating effects, or minimal effective concentration (MEC) for the required period of time to achieve therapeutic efficacy. The MEC will vary for each EZH2 antagonist but can be estimated from in vitro data and animal experiments. Dosages necessary to achieve the MEC will depend on individual characteristics and route of administration. However, high pressure liquid chromatography (HPLC) assays or bioassays can be used to determine plasma concentrations.


[0156] Dosage intervals can also be determined using the MEC value. In certain embodiments, EZH2 antagonists should be administered using a regimen which maintains plasma levels above the MEC for 10-90% of the time, preferably between 30-90% and most preferably between 50-90% until the desired amelioration of symptoms is achieved. In other embodiments, different MEC plasma levels will be maintained for differing amounts of time. In cases of local administration or selective uptake, the effective local concentration of the drug may not be related to plasma concentration.

[0157] One of skill in the art can select from a variety of administration regimens and the amount of EZH2 antagonist administered will, of course, be dependent on the subject being treated, on the subject’s weight, the severity of the affliction, the manner of administration and the judgment of the prescribing physician.

Compounds and Pharmaceutical Compositions

[0158] Aspects of the invention concern compounds which are useful according to the methods of the invention. These compounds are referred to herein as “inhibitors of EZH2” and, equivalently, “EZH2 antagonists”. The compounds can be presented as the compounds per se, pharmaceutically acceptable salts of the compounds, or as pharmaceutical compositions.

[0159] The compounds suitable for use in the method of this invention include compounds of Formula (I):



wherein,

[0160] V¹ is N or CR⁷,

[0161] V² is N or CR², provided when V¹ is N, V² is N,

[0162] X and Z are selected independently from the group consisting of hydrogen, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, unsubstituted or substituted (C₃-C₈)cycloalkyl, unsubstituted or substituted (C₃-C₈)cycloalkyl-(C₁-C₈)alkyl

or $-(C_2-C_8)$ alkenyl, unsubstituted or substituted (C_5-C_8) cycloalkenyl, unsubstituted or substituted (C_5-C_8) cycloalkenyl- (C_1-C_8) alkyl or $-(C_2-C_8)$ alkenyl, (C_6-C_{10}) bicycloalkyl, unsubstituted or substituted heterocycloalkyl, unsubstituted or substituted heterocycloalkyl- (C_1-C_8) alkyl or $-(C_2-C_8)$ alkenyl, unsubstituted or substituted aryl, unsubstituted or substituted aryl- (C_1-C_8) alkyl or $-(C_2-C_8)$ alkenyl, unsubstituted or substituted heteroaryl, unsubstituted or substituted heteroaryl- (C_1-C_8) alkyl or $-(C_2-C_8)$ alkenyl, halo, cyano, $-COR^a$, $-CO_2R^a$, $-CONR^aR^b$, $-CONR^aNR^aR^b$, $-SR^a$, $-SOR^a$, $-SO_2R^a$, $-SO_2NR^aR^b$, nitro, $-NR^aR^b$, $-NR^aC(O)R^b$, $-NR^aC(O)NR^aR^b$, $-NR^aC(O)OR^a$, $-NR^aSO_2R^b$, $-NR^aSO_2NR^aR^b$, $-NR^aNR^aR^b$, $-NR^aNR^aC(O)R^b$, $-NR^aNR^aC(O)NR^aR^b$, $-NR^aNR^aC(O)OR^a$, $-OR^a$, $-OC(O)R^a$, and $-OC(O)NR^aR^b$;

[0163] Y is H or halo;

[0164] R^1 is (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, unsubstituted or substituted (C_3-C_8) cycloalkyl, unsubstituted or substituted (C_3-C_8) cycloalkyl- (C_1-C_8) alkyl or $-(C_2-C_8)$ alkenyl, unsubstituted or substituted (C_5-C_8) cycloalkenyl, unsubstituted or substituted (C_5-C_8) cycloalkenyl- (C_1-C_8) alkyl or $-(C_2-C_8)$ alkenyl, unsubstituted or substituted (C_6-C_{10}) bicycloalkyl, unsubstituted or substituted heterocycloalkyl or $-(C_2-C_8)$ alkenyl, unsubstituted or substituted heterocycloalkyl- (C_1-C_8) alkyl, unsubstituted or substituted aryl, unsubstituted or substituted aryl- (C_1-C_8) alkyl or $-(C_2-C_8)$ alkenyl, unsubstituted or substituted heteroaryl, unsubstituted or substituted heteroaryl- (C_1-C_8) alkyl or $-(C_2-C_8)$ alkenyl, $-COR^a$, $-CO_2R^a$, $-CONR^aR^b$, $-CONR^aNR^aR^b$;

[0165] R^2 is hydrogen, (C_1-C_8) alkyl, trifluoromethyl, alkoxy, or halo, in which said (C_1-C_8) alkyl is optionally substituted with one to two groups selected from amino and (C_1-C_3) alkylamino;

[0166] R^7 is hydrogen, (C_1-C_3) alkyl, or alkoxy;

[0167] R^3 is hydrogen, (C_1-C_8) alkyl, cyano, trifluoromethyl, $-NR^aR^b$, or halo;

[0168] R^6 is selected from the group consisting of hydrogen, halo, (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, unsubstituted or substituted (C_3-C_8) cycloalkyl, unsubstituted or substituted (C_3-C_8) cycloalkyl- (C_1-C_8) alkyl, unsubstituted or substituted (C_5-C_8) cycloalkenyl, unsubstituted or substituted (C_5-C_8) cycloalkenyl- (C_1-C_8) alkyl, (C_6-C_{10}) bicycloalkyl, unsubstituted or substituted heterocycloalkyl, unsubstituted or substituted heterocycloalkyl- (C_1-C_8) alkyl, unsubstituted or substituted aryl, unsubstituted or substituted aryl- (C_1-C_8) alkyl, unsubstituted or substituted heteroaryl, unsubstituted or substituted heteroaryl- (C_1-C_8) alkyl, cyano, $-COR^a$, $-CO_2R^a$, $-CONR^aR^b$, $-CONR^aNR^aR^b$, $-SR^a$, $-SOR^a$, $-SO_2R^a$, $-SO_2NR^aR^b$, nitro, $-NR^aR^b$, $-NR^aC(O)R^b$, $-NR^aC(O)NR^aR^b$, $-NR^aC(O)OR^a$, $-NR^aSO_2R^b$, $-NR^aSO_2NR^aR^b$, $-NR^aNR^aR^b$, $-NR^aNR^aC(O)R^b$, $-NR^aNR^aC(O)NR^aR^b$, $-NR^aNR^aC(O)OR^a$, $-OR^a$, $-OC(O)R^a$, $-OC(O)NR^aR^b$;

[0169] wherein any (C_1-C_8) alkyl, (C_2-C_8) alkenyl, (C_2-C_8) alkynyl, cycloalkyl, cycloalkenyl, bicycloalkyl, heterocycloalkyl, aryl, or heteroaryl group is optionally substituted by 1, 2 or 3 groups independently selected from the group consisting of $-O(C_1-C_6)alkyl(R^c)_{1-2}$, $-S(C_1-C_6)alkyl(R^c)_{1-2}$, $-(C_1-C_6)alkyl(R^c)_{1-2}$, $-(C_1-C_8)alkyl-heterocycloalkyl$, $(C_3-C_8)cycloalkyl-heterocycloalkyl$, halo, $(C_1-C_6)alkyl$, $(C_3-C_8)cycloalkyl$, $(C_5-C_8)cycloalkenyl$, $(C_1-C_6)haloalkyl$, cyano, $-COR^a$, $-CO_2R^a$, $-CONR^aR^b$, $-SR^a$, $-SOR^a$, $-SO_2R^a$, $-SO_2NR^aR^b$, nitro, $-NR^aR^b$, $-NR^aC(O)R^b$, $-NR^aC(O)NR^aR^b$, $-NR^aC(O)OR^a$, $-NR^aSO_2R^b$, $-NR^aSO_2NR^aR^b$, $-OR^a$, $-OC(O)R^a$, $OC(O)NR^aR^b$, heterocycloalkyl, aryl, heteroaryl, aryl- $(C_1-C_4)alkyl$, and heteroaryl- $(C_1-C_4)alkyl$;

[0170] wherein any aryl or heteroaryl moiety of said aryl, heteroaryl, aryl- $(C_1-C_4)alkyl$, or heteroaryl- $(C_1-C_4)alkyl$ is optionally substituted by 1, 2 or 3 groups independently selected from the group consisting of halo, $(C_1-C_6)alkyl$, $(C_3-C_8)cycloalkyl$, $(C_5-C_8)cycloalkenyl$, $(C_1-C_6)haloalkyl$, cyano, $-COR^a$, $-CO_2R^a$, $-CONR^aR^b$, $-SR^a$, $-SOR^a$, $-SO_2R^a$, $-SO_2NR^aR^b$, nitro, $-NR^aR^b$, $-NR^aC(O)R^b$, $-NR^aC(O)NR^aR^b$, $-NR^aC(O)OR^a$, $-NR^aSO_2R^b$, $-NR^aSO_2NR^aR^b$, $-OR^a$, $-OC(O)R^a$, and $-OC(O)NR^aR^b$;

[0171] R^a and R^b are each independently hydrogen, $(C_1-C_8)alkyl$, $(C_2-C_8)alkenyl$, $(C_2-C_8)alkynyl$, $(C_3-C_8)cycloalkyl$, $(C_5-C_8)cycloalkenyl$, $(C_6-C_{10})bicycloalkyl$, heterocycloalkyl, aryl, or heteroaryl, wherein said $(C_1-C_8)alkyl$, $(C_2-C_8)alkenyl$, $(C_2-C_8)alkynyl$, cycloalkyl, cycloalkenyl, bicycloalkyl, heterocycloalkyl, aryl or heteroaryl group is optionally substituted by 1, 2 or 3 groups independently selected from halo, hydroxyl, $(C_1-C_4)alkoxy$, amino, $(C_1-C_4)alkylamino$, $((C_1-C_4)alkyl)((C_1-C_4)alkyl)amino$, $-CO_2H$, $-CO_2(C_1-C_4)alkyl$, $-CONH_2$, $-CONH(C_1-C_4)alkyl$, $-CON((C_1-C_4)alkyl)((C_1-C_4)alkyl)$, $-SO_2(C_1-C_4)alkyl$, $-SO_2NH_2$, $-SO_2NH(C_1-C_4)alkyl$, and $-SO_2N((C_1-C_4)alkyl)((C_1-C_4)alkyl)$;

[0172] or R^a and R^b taken together with the nitrogen to which they are attached represent a 5-8 membered saturated or unsaturated ring, optionally containing an additional heteroatom selected from oxygen, nitrogen, and sulfur, wherein said ring is optionally substituted by 1, 2 or 3 groups independently selected from $(C_1-C_4)alkyl$, $(C_1-C_4)haloalkyl$, amino, $(C_1-C_4)alkylamino$, $((C_1-C_4)alkyl)((C_1-C_4)alkyl)amino$, hydroxyl, oxo, $(C_1-C_4)alkoxy$, and $(C_1-C_4)alkoxy(C_1-C_4)alkyl$, wherein said ring is optionally fused to a $(C_3-C_8)cycloalkyl$, heterocycloalkyl, aryl, or heteroaryl ring;

[0173] or R^a and R^b taken together with the nitrogen to which they are attached represent a 6- to 10-membered bridged bicyclic ring system optionally fused to a $(C_3-C_8)cycloalkyl$, heterocycloalkyl, aryl, or heteroaryl ring;

[0174] each R^c is independently $(C_1-C_4)alkylamino$, $-NR^aSO_2R^b$, $-SOR^a$, $-SO_2R^a$, $-NR^aC(O)OR^a$, $-NR^aR^b$, or $-CO_2R^a$;

[0175] or a salt thereof.

[0176] Subgroups of the compounds encompassed by the general structure of Formula (I) are represented as follows:

[0177] Subgroup A of Formula (I)

[0178] X and Z are selected from the group consisting of $(C_1-C_8)alkyl$, $(C_3-C_8)cycloalkyl$, heterocycloalkyl, aryl, heteroaryl, $-NR^aR^b$, and $-OR^a$;

[0179] Y is H or F;

[0180] R^1 is selected from the group consisting of $(C_1-C_8)alkyl$, $(C_3-C_8)cycloalkyl$, heterocycloalkyl, aryl, and heteroaryl;

[0181] R^2 is hydrogen, $(C_1-C_8)alkyl$, trifluoromethyl, alkoxy, or halo, in which said $(C_1-C_8)alkyl$ is optionally substituted with one to two groups selected from amino and $(C_1-C_3)alkylamino$;

[0182] R^7 is hydrogen, $(C_1-C_3)alkyl$, or alkoxy;

[0183] R^3 is selected from the group consisting of hydrogen, $(C_1-C_8)alkyl$, cyano, trifluoromethyl, $-NR^aR^b$, and halo;

[0184] R^6 is selected from the group consisting of hydrogen, halo, cyano, trifluoromethyl, amino, $(C_1-C_8)alkyl$, $(C_3-C_8)cycloalkyl$, aryl, heteroaryl, acylamino, $(C_2-C_8)alkynyl$, arylalkynyl, heteroarylalkynyl; $-SO_2R^a$; $-SO_2NR^aR^b$ and $-NR^aSO_2R^b$;


[0185] wherein any $(C_1-C_8)alkyl$, $(C_3-C_8)cycloalkyl$, $(C_2-C_8)alkynyl$, arylalkynyl, heteroarylalkynyl group is optionally substituted by 1, 2 or 3 groups independently

selected from $-\text{O}(\text{C}_1\text{-C}_6)\text{alkyl}(\text{R}^a)_{1-2}$, $-\text{S}(\text{C}_1\text{-C}_6)\text{alkyl}(\text{R}^a)_{1-2}$, $-(\text{C}_1\text{-C}_6)\text{alkyl}(\text{R}^a)_{1-2}$, $-(\text{C}_1\text{-C}_8)\text{alkyl}$ -heterocycloalkyl, $(\text{C}_3\text{-C}_8)\text{cycloalkyl}$ -heterocycloalkyl, halo, $(\text{C}_1\text{-C}_6)\text{alkyl}$, $(\text{C}_3\text{-C}_8)\text{cycloalkyl}$, $(\text{C}_5\text{-C}_8)\text{cycloalkenyl}$, $(\text{C}_1\text{-C}_6)\text{haloalkyl}$, cyano, $-\text{COR}^a$, $-\text{CO}_2\text{R}^a$, $-\text{CONR}^a\text{R}^b$, $-\text{SR}^a$, $-\text{SOR}^a$, $-\text{SO}_2\text{R}^a$, $-\text{SO}_2\text{NR}^a\text{R}^b$, nitro, $-\text{NR}^a\text{R}^b$, $-\text{NR}^a\text{C}(\text{O})\text{R}^b$, $-\text{NR}^a\text{C}(\text{O})\text{NR}^a\text{R}^b$, $-\text{NR}^a\text{C}(\text{O})\text{OR}^a$, $-\text{NR}^a\text{SO}_2\text{R}^b$, $-\text{NR}^a\text{SO}_2\text{NR}^a\text{R}^b$, $-\text{OR}^a$, $-\text{OC}(\text{O})\text{R}^a$, $-\text{OC}(\text{O})\text{NR}^a\text{R}^b$, heterocycloalkyl, aryl, heteroaryl, aryl($\text{C}_1\text{-C}_4$)alkyl, and heteroaryl($\text{C}_1\text{-C}_4$)alkyl;

[0186] R^a and R^b are each independently hydrogen, $(\text{C}_1\text{-C}_8)\text{alkyl}$, $(\text{C}_2\text{-C}_8)\text{alkenyl}$, $(\text{C}_2\text{-C}_8)\text{alkynyl}$, $(\text{C}_3\text{-C}_8)\text{cycloalkyl}$, $(\text{C}_5\text{-C}_8)\text{cycloalkenyl}$, $(\text{C}_6\text{-C}_{10})\text{bicycloalkyl}$, heterocycloalkyl, aryl, or heteroaryl, wherein said $(\text{C}_1\text{-C}_8)\text{alkyl}$, $(\text{C}_2\text{-C}_8)\text{alkenyl}$, $(\text{C}_2\text{-C}_8)\text{alkynyl}$, cycloalkyl, cycloalkenyl, bicycloalkyl, heterocycloalkyl, aryl or heteroaryl group is optionally substituted by 1, 2 or 3 groups independently selected from halo, hydroxyl, $(\text{C}_1\text{-C}_4)\text{alkoxy}$, amino, $(\text{C}_1\text{-C}_4)\text{alkylamino}$, $((\text{C}_1\text{-C}_4)\text{alkyl})((\text{C}_1\text{-C}_4)\text{alkyl})\text{amino}$, $-\text{CO}_2\text{H}$, $-\text{CO}_2(\text{C}_1\text{-C}_4)\text{alkyl}$, $-\text{CONH}_2$, $-\text{CONH}(\text{C}_1\text{-C}_4)\text{alkyl}$, $-\text{CON}((\text{C}_1\text{-C}_4)\text{alkyl})((\text{C}_1\text{-C}_4)\text{alkyl})$, $-\text{SO}_2(\text{C}_1\text{-C}_4)\text{alkyl}$, $-\text{SO}_2\text{NH}_2$, $-\text{SO}_2\text{NH}(\text{C}_1\text{-C}_4)\text{alkyl}$, and $-\text{SO}_2\text{N}((\text{C}_1\text{-C}_4)\text{alkyl})((\text{C}_1\text{-C}_4)\text{alkyl})$;

[0187] or R^a and R^b taken together with the nitrogen to which they are attached represent a 5-8 membered saturated or unsaturated ring, optionally containing an additional heteroatom selected from oxygen, nitrogen, and sulfur, wherein said ring is optionally substituted by 1, 2 or 3 groups independently selected from $(\text{C}_1\text{-C}_4)\text{alkyl}$, $(\text{C}_1\text{-C}_4)\text{haloalkyl}$, amino, $(\text{C}_1\text{-C}_4)\text{alkylamino}$, $((\text{C}_1\text{-C}_4)\text{alkyl})((\text{C}_1\text{-C}_4)\text{alkyl})\text{amino}$, hydroxyl, oxo, $(\text{C}_1\text{-C}_4)\text{alkoxy}$, and $(\text{C}_1\text{-C}_4)\text{alkoxy}(\text{C}_1\text{-C}_4)\text{alkyl}$, wherein said ring is optionally fused to a $(\text{C}_3\text{-C}_8)\text{cycloalkyl}$, heterocycloalkyl, aryl, or heteroaryl ring;


[0188] or R^a and R^b taken together with the nitrogen to which they are attached represent a 6- to 10-membered bridged bicyclic ring system optionally fused to a $(\text{C}_3\text{-C}_8)\text{cycloalkyl}$, heterocycloalkyl, aryl, or heteroaryl ring. An aryl or heteroaryl group in this particular subgroup A is selected independently from the group consisting of furan, thiophene, pyrrole, oxazole, thiazole, imidazole, pyrazole, oxadiazole, thiadiazole, triazole, tetrazole, benzofuran, benzothiophene, benzoxazole, benzothiazole, phenyl, pyridine, pyridazine, pyrimidine, pyrazine, triazine, tetrazine, quinoline, cinnoline, quinoxaline, and naphthyridine or another aryl or heteroaryl group as follows:



(1)

wherein in (1),


[0189] A is O, NH, or S; B is CH or N, and C is hydrogen or $\text{C}_1\text{-C}_8$ alkyl; or



(2)

wherein in (2),


[0190] D is N or C optionally substituted by hydrogen or $\text{C}_1\text{-C}_8$ alkyl; or



(3)

wherein in (3),


[0191] E is NH or CH_2 ; F is O or CO; and G is NH or CH_2 ; or



(4)

wherein in (4),

[0192] J is O, S or CO; or



(5)


wherein in (5),

[0193] Q is CH or N;

[0194] M is CH or N; and

[0195] L/(5) is hydrogen, halo, amino, cyano, $(\text{C}_1\text{-C}_8)\text{alkyl}$, $(\text{C}_3\text{-C}_8)\text{cycloalkyl}$, $-\text{COR}^a$, $-\text{CO}_2\text{R}^a$, $-\text{CONR}^a\text{R}^b$, $-\text{CONR}^a\text{NR}^a\text{R}^b$, $-\text{SO}_2\text{R}^a$, $-\text{SO}_2\text{NR}^a\text{R}^b$, $-\text{NR}^a\text{R}^b$, $-\text{NR}^a\text{C}(\text{O})\text{R}^b$, $-\text{NR}^a\text{SO}_2\text{R}^b$, $-\text{NR}^a\text{SO}_2\text{NR}^a\text{R}^b$, $-\text{NR}^a\text{NR}^a\text{R}^b$, $-\text{NR}^a\text{NR}^a\text{C}(\text{O})\text{R}^b$, $-\text{NR}^a\text{NR}^a\text{C}(\text{O})\text{NR}^a\text{R}^b$, or $-\text{OR}^a$,


[0196] wherein any $(\text{C}_1\text{-C}_8)\text{alkyl}$ or $(\text{C}_3\text{-C}_8)\text{cycloalkyl}$ group is optionally substituted by 1, 2 or 3 groups independently selected from $(\text{C}_1\text{-C}_6)\text{alkyl}$, $(\text{C}_3\text{-C}_8)\text{cycloalkyl}$, $(\text{C}_5\text{-C}_8)\text{cycloalkenyl}$, $(\text{C}_1\text{-C}_6)\text{haloalkyl}$, cyano, $-\text{COR}^a$, $-\text{CO}_2\text{R}^a$, $-\text{CONR}^a\text{R}^b$, $-\text{SR}^a$, $-\text{SOR}^a$, $-\text{SO}_2\text{R}^a$, $-\text{SO}_2\text{NR}^a\text{R}^b$, nitro, $-\text{NR}^a\text{R}^b$, $-\text{NR}^a\text{C}(\text{O})\text{R}^b$, $-\text{NR}^a\text{C}(\text{O})\text{NR}^a\text{R}^b$, $-\text{NR}^a\text{C}(\text{O})\text{OR}^a$, $-\text{NR}^a\text{SO}_2\text{R}^b$, $-\text{NR}^a\text{SO}_2\text{NR}^a\text{R}^b$, $-\text{OR}^a$, $-\text{OC}(\text{O})\text{R}^a$, and $-\text{OC}(\text{O})\text{NR}^a\text{R}^b$; wherein R^a and R^b are defined as above; or



(6)

wherein in (6),

[0197] L/(6) is NH or CH_2 ; or




(7)

wherein in 7,

[0198] M/(7) is hydrogen, halo, amino, cyano, (C₁-C₈) alkyl, (C₃-C₈)cycloalkyl, heterocycloalkyl, —COR^a, —CO₂R^a, —CONR^aR^b, —CONR^aNR^aR^b, —SO₂R^a, —SO₂NR^aR^b, —NR^aR^b, —NR^aC(O)R^b, —NR^aSO₂R^b, —NR^aSO₂NR^aR^b, —NR^aNR^aR^b, —NR^aNR^aC(O)R^b, —NR^aNR^aC(O)NR^aR^b, or —OR^a,


[0199] wherein any (C₁-C₈)alkyl, (C₃-C₈)cycloalkyl, or heterocycloalkyl group is optionally substituted by 1, 2 or 3 groups independently selected from (C₁-C₆)alkyl, (C₃-C₈)cycloalkyl, (C₅-C₈)cycloalkenyl, (C₁-C₆)haloalkyl, cyano, —COR^a, —CO₂R^a, —CONR^aR^b, —SR^a, —SOR^a, —SO₂R^a, —SO₂NR^aR^b, nitro, —NR^aR^b, —NR^aC(O)R^b, —NR^aC(O)NR^aR^b, —NR^aC(O)OR^a, —NR^aSO₂R^b, —NR^aSO₂NR^aR^b, —OR^a, —OC(O)R^a, and —OC(O)NR^aR^b; wherein R^a and R^b are defined as above; or



(8)

wherein in (8),

[0200] P is CH₂, NH, O, or S; Q/(8) is CH or N; and n is 0-2; or



(9)

wherein in (9),

[0201] S/(9) and T/(9) is C, or S/(9) is C and T/(9) is N, or S/(9) is N and T/(9) is C;

[0202] R is hydrogen, amino, methyl, trifluoromethyl, or halo;

[0203] U is hydrogen, halo, amino, cyano, nitro, trifluoromethyl, (C₁-C₈)alkyl, (C₃-C₈)cycloalkyl, —COR^a, —CO₂R^a, —CONR^aR^b, —SO₂R^a, —SO₂NR^aR^b, —NR^aR^b, —NR^aC(O)R^b, —NR^aSO₂R^b, —NR^aSO₂NR^aR^b, —NR^aNR^aR^b, —NR^aNR^aC(O)R^b, —OR^a, or 4-(1H-pyrazol-4-yl),

[0204] wherein any (C₁-C₈)alkyl or (C₃-C₈)cycloalkyl group is optionally substituted by 1, 2 or 3 groups independently selected from (C₁-C₆)alkyl, (C₃-C₈)cycloalkyl, (C₅-C₈)cycloalkenyl, (C₁-C₆)haloalkyl, cyano, —COR^a, —CO₂R^a, —CONR^aR^b, —SR^a, —SOR^a, —SO₂R^a, —SO₂NR^aR^b, nitro, —NR^aR^b, —NR^aC(O)R^b, —NR^aC(O)NR^aR^b, —NR^aC(O)OR^a, —NR^aSO₂R^b, —NR^aSO₂NR^aR^b, —OR^a, —OC(O)R^a, and —OC(O)NR^aR^b; wherein R^a and R^b are defined as above.

[0205] Subgroup B of Formula (I)

[0206] X and Z are selected independently from the group consisting of (C₁-C₈)alkyl, (C₃-C₈)cycloalkyl, heterocycloalkyl, aryl, heteroaryl, —NR^aR^b, and —OR^a;

[0207] Y is H;

[0208] R¹ is (C₁-C₈)alkyl, (C₃-C₈)cycloalkyl, or heterocycloalkyl;

[0209] R² is hydrogen, (C₁-C₃)alkyl, or halo, in which said (C₁-C₃)alkyl is optionally substituted with one to two groups selected from amino and (C₁-C₃)alkylamino;

[0210] R⁷ is hydrogen, (C₁-C₃)alkyl, or alkoxy;

[0211] R³ is hydrogen, (C₁-C₈)alkyl or halo;

[0212] R⁶ is hydrogen, halo, cyano, trifluoromethyl, amino, (C₁-C₈)alkyl, (C₃-C₈)cycloalkyl, aryl, heteroaryl, acylamino; (C₂-C₈)alkynyl, arylalkynyl, heteroarylalkynyl, —SO₂R^a, —SO₂NR^aR^b, or


—NR^aSO₂R^b;

[0213] wherein any (C₁-C₈)alkyl, (C₃-C₈)cycloalkyl, (C₂-C₈)alkynyl, arylalkynyl, or heteroarylalkynyl group is optionally substituted by 1, 2 or 3 groups independently selected from halo, (C₁-C₆)alkyl, (C₃-C₈)cycloalkyl, (C₅-C₈)cycloalkenyl, (C₁-C₆)haloalkyl, cyano, —COR^a, —CO₂R^a, —CONR^aR^b, —SOR^a, —SO₂R^a, —SO₂NR^aR^b, nitro, —NR^aR^b, —NR^aC(O)R^b, —NR^aC(O)NR^aR^b, —NR^aC(O)OR^a, —NR^aSO₂R^b, —NR^aSO₂NR^aR^b, —OR^a, —OC(O)R^a, —OC(O)NR^aR^b, heterocycloalkyl, aryl, heteroaryl, aryl(C₁-C₄)alkyl, and heteroaryl(C₁-C₄)alkyl;

[0214] R^a and R^b are each independently hydrogen, (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, (C₃-C₈)cycloalkyl, (C₅-C₈)cycloalkenyl, (C₆-C₁₀)bicycloalkyl, heterocycloalkyl, aryl, or heteroaryl, wherein said (C₁-C₈)alkyl, (C₂-C₈)alkenyl, (C₂-C₈)alkynyl, cycloalkyl, cycloalkenyl, bicycloalkyl, heterocycloalkyl, aryl or heteroaryl group is optionally substituted by 1, 2 or 3 groups independently selected from halo, hydroxyl, (C₁-C₄)alkoxy, amino, (C₁-C₄)alkylamino, ((C₁-C₄)alkyl)((C₁-C₄)alkyl)amino, —CO₂H, —CO₂(C₁-C₄)alkyl, —CONH₂, —CONH(C₁-C₄)alkyl, —CON((C₁-C₄)alkyl)((C₁-C₄)alkyl), —SO₂(C₁-C₄)alkyl, —SO₂NH₂, —SO₂NH(C₁-C₄)alkyl, and —SO₂N((C₁-C₄)alkyl)((C₁-C₄)alkyl);

[0215] or R^a and R^b taken together with the nitrogen to which they are attached represent a 5-8 membered saturated or unsaturated ring, optionally containing an additional heteroatom selected from oxygen, nitrogen, and sulfur, wherein said ring is optionally substituted by 1, 2 or 3 groups independently selected from (C₁-C₄)alkyl, (C₁-C₄)haloalkyl, amino, (C₁-C₄)alkylamino, ((C₁-C₄)alkyl)((C₁-C₄)alkyl) amino, hydroxyl, oxo, (C₁-C₄)alkoxy, and (C₁-C₄)alkoxy (C₁-C₄)alkyl, wherein said ring is optionally fused to a (C₃-C₈)cycloalkyl, heterocycloalkyl, aryl, or heteroaryl ring;


[0216] or R^a and R^b taken together with the nitrogen to which they are attached represent a 6- to 10-membered bridged bicyclic ring system optionally fused to a (C₃-C₈) cycloalkyl, heterocycloalkyl, aryl, or heteroaryl ring. Aryl and heteroaryl in this definition are selected from the group consisting of furan, thiophene, pyrrole, oxazole, thiazole, imidazole, pyrazole, oxadiazole, thiadiazole, triazole, tetrazole, benzofuran, benzothiophene, benzoxazole, benzothiazole, phenyl, pyridine, pyridazine, pyrimidine, pyrazine, triazine, tetrazine, quinoline, cinnoline, quinazoline, quinoxaline, and naphthyridine or a compound of another aryl or heteroaryl group as follows:



(1)

wherein in (1),


[0217] A is O, NH, or S; B is CH or N, and C is hydrogen or C₁-C₈ alkyl; or



(2)

wherein in (2),


[0218] D is N or C optionally substituted by hydrogen or C₁-C₈ alkyl; or



(3)

wherein in (3),

[0219] E is NH or CH₂; F is O or CO; and G is NH or CH₂; or




(4)

wherein

in (4),

[0220] J is O, S or CO; or



(5)

wherein in (5),


[0221] Q is CH or N;

[0222] M is CH or N; and

[0223] L/(5) is hydrogen, halo, amino, cyano, (C₁-C₈)alkyl, (C₃-C₈)cycloalkyl, —COR^a, —CO₂R^a, —CONR^aR^b, —CONR^aNR^aR^b, —SO₂R^a, —SO₂NR^aR^b, —NR^aR^b, —NR^aC(O)R^b, —NR^aSO₂R^b, —NR^aSO₂NR^aR^b, —NR^aNR^aR^b, —NR^aNR^aC(O)R^b, —NR^aNR^aC(O)NR^aR^b, or —OR^a,

[0224] wherein any (C₁-C₈)alkyl, (C₃-C₈)cycloalkyl, group is optionally substituted by 1, 2 or 3 groups independently selected from (C₁-C₆)alkyl, (C₃-C₈)cycloalkyl, (C₅-C₈)cycloalkenyl, (C₁-C₆)haloalkyl, cyano, —COR^a, —CO₂R^a, —CONR^aR^b, —SR^a, —SOR^a, —SO₂R^a, —SO₂NR^aR^b, nitro, —NR^aR^b, —NR^aC(O)R^b, —NR^aC(O)NR^aR^b, —NR^aC(O)OR^a, —NR^aSO₂R^b, —NR^aSO₂NR^aR^b, —OR^a, —OC(O)R^a, and —OC(O)NR^aR^b,


wherein R^a and R^b are defined as above; or



(6)

wherein in (6),

[0225] L/(6) is NH or CH₂; or




(7)

wherein in (7),

[0226] M/(7) is hydrogen, halo, amino, cyano, (C₁-C₈)alkyl, (C₃-C₈)cycloalkyl, heterocycloalkyl, —COR^a, —CO₂R^a, —CONR^aR^b, —CONR^aNR^aR^b, —SO₂R^a, —SO₂NR^aR^b, —NR^aR^b, —NR^aC(O)R^b, —NR^aSO₂R^b, —NR^aSO₂NR^aR^b, —NR^aNR^aR^b, —NR^aNR^aC(O)R^b, —NR^aNR^aC(O)NR^aR^b, or —OR^a,


[0227] wherein any (C₁-C₈)alkyl, (C₃-C₈)cycloalkyl, heterocycloalkyl group is optionally substituted by 1, 2 or 3 groups independently selected from (C₁-C₆)alkyl, (C₃-C₈)cycloalkyl, (C₅-C₈)cycloalkenyl, (C₁-C₆)haloalkyl, cyano, —COR^a, —CO₂R^a, —CONR^aR^b, —SR^a, —SOR^a, —SO₂R^a, —SO₂NR^aR^b, nitro, —NR^aR^b, —NR^aC(O)R^b, —NR^aC(O)NR^aR^b, —NR^aC(O)OR^a, —NR^aSO₂R^b, —NR^aSO₂NR^aR^b, —OR^a, —OC(O)R^a, —OC(O)NR^aR^b; wherein R^a and R^b are defined as above; or



(8)

wherein in (8),

[0228] P is CH₂, NH, O, or S; Q/(8) is CH or N; and n is 0-2; or



(9)

wherein in (9),

[0229] S/(9) and T/(9) is C, or S/(9) is C and T/(9) is N, or S/(9) is N and T/(9) is C;

[0230] R is hydrogen, amino, methyl, trifluoromethyl, halo;

[0231] U is hydrogen, halo, amino, cyano, nitro, trifluoromethyl, (C₁-C₈)alkyl, (C₃-C₈)cycloalkyl, —COR^a, —CO₂R^a, —CONR^aR^b, —SO₂R^a, —SO₂NR^aR^b, —NR^aR^b, —NR^aC(O)R^b, —NR^aSO₂R^b, —NR^aSO₂NR^aR^b, —NR^aNR^aR^b, —NR^aNR^aC(O)R^b, —OR^a, or 4-(1H-pyrazol-4-yl),

[0232] wherein any (C₁-C₈)alkyl, or (C₃-C₈)cycloalkyl group is optionally substituted by 1, 2 or 3 groups independently selected from (C₁-C₆)alkyl, (C₃-C₈)cycloalkyl, (C₅-C₈)cycloalkenyl, (C₁-C₆)haloalkyl, cyano, —COR^a, —CO₂R^a, —CONR^aR^b, —SOR^a, —SO₂R^a, —SO₂NR^aR^b, nitro, —NR^aR^b, —NR^aC(O)

R^b , $—NR^aC(O)NR^aR^b$, $—NR^aC(O)OR^a$, $—NR^aSO_2R^b$, $—NR^aSO_2NR^aR^b$, $—OR^a$, $—OC(O)R^a$, and $—OC(O)NR^aR^b$, wherein R^a and R^b are defined as above.

[0233] Subgroup C of Formula (I)

[0234] X is methyl, ethyl, n-propyl, isopropyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, trifluoromethyl, tetrahydropyran, hydroxymethyl, methoxymethyl, or benzyl;

[0235] Y is H;

[0236] Z is methyl, ethyl, n-propyl, isopropyl, trifluoromethyl, or benzyl;

[0237] R^1 is isopropyl, tert-butyl, cyclobutyl, cyclopentyl, cyclohexyl, (1-methylethyl)cyclopropyl, 1,1-dioxo-tetrahydrothiophene-3-yl, 1-Me-piperidin-4-yl, tetrahydrofuran-3-yl, tetrahydropyran-4-yl, N,N-dimethyl-1-propanaminyl, benzyl, or 4-pyridyl;


[0238] R^2 is hydrogen, (C_1-C_3) alkyl, or halo, in which said (C_1-C_3) alkyl is optionally substituted with one to two groups selected from amino and (C_1-C_3) alkylamino;

[0239] R^7 is hydrogen, (C_1-C_3) alkyl, or alkoxy;

[0240] R^3 is H, methyl, or Br; and

[0241] R^6 is methyl, bis(1,1-dimethylethyl), bis(1-methylethyl), cyclopropyl, propyl, dimethylamino, ethylamino, (2-hydroxyethyl)amino, 2-propen-1-ylamino, 1-piperazinyl, 1-piperidinyl, 4-morpholinyl, 4-piperidinylamino, tetrahydro-2H-pyran-4-ylamino, phenylamino, (phenylmethyl)amino, (4-pyridinylmethyl)amino, [2-(2-pyridinylamino)ethyl]amino, 2-(dimethylamino)ethyl]amino, 4-pyridinylamino, 4-(aminocarbonyl)phenyl]amino, 3-hydroxy-3-methyl-1-butyn-1-yl, 4-pyridinylethynyl, phenylethynyl, 2-furanyl, 3-thienyl; 1H-pyrazol-4-yl, 1H-pyrazol-5-yl, 1H-indazol-6-yl, 3-methyl-1H-indazol-5-yl, 1H-1,2,3-benzotriazol-5-yl, 2-oxo-2,3-dihydro-1H-benzimidazol-5-yl, 2-oxo-2,3-dihydro-1H-indol-5-yl, 2-oxo-2,3-dihydro-1H-indol-6-yl, 2,1,3-benzoxadiazol-5-yl, 2-amino-6-quinazolinyl, 2,4-dioxo-1,2,3,4-tetrahydro-5-pyrimidinyl, 2-amino-5-pyrimidinyl, 7-oxo-1,5,6,7-tetrahydro-1,8-naphthyridin-3-yl, phenyl, 2-methylphenyl, 2-nitrophenyl, 2-phenylethyl, 3-aminophenyl, 4-aminophenyl, 4-chlorophenyl, 4-fluorophenyl, 4-(methyloxy)phenyl, 3-(acetylamino)phenyl, 4-(acetylamino)phenyl, 4-(aminocarbonyl)phenyl, 4-(1H-pyrazol-4-yl)phenyl, 4-(aminosulfonyl)phenyl, 4-(methylsulfonyl)phenyl, 4-[(dimethylamino)sulfonyl]phenyl, 4-[(methylamino)carbonyl]phenyl, 4-[(methylamino)sulfonyl]phenyl, 4-[(methylsulfonyl)amino]phenyl, 3-pyridinyl, 4-pyridinyl, 2-(4-morpholinyl)-4-pyridinyl, 2-amino-4-pyridinyl, 5-(methyloxy)-3-pyridinyl, 5-(methylsulfonyl)-3-pyridinyl, 5-[(cyclopropylsulfonyl)amino]-6-(methyloxy)-3-pyridinyl, 5-[(phenylsulfonyl)amino]-3-pyridinyl, 6-(4-methyl-1-piperazinyl)-3-pyridinyl, 6-(4-morpholinyl)-3-pyridinyl, 6-(acetylamino)-3-pyridinyl, 6-(dimethylamino)-3-pyridinyl, 6-(methyloxy)-3-pyridinyl, 6-[(methylamino)carbonyl]-3-pyridinyl, 6-Rmethylamino)sulfonyl]-3-pyridinyl, 6-methyl-3-pyridinyl, or 4-pyridinyl. (See, e.g., WO 2011/140325; WO 2011/140324; and WO 2012/005805, each of which is incorporated by reference in its entirety.)

[0242] The compounds suitable for use in the method of this invention also include compounds of Formula (II):



wherein

[0243] X_1 is N or CR_{11} ;

[0244] X_2 is N or CR_{13} ;

[0245] Z_1 is NR_7R_8 , OR_{S1} , SR_{S1} , or $CR_7R_8R_{14}$;

[0246] each of R_1 , R_5 , R_9 , and R_{10} , independently, is H or C_1-C_6 alkyl optionally substituted with one or more substituents selected from the group consisting of halo, hydroxyl, $COOH$, $C(O)O-C_1-C_6$ alkyl, cyano, C_1-C_6 alkoxy, amino, mono- C_1-C_6 alkylamino, di- C_1-C_6 alkylamino, C_3-C_8 cycloalkyl, C_6-C_{10} aryl, 4 to 12-membered heterocycloalkyl, and 5- or 6-membered heteroaryl;

[0247] each of R_2 , R_3 , and R_4 , independently, is $-Q_1-T_1$, in which Q_1 is a bond or C_1-C_3 alkyl linker optionally substituted with halo, cyano, hydroxyl or C_1-C_6 alkoxy, and T_1 is H, halo, hydroxyl, $COOH$, cyano, or R_{S1} , in which R_{S1} is C_1-C_3 alkyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl, C_1-C_6 alkoxy, $C(O)O-C_1-C_6$ alkyl, C_3-C_8 cycloalkyl, C_6-C_{10} aryl, amino, mono- C_1-C_6 alkylamino, di- C_1-C_6 alkylamino, 4 to 12-membered heterocycloalkyl, or 5- or 6-membered heteroaryl, and R_{S1} is optionally substituted with one or more substituents selected from the group consisting of halo, hydroxyl, oxo, $COOH$, $C(O)O-C_1-C_6$ alkyl, cyano, C_1-C_6 alkoxy, amino, mono- C_1-C_6 alkylamino, di- C_1-C_6 alkylamino, C_3-C_8 cycloalkyl, C_6-C_{10} aryl, 4 to 12-membered heterocycloalkyl, and 5- or 6-membered heteroaryl;

[0248] R_6 is C_6-C_{10} aryl or 5- or 6-membered heteroaryl, each of which is optionally substituted with one or more $-Q_2-T_2$, wherein Q_2 is a bond or C_1-C_3 alkyl linker optionally substituted with halo, cyano, hydroxyl or C_1-C_6 alkoxy, and T_2 is H, halo, cyano, $—OR_a$, $—NR_aR_b$, $—(NR_aR_bR_c)^+A^-$, $—C(O)R_a$, $—C(O)OR_a$, $—C(O)NR_aR_b$, $—NR_bC(O)R_a$, $—NR_bC(O)OR_a$, $—S(O)_2R_a$, $—S(O)_2NR_aR_b$, or R_{S2} , in which each of R_a , R_b , and R_c , independently, is H or R_{S3} , A^- is a pharmaceutically acceptable anion, each of R_{S2} and R_{S3} , independently, is C_1-C_6 alkyl, C_3-C_8 cycloalkyl, C_6-C_{10} aryl, 4 to 12-membered heterocycloalkyl, or 5- or 6-membered heteroaryl, or R_a and R_b , together with the N atom to which they are attached, form a 4 to 12-membered heterocycloalkyl ring having 0 or 1 additional heteroatom, and each of R_{S2} , R_{S3} , and the 4 to 12-membered heterocycloalkyl ring formed by R_a and R_b , is optionally substituted with one or more $-Q_3-T_3$, wherein Q_3 is a bond or C_1-C_3 alkyl linker each optionally substituted with halo, cyano, hydroxyl or C_1-C_6 alkoxy, and T_3 is selected from the group consisting of halo, cyano, C_1-C_6 alkyl, C_3-C_8 cycloalkyl, C_6-C_{10} aryl, 4 to

12-membered heterocycloalkyl, 5- or 6-membered heteroaryl, OR_d, COOR_d, —S(O)₂R_d, —NR_dR_e, and —C(O)NR_dR_e, each of R_d and R_e independently being H or C₁-C₆ alkyl, or —Q₃-T₃ is oxo; or any two neighboring —Q₂-T₂, together with the atoms to which they are attached form a 5- or 6-membered ring optionally containing 1-4 heteroatoms selected from N, O and S and optionally substituted with one or more substituents selected from the group consisting of halo, hydroxyl, COOH, C(O)O—C₁-C₆ alkyl, cyano, C₁-C₆ alkoxy, amino, mono-C₁-C₆ alkylamino, di-C₁-C₆ alkylamino, C₃-C₈ cycloalkyl, C₆-C₁₀ aryl, 4 to 12-membered heterocycloalkyl, and 5- or 6-membered heteroaryl; provided that —Q₂-T₂ is not H;


[0249] R₇ is —Q₄-T₄, in which Q₄ is a bond, C₁-C₄ alkyl linker, or C₂-C₄ alkenyl linker, each linker optionally substituted with halo, cyano, hydroxyl or C₁-C₆ alkoxy, and T₄ is H, halo, cyano, NR_fR_g, —OR_f, —C(O)R_f, —C(O)OR_f, —C(O)NR_fR_g, —C(O)NR_fOR_g, —NR_fC(O)R_g, —S(O)₂R_f, or R_{S4}, in which each of R_f and R_g, independently is H or R_{S5}, each of R_{S4} and R_{S5}, independently is C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₆-C₁₀ aryl, 4 to 12-membered heterocycloalkyl, or 5- or 6-membered heteroaryl, and each of R_{S4} and R_{S5} is optionally substituted with one or more —Q₅-T₅, wherein Q is a bond, C(O), C(O)NR_k, NR_kC(O), S(O)₂, or C₁-C₃ alkyl linker, R_k being H or C₁-C₆ alkyl, and T₅ is H, halo, C₁-C₆ alkyl, hydroxyl, cyano, C₁-C₆ alkoxy, amino, mono-C₁-C₆ alkylamino, di-C₁-C₆ alkylamino, C₃-C₈ cycloalkyl, C₆-C₁₀ aryl, 4 to 12-membered heterocycloalkyl, 5- or 6-membered heteroaryl, or S(O)_qR_q in which q is 0, 1, or 2 and R_q is C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₆-C₁₀ aryl, 4 to 12-membered heterocycloalkyl, or 5- or 6-membered heteroaryl, and T₅ is optionally substituted with one or more substituents selected from the group consisting of halo, C₁-C₆ alkyl, hydroxyl, cyano, C₁-C₆ alkoxy, amino, mono-C₁-C₆ alkylamino, di-C₁-C₆ alkylamino, C₃-C₈ cycloalkyl, C₆-C₁₀ aryl, 4 to 12-membered heterocycloalkyl, and 5- or 6-membered heteroaryl except when T₅ is H, halo, hydroxyl, or cyano; or —Q₅-T₅ is oxo; provided that R₇ is not H;

[0250] each of R₈, R₁₁, R₁₂, and R₁₃, independently, is H, halo, hydroxyl, COOH, cyano, R_{S6}, OR_{S6}, or COOR_{S6}, in which R_{S6} is C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, 4 to 12-membered heterocycloalkyl, amino, mono-C₁-C₆ alkylamino, or di-C₁-C₆ alkylamino, and R_{S6} is optionally substituted with one or more substituents selected from the group consisting of halo, hydroxyl, COOH, C(O)O—C₁-C₆ alkyl, cyano, C₁-C₆ alkoxy, amino, mono-C₁-C₆ alkylamino, and di-C₁-C₆ alkylamino; or R₇ and R₈, together with the N atom to which they are attached, form a 4 to 11-membered heterocycloalkyl ring having 0 to 2 additional heteroatoms, or R₇ and R₈, together with the C atom to which they are attached, form C₃-C₈ cycloalkyl or a 4 to 11-membered heterocycloalkyl ring having 1 to 3 heteroatoms, and each of the 4 to 11-membered heterocycloalkyl rings or C₃-C₈ cycloalkyl formed by R₇ and R₈ is optionally substituted with one or more —Q₆-T₆, wherein Q₆ is a bond, C(O), C(O)NR_m, NR_mC(O), S(O)₂, or C₁-C₃ alkyl linker, R_m being H or C₁-C₆ alkyl, and T₆ is H, halo, C₁-C₆ alkyl, hydroxyl, cyano, C₁-C₆ alkoxy, amino, mono-C₁-C₆ alkylamino, di-C₁-C₆ alkylamino, C₃-C₈ cycloalkyl, C₆-C₁₀ aryl, 4 to 12-membered heterocycloalkyl, 5- or 6-membered heteroaryl, or S(O)_pR_p in which p is 0, 1, or 2 and R_p is C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₆-C₁₀ aryl, 4 to 12-membered heterocycloalkyl, or 5- or



6-membered heteroaryl, and T₆ is optionally substituted with one or more substituents selected from the group consisting of halo, C₁-C₆ alkyl, hydroxyl, cyano, C₁-C₆ alkoxy, amino, mono-C₁-C₆ alkylamino, di-C₁-C₆ alkylamino, C₃-C₈ cycloalkyl, C₆-C₁₀ aryl, 4 to 12-membered heterocycloalkyl, and 5- or 6-membered heteroaryl except when T₆ is H, halo, hydroxyl, or cyano; or —Q₆-T₆ is oxo; and


[0251] R₁₄ is absent, H, or C₁-C₆ alkyl optionally substituted with one or more substituents selected from the group consisting of halo, hydroxyl, COOH, C(O)O—C₁-C₆ alkyl, cyano, C₁-C₆ alkoxy, amino, mono-C₁-C₆ alkylamino, di-C₁-C₆ alkylamino, C₃-C₈ cycloalkyl, C₆-C₁₀ aryl, 4 to 12-membered heterocycloalkyl, and 5- or 6-membered heteroaryl.

[0252] One subset of the compounds of Formula (II) includes those of Formula (IIa):



[0253] Another subset of the compounds of Formula (II) includes those of Formula (IIb), (IIc), or (IId):





[0254] The compounds of Formulae (II), (IIa), (IIc), and (IIId) can include one or more of the following features:

- [0255] For example, X_1 is CR_{11} and X_2 is CR_{13} .
 [0256] For example, X_1 is CR_{11} and X_2 is N.
 [0257] For example, X_1 is N and X_2 is CR_{13} .
 [0258] For example, X_1 is N and X_2 is N.
 [0259] For example, Z_1 is NR_7R_8 .
 [0260] For example, Z_1 is $CR_7R_8R_{14}$.
 [0261] For example, Z_1 is OR_7 .
 [0262] For example, Z_1 is SR_7 .
 [0263] For example, R_6 is phenyl substituted with one or more $-Q_2-T_2$.
 [0264] For example, R_6 is 5 to 6-membered heteroaryl containing 1-3 additional heteroatoms selected from N, O, and S and optionally substituted with one or more $-Q_2-T_2$.
 [0265] For example, R_6 is pyridinyl, pyrazolyl, pyrimidinyl, quinolinyl, tetrazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, furyl, or thienyl, each of which is optionally substituted with one or more $-Q_2-T_2$.
 [0266] For example, Q_2 is a bond.
 [0267] For example, Q_2 is an unsubstituted C_1-C_3 alkyl linker.
 [0268] For example, T_2 is C_1-C_6 alkyl or C_6-C_{10} aryl, each optionally substituted with one or more $-Q_3-T_3$.
 [0269] For example, T_2 is an unsubstituted substituted straight chain C_1-C_6 or branched C_3-C_6 alkyl, including but not limited to, methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl, t-butyl, n-pentyl, s-pentyl and n-hexyl.
 [0270] For example, T_2 is phenyl.
 [0271] For example, T_2 is halo (e.g., fluorine, chlorine, bromine, and iodine).
 [0272] For example, T_2 is 4 to 7-membered heterocycloalkyl (e.g., azetidiny, oxetanyl, thietanyl, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, triazolidinyl, tetrahydropyranyl, piperidinyl, 1,2,3,6-tetrahydropyridinyl, piperazinyl, tetrahydro-2H-pyranyl, 3,6-dihydro-2H-pyranyl, tetrahydro-2H-thiopyran, morpholinyl, 1,4-diazepanyl, 1,4-oxazepanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 2,5-diazabicyclo[2.2.1]heptanyl, and the like) optionally substituted with one or more $-Q_3-T_3$.
 [0273] For example, T_2 is OR_a , NR_aR_b , $-(NR_aR_bR_c)^+A^-$, $-C(O)R_a$, $-C(O)OR_a$, $-C(O)NR_aR_b$, $NR_bC(O)R_a$, $NR_bC(O)OR_a$, $-S(O)_2R_a$, or $-S(O)_2NR_aR_b$.
 [0274] For example, T_2 is NR_aR_b or $-C(O)NR_aR_b$, in which each of R_a and R_b , independently is H or C_1-C_6 alkyl, or R_a and R_b , together with the N atom to which they are attached, form a 4 to 7-membered heterocycloalkyl ring having 0 or 1 additional heteroatom, the C_1-C_6 alkyl and the 4 to 7-membered heterocycloalkyl ring being optionally substituted with one or more $-Q_3-T_3$.

[0275] For example, Q_2 is C_1-C_3 alkyl linker optionally substituted with halo or hydroxyl.

[0276] For example, Q_2 is a bond or methyl linker and T_2 is H, halo, OR_a , NR_aR_b , $-(NR_aR_bR_c)^+A^-$, or $-S(O)_2NR_aR_b$.

[0277] For example, each of R_a , R_b , and R_c , independently is H or C_1-C_6 alkyl optionally substituted with one or more $-Q_3-T_3$.

[0278] For example, one of R_a , R_b , and R_c is H.

[0279] For example, R_a and R_b , together with the N atom to which they are attached, form a 4 to 7-membered heterocycloalkyl ring having 0 or 1 additional heteroatoms to the N atom (e.g., azetidiny, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, triazolidinyl, piperidinyl, 1,2,3,6-tetrahydropyridinyl, piperazinyl, morpholinyl, 1,4-diazepanyl, 1,4-oxazepanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 2,5-diazabicyclo[2.2.1]heptanyl, and the like) and the ring is optionally substituted with one or more $-Q_3-T_3$.

[0280] For example, $-Q_3-T_3$ is oxo.

[0281] For example, T_2 is 4 to 7-membered heterocycloalkyl or C_3-C_8 cycloalkyl and one or more $-Q_3-T_3$ are oxo.

[0282] For example, Q_3 is a bond or unsubstituted or substituted C_1-C_3 alkyl linker.

[0283] For example, T_3 is H, halo, 4 to 7-membered heterocycloalkyl, C_1-C_3 alkyl, OR_d , $COOR_d$, $-S(O)_2R_d$, or NR_dR_e .

[0284] For example, one of R_d and R_e is H.

[0285] For example, R_7 is $-C(O)R_f$.

[0286] For example, R_7 is $-C(O)R_f$ in which R_f is C_3-C_8 cycloalkyl.

[0287] For example, R_7 is C_6-C_{10} aryl substituted with one or more $-Q_5-T_5$.

[0288] For example, R_7 is phenyl optionally substituted with one or more $-Q_5-T_5$.

[0289] For example, R_7 is C_1-C_6 alkyl optionally substituted with one or more $-Q_5-T_5$.

[0290] For example, R_7 is C_3-C_8 cycloalkyl optionally substituted with one or more $-Q_5-T_5$.

[0291] For example, R_7 is 4 to 7-membered heterocycloalkyl (e.g., azetidiny, oxetanyl, thietanyl, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, triazolidinyl, tetrahydropyranyl, piperidinyl, 1,2,3,6-tetrahydropyridinyl, piperazinyl, tetrahydro-2H-pyranyl, 3,6-dihydro-2H-pyranyl, tetrahydro-2H-thiopyran, and morpholinyl, and the like) optionally substituted with one or more $-Q_5-T_5$.

[0292] For example, R_7 is 5 to 6-membered heterocycloalkyl optionally substituted with one or more $-Q_5-T_5$.

[0293] For example, R_7 is isopropyl.

[0294] For example, R_7 is pyrrolidinyl, piperidinyl, tetrahydropyran, tetrahydro-2H-thiopyran, cyclopentyl, or cyclohexyl, each optionally substituted with one or more $-Q_5-T_5$.

[0295] For example, R_7 is cyclopentyl cyclohexyl or tetrahydro-2H-thiopyran, each optionally substituted with one or more $-Q_5-T_5$.

[0296] For example, one or more $-Q_5-T_5$ are oxo.

[0297] For example, R_7 is 1-oxide-tetrahydro-2H-thiopyran or 1,1-dioxide-tetrahydro-2H-thiopyran.

[0298] For example, Q is a bond and T_5 is amino, mono- C_1-C_6 alkylamino, di- C_1-C_6 alkylamino.

[0299] For example, Q_5 is $NHC(O)$ and T_5 is C_1-C_6 alkyl or C_1-C_6 alkoxy.

[0300] For example, T_4 is 4 to 7-membered heterocycloalkyl or C_3-C_8 cycloalkyl and one or more $-Q_5-T_5$ are oxo.

[0301] For example, T_5 is H, halo, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_3 - C_8 cycloalkyl, C_6 - C_{10} aryl, or 4 to 7-membered heterocycloalkyl.

[0302] For example, Q is a bond and T_5 is C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, or 4 to 7-membered heterocycloalkyl.

[0303] For example, Q_5 is CO, $S(O)_2$, or $NHC(O)$; and T_5 is C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_3 - C_8 cycloalkyl, or 4 to 7-membered heterocycloalkyl.

[0304] For example, T_5 is C_1 - C_6 alkyl or C_1 - C_6 alkoxy, each optionally substituted with halo, hydroxyl, cyano, C_1 - C_6 alkoxy, amino, mono- C_1 - C_6 alkylamino, di- C_1 - C_6 alkylamino, or C_3 - C_8 cycloalkyl.

[0305] For example, Q_5 is C_1 - C_3 alkyl linker and T_5 is H or C_6 - C_{10} aryl.

[0306] For example, Q is C_1 - C_3 alkyl linker and T_5 is C_3 - C_8 cycloalkyl, 4 to 7-membered heterocycloalkyl, or $S(O)_q R_q$.

[0307] For example, R_{11} is H.

[0308] For example, each of R_2 and R_4 , independently, is H or C_1 - C_6 alkyl optionally substituted with amino, mono- C_1 - C_6 alkylamino, di- C_1 - C_6 alkylamino, or C_6 - C_{10} aryl.

[0309] For example, each of R_2 and R_4 , independently is C_1 - C_3 alkyl optionally substituted with C_1 - C_6 alkoxy.

[0310] For example, each of R_2 and R_4 is methyl.

[0311] For example, R_1 is H.

[0312] For example, R_{12} is H, methyl, ethyl, ethenyl, or halo.

[0313] For example, R_{12} is methyl.

[0314] For example, R_{12} is ethyl.

[0315] For example, R_{12} is ethenyl.

[0316] For example, R_8 is H, methyl, ethyl, or ethenyl.

[0317] For example, R_8 is methyl.

[0318] For example, R_8 is ethyl.

[0319] For example, R_8 is 4 to 7-membered heterocycloalkyl (e.g., azetidiny, oxetanyl, thietanyl, pyrrolidiny, imidazolidiny, pyrazolidiny, oxazolidiny, isoxazolidiny, triazolidiny, tetrahydrofuranyl, piperidiny, 1,2,3,6-tetrahydropyridiny, piperaziny, tetrahydro-2H-pyran, 3,6-dihydro-2H-pyran, tetrahydro-2H-thiopyran, morpholinyl, 1,4-diazepanyl, 1,4-oxazepanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 2,5-diazabicyclo[2.2.1]heptanyl, and the like).

[0320] For example, R_8 is tetrahydropyran.

[0321] For example, R_8 is tetrahydropyran and R_7 is $-Q_4-T_4$, in which Q_4 is a bond or C_1 - C_4 alkyl linker and T_4 is H, C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl or 4 to 7-membered heterocycloalkyl.

[0322] For example, Z_1 is NR_7R_8 or $CR_7R_8R_{14}$ wherein R_7 and R_8 , together with the atom to which they are attached, form a 4 to 11-membered heterocycloalkyl ring having 1 to 3 heteroatoms (e.g., azetidiny, oxetanyl, thietanyl, pyrrolidiny, imidazolidiny, pyrazolidiny, oxazolidiny, isoxazolidiny, triazolidiny, tetrahydrofuranyl, piperidiny, 1,2,3,6-tetrahydropyridiny, piperaziny, tetrahydro-2H-pyran, 3,6-dihydro-2H-pyran, tetrahydro-2H-thiopyran, and morpholinyl, and the like) or C_3 - C_8 cycloalkyl, each optionally substituted with one or more $-Q_6-T_6$.

[0323] For example, the ring formed by R_7 and R_8 is selected from the group consisting of azetidiny, pyrrolidiny, piperidiny, morpholinyl, piperaziny, and cyclohexenyl, each optionally substituted with one $-Q_6-T_6$.

[0324] For example, $-Q_6-T_6$ is oxo.

[0325] For example, T_6 is H, halo, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_3 - C_8 cycloalkyl, C_6 - C_{10} aryl, or 4 to 7-membered heterocycloalkyl.

[0326] For example, Q_6 is a bond and T_6 is C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, or 4 to 7-membered heterocycloalkyl.

[0327] For example, Q_6 is CO, $S(O)_2$, or $NHC(O)$; and T_6 is C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_3 - C_8 cycloalkyl, or 4 to 7-membered heterocycloalkyl.

[0328] For example, T_6 is C_1 - C_6 alkyl or C_1 - C_6 alkoxy, each optionally substituted with halo, hydroxyl, cyano, C_1 - C_6 alkoxy, amino, mono- C_1 - C_6 alkylamino, di- C_1 - C_6 alkylamino, or C_3 - C_8 cycloalkyl.

[0329] For example, Q_6 is C_1 - C_3 alkyl linker and T_6 is H or C_6 - C_{10} aryl.

[0330] For example, Q_6 is C_1 - C_3 alkyl linker and T_6 is C_3 - C_8 cycloalkyl, 4 to 7-membered heterocycloalkyl, or $S(O)R_p$.

[0331] For example, each of R_p and R_q , independently, is C_1 - C_6 alkyl.

[0332] For example, R_{13} is H or methyl.

[0333] For example, R_{13} is H.


[0334] For example, R_3 is H.

[0335] For example, A^- is Br^- .

[0336] For example, each of R_5 , R_9 , and R_{10} is H.

[0337] Another subset of the compounds of Formula (II) includes those of Formula (He):

(IIe)



[0338] The compounds of Formula (He) can include one or more of the following features:

[0339] For example, each of R_a and R_b , independently is H or C_1 - C_6 alkyl optionally substituted with one or more $-Q_3-T_3$.

[0340] For example, one of R_a and R_b is H.

[0341] For example, R_a and R_b , together with the N atom to which they are attached, form a 4 to 7-membered heterocycloalkyl ring having 0 or 1 additional heteroatoms to the N atom (e.g., azetidiny, pyrrolidiny, imidazolidiny, pyrazolidiny, oxazolidiny, isoxazolidiny, triazolidiny, piperidiny, 1,2,3,6-tetrahydropyridiny, piperaziny, morpholinyl, 1,4-diazepanyl, 1,4-oxazepanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 2,5-diazabicyclo[2.2.1]heptanyl, and the like) and the ring is optionally substituted with one or more $-Q_3-T_3$.

[0342] For example, R_a and R_b , together with the N atom to which they are attached, form azetidiny, pyrrolidiny, imidazolidiny, pyrazolidiny, oxazolidiny, isoxazolidiny, triazolidiny, tetrahydrofuranyl, piperidiny, 1,2,3,6-tetrahydropyridiny, piperaziny, or morpholinyl, and the ring is optionally substituted with one or more $-Q_3-T_3$.

[0343] For example, one or more $-Q_3-T_3$ are oxo.

[0344] For example, Q_3 is a bond or unsubstituted or substituted C_1 - C_3 alkyl linker.

[0345] For example, T_3 is H, halo, 4 to 7-membered heterocycloalkyl, C_1 - C_3 alkyl, OR_d , $COOR_d$, $-S(O)_2R_d$, or $-NR_dR_c$.

[0346] For example, one of R_d and R_e is H.

[0347] For example, R_7 is C_3 - C_8 cycloalkyl or 4 to 7-membered heterocycloalkyl, each optionally substituted with one or more $-Q_5$ - T_5 .

[0348] For example, R_7 is piperidinyl, tetrahydropyran, tetrahydro-2H-thiopyranyl, cyclopentyl, cyclohexyl, pyrrolidinyl, or cycloheptyl, each optionally substituted with one or more $-Q_5$ - T_5 .

[0349] For example, R_7 is cyclopentyl cyclohexyl or tetrahydro-2H-thiopyranyl, each optionally substituted with one or more $-Q_5$ - T_5 .

[0350] For example, Q_5 is $NHC(O)$ and T_5 is C_1 - C_6 alkyl or C_1 - C_6 alkoxy.

[0351] For example, one or more $-Q_5$ - T_5 are oxo.


[0352] For example, R_7 is 1-oxide-tetrahydro-2H-thiopyranyl or 1,1-dioxide-tetrahydro-2H-thiopyranyl.

[0353] For example, Q_5 is a bond and T_5 is amino, mono- C_1 - C_6 alkylamino, di- C_1 - C_6 alkylamino.

[0354] For example, Q_5 is CO , $S(O)_2$, or $NHC(O)$; and T_5 is C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_3 - C_8 cycloalkyl, or 4 to 7-membered heterocycloalkyl.

[0355] For example, R_8 is H, methyl, or ethyl.

[0356] The compounds suitable for use in the method of this invention also include compounds of Formula (III):



or a pharmaceutically acceptable salt or ester thereof. In Formula (III):

[0357] X_1' is N or CR_{11}' ;

[0358] X_2' is N or CR_{13}' ;

[0359] X_3 is N or C, and when X_3 is N, R_6' is absent;

[0360] Z_2 is $NR_7'R_8'$, OR_5' , $S(O)_2R_7'$, or $CR_7'R_8'R_{14}'$, in which a' is 0, 1, or 2;

[0361] each of R_1' , R_5' , R_9' , and R_{10}' , independently, is H or C_1 - C_6 alkyl optionally substituted with one or more substituents selected from the group consisting of halo, hydroxyl, $COOH$, $C(O)O$ - C_1 - C_6 alkyl, cyano, C_1 - C_6 alkoxy, amino, mono- C_1 - C_6 alkylamino, di- C_1 - C_6 alkylamino, C_3 - C_8 cycloalkyl, C_6 - C_{10} aryl, 4 to 12-membered heterocycloalkyl, and 5- or 6-membered heteroaryl;

[0362] each of R_2' , R_3' , and R_4' , independently, is $-Q_1$ - T_1 , in which Q_1 is a bond or C_1 - C_3 alkyl linker optionally substituted with halo, cyano, hydroxyl or C_1 - C_6 alkoxy, and T_1 is H, halo, hydroxyl, $COOH$, cyano, azido, or R_{S1}' , in which R_{S1}' is C_1 - C_3 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_1 - C_6 alkoxy, $C(O)O$ - C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, C_6 - C_{10} aryl, amino,

mono- C_1 - C_6 alkylamino, di- C_1 - C_6 alkylamino, 4 to 12-membered heterocycloalkyl, or 5- or 6-membered heteroaryl, and R_{S1}' is optionally substituted with one or more substituents selected from the group consisting of halo, hydroxyl, oxo, $COOH$, $C(O)O$ - C_1 - C_6 alkyl, cyano, C_1 - C_6 alkoxy, amino, mono- C_1 - C_6 alkylamino, di- C_1 - C_6 alkylamino, C_3 - C_8 cycloalkyl, C_6 - C_{10} aryl, 4 to 12-membered heterocycloalkyl, and 5- or 6-membered heteroaryl;

[0363] R_6' is H, halo, cyano, azido, OR_d' , $-C(O)R_a'$, $-C(O)OR_a'$, $-C(O)NR_a'R_b'$, $-NR_b'C(O)R_a'$, $-S(O)_bR_a'$, $-S(O)_bNR_a'R_b'$, or R_{S2}' , in which R_{S2}' is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_3 - C_8 cycloalkyl, or 4 to 12-membered heterocycloalkyl, b' is 0, 1, or 2, each of R_a' and R_b' , independently is H or R_{S3}' , and R_{S3}' is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_3 - C_8 cycloalkyl, C_6 - C_{10} aryl, 4 to 12-membered heterocycloalkyl, or 5- or 6-membered heteroaryl; or R_a' and R_b' , together with the N atom to which they are attached, form a 4 to 12-membered heterocycloalkyl ring having 0 or 1 additional heteroatom; and each of R_{S2}' , R_{S3}' , and the 4 to 12-membered heterocycloalkyl ring formed by R_a' and R_b' , is optionally substituted with one or more $-Q_2$ - T_2 , wherein Q_2 is a bond or C_1 - C_3 alkyl linker each optionally substituted with halo, cyano, hydroxyl or C_1 - C_6 alkoxy, and T_2 is H, halo, cyano, $-OR_c'$, $-NR_c'R_d'$, $-C(O)R_c'$, $-C(O)OR_c'$, $-C(O)NR_c'R_d'$, $-NR_d'C(O)R_c'$, $-NR_d'C(O)OR_c'$, $-S(O)_2R_c'$, $-S(O)_2NR_c'R_d'$, or R_{S4}' , in which each of R_c' and R_d' , independently is H or R_{S5}' , each of R_{S4}' and R_{S5}' , independently, is C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, C_6 - C_{10} aryl, 4 to 12-membered heterocycloalkyl, or 5- or 6-membered heteroaryl, or R_c' and R_d' , together with the N atom to which they are attached, form a 4 to 12-membered heterocycloalkyl ring having 0 or 1 additional heteroatom, and each of R_{S4}' , R_{S5}' , and the 4 to 12-membered heterocycloalkyl ring formed by R_c' and R_d' , is optionally substituted with one or more $-Q_3$ - T_3 , wherein Q_3 is a bond or C_1 - C_3 alkyl linker each optionally substituted with halo, cyano, hydroxyl or C_1 - C_6 alkoxy, and T_3 is selected from the group consisting of halo, cyano, C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, C_6 - C_{10} aryl, 4 to 12-membered heterocycloalkyl, 5- or 6-membered heteroaryl, OR_e' , $COOR_e'$, $-S(O)_2R_e'$, $-NR_e'R_f'$, and $-C(O)NR_e'R_f'$, each of R_e' and R_f' independently being H or C_1 - C_6 alkyl, or $-Q_3$ - T_3 is oxo; or $-Q_2$ - T_2 is oxo; provided that $-Q_2$ - T_2 is not H;


[0364] R_7' is $-Q_4$ - T_4 , in which Q_4 is a bond, C_1 - C_4 alkyl linker, or C_2 - C_4 alkenyl linker, each linker optionally substituted with halo, cyano, hydroxyl or C_1 - C_6 alkoxy, and T_4 is H, halo, cyano, $NR_g'R_h'$, $-OR_g'$, $-C(O)R_g'$, $-C(O)OR_g'$, $-C(O)NR_g'R_h'$, $-C(O)NR_g'OR_h'$, $-NR_g'C(O)R_h'$, $-S(O)_2R_g'$, or R_{S6}' , in which each of R_g' and R_h' , independently is H or R_{S7}' , each of R_{S6}' and R_{S7}' , independently is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_3 - C_8 cycloalkyl, C_6 - C_{10} aryl, 4 to 12-membered heterocycloalkyl, or 5- or 6-membered heteroaryl, and each of R_{S6}' and R_{S7}' is optionally substituted with one or more $-Q_5$ - T_5 , wherein Q' is a bond, $C(O)$, $C(O)NR_i'$, $NR_i'C(O)$, $S(O)_2$, or C_1 - C_3 alkyl linker, R_i' being H or C_1 - C_6 alkyl, and T_5 is H, halo, C_1 - C_6 alkyl, hydroxyl, cyano, C_1 - C_6 alkoxy, amino, mono- C_1 - C_6 alkylamino, di- C_1 - C_6 alkylamino, C_3 - C_8 cycloalkyl, C_6 - C_{10} aryl, 4 to 12-membered heterocycloalkyl, 5- or 6-membered heteroaryl, or $S(O)_qR_q'$ in which q' is 0, 1, or 2 and R_q' is C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_3 - C_8 cycloalkyl, C_6 - C_{10} aryl, 4 to 12-membered heterocycloalkyl, or 5- or 6-membered heteroaryl, and T_5 is optionally substituted with one or more substituents selected from the group consisting of halo, C_1 - C_6

alkyl, hydroxyl, cyano, C₁-C₆ alkoxy, amino, mono-C₁-C₆ alkylamino, di-C₁-C₆ alkylamino, C₃-C₈ cycloalkyl, C₆-C₁₀ aryl, 4 to 12-membered heterocycloalkyl, and 5- or 6-membered heteroaryl except when T₅ is H, halo, hydroxyl, or cyano; or -Q₅'-T₅' is oxo; provided that R₇' is not H;




[0365] each of R₈', R₁₁', R₁₂', and R₁₃', independently, is H, halo, hydroxyl, COOH, cyano, R_{SS}', OR_{SS}', or COOR_{SS}', in which R_{SS}' is C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, 4 to 12-membered heterocycloalkyl, amino, mono-C₁-C₆ alkylamino, or di-C₁-C₆ alkylamino, and R_{SS}' is optionally substituted with one or more substituents selected from the group consisting of halo, hydroxyl, COOH, C(O)O—C₁-C₆ alkyl, cyano, C₁-C₆ alkoxy, amino, mono-C₁-C₆ alkylamino, and di-C₁-C₆ alkylamino; or R₇' and R₈', together with the N atom to which they are attached, form a 4 to 12-membered heterocycloalkyl ring having 0 to 2 additional heteroatoms, or R₇' and R₈', together with the C atom to which they are attached, form C₃-C₈ cycloalkyl or a 4 to 12-membered heterocycloalkyl ring having 1 to 3 heteroatoms, and each of the 4 to 12-membered heterocycloalkyl rings or C₃-C₈ cycloalkyl formed by R₇' and R₈' is optionally substituted with one or more -Q₆'-T₆', wherein Q₆' is a bond, C(O), C(O)NR_m', NR_m'C(O), S(O)₂, or C₁-C₃ alkyl linker, R_m' being H or C₁-C₆ alkyl, and T₆' is H, halo, C₁-C₆ alkyl, hydroxyl, cyano, C₁-C₆ alkoxy, amino, mono-C₁-C₆ alkylamino, di-C₁-C₆ alkylamino, C₃-C₈ cycloalkyl, C₆-C₁₀ aryl, 4 to 12-membered heterocycloalkyl, 5- or 6-membered heteroaryl, or S(O)_pR_p' in which p' is 0, 1, or 2 and R_p' is C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₃-C₈ cycloalkyl, C₆-C₁₀ aryl, 4 to 12-membered heterocycloalkyl, or 5- or 6-membered heteroaryl, and T₆' is optionally substituted with one or more substituents selected from the group consisting of halo, C₁-C₆ alkyl, hydroxyl, cyano, C₁-C₆ alkoxy, amino, mono-C₁-C₆ alkylamino, di-C₁-C₆ alkylamino, C₃-C₈ cycloalkyl, C₆-C₁₀ aryl, 4 to 12-membered heterocycloalkyl, and 5- or 6-membered heteroaryl except when T₆' is H, halo, hydroxyl, or cyano; or -Q₆'-T₆' is oxo; and

[0366] R₁₄' is absent, H, or C₁-C₆ alkyl optionally substituted with one or more substituents selected from the group consisting of halo, hydroxyl, COOH, C(O)O—C₁-C₆ alkyl, cyano, C₁-C₆ alkoxy, amino, mono-C₁-C₆ alkylamino, di-C₁-C₆ alkylamino, C₃-C₈ cycloalkyl, C₆-C₁₀ aryl, 4 to 12-membered heterocycloalkyl, and 5- or 6-membered heteroaryl.

[0367] One subset of the compounds of Formula (III) includes those of Formula (Ma):



[0368] Another subset of the compounds of Formula (III) includes those of Formula (IIIb), (IIIc), or (IIId):



[0369] The compounds of Formulae (III), (IIIa), (IIIb), (IIIc), and (IIId) can include one or more of the following features:

[0370] For example, the compounds of Formula (III) are not

[0371] N-(5-(((4,6-dimethyl-2-oxo-1,2-dihydropyridin-3-yl)methyl)carbamoyl)-2-methylphenyl)furan-2-carboxamide,

[0372] N,N'-(5-(((4,6-dimethyl-2-oxo-1,2-dihydropyridin-3-yl)methyl)carbamoyl)-1,3-phenylene)diacetamide,

[0373] N-(((4,6-dimethyl-2-oxo-1,2-dihydropyridin-3-yl)methyl)-3-pivalamidobenzamide,

[0374] 3-(3,4-dihydro-2H-benzo[b][1,4]dioxepine-7-sulfonamido)-N-(((4,6-dimethyl-2-oxo-1,2-dihydropyridin-3-yl)methyl)benzamide,

[0375] N-(((4,6-dimethyl-2-oxo-1,2-dihydropyridin-3-yl)methyl)-3,5-dimethoxybenzamide,

[0376] N-(((4,6-dimethyl-2-oxo-1,2-dihydropyridin-3-yl)methyl)-3,4,5-trimethoxybenzamide,

[0377] 3-allyl-N-(((4,6-dimethyl-2-oxo-1,2-dihydropyridin-3-yl)methyl)-4,5-dimethoxybenzamide,

[0378] 4-(2-amino-2-oxoethoxy)-3-chloro-N-((4,6-dimethyl-2-oxo-1,2-dihydropyridin-3-yl)methyl)-5-methoxybenzamide,

[0379] 3-chloro-N-((4,6-dimethyl-2-oxo-1,2-dihydropyridin-3-yl)methyl)-4-hydroxy-5-methoxybenzamide, or

[0380] 3-bromo-N-((4,6-dimethyl-2-oxo-1,2-dihydropyridin-3-yl)methyl)-5-methoxy-4-propoxybenzamide.

[0381] For example, X_1' is CR_{11}' and X_2' is CR_{13}' .

[0382] For example, X_1' is CR_{11}' and X_2' is N.

[0383] For example, X_1' is N and X_2' is CR_{13}' .

[0384] For example, X_1' is N and X_2' is N.

[0385] For example, X_3 is C.

[0386] For example, X_3 is N and R_6' is absent.

[0387] For example, Z_2 is $NR_7'R_8'$.

[0388] For example, Z_2 is $CR_7'R_8'R_{14}'$.

[0389] For example, Z_2 is OR_7' .

[0390] For example, Z_2 is $S(O)_a'R_7'$, in which a' is 0, 1, or 2.

[0391] For example, R_6' is H.

[0392] For example, R_6' is halo (e.g., fluorine, chlorine, bromine, and iodine).

[0393] For example, R_6' is C_1 - C_3 alkyl optionally substituted with one or more $-Q_2'-T_2'$.

[0394] For example, R_6' is CF_3 .

[0395] For example, R_6' is C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, or C_3 - C_6 cycloalkyl each optionally substituted with one or more $-Q_2'-T_2'$.

[0396] For example, R_6' is ethenyl.

[0397] For example, R_6' is ethynyl.

[0398] For example, R_6' is ethynyl substituted with one or more $-Q_2'-T_2'$, in which Q_2' is a bond or C_1 - C_3 alkyl linker and T_2' is C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, or 4 to 7-membered heterocycloalkyl (e.g., azetidiny, oxetanyl, thietanyl, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, triazolidinyl, tetrahydrofuranly, piperidinyl, 1,2,3,6-tetrahydropyridinyl, piperazinyl, tetrahydro-2H-pyranly, 3,6-dihydro-2H-pyranly, tetrahydro-2H-thiopyranly, 1,4-diazepanyl, 1,4-oxazepanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 2,5-diazabicyclo[2.2.1]heptanyl, and morpholinyl, and the like) optionally substituted with one or more $-Q_3'-T_3'$.

[0399] For example, R_6' is cyano.

[0400] For example, R_6' is azido.

[0401] For example, R_6' is $C(O)H$.

[0402] For example, R_6' is OR_a' or $-C(O)R_a'$.

[0403] For example, R_a' is C_3 - C_6 alkyl or 4 to 7-membered heterocycloalkyl (e.g., azetidiny, oxetanyl, thietanyl, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, triazolidinyl, tetrahydrofuranly, piperidinyl, 1,2,3,6-tetrahydropyridinyl, piperazinyl, tetrahydro-2H-pyranly, 3,6-dihydro-2H-pyranly, tetrahydro-2H-thiopyranly, 1,4-diazepanyl, 1,4-oxazepanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 2,5-diazabicyclo[2.2.1]heptanyl, and morpholinyl, and the like), which is optionally substituted with one or more $-Q_2'-T_2'$.

[0404] For example, R_6' is 4 to 7-membered heterocycloalkyl (e.g., azetidiny, oxetanyl, thietanyl, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, triazolidinyl, tetrahydrofuranly, piperidinyl, 1,2,3,6-tetrahydropyridinyl, piperazinyl, tetrahydro-2H-pyranly, 3,6-dihydro-2H-pyranly, tetrahydro-2H-thiopyranly, 1,4-diazepanyl, 1,4-oxazepanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 2,5-diazabicyclo[2.2.1]heptanyl, and morpholinyl, and the like) optionally substituted with one or more $-Q_2'-T_2'$.

[0405] For example, R_6' is piperidinyl, 2,2,6,6-tetramethylpiperidinyl, 1,2,3,6-tetrahydropyridinyl, 2,2,6,6-tetramethyl-

1,2,3,6-tetrahydropyridinyl, piperazinyl, morpholinyl, tetrahydro-2H-pyranly, 3,6-dihydro-2H-pyranly, or pyrrolidinyl, each of which is optionally substituted with one or more $-Q_2'-T_2'$.

[0406] For example, R_6' is 4 to 7-membered heterocycloalkyl optionally substituted with one or more $-Q_2'-T_2'$, and $-Q_2'-T_2'$ is oxo or Q_2' is a bond and T_2' is $-OR_c'$, $-NR_c'R_d'$, $-C(O)R_c'$, $-C(O)OR_c'$, $-S(O)_2R_c'$, C_1 - C_6 alkyl, or 4 to 7-membered heterocycloalkyl, each of which is optionally substituted with one or more $-Q_3'-T_3'$ when R_c' or R_d' is not H.

[0407] For example, R_6' is $-C(O)R_c'$, $-C(O)OR_a'$, $-C(O)NR_a'R_b'$, $-NR_b'C(O)R_a'$, $-SR_a'$, $-S(O)_2R_a'$, or $-S(O)_2NR_a'R_b'$.

[0408] For example, each of R_a' and R_b' , independently is H, C_1 - C_6 alkyl, or C_3 - C_8 cycloalkyl optionally substituted with one or more $-Q_2'-T_2'$.

[0409] For example, one of R_a' and R_b' is H.

[0410] For example, R_a' and R_b' , together with the N atom to which they are attached, form a 4 to 7-membered heterocycloalkyl ring having 0 or 1 additional heteroatoms to the N atom (e.g., azetidiny, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, triazolidinyl, piperidinyl, 1,2,3,6-tetrahydropyridinyl, piperazinyl, 1,4-diazepanyl, 1,4-oxazepanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 2,5-diazabicyclo[2.2.1]heptanyl, and morpholinyl, and the like) and the ring is optionally substituted with one or more $-Q_2'-T_2'$.

[0411] For example, $-Q_2'-T_2'$ is oxo.

[0412] For example, Q_2' is a bond.

[0413] For example, Q_2' is an unsubstituted C_1 - C_3 alkyl linker.

[0414] For example, T_2' is C_1 - C_6 alkyl or C_6 - C_{10} aryl, each optionally substituted with one or more $-Q_3'-T_3'$.

[0415] For example, T_2' is an unsubstituted substituted straight chain C_1 - C_6 or branched C_3 - C_6 alkyl, including but not limited to, methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl, t-butyl, n-pentyl, s-pentyl and n-hexyl.

[0416] For example, T_2' is phenyl.

[0417] For example, T_2' is halo (e.g., fluorine, chlorine, bromine, and iodine).

[0418] For example, T_2' is 4 to 7-membered heterocycloalkyl (e.g., azetidiny, oxetanyl, thietanyl, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, triazolidinyl, tetrahydrofuranly, piperidinyl, 1,2,3,6-tetrahydropyridinyl, piperazinyl, tetrahydro-2H-pyranly, 3,6-dihydro-2H-pyranly, tetrahydro-2H-thiopyranly, 1,4-diazepanyl, 1,4-oxazepanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 2,5-diazabicyclo[2.2.1]heptanyl, and morpholinyl, and the like) optionally substituted with one or more $-Q_3'-T_3'$.

[0419] For example, T_2' is $-OR_c'$, $-C(O)R_c'$, $-C(O)OR_c'$, or $-S(O)_2R_c'$.

[0420] For example, is C_1 - C_6 alkyl or 4 to 7-membered heterocycloalkyl (e.g., azetidiny, oxetanyl, thietanyl, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, triazolidinyl, tetrahydrofuranly, piperidinyl, 1,2,3,6-tetrahydropyridinyl, piperazinyl, tetrahydro-2H-pyranly, 3,6-dihydro-2H-pyranly, tetrahydro-2H-thiopyranly, 1,4-diazepanyl, 1,4-oxazepanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 2,5-diazabicyclo[2.2.1]heptanyl, and morpholinyl, and the like), which is optionally substituted with one or more $-Q_3'-T_3'$.

[0421] For example, each of R_c' and R_d' , independently is H or C_1 - C_6 alkyl optionally substituted with one or more $-Q_3'-T_3'$.

[0422] For example, R_c' is H.

[0423] For example, R_d' is H.

[0424] For example, R_c' and R_d' , together with the N atom to which they are attached, form a 4 to 7-membered heterocycloalkyl ring having 0 or 1 additional heteroatoms to the N atom (e.g., azetidiny, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, triazolidinyl, piperidinyl, 1,2,3,6-tetrahydropyridinyl, piperazinyl, 1,4-diazepanyl, 1,4-oxazepanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 2,5-diazabicyclo[2.2.1]heptanyl, and morpholinyl, and the like) and the ring is optionally substituted with one or more $-Q_3'-T_3'$.

[0425] For example, Q_2' is a bond and T_2' is $-OR_c'$, $-NR_d'R_e'$, $-C(O)R_c'$, $-C(O)OR_c'$, $-S(O)_2R_c'$, C_1-C_6 alkyl, or 4 to 7-membered heterocycloalkyl, each of which is optionally substituted with one or more $-Q_3'-T_3'$ when R_c' or R_d' is not H.

[0426] For example, $-Q_3'-T_3'$ is oxo.

[0427] For example, T_2' is 4 to 7-membered heterocycloalkyl or C_3-C_8 cycloalkyl and one or more $-Q_3'-T_3'$ are oxo.

[0428] For example, Q_3' is a bond or unsubstituted or substituted C_1-C_3 alkyl linker.

[0429] For example, T_3' is H, halo, 4 to 7-membered heterocycloalkyl, C_1-C_3 alkyl, OR_e' , $COOR_e'$, $-S(O)_2R_e'$, $-NR_e'R_f'$, or $-C(O)NR_e'R_f'$.

[0430] For example, one of R_d' and R_e' is H.

[0431] For example, Q_3' is a bond or C_1-C_3 alkyl linker and T_3' is selected from the group consisting of C_1-C_3 alkyl, halo, OR_e' , $-S(O)_2R_e'$, $-NR_e'R_f'$, and $-C(O)NR_e'R_f'$.

[0432] For example, Q_3' is a bond or C_1-C_3 alkyl linker and T_3' is selected from the group consisting of C_1-C_3 alkyl, OR_e' , $-S(O)_2R_e'$, or $-NR_e'R_f'$.

[0433] For example, R_e' is H.

[0434] For example, R_f' is H.

[0435] For example, R_7' is $-C(O)R_g'$.

[0436] For example, R_7' is $-C(O)R_g'$, in which R_g' is C_3-C_8 cycloalkyl, 4 to 7-membered heterocycloalkyl, C_3-C_8 cycloalkyl.

[0437] For example, R_7' is C_6-C_{10} aryl substituted with one or more $-Q_5'-T_5'$.

[0438] For example, R_7' is phenyl optionally substituted with one or more $-Q_5'-T_5'$.

[0439] For example, R_7' is C_1-C_6 alkyl optionally substituted with one or more $-Q_5'-T_5'$.

[0440] For example, R_7' is C_3-C_8 cycloalkyl optionally substituted with one or more $-Q_5'-T_5'$.

[0441] For example, R_7' is 4 to 7-membered heterocycloalkyl (e.g., azetidiny, oxetanyl, thietanyl, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, triazolidinyl, tetrahydrofuranly, piperidinyl, 1,2,3,6-tetrahydropyridinyl, piperazinyl, tetrahydro-2H-pyranly, 3,6-dihydro-2H-pyranly, tetrahydro-2H-thiopyranly, 1,4-diazepanyl, 1,4-oxazepanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 2,5-diazabicyclo[2.2.1]heptanyl, and morpholinyl, and the like) optionally substituted with one or more $-Q_5'-T_5'$.

[0442] For example, R_7' is 5 to 6-membered heterocycloalkyl optionally substituted with one or more $-Q_5'-T_5'$.

[0443] For example, R_7' is isopropyl.

[0444] For example, R_7' is pyrrolidinyl, piperidinyl, tetrahydropyran, cyclopentyl, or cyclohexyl, cycloheptyl, each optionally substituted with one $-Q_5'-T_5'$.

[0445] For example, R_7' is cyclopentyl or cyclohexyl, each optionally substituted with one $-Q_5'-T_5'$.

[0446] For example, Q_5' is $NHC(O)$ and T_5' is C_1-C_6 alkyl or C_1-C_6 alkoxy.

[0447] For example, $-Q_5'-T_5'$ is oxo.

[0448] For example, T_4' is 4 to 7-membered heterocycloalkyl, C_3-C_8 cycloalkyl, or C_6-C_{10} aryl, and one or more $-Q_5'-T_5'$ are oxo.

[0449] For example, R_7' is 1-oxide-tetrahydro-2H-thiopyranly or 1,1-dioxide-tetrahydro-2H-thiopyranly.

[0450] For example, R_7' is cyclohexanonyl, e.g., cyclohexanon-4-yl.

[0451] For example, T_5' is H, halo, C_1-C_6 alkyl, C_1-C_6 alkoxy, C_3-C_8 cycloalkyl, C_6-C_{10} aryl, or 4 to 7-membered heterocycloalkyl.

[0452] For example, Q_5' is a bond and T_5' is C_1-C_6 alkyl, C_3-C_8 cycloalkyl, or 4 to 7-membered heterocycloalkyl.

[0453] For example, Q_5' is a bond and T_5' is 5- or 6-membered heteroaryl, amino, mono- C_1-C_6 alkylamino, di- C_1-C_6 alkylamino, T_5' being optionally substituted with one or more substituents selected from the group consisting of halo, hydroxyl, C_1-C_6 alkoxy, or C_3-C_8 cycloalkyl.

[0454] For example, Q_5' is CO , $S(O)_2$, or $NHC(O)$; and T_5' is C_1-C_6 alkyl, C_1-C_6 alkoxy, C_3-C_8 cycloalkyl, or 4 to 7-membered heterocycloalkyl.

[0455] For example, T_5' is C_1-C_6 alkyl or C_1-C_6 alkoxy, each optionally substituted with halo, hydroxyl, cyano, C_1-C_6 alkoxy, amino, mono- C_1-C_6 alkylamino, di- C_1-C_6 alkylamino, or C_3-C_8 cycloalkyl.

[0456] For example, Q_5' is C_1-C_3 alkyl linker and T_5' is H or C_6-C_{10} aryl.

[0457] For example, Q_5' is C_1-C_3 alkyl linker and T_5' is C_3-C_8 cycloalkyl, 4 to 7-membered heterocycloalkyl, or $S(O)_2R_g'$.

[0458] For example, R_6' is halo (e.g., fluorine, chlorine, bromine, and iodine) and Z_2 is $S(O)_aR_7'$, in which a is 0, 1, or 2 and R_7' is C_1-C_6 alkyl (e.g., methyl, ethyl, n-propyl, i-propyl, butyl, or t-butyl), C_3-C_8 cycloalkyl (e.g., cyclopentyl, cyclohexyl, or cycloheptyl) or 4 to 7-membered heterocycloalkyl (e.g., azetidiny, oxetanyl, thietanyl, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, triazolidinyl, tetrahydrofuranly, piperidinyl, 1,2,3,6-tetrahydropyridinyl, piperazinyl, tetrahydro-2H-pyranly, 3,6-dihydro-2H-pyranly, tetrahydro-2H-thiopyranly, 1,4-diazepanyl, 1,4-oxazepanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 2,5-diazabicyclo[2.2.1]heptanyl, and morpholinyl, and the like) and R_7' is optionally substituted with one or more $-Q_5'-T_5'$.

[0459] For example, R_6' is halo (e.g., fluorine, chlorine, bromine, and iodine) and Z_2 is OR_7' in which R_7' is 4 to 7-membered heterocycloalkyl (e.g., azetidiny, oxetanyl, thietanyl, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, triazolidinyl, tetrahydrofuranly, piperidinyl, 1,2,3,6-tetrahydropyridinyl, piperazinyl, tetrahydro-2H-pyranly, 3,6-dihydro-2H-pyranly, tetrahydro-2H-thiopyranly, 1,4-diazepanyl, 1,4-oxazepanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 2,5-diazabicyclo[2.2.1]heptanyl, and morpholinyl, and the like) and R_7' is optionally substituted with one or more $-Q_5'-T_5'$.

[0460] For example, R_{11}' is H.

[0461] For example, each of R_2' and R_4' , independently, is H or C_1-C_6 alkyl optionally substituted with azido, halo, amino, mono- C_1-C_6 alkylamino, di- C_1-C_6 alkylamino, or C_6-C_{10} aryl.

[0462] For example, each of R_2' and R_4' , independently, is C_1-C_3 alkyl optionally substituted with C_1-C_6 alkoxy.

[0463] For example, each of R_2' and R_4' is methyl.

[0464] For example, R_1' is H.

[0465] For example, R_1' is C_1 - C_6 alkyl optionally substituted with azido, halo, amino, mono- C_1 - C_6 alkylamino, di- C_1 - C_6 alkylamino, or C_6 - C_{10} aryl.

[0466] For example, R_{12}' is H, methyl, ethyl, ethenyl, or halo.

[0467] For example, R_{12}' is methyl.

[0468] For example, R_{12}' is ethyl.

[0469] For example, R_{12}' is ethenyl or propenyl.

[0470] For example, R_{12}' is methoxyl.

[0471] For example, R_8' is H, methyl, ethyl, or ethenyl.

[0472] For example, R_8' is methyl.

[0473] For example, R_8' is ethyl.

[0474] For example, R_8' is propyl.

[0475] For example, R_8' is ethenyl or propenyl.

[0476] For example, R_8' is C_1 - C_6 alkyl substituted with one or more substituents selected from the group consisting of halo (e.g., F, Cl, or Br), hydroxyl, or C_1 - C_6 alkoxy.

[0477] For example, R_8' is 4 to 7-membered optionally substituted heterocycloalkyl (e.g., azetidiny, oxetanyl, thietanyl, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, triazolidinyl, tetrahydrofuranyl, piperidinyl, 1,2,3,6-tetrahydropyridinyl, piperazinyl, tetrahydro-2H-pyranyl, 3,6-dihydro-2H-pyranyl, tetrahydro-2H-thiopyranyl, 1,4-diazepanyl, 1,4-oxazepanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 2,5-diazabicyclo[2.2.1]heptanyl, and morpholinyl, and the like).

[0478] For example, R_8' is piperidinyl.

[0479] For example, R_8' is 4 to 7-membered optionally substituted heterocycloalkyl and R_7' is $-Q_4'-T_4'$, in which Q_4' is a bond or C_1 - C_4 alkyl linker and T_4' is H, C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl or 4 to 7-membered heterocycloalkyl.

[0480] For example, Z_2 is $NR_7'R_8'$ or $CR_7'R_8'R_{14}'$ wherein R_7' and R_8' , together with the atom to which they are attached, form a 4 to 11-membered heterocycloalkyl ring having 1 to 3 heteroatoms (e.g., azetidiny, oxetanyl, thietanyl, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, triazolidinyl, tetrahydrofuranyl, piperidinyl, 1,2,3,6-tetrahydropyridinyl, piperazinyl, tetrahydro-2H-pyranyl, 3,6-dihydro-2H-pyranyl, tetrahydro-2H-thiopyranyl, 1,4-diazepanyl, 1,4-oxazepanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 2,5-diazabicyclo[2.2.1]heptanyl, morpholinyl, and the like) or C_3 - C_8 cycloalkyl, each optionally substituted with one or more $-Q_6'-T_6'$.

[0481] For example, the ring formed by R_7' and R_8' is selected from the group consisting of azetidiny, pyrrolidinyl, piperidinyl, morpholinyl, piperazinyl, and cyclohexenyl, each optionally substituted with one $-Q_6'-T_6'$.

[0482] For example, $-Q_6'-T_6'$ is oxo.

[0483] For example, T_6' is H, halo, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_3 - C_8 cycloalkyl, C_6 - C_{10} aryl, or 4 to 7-membered heterocycloalkyl.

[0484] For example, Q_6' is a bond and T_6' is C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, or 4 to 7-membered heterocycloalkyl.

[0485] For example, Q_6' is CO, $S(O)_2$, or $NHC(O)$; and T_6' is C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_3 - C_8 cycloalkyl, or 4 to 7-membered heterocycloalkyl.

[0486] For example, T_6' is C_1 - C_6 alkyl or C_1 - C_6 alkoxy, each optionally substituted with halo, hydroxyl, cyano, C_1 - C_6 alkoxy, amino, mono- C_1 - C_6 alkylamino, di- C_1 - C_6 alkylamino, or C_3 - C_8 cycloalkyl.

[0487] For example, Q_6' is C_1 - C_3 alkyl linker and T_6' is H or C_6 - C_{10} aryl.

[0488] For example, Q_6' is C_1 - C_3 alkyl linker and T_6' is C_3 - C_8 cycloalkyl, 4 to 7-membered heterocycloalkyl, or $S(O)_pR_p'$.

[0489] For example, each of R_p' and R_q' , independently, is C_1 - C_6 alkyl.

[0490] For example, R_6' is $-S(O)_bR_a'$ or azido, in which b' is 0, 1, or 2 and R_a' is C_1 - C_6 alkyl or C_3 - C_8 cycloalkyl; and Z_2 is $NR_7'R_8'$, in which R_7' is C_3 - C_8 cycloalkyl (e.g., cyclopentyl, cyclohexyl, or cycloheptyl) or 4 to 7-membered heterocycloalkyl (e.g., azetidiny, oxetanyl, thietanyl, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, triazolidinyl, tetrahydrofuranyl, piperidinyl, 1,2,3,6-tetrahydropyridinyl, piperazinyl, tetrahydro-2H-pyranyl, 3,6-dihydro-2H-pyranyl, tetrahydro-2H-thiopyranyl, 1,4-diazepanyl, 1,4-oxazepanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 2,5-diazabicyclo[2.2.1]heptanyl, and morpholinyl, and the like), each optionally substituted with one or more $-Q_5'-T_5'$ and R_8' is H or C_1 - C_6 alkyl (e.g., methyl, ethyl, n-propyl, i-propyl, butyl, or t-butyl).

[0491] For example, R_6' is halo (e.g., fluorine, chlorine, bromine, and iodine) and Z_2 is $NR_7'R_8'$ or $CR_7'R_8'R_{14}'$ wherein R_7' and R_8' , together with the atom to which they are attached, form a 4 to 11-membered heterocycloalkyl ring having 1 to 3 heteroatoms (e.g., azetidiny, oxetanyl, thietanyl, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, triazolidinyl, tetrahydrofuranyl, piperidinyl, 1,2,3,6-tetrahydropyridinyl, piperazinyl, tetrahydro-2H-pyranyl, 3,6-dihydro-2H-pyranyl, tetrahydro-2H-thiopyranyl, 1,4-diazepanyl, 1,4-oxazepanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 2,5-diazabicyclo[2.2.1]heptanyl, morpholinyl, and the like) or C_3 - C_8 cycloalkyl, each optionally substituted with one or more $-Q_6'-T_6'$.

[0492] For example, R_{13}' is H or methyl.


[0493] For example, R_{13}' is H.

[0494] For example, R_3' is H.

[0495] For example, each of R_5' , R_9' , and R_{10}' is H.

[0496] Other compounds suitable for the methods of the invention are described in PCT/US2012/026953, filed on Feb. 28, 2012; U.S. Provisional Application Ser. Nos. 61/474,821, filed on Apr. 13, 2011; 61/474,825, filed on Apr. 13, 2011; 61/499,595, filed on Jun. 21, 2011; and 61/505,676, filed on Jul. 8, 2011, the contents of which are hereby incorporated by reference in their entireties.

[0497] Exemplary EZH2 inhibitor compounds of the present invention are shown in Table 1. In the table below, each occurrence of



should be construed as





TABLE 1

Compound Number	Structure
1	<p>(“SAH”)</p>
2	<p>(“SAH”)</p>
3	<p>(“SAH”)</p>
4	<p>(“SAH”)</p>

TABLE 1-continued


Compound Number	Structure
5	<p>(“SAH”)</p>
6	<p>(“SAH”)</p>
7	<p>(“SAH”)</p>

TABLE 1-continued

Compound Number	Structure
8	

CCN(CC1CCOCC1)c2cc(cc(c2C)C(=O)NCc3cc(C)c(C)cn3C)c4ccc(cc4CNCC5CCOCC5)C6=CC=CC=C6

10



Chemical structure 10 is a complex molecule featuring a central benzene ring. This central ring is substituted with a methyl group at the 2-position, a 1-ethyl-4-(tetrahydro-2H-pyran-2-yl)pyrrolidin-3-yl group at the 3-position, and a 4-(2-(2,6-dimethyl-4-oxo-1,2,3,4-tetrahydropyridin-4-yl)ethyl)phenyl group at the 4-position. The 4-(2-(2,6-dimethyl-4-oxo-1,2,3,4-tetrahydropyridin-4-yl)ethyl)phenyl group consists of a phenyl ring connected via a methylene bridge to a nitrogen atom, which is part of a 7-membered ring containing another nitrogen atom.

TABLE 1-continued

Compound Number	Structure
11	

12

Chemical structure of a substituted benzamide derivative. The central benzene ring has a chlorine atom at the top position. Moving clockwise, there is a carbonyl group (C=O) at the 2-position, an amide group (-NH-CH₂-) at the 3-position, a nitrogen atom bonded to an ethyl group and a cyclohexyl ring at the 4-position, and a methyl group at the 5-position. The cyclohexyl ring is further substituted with a dimethylamino group (-N(CH₃)₂) at the 1-position.

13

Chemical structure 13 is a complex molecule. It features a central benzene ring substituted with three groups: a morpholine-4-ylmethyl group, a 2-ethyl-1-(cyclohexyl)pyrrolidin-1-yl group, and a 2-((2-methyl-4-oxo-1,2,3,4-tetrahydropyridin-3-yl)methyl)amino group.

TABLE 1-continued

Compound Number	Structure
14	
15	
16	

TABLE 1-continued

Compound Number	Structure
17	
18	
19	

TABLE 1-continued




Compound Number	Structure
20	
21	
22	

TABLE 1-continued




Compound Number	Structure
23	
24	
25	

TABLE 1-continued




Compound Number	Structure
26	
27	
28	

TABLE 1-continued




Compound Number	Structure
29	
30	
31	

TABLE 1-continued




Compound Number	Structure
32	
33	
34	

TABLE 1-continued







Compound Number	Structure
35	
36	
37	
38	

TABLE 1-continued



40



41






TABLE 1-continued



43



44





TABLE 1-continued

Compound Number	Structure
-----------------	-----------

45



46






TABLE 1-continued

Compound Number	Structure
-----------------	-----------

47



48



49






TABLE 1-continued

Compound Number	Structure
--------------------	-----------

50



51



52






TABLE 1-continued

Compound Number	Structure
--------------------	-----------

53



54



55





TABLE 1-continued

Compound Number	Structure
--------------------	-----------

56



57





TABLE 1-continued

Compound Number	Structure
--------------------	-----------

58



59




TABLE 1-continued

Compound Number	Structure
60	
61	
62	

[0498] Unless otherwise indicated, the term “substituted” means substituted by one or more defined groups. In the case where groups may be selected from a number of alternative groups the selected groups may be the same or different.

[0499] The term “independently” means that where more than one substituent is selected from a number of possible substituents, those substituents may be the same or different.

[0500] An “effective amount” means that amount of a drug or pharmaceutical agent that will elicit the biological or medical response of a tissue, system, animal or human that is being sought, for instance, by a researcher or clinician. Furthermore, the term “therapeutically effective amount” means any amount which, as compared to a corresponding subject who

has not received such amount, results in improved treatment, healing, prevention, or amelioration of a disease, disorder, or side effect, or a decrease in the rate of advancement of a disease or disorder. The term also includes within its scope amounts effective to enhance normal physiological function.

[0501] As used herein, “alkyl”, “C₁, C₂, C₃, C₄, C₅ or C₆ alkyl” or “C₁-C₆ alkyl” is intended to include C₁, C₂, C₃, C₄, C₅ or C₆ straight chain (linear) saturated aliphatic hydrocarbon groups and C₃, C₄, C₅ or C₆ branched saturated aliphatic hydrocarbon groups. For example, C₁-C₆ alkyl is intended to include C₁, C₂, C₃, C₄, C₅ and C₆ alkyl groups. Examples of alkyl include, moieties having from one to six carbon atoms, such as, but not limited to, methyl, ethyl, n-propyl, i-propyl, n-butyl, s-butyl, t-butyl, n-pentyl, s-pentyl or n-hexyl.

[0502] In certain embodiments, a straight chain or branched alkyl has six or fewer carbon atoms (e.g., C₁-C₆ for straight chain, C₃-C₆ for branched chain), and in another embodiment, a straight chain or branched alkyl has four or fewer carbon atoms.

[0503] As used herein, the term “cycloalkyl” refers to a saturated or unsaturated nonaromatic hydrocarbon mono- or multi-ring (e.g., fused, bridged; or spiro rings) system having 3 to 30 carbon atoms (e.g., C₃-C₁₀). Examples of cycloalkyl include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, and adamantyl. The term “heterocycloalkyl” refers to a saturated or unsaturated nonaromatic 3-8 membered monocyclic, 7-12 membered bicyclic (fused, bridged, or spiro rings), or 11-14 membered tricyclic ring system (fused, bridged, or spiro rings) having one or more heteroatoms (such as O, N, S, or Se), unless specified otherwise. Examples of heterocycloalkyl groups include, but are not limited to, piperidinyl, piperazinyl, pyrrolidinyl, dioxanyl, tetrahydrofuranyl, isoindolinyl, indolinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, triazolidinyl, tetrahydrofuranyl, oxiranyl, azetidiny, oxetanyl, thietanyl, 1,2,3,6-tetrahydropyridinyl, tetrahydropyran, dihydropyran, pyran, morpholinyl, 1,4-diazepanyl, 1,4-oxazepanyl, 2-oxa-5-azabicyclo[2.2.1]heptanyl, 2,5-diazabicyclo[2.2.1]heptanyl, and the like.

[0504] The term “optionally substituted alkyl” refers to unsubstituted alkyl or alkyl having designated substituents replacing one or more hydrogen atoms on one or more carbons of the hydrocarbon backbone. Such substituents can include, for example, alkyl, alkenyl, alkynyl, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxy, carbonyloxy, aryloxy, aryloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxy, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxy, phosphate, phosphonate, phosphinate, amino (including alkylamino, dialkylamino, arylamino, diarylamino and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfinyl, alkylthio, arylthio, thiocarbonyloxy, sulfates, alkylsulfinyl, sulfonate, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety.

[0505] An “arylalkyl” or an “aralkyl” moiety is an alkyl substituted with an aryl (e.g., phenylmethyl (benzyl)). An “alkylaryl” moiety is an aryl substituted with an alkyl (e.g., methylphenyl).

[0506] As used herein, “alkyl linker” is intended to include C₁, C₂, C₃, C₄, C₅ or C₆ straight chain (linear) saturated divalent aliphatic hydrocarbon groups and C₃, C₄, C₅ or C₆

branched saturated aliphatic hydrocarbon groups. For example, C₁-C₆ alkyl linker is intended to include C₁, C₂, C₃, C₄, C₅ and C₆ alkyl linker groups. Examples of alkyl linker include, moieties having from one to six carbon atoms, such as, but not limited to, methyl (—CH₂—), ethyl (—CH₂CH₂—), n-propyl (—CH₂CH₂CH₂—), i-propyl (—CHCH₃CH₂—), n-butyl (—CH₂CH₂CH₂CH₂—), s-butyl (—CHCH₃CH₂CH₂—), i-butyl (—C(CH₃)₂CH₂—), n-pentyl (—CH₂CH₂CH₂CH₂CH₂—), s-pentyl (—CHCH₃CH₂CH₂CH₂—) or n-hexyl (—CH₂CH₂CH₂CH₂CH₂CH₂—).

[0507] “Alkenyl” includes unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but that contain at least one double bond. For example, the term “alkenyl” includes straight chain alkenyl groups (e.g., ethenyl, propenyl, butenyl, pentenyl, hexenyl, heptenyl, octenyl, nonenyl, decenyl), and branched alkenyl groups. In certain embodiments, a straight chain or branched alkenyl group has six or fewer carbon atoms in its backbone (e.g., C₂-C₆ for straight chain, C₃-C₆ for branched chain). The term “C₂-C₆” includes alkenyl groups containing two to six carbon atoms. The term “C₃-C₆” includes alkenyl groups containing three to six carbon atoms.

[0508] The term “optionally substituted alkenyl” refers to unsubstituted alkenyl or alkenyl having designated substituents replacing one or more hydrogen atoms on one or more hydrocarbon backbone carbon atoms. Such substituents can include, for example, alkyl, alkenyl, alkynyl, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxy carbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxy, phosphate, phosphonate, phosphinato, amino (including alkylamino, dialkylamino, arylamino, diarylamino and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonate, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety.

[0509] “Alkynyl” includes unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but which contain at least one triple bond. For example, “alkynyl” includes straight chain alkynyl groups (e.g., ethynyl, propynyl, butynyl, pentynyl, hexynyl, heptynyl, octynyl, nonynyl, decynyl), and branched alkynyl groups. In certain embodiments, a straight chain or branched alkynyl group has six or fewer carbon atoms in its backbone (e.g., C₂-C₆ for straight chain, C₃-C₆ for branched chain). The term “C₂-C₆” includes alkynyl groups containing two to six carbon atoms. The term “C₃-C₆” includes alkynyl groups containing three to six carbon atoms.

[0510] The term “optionally substituted alkynyl” refers to unsubstituted alkynyl or alkynyl having designated substituents replacing one or more hydrogen atoms on one or more hydrocarbon backbone carbon atoms. Such substituents can include, for example, alkyl, alkenyl, alkynyl, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxy carbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxy, phosphate, phosphonate, phosphinato, amino (including alkylamino, dialkylamino, arylamino, diarylamino and alkylarylamino), acylamino (including alkylcarbony-

lamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonate, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety.

[0511] Other optionally substituted moieties (such as optionally substituted cycloalkyl, heterocycloalkyl, aryl, or heteroaryl) include both the unsubstituted moieties and the moieties having one or more of the designated substituents. For example, substituted heterocycloalkyl includes those substituted with one or more alkyl groups, such as 2,2,6,6-tetramethyl-piperidiny and 2,2,6,6-tetramethyl-1,2,3,6-tetrahydropyridiny.

[0512] “Aryl” includes groups with aromaticity, including “conjugated,” or multicyclic systems with at least one aromatic ring and do not contain any heteroatom in the ring structure. Examples include phenyl, benzyl, 1,2,3,4-tetrahydronaphthalenyl, etc.

[0513] “Heteroaryl” groups are aryl groups, as defined above, except having from one to four heteroatoms in the ring structure, and may also be referred to as “aryl heterocycles” or “heteroaromatics.” As used herein, the term “heteroaryl” is intended to include a stable 5-, 6-, or 7-membered monocyclic or 7-, 8-, 9-, 10-, 11- or 12-membered bicyclic aromatic heterocyclic ring which consists of carbon atoms and one or more heteroatoms, e.g., 1 or 1-2 or 1-3 or 1-4 or 1-5 or 1-6 heteroatoms, or e.g., 2, 3, 4, 5, or 6 heteroatoms, independently selected from the group consisting of nitrogen, oxygen and sulfur. The nitrogen atom may be substituted or unsubstituted (i.e., N or NR wherein R is H or other substituents, as defined). The nitrogen and sulfur heteroatoms may optionally be oxidized (i.e., N→O and S(O)_p, where p=1 or 2). It is to be noted that total number of S and O atoms in the aromatic heterocycle is not more than 1.

[0514] Examples of heteroaryl groups include pyrrole, furan, thiophene, thiazole, isothiazole, imidazole, triazole, tetrazole, pyrazole, oxazole, isoxazole, pyridine, pyrazine, pyridazine, pyrimidine, and the like.

[0515] Furthermore, the terms “aryl” and “heteroaryl” include multicyclic aryl and heteroaryl groups, e.g., tricyclic, bicyclic, e.g., naphthalene, benzoxazole, benzodioxazole, benzothiazole, benzoimidazole, benzothienophene, methylenedioxyphenyl, quinoline, isoquinoline, naphthrydine, indole, benzofuran, purine, benzofuran, deazapurine, indolizirine.

[0516] In the case of multicyclic aromatic rings, only one of the rings needs to be aromatic (e.g., 2,3-dihydroindole), although all of the rings may be aromatic (e.g., quinoline). The second ring can also be fused or bridged.

[0517] The cycloalkyl, heterocycloalkyl, aryl, or heteroaryl ring can be substituted at one or more ring positions (e.g., the ring-forming carbon or heteroatom such as N) with such substituents as described above, for example, alkyl, alkenyl, alkynyl, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, arylcarbonyloxy, alkoxy carbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, alkylaminocarbonyl, aralkylaminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, aralkylcarbonyl, alkenylcarbonyl, alkoxy carbonyl, aminocarbonyl, alkylthiocarbonyl, phosphate, phosphonate, phosphinato, amino (including alkylamino, dialkylamino, arylamino, diarylamino and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonate, sulfamoyl, sulfonamido, nitro, trifluoromethyl,

cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety. Aryl and heteroaryl groups can also be fused or bridged with alicyclic or heterocyclic rings, which are not aromatic so as to form a multicyclic system (e.g., tetralin, methylenedioxyphenyl).

[0518] As used herein, “carbocycle” or “carbocyclic ring” is intended to include any stable monocyclic, bicyclic or tricyclic ring having the specified number of carbons, any of which may be saturated, unsaturated, or aromatic. Carbocycle includes cycloalkyl and aryl. For example, a C₃-C₁₄ carbocycle is intended to include a monocyclic, bicyclic or tricyclic ring having 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13 or 14 carbon atoms. Examples of carbocycles include, but are not limited to, cyclopropyl, cyclobutyl, cyclobutenyl, cyclopentyl, cyclopentenyl, cyclohexyl, cycloheptenyl, cycloheptyl, cycloheptenyl, adamantyl, cyclooctyl, cyclooctenyl, cyclooctadienyl, fluorenyl, phenyl, naphthyl, indanyl, adamantyl and tetrahydronaphthyl. Bridged rings are also included in the definition of carbocycle, including, for example, [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane and [2.2.2]bicyclooctane. A bridged ring occurs when one or more carbon atoms link two non-adjacent carbon atoms. In one embodiment, bridge rings are one or two carbon atoms. It is noted that a bridge always converts a monocyclic ring into a tricyclic ring. When a ring is bridged, the substituents recited for the ring may also be present on the bridge. Fused (e.g., naphthyl, tetrahydronaphthyl) and Spiro rings are also included.

[0519] As used herein, “heterocycle” or “heterocyclic group” includes any ring structure (saturated, unsaturated, or aromatic) which contains at least one ring heteroatom (e.g., N, O or S). Heterocycle includes heterocycloalkyl and heteroaryl. Examples of heterocycles include, but are not limited to, morpholine, pyrrolidine, tetrahydrothiophene, piperidine, piperazine, oxetane, pyran, tetrahydropyran, azetidine, and tetrahydrofuran.

[0520] Examples of heterocyclic groups include, but are not limited to, acridinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzoxazolyl, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazolyl, carbazolyl, 4aH-carbazolyl, carbolinyl, chromanyl, chromenyl, cinnolinyl, decahydroquinolinyl, 2H,6H-1,5,2-dithiazinyl, dihydrofuro[2,3-b]tetrahydrofuran, furanyl, furazanyl, imidazolidinyl, imidazolyl, imidazolyl, 1H-indazolyl, indolyl, indolinyl, indolizyl, indolyl, 3H-indolyl, isatinoyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindolinyl, isoindolyl, isoquinolinyl, isothiazolyl, isoxazolyl, methylenedioxyphenyl, morpholinyl, naphthyridinyl, octahydroisoquinolinyl, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,4-oxadiazolyl (4H)-one, oxazolidinyl, oxazolyl, oxindolyl, pyrimidinyl, phenanthridinyl, phenanthrolinyl, phenazinyl, phenothiazinyl, phenoxathinyl, phenoxazinyl, phthalazinyl, piperazinyl, piperidinyl, piperidonyl, 4-piperidonyl, piperonyl, pteridinyl, purinyl, pyranyl, pyrazinyl, pyrazolidinyl, pyrazolinyl, pyrazolyl, pyridazinyl, pyridooxazole, pyridoimidazole, pyridothiazole, pyridinyl, pyridyl, pyrimidinyl, pyrrolidinyl, pyrrolinyl, 2H-pyrrolyl, pyrrolyl, quinazolinyl, quinolinyl, 4H-quinolinyl, quinoxalinyl, quinuclidinyl, tetrahydrofuranyl, tetrahydroisoquinolinyl, tetrahydroquinolinyl, tetrazolyl, 6H-1,2,5-thiadiazinyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, thianthrenyl, thiazolyl, thienyl, thienothiazolyl, thienooxazolyl, thienoimi-

dazolyl, thiophenyl, triazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl and xanthenyl.

[0521] The term “substituted,” as used herein, means that any one or more hydrogen atoms on the designated atom is replaced with a selection from the indicated groups, provided that the designated atom’s normal valency is not exceeded, and that the substitution results in a stable compound. When a substituent is pxo or keto (i.e., =O), then 2 hydrogen atoms on the atom are replaced. Keto substituents are not present on aromatic moieties. Ring double bonds, as used herein, are double bonds that are formed between two adjacent ring atoms (e.g., C=C, C=N or N=N). “Stable compound” and “stable structure” are meant to indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

[0522] When a bond to a substituent is shown to cross a bond connecting two atoms in a ring, then such substituent may be bonded to any atom in the ring. When a substituent is listed without indicating the atom via which such substituent is bonded to the rest of the compound of a given formula, then such substituent may be bonded via any atom in such formula. Combinations of substituents and/or variables are permissible, but only if such combinations result in stable compounds.

[0523] When any variable (e.g., R₁) occurs more than one time in any constituent or formula for a compound, its definition at each occurrence is independent of its definition at every other occurrence. Thus, for example, if a group is shown to be substituted with 0-2 R₁ moieties, then the group may optionally be substituted with up to two R₁ moieties and R₁ at each occurrence is selected independently from the definition of R₁. Also, combinations of substituents and/or variables are permissible, but only if such combinations result in stable compounds.

[0524] The term “hydroxy” or “hydroxyl” includes groups with an —OH or —O[−].

[0525] As used herein, “halo” or “halogen” refers to fluoro, chloro, bromo and iodo. The term “perhalogenated” generally refers to a moiety wherein all hydrogen atoms are replaced by halogen atoms. The term “haloalkyl” or “haloalkoxy” refers to an alkyl or alkoxy substituted with one or more halogen atoms.

[0526] The term “carbonyl” includes compounds and moieties which contain a carbon connected with a double bond to an oxygen atom. Examples of moieties containing a carbonyl include, but are not limited to, aldehydes, ketones, carboxylic acids, amides, esters, anhydrides, etc.

[0527] The term “carboxyl” refers to —COOH or its C₁-C₆ alkyl ester.

[0528] “Acyl” includes moieties that contain the acyl radical (R—C(O)—) or a carbonyl group. “Substituted acyl” includes acyl groups where one or more of the hydrogen atoms are replaced by, for example, alkyl groups, alkynyl groups, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy, aryloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxy, phosphate, phosphonate, phosphinate, amino (including alkylamino, dialkylamino, arylamino, diarylamino and alkylaryl amino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfanyl, sulfonate, sulfamoyl, sulfona-

mido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety.

[0529] “Aroyl” includes moieties with an aryl or heteroaromatic moiety bound to a carbonyl group. Examples of aroyl groups include phenylcarboxy, naphthyl carboxy, etc.

[0530] “Alkoxyalkyl,” “alkylaminoalkyl,” and “thioalkoxyalkyl” include alkyl groups, as described above, wherein oxygen, nitrogen, or sulfur atoms replace one or more hydrocarbon backbone carbon atoms.

[0531] The term “alkoxy” or “alkoxy” includes substituted and unsubstituted alkyl, alkenyl and alkynyl groups covalently linked to an oxygen atom. Examples of alkoxy groups or alkoxy radicals include, but are not limited to, methoxy, ethoxy, isopropoxy, propoxy, butoxy and pentoxy groups. Examples of substituted alkoxy groups include halogenated alkoxy groups. The alkoxy groups can be substituted with groups such as alkenyl, alkynyl, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxy carbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxy, phosphate, phosphonate, phosphinato, amino (including alkylamino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonate, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moieties. Examples of halogen substituted alkoxy groups include, but are not limited to, fluoromethoxy, difluoromethoxy, trifluoromethoxy, chloromethoxy, dichloromethoxy and trichloromethoxy.

[0532] The term “ether” or “alkoxy” includes compounds or moieties which contain an oxygen bonded to two carbon atoms or heteroatoms. For example, the term includes “alkoxyalkyl,” which refers to an alkyl, alkenyl, or alkynyl group covalently bonded to an oxygen atom which is covalently bonded to an alkyl group.

[0533] The term “ester” includes compounds or moieties which contain a carbon or a heteroatom bound to an oxygen atom which is bonded to the carbon of a carbonyl group. The term “ester” includes alkoxy carbonyl groups such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, pentoxycarbonyl, etc.

[0534] The term “thioalkyl” includes compounds or moieties which contain an alkyl group connected with a sulfur atom. The thioalkyl groups can be substituted with groups such as alkyl, alkenyl, alkynyl, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxy carbonyloxy, aryloxy carbonyloxy, carboxylate, carboxylic acid, alkylcarbonyl, arylcarbonyl, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxy, amino (including alkylamino, dialkylamino, arylamino, diarylamino and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonate, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moieties.

[0535] The term “thiocarbonyl” or “thiocarboxy” includes compounds and moieties which contain a carbon connected with a double bond to a sulfur atom.

[0536] The term “thioether” includes moieties which contain a sulfur atom bonded to two carbon atoms or heteroatom

oms. Examples of thioethers include, but are not limited to althioalkyls, althioalkenyls, and althioalkynyls. The term “althioalkyls” include moieties with an alkyl, alkenyl, or alkynyl group bonded to a sulfur atom which is bonded to an alkyl group. Similarly, the term “althioalkenyls” refers to moieties wherein an alkyl, alkenyl or alkynyl group is bonded to a sulfur atom which is covalently bonded to an alkenyl group; and althioalkynyls” refers to moieties wherein an alkyl, alkenyl or alkynyl group is bonded to a sulfur atom which is covalently bonded to an alkynyl group.

[0537] As used herein, “amine” or “amino” refers to unsubstituted or substituted —NH_2 . “Alkylamino” includes groups of compounds wherein nitrogen of —NH_2 is bound to at least one alkyl group. Examples of alkylamino groups include benzylamino, methylamino, ethylamino, phenethylamino, etc. “Dialkylamino” includes groups wherein the nitrogen of —NH_2 is bound to at least two additional alkyl groups. Examples of dialkylamino groups include, but are not limited to, dimethylamino and diethylamino. “Arylamino” and “diarylamino” include groups wherein the nitrogen is bound to at least one or two aryl groups, respectively. “Aminoaryl” and “aminoaryloxy” refer to aryl and aryloxy substituted with amino. “Alkylarylamino,” “alkylaminoaryl” or “arylaminoalkyl” refers to an amino group which is bound to at least one alkyl group and at least one aryl group. “Alkaminoalkyl” refers to an alkyl, alkenyl, or alkynyl group bound to a nitrogen atom which is also bound to an alkyl group. “Acylamino” includes groups wherein nitrogen is bound to an acyl group. Examples of acylamino include, but are not limited to, alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido groups.

[0538] The term “amide” or “aminocarboxy” includes compounds or moieties that contain a nitrogen atom that is bound to the carbon of a carbonyl or a thiocarbonyl group. The term includes “alkaminocarboxy” groups that include alkyl, alkenyl or alkynyl groups bound to an amino group which is bound to the carbon of a carbonyl or thiocarbonyl group. It also includes “arylamino carboxy” groups that include aryl or heteroaryl moieties bound to an amino group that is bound to the carbon of a carbonyl or thiocarbonyl group. The terms “alkylaminocarboxy”; “alkenylaminocarboxy”, “alkynylaminocarboxy” and “arylamino carboxy” include moieties wherein alkyl, alkenyl, alkynyl and aryl moieties, respectively, are bound to a nitrogen atom which is in turn bound to the carbon of a carbonyl group. Amides can be substituted with substituents such as straight chain alkyl, branched alkyl, cycloalkyl, aryl, heteroaryl or heterocycle. Substituents on amide groups may be further substituted.

[0539] The term “optionally” means that the subsequently described event(s) may or may not occur, and includes both event(s), which occur, and events that do not occur.

[0540] Herein, the term “pharmaceutically-acceptable salts” refers to salts that retain the desired biological activity of the subject compound and exhibit minimal undesired toxicological effects. These pharmaceutically-acceptable salts may be prepared in situ during the final isolation and purification of the compound, or by separately reacting the purified compound in its free acid or free base form with a suitable base or acid, respectively.

[0541] By the term “co-administering” and derivatives thereof as used herein is meant either simultaneous administration or any manner of separate sequential administration of one or more additional pharmaceutically active compounds, whether for treating cancer, the side effects of cancer or

cancer therapy, or some other disease. Preferably, if the administration is not simultaneous, the compounds are administered in a close time proximity to each other. Furthermore, it does not matter if the compounds are administered in the same dosage form, e.g. one compound may be administered topically and another compound may be administered orally.

[0542] The compounds of Formulae (I)-(III) include the compounds themselves, as well as their salts, their solvates, their N-oxides, and their prodrugs, if applicable. In certain embodiments, compounds according to Formulae (I)-(III) may contain an acidic functional group, one acidic enough to form salts. Representative salts include pharmaceutically acceptable metal salts such as sodium, potassium, lithium, calcium, magnesium, aluminum, and zinc salts; carbonates and bicarbonates of a pharmaceutically-acceptable metal salts such as sodium, potassium, lithium, calcium, magnesium, aluminum, and zinc; pharmaceutically-acceptable organic primary, secondary, and tertiary amines including aliphatic amines, aromatic amines, aliphatic diamines, and hydroxy alkylamines such as methylamine, ethylamine, 2-hydroxyethylamine, diethylamine, triethylamine, ethylenediamine, ethanolamine, diethanolamine, and cyclohexylamine.

[0543] In certain embodiments, compounds according to Formulae (I)-(III) may contain a basic functional group and are therefore capable of forming pharmaceutically-acceptable acid addition salts by treatment with a suitable acid. Suitable acids include pharmaceutically-acceptable inorganic acids and pharmaceutically-acceptable organic acids. Representative pharmaceutically acceptable acid addition salts include hydrochloride, hydrobromide, nitrate, methylnitrate, sulfate, bisulfate, sulfamate, phosphate, acetate, hydroxyacetate, phenylacetate, propionate, butyrate, isobutyrate, valerate, maleate, hydroxymaleate, acrylate, fumarate, malate, tartrate, citrate, salicylate, p-aminosalicylate, glycolate, lactate, heptanoate, phthalate, oxalate, succinate, benzoate, o-acetoxybenzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, mandelate, tannate, formate, stearate, ascorbate, palmitate, oleate, pyruvate, pantoate, malonate, laurate, glutarate, glutamate, estolate, methanesulfonate (mesylate), ethanesulfonate (esylate), 2-hydroxyethanesulfonate, benzenesulfonate (besylate), p-aminobenzenesulfonate, p-toluenesulfonate (tosylate) and naphthalene-2-sulfonate.

[0544] Compounds of the present invention that contain nitrogens can be converted to N-oxides by treatment with an oxidizing agent (e.g., 3-chloroperoxybenzoic acid (mCPBA) and/or hydrogen peroxides) to afford other compounds of the present invention. Thus, all shown and claimed nitrogen-containing compounds are considered, when allowed by valency and structure, to include both the compound as shown and its N-oxide derivative (which can be designated as $N \rightarrow O$ or $N^+ - O^-$). Furthermore, in other instances, the nitrogens in the compounds of the present invention can be converted to N-hydroxy or N-alkoxy compounds. For example, N-hydroxy compounds can be prepared by oxidation of the parent amine by an oxidizing agent such as m-CPBA. All shown and claimed nitrogen-containing compounds are also considered, when allowed by valency and structure, to cover both the compound as shown and its N-hydroxy (i.e., $N-OH$) and N-alkoxy (i.e., $N-OR$, wherein R is substituted or unsubstituted C_1-C_6 alkyl, C_1-C_6 alkenyl, C_1-C_6 alkynyl, 3-14-membered carbocycle or 3-14-membered heterocycle) derivatives.

[0545] In the present specification, the structural formula of the compound represents a certain isomer for convenience in some cases, but the present invention includes all isomers, such as geometrical isomers, optical isomers based on an asymmetrical carbon, stereoisomers, tautomers, and the like. In addition, a crystal polymorphism may be present for the compounds represented by the formula. It is noted that any crystal form, crystal form mixture, or anhydride or hydrate thereof is included in the scope of the present invention. Furthermore, so-called metabolite which is produced by degradation of the present compound in vivo is included in the scope of the present invention.

[0546] "Isomerism" means compounds that have identical molecular formulae but differ in the sequence of bonding of their atoms or in the arrangement of their atoms in space. Isomers that differ in the arrangement of their atoms in space are termed "stereoisomers." Stereoisomers that are not mirror images of one another are termed "diastereoisomers," and stereoisomers that are non-superimposable mirror images of each other are termed "enantiomers" or sometimes optical isomers. A mixture containing equal amounts of individual enantiomeric forms of opposite chirality is termed a "racemic mixture."

[0547] A carbon atom bonded to four nonidentical substituents is termed a "chiral center."

[0548] "Chiral isomer" means a compound with at least one chiral center. Compounds with more than one chiral center may exist either as an individual diastereomer or as a mixture of diastereomers, termed "diastereomeric mixture." When one chiral center is present, a stereoisomer may be characterized by the absolute configuration (R or S) of that chiral center. Absolute configuration refers to the arrangement in space of the substituents attached to the chiral center. The substituents attached to the chiral center under consideration are ranked in accordance with the Sequence Rule of Cahn, Ingold and Prelog. (Cahn et al., *Angew. Chem. Inter. Edit.* 1966, 5, 385; errata 511; Cahn et al., *Angew. Chem.* 1966, 78, 413; Cahn and Ingold, *J. Chem. Soc.* 1951 (London), 612; Cahn et al., *Experientia* 1956, 12, 81; Cahn, *J. Chem. Educ.* 1964, 41, 116).

[0549] "Geometric isomer" means the diastereomers that owe their existence to hindered rotation about double bonds or a cycloalkyl linker (e.g., 1,3-cyclobutyl). These configurations are differentiated in their names by the prefixes cis and trans, or Z and E, which indicate that the groups are on the same or opposite side of the double bond in the molecule according to the Cahn-Ingold-Prelog rules.


[0550] It is to be understood that the compounds of the present invention may be depicted as different chiral isomers or geometric isomers. It should also be understood that when compounds have chiral isomeric or geometric isomeric forms, all isomeric forms are intended to be included in the scope of the present invention, and the naming of the compounds does not exclude any isomeric forms.

[0551] Furthermore, the structures and other compounds discussed in this invention include all atropic isomers thereof. "Atropic isomers" are a type of stereoisomer in which the atoms of two isomers are arranged differently in space. Atropic isomers owe their existence to a restricted rotation caused by hindrance of rotation of large groups about a central bond. Such atropic isomers typically exist as a mixture, however as a result of recent advances in chromatography techniques, it has been possible to separate mixtures of two atropic isomers in select cases.

[0552] “Tautomer” is one of two or more structural isomers that exist in equilibrium and is readily converted from one isomeric form to another. This conversion results in the formal migration of a hydrogen atom accompanied by a switch of adjacent conjugated double bonds. Tautomers exist as a mixture of a tautomeric set in solution. In solutions where tautomerization is possible, a chemical equilibrium of the tautomers will be reached. The exact ratio of the tautomers depends on several factors, including temperature, solvent and pH. The concept of tautomers that are interconvertible by tautomerizations is called tautomerism.

[0553] Of the various types of tautomerism that are possible, two are commonly observed. In keto-enol tautomerism a simultaneous shift of electrons and a hydrogen atom occurs. Ring-chain tautomerism arises as a result of the aldehyde group (—CHO) in a sugar chain molecule reacting with one of the hydroxy groups (—OH) in the same molecule to give it a cyclic (ring-shaped) form as exhibited by glucose.

[0554] Common tautomeric pairs are: ketone-enol, amide-nitrile, lactam-lactim, amide-imidic acid tautomerism in heterocyclic rings (e.g., in nucleobases such as guanine, thymine and cytosine), imine-enamine and enamine-enamine. An example of keto-enol equilibria is between pyridin-2(1H)-ones and the corresponding pyridin-2-ols, as shown below.



[0555] It is to be understood that the compounds of the present invention may be depicted as different tautomers. It should also be understood that when compounds have tautomeric forms, all tautomeric forms, including mixtures thereof, are intended to be included in the scope of the present invention, and the naming of the compounds does not exclude any tautomer form.

[0556] The compounds of Formulae (I)-(III) may be prepared in crystalline or non-crystalline form, and, if crystalline, may optionally be solvated, e.g., as the hydrate. This invention includes within its scope stoichiometric solvates (e.g., hydrates) as well as compounds containing variable amounts of solvent (e.g., water).

[0557] Certain of the compounds described herein may contain one or more chiral atoms, or may otherwise be capable of existing as two enantiomers. The compounds claimed below include mixtures of enantiomers as well as purified enantiomers or enantiomerically enriched mixtures. Also included within the scope of the invention are the individual isomers of the compounds represented by Formulae (I)-(III), or claimed below, as well as any wholly or partially equilibrated mixtures thereof. The present invention also covers the individual isomers of the claimed compounds as mixtures with isomers thereof in which one or more chiral centers are inverted.

[0558] Where there are different isomeric forms they may be separated or resolved one from the other by conventional methods, or any given isomer may be obtained by conventional synthetic methods or by stereospecific or asymmetric syntheses.

[0559] While it is possible that, for use in therapy, a compound of Formulae (I)-(III), as well as salts, solvates, N-oxides, and the like, may be administered as a neat preparation, i.e., no additional carrier, the more usual practice is to present the active ingredient confectioned with a carrier or diluent. Accordingly, the invention further provides pharmaceutical compositions, which includes a compound of Formulae (I)-(III) and salts, solvates and the like, and one or more pharmaceutically acceptable carriers, diluents, or excipients. The compounds of Formulae (I)-(III) and salts, solvates, N-oxides, etc. are as described above. The carrier(s), diluent, or excipient(s) must be acceptable in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof. In accordance with another aspect of the invention there is also provided a process for the preparation of a pharmaceutical formulation including admixing a compound of the Formulae (I)-(III), or salts, solvates, N-oxides, etc. with one or more pharmaceutically acceptable carriers, diluents or excipients.

[0560] It will be appreciated by those skilled in the art that certain protected derivatives of compounds of Formulae (I)-(III), which may be made prior to a final deprotection stage, may not possess pharmacological activity as such, but may, in certain instances, be administered orally or parenterally and thereafter metabolized in the body to form compounds of the invention which are pharmacologically active. Such derivatives may therefore be described as “prodrugs”. Further, certain compounds of the invention may act as prodrugs of other compounds of the invention. All protected derivatives and prodrugs of compounds of the invention are included within the scope of the invention. It will further be appreciated by those skilled in the art, that certain moieties, known to those skilled in the art as “pro-moieties” may be placed on appropriate functionalities when such functionalities are present within compounds of the invention. Preferred prodrugs for compounds of the invention include: esters, carbonate esters, hemi-esters, phosphate esters, nitro esters, sulfate esters, sulfoxides, amides, carbamates, azo-compounds, phosphamides, glycosides, ethers, acetals, and ketals.

[0561] Additionally, the compounds of the present invention, for example, the salts of the compounds, can exist in either hydrated or unhydrated (the anhydrous) form or as solvates with other solvent molecules. Nonlimiting examples of hydrates include monohydrates, dihydrates, etc. Nonlimiting examples of solvates include ethanol solvates, acetone solvates, etc.

[0562] “Solvate” means solvent addition forms that contain either stoichiometric or non stoichiometric amounts of solvent. Some compounds have a tendency to trap a fixed molar ratio of solvent molecules in the crystalline solid state, thus forming a solvate. If the solvent is water the solvate formed is a hydrate; and if the solvent is alcohol, the solvate formed is an alcoholate. Hydrates are formed by the combination of one or more molecules of water with one molecule of the substance in which the water retains its molecular state as H_2O .


[0563] As used herein, the term “analog” refers to a chemical compound that is structurally similar to another but differs slightly in composition (as in the replacement of one atom by an atom of a different element or in the presence of a particular functional group, or the replacement of one functional group by another functional group). Thus, an analog is a compound that is similar or comparable in function and appearance, but not in structure or origin to the reference compound.

[0564] As defined herein, the term “derivative” refers to compounds that have a common core structure, and are substituted with various groups as described herein. For example, all of the compounds represented by Formula (I) are aryl- or heteroaryl-substituted benzene compounds, and have Formula (I) as a common core.

[0565] The term “bioisostere” refers to a compound resulting from the exchange of an atom or of a group of atoms with another, broadly similar, atom or group of atoms. The objective of a bioisosteric replacement is to create a new compound with similar biological properties to the parent compound. The bioisosteric replacement may be physicochemically or topologically based. Examples of carboxylic acid bioisosteres include, but are not limited to, acyl sulfonimides; tetrazoles, sulfonates and phosphonates. See, e.g., Patani and LaVoie, *Chem. Rev.* 96, 3147-3176, 1996.


[0566] The present invention is intended to include all isotopes of atoms occurring in the present compounds. Isotopes include those atoms having the same atomic number but different mass numbers. By way of general example and without limitation, isotopes of hydrogen include tritium and deuterium, and isotopes of carbon include C-13 and C-14.

[0567] Such compounds also include Compound 75



and pharmaceutically acceptable salts thereof.

[0568] The invention further includes a pharmaceutical composition comprising Compound 75



or a pharmaceutically acceptable salt thereof.

[0569] An EZH2 antagonist and optionally other therapeutics can be administered per se (neat) or in the form of a

pharmaceutically acceptable salt. When used in medicine the salts should be pharmaceutically acceptable, but non-pharmaceutically acceptable salts can conveniently be used to prepare pharmaceutically acceptable salts thereof.

[0570] Compounds useful in accordance with the invention may be provided as salts with pharmaceutically compatible counterions (i.e., pharmaceutically acceptable salts). A “pharmaceutically acceptable salt” means any non-toxic salt that, upon administration to a recipient, is capable of providing, either directly or indirectly, a compound or a prodrug of a compound useful in accordance with this invention. A “pharmaceutically acceptable counterion” is an ionic portion of a salt that is not toxic when released from the salt upon administration to a subject. Pharmaceutically compatible salts may be formed with many acids, including but not limited to hydrochloric, sulfuric, acetic, lactic, tartaric, malic, and succinic acids. Salts tend to be more soluble in water or other protic solvents than their corresponding free base forms. The present invention includes the use of such salts.

[0571] Pharmaceutically acceptable acid addition salts include those formed with mineral acids such as hydrochloric acid and hydrobromic acid, and also those formed with organic acids such as maleic acid. For example, acids commonly employed to form pharmaceutically acceptable salts include inorganic acids such as hydrogen bisulfide, hydrochloric, hydrobromic, hydroiodic, sulfuric and phosphoric acid, as well as organic acids such as para-toluenesulfonic, salicylic, tartaric, bitartaric, ascorbic, maleic, besylic, fumaric, gluconic, glucuronic, formic, glutamic, methanesulfonic, ethanesulfonic, benzenesulfonic, lactic, oxalic, para-bromophenylsulfonic, carbonic, succinic, citric, benzoic and acetic acid, and related inorganic and organic acids. Such pharmaceutically acceptable salts thus include sulfate, pyrosulfate, bisulfate, sulfite, bisulfite, phosphate, monohydrogenphosphate, dihydrogenphosphate, metaphosphate, pyrophosphate, chloride, bromide, iodide, acetate, propionate, decanoate, caprylate, acrylate, formate, isobutyrate, caprate, heptanoate, propiolate, oxalate, malonate, succinate, suberate, sebacate, fumarate, maleate, butyne-1,4-dioate, hexyne-1,6-dioate, benzoate, chlorobenzoate, methylbenzoate, dinitrobenzoate, hydroxybenzoate, methoxybenzoate, phthalate, terephthalate, sulfonate, xylenesulfonate, phenylacetate, phenylpropionate, phenylbutyrate, citrate, lactate, β -hydroxybutyrate, glycolate, maleate, tartrate, methanesulfonate, propanesulfonate, naphthalene-1-sulfonate, naphthalene-2-sulfonate, mandelate and the like.

[0572] Suitable bases for forming pharmaceutically acceptable salts with acidic functional groups include, but are not limited to, hydroxides of alkali metals such as sodium, potassium, and lithium; hydroxides of alkaline earth metal such as calcium and magnesium; hydroxides of other metals, such as aluminum and zinc; ammonia, and organic amines, such as unsubstituted or hydroxy-substituted mono-, di-, or trialkylamines; dicyclohexylamine; tributyl amine; pyridine; N-methyl,N-ethylamine; diethylamine; triethylamine; mono-, bis-, or tris-(2-hydroxy-lower alkyl amines), such as mono-, bis-, or tris-(2-hydroxyethyl)amine, 2-hydroxy-tert-butylamine, or tris-(hydroxymethyl)methylamine, N,N-di alkyl-N-(hydroxy alkyl)-amines, such as N,N-dimethyl-N-(2-hydroxyethyl)amine, or tri-(2-hydroxyethyl)amine; N-methyl-D-glucamine; and amino acids such as arginine, lysine, and the like.

[0573] Certain compounds useful in accordance with the invention and their salts may exist in more than one crystal-

line form (i.e., polymorph); the present invention includes the use of each of the crystal forms and mixtures thereof.

[0574] Certain compounds useful in accordance with the invention may contain one or more chiral centers, and exist in different optically active forms. When compounds useful in accordance with the invention contain one chiral center, the compounds exist in two enantiomeric forms and the present invention includes the use of both enantiomers and mixtures of enantiomers, such as racemic mixtures thereof. The enantiomers may be resolved by methods known to those skilled in the art; for example, enantiomers may be resolved by formation of diastereoisomeric salts which may be separated, for example, by crystallization; formation of diastereoisomeric derivatives or complexes which may be separated, for example, by crystallization, gas-liquid or liquid chromatography; selective reaction of one enantiomer with an enantiomer-specific reagent, for example, via enzymatic esterification; or gas-liquid or liquid chromatography in a chiral environment, for example, on a chiral support (e.g., silica with a bound chiral ligand) or in the presence of a chiral solvent. Where the desired enantiomer is converted into another chemical entity by one of the separation procedures described above, a further step may be used to liberate the desired purified enantiomer. Alternatively, specific enantiomers may be synthesized by asymmetric synthesis using optically active reagents, substrates, catalysts or solvents, or by converting one enantiomer into the other by asymmetric transformation.

[0575] When a compound useful in accordance with the invention contains more than one chiral center, it may exist in diastereoisomeric forms. The diastereoisomeric compounds may be separated by methods known to those skilled in the art (for example, chromatography or crystallization) and the individual enantiomers may be separated as described above. The present invention includes the use of various diastereoisomers of compounds useful in accordance with the invention, and mixtures thereof. Compounds useful in accordance with the invention may exist in different tautomeric forms or as different geometric isomers, and the present invention includes the use of each tautomer and/or geometric isomer of compounds useful in accordance with the invention, and mixtures thereof. Compounds useful in accordance with the invention may exist in zwitterionic form. The present invention includes the use of each zwitterionic form of compounds useful in accordance with the invention, and mixtures thereof.

Kits

[0576] An EZH2 antagonist may, if desired, be presented in a kit (e.g., a pack or dispenser device) which may contain one or more unit dosage forms containing the EZH2 antagonist. The pack may for example comprise metal or plastic foil, such as a blister pack. The pack or dispenser device may be accompanied by instructions for administration. Compositions comprising an EZH2 antagonist of the invention formulated in a compatible pharmaceutical carrier may also be prepared, placed in an appropriate container, and labeled for treatment of an indicated condition. Instructions for use may also be provided.

[0577] Also provided herein are kits comprising a plurality of methylation detection reagents that detect the methylated H3-K27. For example, the kit includes mono-methylated H3-K27, di-methylated H3-K27 and tri-methylated H3-K27 detection reagents. The detection reagent is for example antibodies or fragments thereof, polypeptide or aptamers.

[0578] A kit may also include an EZH2 mutant detection reagent, e.g., nucleic acids that specifically identify a mutant EZH2 nucleic acid sequence by having homologous nucleic acid sequences, such as oligonucleotide sequences, complementary to a portion of the mutant EZH2 nucleic acid sequence or antibodies to proteins encoded by the mutant EZH2 nucleic acids packaged together in the form of a kit. The oligonucleotides can be fragments of the EZH2 gene. For example the oligonucleotides can be 200, 150, 100, 50, 25, 10 or less nucleotides in length. The kit may contain in separate containers an aptamer or an antibody, control formulations (positive and/or negative), and/or a detectable label such as fluorescein, green fluorescent protein, rhodamine, cyanine dyes, Alexa dyes, luciferase, radiolabels, among others. Instructions (e.g., written, tape, VCR, CD-ROM, etc.) for carrying out the assay may be included in the kit. The assay may for example be in the form of a Western Blot analysis, Immunohistochemistry (IHC), immunofluorescence (IF), sequencing and Mass spectrometry (MS) as known in the art.

DEFINITIONS

[0579] For convenience, certain terms employed in the specification, examples, and appended claims are collected here. All definitions, as defined and used herein, supersede dictionary definitions, definitions in documents incorporated by reference, and/or ordinary meanings of the defined terms.

[0580] The articles “a” and “an” are used herein to refer to one or to more than one (i.e., to at least one) of the grammatical object of the article. By way of example, “an element” means one element or more than one element.

[0581] The phrase “and/or,” as used herein in the specification and in the claims, should be understood to mean “either or both” of the elements so conjoined, i.e., elements that are conjunctively present in some cases and disjunctively present in other cases. Multiple elements listed with “and/or” should be construed in the same fashion, i.e., “one or more” of the elements so conjoined. Other elements may optionally be present other than the elements specifically identified by the “and/or” clause, whether related or unrelated to those elements specifically identified. Thus, as a non-limiting example, a reference to “A and/or B”, when used in conjunction with open-ended language such as “comprising” can refer, in one embodiment, to A only (optionally including elements other than B); in another embodiment, to B only (optionally including elements other than A); in yet another embodiment, to both A and B (optionally including other elements); etc.

[0582] As used herein in the specification and in the claims, “or” should be understood to have the same meaning as “and/or” as defined above. For example, when separating items in a list, “or” or “and/or” shall be interpreted as being inclusive, i.e., the inclusion of at least one, but also including more than one, of a number or list of elements, and, optionally, additional unlisted items. Only terms clearly indicated to the contrary, such as “only one of” or “exactly one of,” or, when used in the claims, “consisting of,” will refer to the inclusion of exactly one element of a number or list of elements. In general, the term “or” as used herein shall only be interpreted as indicating exclusive alternatives (i.e., “one or the other but not both”) when preceded by terms of exclusivity, such as “either,” “one of,” “only one of,” or “exactly one of.” “Consisting essentially of,” when used in the claims, shall have its ordinary meaning as used in the field of patent law.

[0583] As used herein in the specification and in the claims, the phrase “at least one,” in reference to a list of one or more elements, should be understood to mean at least one element selected from any one or more of the elements in the list of elements, but not necessarily including at least one of each and every element specifically listed within the list of elements and not excluding any combinations of elements in the list of elements. This definition also allows that elements may optionally be present other than the elements specifically identified within the list of elements to which the phrase “at least one” refers, whether related or unrelated to those elements specifically identified. Thus, as a non-limiting example, “at least one of A and B” (or, equivalently, “at least one of A or B;” or, equivalently “at least one of A and/or B”) can refer, in one embodiment, to at least one, optionally including more than one, A, with no B present (and optionally including elements other than B); in another embodiment, to at least one, optionally including more than one, B, with no A present (and optionally including elements other than A); in yet another embodiment, to at least one, optionally including more than one, A, and at least one, optionally including more than one, B (and optionally including other elements); etc.

[0584] It should also be understood that, unless clearly indicated to the contrary, in any methods claimed herein that include more than one step or act, the order of the steps or acts of the method is not necessarily limited to the order in which the steps or acts of the method are recited.

[0585] In the claims, as well as in the specification above, all transitional phrases such as “comprising,” “including,” “carrying,” “having,” “containing,” “involving,” “holding,” “composed of,” and the like are to be understood to be open-ended, i.e., to mean including but not limited to. Only the transitional phrases “consisting of” and “consisting essentially of” shall be closed or semi-closed transitional phrases, respectively, as set forth in the United States Patent Office Manual of Patent Examining Procedures, Section 2111.03.

[0586] The terms “co-administration” and “co-administering” refer to both concurrent administration (administration of two or more therapeutic agents at the same time) and time varied administration (administration of one or more therapeutic agents at a time different from that of the administration of an additional therapeutic agent or agents), as long as the therapeutic agents are present in the patient to some extent at the same time.

[0587] The term “precancerous condition” or “pre-malignant condition” refers to a disease, syndrome, or finding that, if left untreated, may lead to cancer. It is a generalized state associated with a significantly increased risk of cancer.

[0588] The term “treating” as used herein refers to alleviate of at least one symptom of the disease, disorder or condition. The term encompasses the administration and/or application of one or more compounds described herein, to a subject, for the purpose of providing management of, or remedy for a condition. “Treatment” for the purposes of this disclosure, may, but does not have to, provide a cure; rather, “treatment” may be in the form of management of the condition. When the compounds described herein are used to treat unwanted proliferating cells, including cancers, “treatment” includes partial or total destruction of the undesirable proliferating cells with minimal destructive effects on normal cells. A desired mechanism of treatment of unwanted rapidly proliferating cells, including cancer cells, at the cellular level is apoptosis.

[0589] The term “preventing” as used herein includes either preventing or slowing the onset of a clinically evident disease

progression altogether or preventing or slowing the onset of a preclinically evident stage of a disease in individuals at risk. This includes prophylactic treatment of those at risk of developing a disease.

[0590] The term “subject” as used herein for purposes of treatment includes any human subject who has been diagnosed with, has symptoms of, or is at risk of developing a cancer or a precancerous condition. For methods of prevention the subject is any human subject. To illustrate, for purposes of prevention, a subject may be a human subject who is at risk of or is genetically predisposed to obtaining a disorder characterized by unwanted, rapid cell proliferation, such as cancer. The subject may be at risk due to exposure to carcinogenic agents, being genetically predisposed to disorders characterized by unwanted, rapid cell proliferation, and so on.

[0591] Except as otherwise indicated, standard methods can be used for the production of recombinant and synthetic polypeptides, fusion proteins, antibodies or antigen-binding fragments thereof, manipulation of nucleic acid sequences, production of transformed cells, and the like. Such techniques are known to those skilled in the art. See, e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 3rd Ed. (Cold Spring Harbor, N.Y., 2001); F. M. Ausubel et al. *Current Protocols in Molecular Biology* (Green Publishing Associates, Inc. and John Wiley & Sons, Inc., New York).

[0592] The term “EZH2 polypeptide” encompasses functional fragments of the full-length polypeptides and functional equivalents of either of the foregoing that have substantially similar or substantially identical amino acid sequences (at least about 75%, 80%, 85%, 90%, 95% 98% or more amino acid sequence similarity or identity), where the functional fragment or functional equivalent retains one or more of the functional properties of the native polypeptide.

[0593] By “functional” it is meant that the polypeptide (or nucleic acid) has the same or substantially similar activity with respect to one or more of the biological properties of the native polypeptide (or nucleic acid), e.g., at least about 50%, 75%, 85%, 90%, 95% or 98% or more of the activity of the native polypeptide (or nucleic acid).

[0594] The term “modulate” (and grammatical equivalents) refers to an increase or decrease in activity. In particular embodiments, the term “increase” or “enhance” (and grammatical equivalents) means an elevation by at least about 25%, 50%, 75%, 2-fold, 3-fold, 5-fold, 10-fold, 15-fold, 20-fold or more. In particular embodiments, the terms “decrease” or “reduce” (and grammatical equivalents) means a diminishment by at least about 25%, 40%, 50%, 60%, 75%, 80%, 85%, 90%, 95%, 98% or more. In some embodiments, the indicated activity, substance or other parameter is not detectable. Specifically provided are antagonists of EZH2.

[0595] The term “pharmacodynamic marker” refers to a molecular marker of drug response that can be measured in patients receiving the drug. The marker should be a direct measure of modulation of the drug target and be able to show quantitative changes in response to dose. A potential pharmacodynamic marker for EZH2 antagonists could be levels of histone H3-K27 methylation in disease or surrogate tissue.

[0596] As used herein, the term “responsiveness” is interchangeable with terms “responsive”, “sensitive”, and “sensitivity”, and it is meant that a subject showing therapeutic response when administered an EZH inhibitor, e.g., tumor cells or tumor tissues of the subject undergo apoptosis and/or necrosis, and/or display reduced growing, dividing, or proliferation.

[0597] The term “control” or “reference” refers to methylation levels (e.g., monomethylation level, dimethylation level or trimethylation level) detected in an adjacent non-tumor tissue isolated from the subject, detected in a healthy tissue from a healthy subject, or established by a pathologist with standard methods in the art.

[0598] By “sample” it means any biological sample derived from the subject, includes but is not limited to, cells, tissues samples and body fluids (including, but not limited to, mucus, blood, plasma, serum, urine, saliva, and semen).

[0599] Throughout the description, where compositions are described as having, including, or comprising specific components, it is contemplated that compositions also consist essentially of, or consist of, the recited components. Similarly, where methods or processes are described as having, including, or comprising specific process steps, the processes also consist essentially of, or consist of, the recited processing steps. Further, it should be understood that the order of steps or order for performing certain actions is immaterial so long as the invention remains operable. Moreover, two or more steps or actions can be conducted simultaneously.

[0600] The synthetic processes of the invention can tolerate a wide variety of functional groups; therefore various substituted starting materials can be used. The processes generally provide the desired final compound at or near the end of the overall process, although it may be desirable in certain instances to further convert the compound to a pharmaceutically acceptable salt, ester or prodrug thereof.

[0601] Compounds of the present invention can be prepared in a variety of ways using commercially available starting materials, compounds known in the literature, or from readily prepared intermediates, by employing standard synthetic methods and procedures either known to those skilled in the art, or which will be apparent to the skilled artisan in light of the teachings herein. Standard synthetic methods and procedures for the preparation of organic molecules and functional group transformations and manipulations can be obtained from the relevant scientific literature or from standard textbooks in the field. Although not limited to any one or several sources, classic texts such as Smith, M. B., March, J., *March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure*, 5th edition, John Wiley & Sons: New York, 2001; and Greene, T. W., Wuts, P. G. M., *Protective Groups in Organic Synthesis*, 3rd edition, John Wiley & Sons: New York, 1999, incorporated by reference herein, are useful and recognized reference textbooks of organic synthesis known to those in the art. The following descriptions of synthetic methods are designed to illustrate, but not to limit, general procedures for the preparation of compounds of the present invention.

[0602] Compounds of the present invention can be conveniently prepared by a variety of methods familiar to those skilled in the art. The compounds of this invention with each of the formulae described herein may be prepared according to the following procedures from commercially available starting materials or starting materials which can be prepared using literature procedures. These procedures show the preparation of representative compounds of this invention.

[0603] Compounds designed, selected and/or optimized by methods described above, once produced, can be characterized using a variety of assays known to those skilled in the art to determine whether the compounds have biological activity. For example, the molecules can be characterized by conventional assays, including but not limited to those assays

described below, to determine whether they have a predicted activity, binding activity and/or binding specificity.

[0604] Furthermore, high-throughput screening can be used to speed up analysis using such assays. As a result, it can be possible to rapidly screen the molecules described herein for activity, using techniques known in the art. General methodologies for performing high-throughput screening are described, for example, in Devlin (1998) *High Throughput Screening*, Marcel Dekker; and U.S. Pat. No. 5,763,263. High-throughput assays can use one or more different assay techniques including, but not limited to, those described below.

[0605] All publications and patent documents cited herein are incorporated herein by reference as if each such publication or document was specifically and individually indicated to be incorporated herein by reference. Citation of publications and patent documents is not intended as an admission that any is pertinent prior art, nor does it constitute any admission as to the contents or date of the same. The invention having now been described by way of written description, those of skill in the art will recognize that the invention can be practiced in a variety of embodiments and that the foregoing description and examples below are for purposes of illustration and not limitation of the claims that follow.

EXAMPLES

[0606] The invention now being generally described, it will be more readily understood by reference to the following examples, which are included merely for purposes of illustration of certain aspects and embodiments of the present invention, and are not intended to limit the invention.

Example 1

Recombinant Five-Component PRC2 Complex

[0607] Wild-type EZH2 (GenBank Accession No. NM_004456) or Tyr641 mutants were co-expressed with wild-type AEBP2 (GenBank Accession No. NM_153207), EED (GenBank Accession No. NM_003797), SUZ₁₂ (GenBank Accession No. NM_015355) and RbAp48 (GenBank Accession No. NM_005610) in *Spodoptera frugiperda* (Sf9) cells using a baculovirus expression system. An N-terminal FLAG tag on the EED was used to purify active PRC2 complex from cell lysates (BPS Bioscience, catalog number 51004). The purity of the final PRC2 preparations was assessed by SDS-PAGE with Coomassie blue staining.

Example 2

H3, H4 Peptide Panel

[0608] A library consisting of 44 peptides of 15 amino acids each was synthesized by 21st Century Biochemicals (Marlboro, Mass.). This peptide panel encompassed all of the amino acids of human histones H3 and H4 with 5 residue overlaps between consecutive peptide sequences. The N-terminus of each peptide was appended with biotin, and the C-termini were represented as the amide. Purity (>95%) and identity were confirmed by liquid chromatography/mass spectral analysis.

[0609] For study of the H3-K27 methylation status dependence of enzyme activity, peptides were synthesized representing the amino acid sequence of human H3 from residues 21-44 (H3:21-44) with lysine 27 represented as the unmodi-

fied, mono-methylated, di-methylated or tri-methylated side chain amine. These peptides were purchased from New England Peptide (Gardner, Mass.) with biotin appended to the C-terminus of each peptide.

Example 3

Evaluation of H3-K27 Methylation Status in Cells

[0610] The cell lines OCI-LY19 (ACC 528), KARPAS-422 (ACC 32), and WSU-DLCL2 (ACC 575) were obtained from DSMZ. The cell lines DB (CRL-2289) and SU-DHL6 (CRL-2959) were obtained from ATCC. OCI-LY19, WSU-DLCL2, and DB cell lines were grown in RPMI-1640 with 10% FBS, and KARPAS-422 and SU-DHL6 cell lines were grown in RPMI-1640 plus 20% FBS. Cells were grown to a density of 1.5×10^6 cells/mL and 1×10^7 cells were harvested by centrifugation at $264 \times g$, washed in ice cold PBS and lysed by resuspension in a $10 \times$ pellet volume of RIPA lysis buffer containing 50 mM Tris-HCl, 150 mM NaCl, 0.25% DOC, 1% NP-40, and 1 mM EDTA (Millipore #20-188), plus 0.1% SDS and protease inhibitor tablets (Roche #1836153). Lysates were sonicated by 2 rounds of 10 1-second bursts at setting 3 with a Misonix XL-2000 to ensure efficient histone extraction, and cleared by centrifugation at 4°C using a bench top centrifuge at 14,000 rpm for 10 minutes. Protein concentration was determined by BCA assay (Pierce). Four micrograms of each lysate was fractionated on 4-20% Tris-Glycine gel (Invitrogen), transferred to PVDF, and probed with the following antibodies in Odyssey blocking buffer: mouse anti-EZH2 (CST 3147; 1:2000 dilution), rabbit anti-H3-K27me3 (CST 9733; 1:10000 dilution), rabbit anti-H3-K27me2 (CST 9755; 1:5000 dilution), rabbit anti-H3-K27me1 (Active Motif 39377; 1:5000 dilution), and mouse anti-Total H3 (CST 3638; 1:20000 dilution). Following primary Ab incubation, membranes were probed with IRDye 800CW donkey-anti-mouse IgG (LiCOR #926-32212) or Alexa Fluor 680 goat-anti-rabbit IgG (Invitrogen #A-21076) secondary Ab and imaged using the LiCOR Odyssey system.

Example 4

Enzymology

[0611] As noted above, it had previously been concluded that the disease-associated changes at Tyr641 resulted in loss of function with respect to EZH2-catalyzed H3-K27 methylation. However, a presumptive reduction in the rate of H3-K27 methylation due to enzyme heterozygosity was difficult to rationalize as the basis for a malignant phenotype, especially in light of previous data indicating that overexpression of EZH2, loss-of-function mutations in the corresponding H3-K27 demethylase UTX, or overexpression of components of the PRC2, such as PHF19/PCL3, involved in increased H3-K27 trimethylation, all result in malignant phenotypes in specific human cancers. Morin et al. (2010) *Nat Genet* 42:181-5; Martinez-Garcia et al. (2010) *Nat Genet* 42:100-1; Bracken et al. (2003) *EMBO J* 22:5323-35; Kleer et al. (2003) *Proc Natl Acad Sci USA* 100:11606-11; Varambally et al. (2002) *Nature* 419:624-9; Simon et al. (2008) *Mutat Res* 647:21-9; van Haaften et al. (2009) *Nat Genet.* 41:521-3; Wang et al. (2004) *Gene* 343:69-78; Cao et al. (2008) *Mol Cell Biol* 28:1862-72; and Sarma et al. (2008) *Mol Cell Biol* 28:2718-31). Therefore, the enzymology of these mutations was explored in greater detail.

[0612] Recombinant PRC2 complexes were prepared with WT and Tyr641 mutant versions of human EZH2 (see Example 1 above; Cao et al. (2004) *Mol Cell* 15:57-67). Equal

concentrations (nominally 8 nM, based on protein determinations) of each complex were initially tested for the ability to catalyze ^3H -methyl transfer from labeled S-adenosyl methionine (SAM) to an unmodified peptide representing the amino acid sequence surrounding H3-K27 (H3:21-44) or to native avian erythrocyte oligonucleosomes. As previously reported (Morin et al. (2010) *Nat Genet* 42:181-5), it was found that the WT enzyme displayed robust activity for methyl transfer to this unmethylated peptidic substrate, but that none of the mutant enzymes displayed significant methyltransferase activity (FIG. 1A). In contrast to the previously reported data and that in FIG. 1A, it was found that all of the mutant EZH2 constructs were active methyltransferases against the avian nucleosome substrate (FIG. 1B). The nucleosomes isolated from the avian natural source represent an admixture of states of histone modification, including various states of H3-K27 methylation as judged by Western blotting with H3-K27 methylation-specific antibodies.

[0613] There are several potential explanations for the discordant activity of the mutant PRC2 complexes on peptide and nucleosome substrates. One possibility is that substrate recognition sites distal to the enzyme active site (i.e., exosites) are important determinants of substrate binding and turnover; these sites would engage complementary recognition elements on the nucleosome that are not available on small peptidic substrates. However, when *E. coli*-expressed, recombinant human histone H3 was tested as a substrate for the WT and mutant PRC2 complexes, the resulting pattern of activity was identical to that seen for the peptide substrate; that is, the WT enzyme demonstrated robust methyltransferase activity against the H3 substrate, the Y641F mutant showed 7% the activity of WT complex, and all other mutants displayed $\leq 1\%$ the activity of WT complex. Hence, exosite engagement seems an unlikely explanation for the current results. The nucleosome presents many lysine residues beyond H3-K27 as potential sites of methylation that would not be present in the small peptidic substrate. Thus, another possibility is that mutation of Y641 alters the substrate specificity of EZH2 to result in methylation of lysine residues other than H3-K27. This possibility is unlikely given the excellent agreement between mutant activity on small peptide and recombinant H3 protein substrates.

[0614] The apparent discordance between the present results and those previously reported was resolved when the enzymatic activity of the WT and mutant PRC2 complexes were tested against a panel of peptidic substrates that represent all possible lysine (K) residues of histone H3 and histone H4 (see Example 2 above). All of the enzyme forms showed significant activity only against peptides containing the equivalent of residue H3-K27. The specific activity of the mutants, however, was greatly reduced relative to WT in the order WT >> Y641F > Y641S > Y641H > Y641N, again consistent with previous reported findings.

Example 5

Enzymology

[0615] To understand further the enzymatic activity of these mutants, and to reconcile the apparent discrepancy between activity against peptidic and nucleosome substrates, the ability of the enzyme forms to catalyze further methylation of various H3-K27 methylation states in the context of the H3:21-44 peptide was studied. As stated above, it was found that all of the mutant enzymes were deficient catalysts of unmodified H3-K27 peptide methylation, relative to the WT enzyme. Remarkably, however, all of the mutant enzymes were found to be superior to WT enzyme in cata-

lyzing further methylation of the mono- and especially the di-methylated H3-K27 peptides (FIG. 2). Thus, the data suggest that the WT enzyme is most efficient in catalyzing the zero- to mono-methylation reaction. The mutant enzymes are defective in catalyzing this initial step, but are more efficient than the WT enzyme in catalyzing the subsequent steps leading from mono-methyl to di- and tri-methyl H3-K27.

[0616] The origins of the differential substrate specificities of WT and mutant EZH2 were explored through steady state enzyme kinetics. As summarized in Table 2, the mutations have minimal effects on ground-state substrate recognition, as demonstrated by the similar values of K_m for nucleosome and of $K_{1/2}$ for peptide substrates. In all cases the peptidic substrates displayed sigmoidal binding behavior; hence the concentration of peptide resulting in half-maximal velocity is reported here as $K_{1/2}$ instead of the more common Michaelis constant, K_m . Copeland (2005) *Evaluation of Enzyme Inhibitors in Drug Discovery: A Guide to Medicinal Chemists and Pharmacologists*, Wiley. The SAM K_m likewise displayed minimal variation among the enzyme forms, ranging from 208 ± 50 to 304 ± 64 nM. Instead, the differences in substrate utilization appear to have their origin in transition state recognition, as demonstrated by differences in k_{cat} values among the enzymes for various substrates (Table 2). As a result, the catalytic efficiency, quantified as the ratio k_{cat}/K (where K is either K_m or $K_{1/2}$, depending on substrate identity; vide supra), varies between the WT and mutant enzymes for different states of H3-K27 methylation (Table 2).

TABLE 2

Steady state kinetic parameters for methylation reactions catalyzed by PRC2 containing wild-type or Y641 mutants of EZH2.				
Enzyme	Substrate H3-K27 Methylation Status	K (nM)	k_{cat} ($h^{-1} \times 10^{-2}$)	k_{cat}/K ($h^{-1} \cdot nM^{-1} \times 10^{-4}$)
WT	0	184 ± 10	84.0 ± 3.0	45.7 ± 3.0
	1	436 ± 42	65.4 ± 5.8	15.0 ± 2.0
	2	178 ± 16	6.0 ± 0.3	3.4 ± 0.3
	Nucleosome	141 ± 31	42.6 ± 2.6	30.2 ± 6.9
Y641F	0	240 ± 19	4.8 ± 0.3	2.0 ± 0.2
	1	404 ± 124	15.0 ± 4.3	3.7 ± 1.6
	2	191 ± 10	84.0 ± 2.8	44.0 ± 2.7
	Nucleosome	176 ± 19	65.4 ± 2.0	37.2 ± 4.2
Y641H	0	— ^a	—	—
	1	319 ± 57	28.2 ± 3.7	8.8 ± 2.0
	2	148 ± 9	22.8 ± 0.9	15.4 ± 1.1
	Nucleosome	140 ± 22	23.4 ± 1.0	16.7 ± 2.7
Y641N	0	—	—	—
	1	280 ± 11	23.4 ± 0.8	8.4 ± 0.4
	2	157 ± 11	96.0 ± 4.0	61.1 ± 5.0
	Nucleosome	191 ± 34	23.4 ± 1.3	12.3 ± 2.3
Y641S	0	—	—	—
	1	249 ± 8	27.6 ± 0.8	11.1 ± 0.5
	2	136 ± 8	59.4 ± 2.0	43.7 ± 3.0
	Nucleosome	137 ± 28	23.4 ± 1.4	17.1 ± 3.6

^aActivity too low to measure.

Example 6

Enzymology

[0617] The steady state kinetic parameters listed in Table 2 made it possible to calculate the expected levels of different H3-K27 methylation states for cells heterozygous for the various mutant EZH2 forms, relative to cells homozygous for the WT enzyme. To perform these simulations, a number of simplifying assumptions were made: (1) that steady state

enzyme kinetics are relevant to PRC2-catalyzed H3-K27 methylation in the cellular context and that all measurements are made at the same time point in cell growth; (2) that the mutant and WT enzyme are expressed at equal levels in heterozygous cells and that the total EZH2 level is equal in all cells; (3) that the cellular concentration of SAM, relative to its K_m , is saturating and does not change among the cells; (4) that the cellular concentration of nucleosome, is similar to its K_m and likewise does not change among cells; (5) that EZH1 catalyzed methylation of H3-K27 was insignificant and constant among the cells; and (6) that any H3-K27 demethylase activity was also constant among the cells.

[0618] With these assumptions in place, the predictions illustrated in FIG. 3A were obtained for relative levels of H3-K27me3 (top panel), H3-K27me2 (middle panel) and H3-K27me1 (bottom panel). A clear pattern emerges from these simulations. The level of H3-K27me3 increases relative to WT cells for all mutant-harboring cells, ranging from a 30% increase for the Y641H mutant to >400% for the Y641N mutant. At the same time, the levels of H3-K27me2 decreases to <50% of WT for all of the mutants, and the levels of H3-K27me1 are reduced by approximately half for all mutants, relative to WT.

[0619] The relative levels of the H3-K27 methylation states in B-cell lymphoma cell lines that are known to be homozygous for WT EZH2 (OCI-LY19) or heterozygous for EZH2 Y641N (DB, KARPAS 422, and SU-DHL-6) or EZH2 Y641F (WSU-DLCL2) were then measured by Western blotting (FIG. 3B). The pattern of relative H3-K27 methylation states seen in FIG. 3b is in excellent agreement with the results of the simulations based on in vitro steady state kinetic parameters, despite the assumptions used in the simulations and the use of a non-physiological peptide surrogate as substrate.

[0620] Thus, increased H3-K27me3 was observed for all Y641 mutant-harboring cells relative to WT, decreased H3-K27me2 was observed for all Y641 mutant-harboring cells relative to WT, and decreased H3-K27me1 was observed for at least two of the four mutant cell lines. The near-comparable levels of H3-K27me1 in WT and KARPAS 422 and SU-DHL-6 cells may reflect different expression levels of WT and mutant EZH2, different contributions of EZH1, or other factors not accounted for in the simulations. Nevertheless, the concordance between the predicted and experimental patterns of H3-K27 methylation status is remarkable and supports the view that enzymatic coupling between WT and mutant EZH2 leads to increased H3-K27me3, thus resulting in the malignant phenotype of cells that are heterozygous for these mutants.

Example 7

In Vitro Assays of PRC2 Methyltransferase Activity

[0621] Flashplate Assay with Peptide Substrate.

[0622] For initial comparison of WT and Y641 mutants of EZH2, biotinylated histone H3:21-44 peptide containing unmethylated K27 (New England Peptide), monomethylated K27 (Millipore) or dimethylated K27 (Millipore) at a concentration of 800 nM was combined with a mixture of S-adenosylmethionine-Cl (SAM) at 1,700 nM, and 300 nM tritiated SAM (Perkin Elmer). This substrate combination was then added to the PRC2 in assay buffer (20 mM BICINE, 1 mM DTT, 0.002% Tween 20, 0.005% bovine skin gelatin (BSG), pH 7.6). Reactions were allowed to proceed for the indicated time interval and then quenched by addition of excess cold SAM (600 μ M final concentration). Quenched reaction mixtures were transferred to a streptavidin-coated Flashplate (Perkin Elmer, catalog number SMP410), allowed

to bind for one hour, and then detected on a TopCount NXT HTS scintillation and luminescence counter (Perkin Elmer). Each time point represented the average of six individual reactions. Steady state kinetic parameters were determined under identical reaction conditions except that the concentration of peptide or SAM was varied, while at saturating conditions of the other substrate. Velocity was plotted as a function of varied substrate concentration and the data were fitted to the untransformed version of the Michaelis-Menten equation or the untransformed version of a sigmoidal kinetic equation to calculate values of K and k_{cat} . Standard errors of fitted parameters are listed in Table 2 and were used to construct the error bars illustrated in FIG. 2 panels B and C. Error associated with k_{cat}/K (Table 2) were calculated according to standard methods of error propagation; the fractional error of k_{cat}/K was determined as:

$$\mu \frac{k_{cat}}{K} = \sqrt{\left(\frac{\mu k_{cat}}{k_{cat}}\right)^2 + \left(\frac{\mu K}{K}\right)^2} \quad (1)$$

where μk_{cat} is the standard error of k_{cat} and μK is the standard error of K .

[0623] Filterplate Assay with Oligonucleosome.


[0624] Chicken erythrocyte oligonucleosomes were purified as previously described. Fang et al. (2004) *Methods Enzymol* 377:213-26. Nucleosomes were combined with a mixture of SAM and tritiated SAM, and added to PRC2 in assay buffer (20 mM BICINE, 100 mM KCl, 1 mM DTT, 0.002% Tween 20, 0.005% BSG, pH 7.6). Reactions were run and quenched as above. Quenched reaction mixture was transferred to a glass fiber filterplate (Millipore, catalog number MSFBN6B) and washed three times with 10% trichloroacetic acid and allowed to dry. Microscint Zero (30 μ L) was added and tritium incorporation was detected on a TopCount scintillation and luminescence counter. Steady state parameters were determined under identical reaction conditions except that the concentration of nucleosome or SAM was varied while at saturating conditions of the other substrate. Velocity was plotted as a function of varied substrate concentration and fitted to the untransformed version of the Michaelis-Menten equation to derive the values of K_m and k_{cat} as described above.

Example 8

Compound Preparation

A. Preparation of Compound 63

[0625]




[0626] To a solution of 9-((3aR,4R,6R,6aR)-6-(aminomethyl)-2,2-dimethyltetrahydrofuro[3,4-d][1,3]dioxol-4-yl)-

9H-purin-6-amine (Townsend, A. P. et al. (2009) *Org. Lett.* 11:2976-2979) (3.05 g, 9.96 mmol) in DCE (250 mL) was added (9H-fluoren-9-yl)methyl (2-oxoethyl)carbamate (2.8 g, 9.96 mmol) and NaB(OAc)₃H (2.96 g, 13.95 mmol), the mixture stirred for 4 h at room temperature. K₂CO₃ solution was added to pH at 8-9. DCM was added, the organic layer was dried with Na₂SO₄, concentrated and purified by SGC (DCM:MeOH=30:1) to give 63 (2.9 g, yield: 50.9%).

B. Preparation of Compound 65


[0627]




[0628] To a solution of 63 (2.9 g, 5.08 mmol) in DCE (250 mL), (S)-benzyl 2-((tert-butoxycarbonyl)amino)-4-oxobutanoate (1.56 g, 5.08 mmol) and NaB(OAc)₃H (1.51 g, 7.11 mmol) were added, the mixture stirred for 4 h at room temperature. K₂CO₃ solution was added to pH at 8-9. DCM was added, the organic layer was dried with Na₂SO₄, concentrated and purified by SGC (DCM: MeOH=100:1) to give 65 (2.8 g, yield: 63.9%).

C. Preparation of Compound 75

[0629]



-continued



temperature under H_2 overnight. The mixture was filtered and the filtrate was concentrated to give crude product. The crude product was purified by preparative TLC (DCM/MeOH=8:1) to afford 74 (20 mg, yield: 69%) as white powder. LC/MS (m/z): 689.7 [M+1]⁺.

[0633] Step 4. A solution of 74 (20 mg, 0.028 mmol) in 90% TFA (1 mL) was stirred at room temperature for 1 h, and concentrated as a solid to remove. TFA to give the compound 75 (TFA salt) as a colorless oil without purification. LC/MS (m/z): 549.7 [M+1]⁺.

Example 9

Inhibition of EZH2 Wild-Type and Y641 Mutants by SAH

[0634] S-Adenosyl-L-homocysteine (SAH) was serially diluted 3 fold in DMSO for 10 points and 1 μ L was plated in a 384 well microtiter plate. Positive control (100% inhibition standard) was 100 μ M final concentration of SAH and negative control (0% inhibition standard) contained 1 μ L of DMSO. SAH was then incubated for 30 minutes with 40 μ L per well of EZH2 wild-type and mutants at 8 nM in pH 7.6 assay buffer (20 mM BICINE, 100 mM KCl, 1 mM DTT, 0.002% Tween 20, 0.005% BSG). A substrate mix at 10 μ L per well was added which contained S-adenosylmethionine-C1 (SAM) at 150 nM and tritiated SAM at 100 nM, and biotinylated oligonucleosome at 150 nM in pH 7.6 assay buffer. Quenched enzyme reaction was transferred to a streptavidin-coated Flashplate (Perkin Elmer, catalog number SMP410), allowed to bind for one hour, and detected on a TopCount NXT HTS (Perkin Elmer).

[0635] Results are shown in FIG. 7. IC₅₀ values are shown in Table 3.

TABLE 3

Inhibition of WT EZH2 and Y641 mutants of EZH2 by SAH.					
	WT	Y641H	Y641S	Y641N	Y641F
IC ₅₀ , μ M	0.467	0.263	0.283	0.380	4.80

Example 10

Inhibition of EZH2 Wild-Type and Y641 Mutants by Compound 75

[0636] Compound 75 was serially diluted 3 fold in DMSO for 10 points and 1 μ L was plated in a 384 well microtiter plate. Positive control (100% inhibition standard) was 100 μ M final concentration of SAH and negative control (0% inhibition standard) contained 1 μ L of DMSO. Compound 75 was then incubated for 30 minutes with 40 μ L per well of EZH2 wild-type and mutants at 8 nM in pH 7.6 assay buffer (20 mM BICINE, 100 mM KCl, 1 mM DTT, 0.002% Tween 20, 0.005% BSG). A substrate mix at 10 μ L per well was added which contained S-adenosylmethionine-C1 (SAM) at 150 nM and tritiated SAM at 100 nM, and biotinylated oligonucleosome at 150 nM in pH 7.6 assay buffer. Quenched enzyme reaction was transferred to a streptavidin-coated Flashplate (Perkin Elmer, catalog number SMP410), allowed to bind for one hour, and detected on a TopCount NXT HTS (Perkin Elmer).

[0630] Step 1. To a solution of 65B (2.2 g, 2.55 mmol) in DCM (10 mL), Et₃NH (1.1 g, 15.3 mmol) were added, the mixture stirred for 4 h at room temperature. The mixture was concentrated to give crude 72 (2.2 g).

[0631] Step 2. To a stirred solution of 72 (167 mg, 0.26 mmol) in MeOH (4 mL), 2-(4-chlorophenyl)acetaldehyde (40 mg, 0.26 mmol) was added and stirred at room temperature for 20 min. Then Na(OAc)₃BH (83 mg, 0.39 mmol) and HOAc (0.4 mL) was added and stirred overnight. Then NaHCO₃ (aq) was added and extracted with DCM (25 mL×3), washed with brine, dried with Na₂SO₄ and concentrated. The crude product was purified by preparative TLC (DCM/MeOH=10:1) to afford 73 (30 mg, yield: 14%) as white powder. LC/MS (m/z): 779.7 [M+1]⁺.

[0632] Step 3. A mixture of 73 (30 mg, 0.038 mmol) and 10% Pd/C (15 mg) in MeOH (2 mL) was stirred at room

[0637] Results are shown in FIG. 8. IC50 values are shown in Table 4.

TABLE 4

Inhibition of WT EZH2 and Y641 mutants of EZH2 by Compound 75.					
	WT	Y641S	Y641N	Y641F	Y641H
IC50, μM	8.95	2.50	4.10	7.18	7.56

Example 11

H3-K27me2/me3 Ratios Predict Sensitivity to an EZH2 Inhibitor

[0638] Tumor cell lines heterozygous for the EZH2 (Y641) mutation display increased levels of H3-K27me3, the methylation state of H3-K27 thought to be important in tumorigenesis. Levels of the mono (H3-K27me1), di (H3-K27me2), or trimethylated (H3-K27me3) forms of H3-K27 in a panel of cell lines that were WT for EZH2, or heterozygous for EZH2 (Y641) mutations were evaluated. Cell lines used are listed in Table 5. The majority of lines are B-cell lymphoma lines, however two melanoma lines were also included. IGR1 is a melanoma line that has recently been found to contain a Y641N mutation in EZH2, and A375 cells were included as a WT EZH2 melanoma control line. FIGS. 9A and B show the results of western blot analysis of histones isolated from this cell line panel probed with antibodies recognizing H3-K27me1, H3-K27me2, or H3-K27me3. In general, global H3-K27me3 levels are higher in Y641 mutant containing cell lines than in cell lines expressing WT EZH2 exclusively. The exception is Farage cells, where H3-K27me3 levels were similar to those in WT lines. More striking are the dramatically lower levels of H3-K27me2 in EZH2 Y641 mutant cell lines relative to wild type cell lines. Little or no H3-K27me2 signal was observed in western blot of histones extracted from Y641 mutant cell lines, whereas the signal observed with the same antibody in WT cell lines was more intense than that observed with the antibody specific for H3-K27me3. Overall, in WT cell lines the western blot signal with an HK27me2 antibody was higher than the signal observed with the H3-K27me3 antibody, whereas the opposite was true in Y641 mutant cell lines. Thus the ratio of H3-K27me3/me2 signal in Y641 lines is higher than that observed in WT lines.

[0639] The H3-K27 methylation state can also be examined by Mass spectrometry (MS), an independent method that does not rely on antibody reagents. The MS analysis demonstrated that H3-K27me3 levels are higher in Y641 mutant and Pfeiffer lines (A677G) than in the other WT lines, whereas the opposite is true for H3-K27me2 levels. In the Y641 mutant and Pfeiffer lines (A677G), H3-K27me3 levels were higher than H3-K27me2 levels, whereas the opposite was true in the other WT lines. These results are consistent with those observed by western blot analysis in FIGS. 9A and B.

[0640] The differences in H3-K27 methylation state was also detected by immunocytochemistry using antibodies to H3-K27me2 or H3-K27me3. This immunohistochemistry assay is used for detecting aberrant H3-K27me2/3 ratios associated with Y641 mutant EZH2 in formalin fixed paraffin embedded patient tumor tissue samples. A panel of five WT and five Y641 mutant lymphoma cell line pellets were fixed and embedded in paraffin blocks and stained with anti-H3-

K27me2 or H3-K27me3 antibodies. An antibody to histone H3 was included as a positive control, since all cells should contain nuclear histone H3. FIG. 10 shows that all cell lines were positive in 100% of cells for both H3-K27me3 and H3 staining. Under these conditions, no clear difference in H3-K27me3 staining intensity was observed between WT and Y641 mutant cell lines. This may reflect the limited dynamic range of chromogenic immunocytochemistry staining compared to other methods of detection. However, as shown in FIG. 11, cell lines could be clearly segregated into those staining positive or negative for H3-K27me2. All WT cell lines stained positive for H3-K27me2, whereas all Y641 mutant cell lines and Pfeiffer cells (A677G) showed no staining with the H3-K27me2 antibody. These results are consistent with those obtained by western and MS analysis.

[0641] Without wishing to be bound by theory, the increased levels of H3-K27me3 associated with the gain of function EZH2 (Y641) mutations may render cells bearing EZH2 mutations more sensitive to small molecule EZH2 inhibitors. To evaluate whether the increased H3-K27me3 and/or decreased H3-K27me2 levels observed in Pfeiffer cells in the absence of an EZH2 Y641 mutation would also correlate with sensitivity to EZH2 inhibitors, two compounds that demonstrate potent inhibition of EZH2 in biochemical assays with IC50s of 85 and 16 nM respectively were tested. Treatment of WSU-DLCL2 cells with either compound led to inhibition of global H3-K27me3 levels, confirming their ability to enter cells and inhibit cellular EZH2 methyltransferase activity (FIG. 12).

[0642] The sensitivity of a panel of WT and Y641 mutant cell lines to each compound was evaluated in proliferation assays. Because the anti-proliferative activity of EZH2 inhibitors takes several days to manifest, compounds were assessed in 11-day proliferation assays. FIG. 13 shows representative growth curves for WT (OCI-LY19), or Y641 mutant (WSU-DLCL2) cell lines treated with the test compounds. Both compounds demonstrated anti-proliferative activity against WSU-DLCL2 cells, but little activity against OCI-LY19 cells. Inhibitor A was a more potent inhibitor of WSU-DLCL2 proliferation than Inhibitor B and this is consistent with Inhibitor A being a more potent inhibitor of EZH2 in biochemical assays. Proliferation assays were performed in a panel of WT and Y641 mutant lymphoma cell lines, with Inhibitor B, and day 11 IC90 values were derived. FIG. 14A shows IC90 values of lymphoma cell lines grouped by EZH2 Y641 status. Overall, Y641 mutant cell lines demonstrated increased sensitivity to EZH2 inhibitors relative to WT cell lines, although RL and SUDHL4 cells were significantly less sensitive than other mutant lines. Pfeiffer cells (A677G) demonstrate high H3-K27me3 and low H3-K27me2 levels, and so grouping cell lines according to high H3-K27me3 and low H3-K27me2 gives better discrimination of EZH2 inhibitor sensitivity as shown for Inhibitor B in FIG. 14B. Thus, high H3-K27me3 and low H3-K27me2 levels can be used to predict sensitivity to EZH2 inhibitors, independent of knowledge of mutational status.

[0643] These results demonstrates that identifying EZH2 Y641 mutations in patient tumors and/or detecting low levels of H3-K27me2 relative to H3-K27me3 through use of techniques such as western blot, MS or IHC in a patient can be used to identify which patient will respond to EZH2 inhibitor treatment.

TABLE 5

Cell lines used in this study.		
Cancer	EZH2 Status	Cell Line
Lymphoma: DLBCL (Diffuse Large Cell B Cell Lymphoma) and other B-cell Lymphoma	Wild Type	OCI-LY19
		HT
		MC116
		BC-1
		BC-3
		Toledo
		DOHH-2
		Farage
		SR
		NU-DHL-1
	Y641 Mutation	NU-DUL-1
		SU-DHL-10 (Y641F)
		DB (Y641N)
		KARPAS 422 (Y641N)
		SU-DHL-6 (Y641N)
Melanoma	Wild Type	WSU-DLCL-2 (Y641F)
	Y641 Mutation	RL (Y641N)
		SU-DHL-4 (Y641S)
		A375
		IGR-1 (Y641N)

Example 12

Recombinant 4-Component PRC2 Complexes

[0644] Wild-type (WT) EZH2 (GenBank Accession No. NM_004456) or A677G and A687V mutants were co-expressed with wild-type EED (GenBank Accession No. NM_003797), SUZ12 (GenBank Accession No. NM_015355) and RbAp48 (GenBank Accession No. NM005610) in *Spodoptera frugiperda* (Sf9) cells using a baculovirus expression system. An N-terminal FLAG tag on the EED was used to purify active PRC2 complex from cell lysates. The purity of the final PRC2 preparations was assessed by SDS-PAGE with Coomassie blue staining and protein concentration was determined using a bovine serum albumin standard curve in a Bradford assay.

Example 13

In Vitro Assays of PRC2 Methyltransferase Activity

[0645] Standard Procedure for Flashplate Assay with Peptide Substrates.

[0646] Activity of the wild-type or mutant EZH2-containing PRC2 complexes was investigated using a series of four peptides representing the amino acid sequence of human H3 from residues 21-44 (H3:21-44) with lysine 27 represented as the unmodified, monomethylated, dimethylated or trimethylated side chain amine, consisting of the following sequence, with the H3-K27 lysine subject to modification underlined, ATKAAAR
KSAPATGGVKKPHRYRPGG[K-Ahx-Biot]-amide (SEQ ID NO: 20). Biotin (Biot) was appended to a C-terminal lysine (K) residue through an aminohexyl linker (AHX) attached to the lysine side chain amine (21st Century Biochemicals). For comparison of mutant of WT and mutant EZH2 activity, biotinylated histone H3:21-44 peptides were combined with a mixture of S-adenosylmethionine (SAM; Sigma-Aldrich) and tritiated SAM (³H-SAM; American Radiolabeled Chemicals) and recombinant 4-component PRC2 in assay buffer (20 mM BICINE, 1 mM DTT, 0.002% Tween 20, 0.005% bovine skin gelatin (BSG), pH 7.6). Reac-

tions were allowed to proceed for the indicated time interval and then quenched by addition of excess cold SAM (100 μ M final concentration). Quenched reaction mixtures were transferred to a streptavidin-coated Flashplate (Perkin Elmer, catalog number SMP410), and allowed to bind for one hour before the plates were washed in a Biotek EL-405 \times plate-washer and read on a TopCount NXT HTS scintillation and luminescence counter (PerkinElmer).

Example 14

Enzymology

[0647] Recombinant 4-component PRC2 complexes were prepared with wild-type and either A677G or A687V mutant versions of EZH2 (see Example 12 above; Cao et al. (2004) *Mol Cell* 15:57-67). Each complex was initially tested for the ability to catalyze ³H-methyl transfer from labeled S-adenosyl methionine (SAM) to each of the four H3:21-44 peptides. Enzyme was serially diluted, and a mixture of peptide (200 nM) and SAM (200 nM ³H-SAM and 800 nM unlabeled SAM) was added. Reactions were quenched at 15 minute intervals by the addition of an excess of unlabeled SAM and reaction velocity was calculated based on the linear regression of raw counts per minute (CPM) vs. time. As shown in FIG. 15, the mutant enzymes displayed a different pattern of activity than the wild-type enzyme. The A677G and A687V mutants had robust activity on all of the unmethylated, monomethylated and dimethylated H3:21-44 peptides, whereas the wild-type enzyme only showed robust activity on the unmethylated and monomethylated peptides. The control peptide, containing fully trimethylated H3-K27 was not methylated in the assay, indicating that H3-K27 was the target lysine.

Example 15

Enzymology

[0648] To further understand the enzymatic activity of these mutants, the origins of the differential substrate specificities of wild-type and mutant EZH2 were explored through steady-state enzyme kinetics. Reactions containing a titration of the H3-K27 peptides with fixed enzyme (4 nM) and SAM (200 nM ³H-SAM and 800 nM unlabeled SAM) were performed. In all cases the peptidic substrates displayed sigmoidal binding behavior; hence the concentration of peptide resulting in half-maximal velocity is reported here as $K_{1/2}$ instead of the more common Michaelis-Menten constant, K_m (Copeland (2005) *Evaluation of Enzyme Inhibitors in Drug Discovery: A Guide to Medicinal Chemists and Pharmacologists*, Wiley). As summarized in Table 6, the mutations have an effect on ground-state substrate recognition as demonstrated by lower $K_{1/2}$ for the unmethylated peptide substrates and higher $K_{1/2}$ values for the dimethylated peptide substrates. Additionally, the maximal velocities of the enzymes are affected by the mutations. The A677G mutation leads to 2.9-, 3.7- and 22-fold increases in k_{cat} on respective un-, mono-, and dimethyl H3-K27 peptide substrates, while the A687V mutation results in a 3-fold reduction in k_{cat} on the unmethyl H3-K27 peptide, but produces respective 3.5- and 2.5-fold increases in k_{cat} on the mono- and dimethyl H3-K27 peptides. The SAM K_m displayed minimal variation among the enzyme forms, 403 \pm 64 nM to 899 \pm 89 nM on the substrate containing the preferred methylation state at the H3-K27 residue.

TABLE 6

Summary of Steady-State Enzyme Kinetics for WT and Mutant EZH2 Enzymes				
Enzyme	Peptide Substrate H3- K27 Methylation Status	K _{1/2} (nM)	k _{cat} (h ⁻¹)	k _{cat} /K _{1/2} (h ⁻¹ · nM ⁻¹ × 10 ⁻⁴)
*WT	0	154 ± 12	4.80 ± 0.20	305 ± 26
	1	337 ± 26	3.33 ± 0.21	99 ± 9
	2	144 ± 11	1.08 ± 0.04	75 ± 6
A677G	0	88 ± 6	14.05 ± 0.54	1590 ± 120
	1	222 ± 51	12.25 ± 1.67	570 ± 150
	2	522 ± 117	23.85 ± 3.60	450 ± 120
A687V	0	43 ± 3	1.58 ± 0.08	370 ± 30
	1	176 ± 13	11.49 ± 0.50	650 ± 60
	2	352 ± 155	2.70 ± 0.65	80 ± 40

*Wild-type data was previously published in Wigle et al., *Febs Lett* (2011) Oct 3; 585(19): 3011-4 using 4-component EZH2.

Example 16

Inhibition of Wild-Type EZH2 and EZH2 Mutants
by EZH2 Inhibitors

[0649] Test compounds were serially diluted 3-fold in DMSO in a 10 point-curve and 1 µL was spotted into a 384-well microplate in duplicate using a Platemate Plus equipped with 384-channel head (Thermo Scientific). The final top concentration of test compounds in the assay was 10 µM. Positive control (100% inhibition standard) was 1 mM final concentration of SAH and negative control (0% inhibition standard) contained 1 µL of DMSO. Test compounds were then incubated for 30 minutes with 40 µL per well of wild-type EZH2 (final concentration was 4 nM), Y641F EZH2 (final concentration was 0.1 nM) and A677G and A687V EZH2 (for each, final concentration was 2 nM) and peptide in 1× assay buffer (20 mM BICINE pH=7.6, 1 mM DTT, 0.002% Tween 20, 0.005% BSG). For the wild-type EZH2 and A677G EZH2 assays, biotinylated peptide H3:21-44 with unmethylated K27 was present at a final concentration of 200 nM, while in the A687V EZH2 assay, biotinylated peptide H3:21-44 with monomethylated K27 was present at a final concentration of 200 nM and in the Y641F EZH2 assay biotinylated peptide H3:21-44 with dimethylated K27 was present at a final concentration of 200 nM. To initiate the reaction containing the wild-type EZH2 enzyme, a substrate mix of 10 µL per well was added that contained unlabeled SAM (final concentration was 1800 nM) and ³H-SAM (final concentration was 200 nM) in 1× assay buffer. To initiate the reaction containing the Y641F EZH2 enzyme a substrate mix of 10 µL per well was added that contained unlabeled SAM (final concentration was 700 nM), and ³H-SAM (final concentration was 300 nM) in 1× assay buffer. To initiate the reactions containing the A677G or A687V EZH2, a substrate mix of 10 µL per well was added which contained unlabeled SAM (final concentration was 400 nM) and ³H-SAM (final concentration was 100 nM). Reactions proceeded for 90 min, then were quenched with excess unlabeled SAM (167 µM), then were transferred to a streptavidin-coated Flashplate (PerkinElmer, catalog number SMP410), allowed to bind for one hour, and detected on a TopCount NXT HTS (PerkinElmer). The IC₅₀ values are obtained from 4-parameter fits of the % inhibition of enzyme activity and are tabulated in Table 7. The formulae used to derive IC₅₀ values are indicated below.

% Inhibition Calculation

[0650]

$$\% inh = 100 - \left(\frac{dpm_{cmpd} - dpm_{min}}{dpm_{max} - dpm_{min}} \right) \times 100$$

Where dpm=disintegrations per minute, cmpd=signal in assay well, and min and max are the respective minimum and maximum signal controls.

Four-Parameter 1050 Fit

[0651]

$$Y = \text{Bottom} + \frac{(\text{Top} - \text{Bottom})}{\left(1 + \left(\frac{X}{IC_{50}} \right)^{\text{HillCoefficient}} \right)}$$

[0652] Where top and bottom are the normally allowed to float, but may be fixed at 100 or 0 respectively in a 3-parameter fit. The Hill Coefficient normally allowed to float but may also be fixed at 1 in a 3-parameter fit. Y is the % inhibition and X is the compound concentration.

TABLE 7

Inhibition of wild-type and mutant EZH2 by EZH2 inhibitors					
Compound Number	EZH2 Inhibitor	WT (uM)	Y641F (uM)	A677G (uM)	A687V (uM)
1.	SAH	6.9082	16.6193	6.2379	5.9034
2.	EPZ004710	2.9758	3.7887	0.3187	0.5378
3.	EPZ004744	1.5203	0.7432	0.1128	0.1354
4.	EPZ005030	0.2600	0.1846	0.0418	0.0504
5.	EPZ005100	0.3579	0.2923	0.0316	0.0599
6.	EPZ005260	2.3755	1.5781	0.2130	0.3189
7.	EPZ005687	0.0950	0.0750	0.0113	0.0082
8.	EPZ006089	0.2300	0.3110	0.0190	0.0277
9.	EPZ006438	0.0062	0.0111	0.0022	0.0015
10.	EPZ006632	0.0025	0.0034	0.0040	0.0032
11.	EPZ007038	0.0099	0.0123	0.0060	0.0032
12.	EPZ007209	0.0065	0.0074	0.0086	0.0068
13.	EPZ007210	0.0043	0.0044	0.0038	0.0036
14.	EPZ007227	0.0143	0.0207	0.0122	0.0128
15.	EPZ007426	0.0181	0.0088	0.0034	0.0040
16.	EPZ007428	0.0014	0.0055	0.0019	0.0021
17.	EPZ007478	0.0088	0.0114	0.0042	0.0071
18.	EPZ007648	0.0025	0.0079	0.0058	0.0067
19.	EPZ007649	0.0094	0.0092	0.0096	0.0082
20.	EPZ007655	0.0125	0.0104	0.0192	0.0171
21.	EPZ007692	0.0100	0.0117	0.0116	0.0103
22.	EPZ007789	0.0108	0.0114	0.0048	0.0051
23.	EPZ007790	0.0169	0.0158	0.0073	0.0065
24.	EPZ008205	0.0129	0.0093	0.0112	0.0101
25.	EPZ008277	0.0333	0.0092	0.0023	0.0055
26.	EPZ008278	0.0384	0.0106	0.0093	0.0191
27.	EPZ008279	0.0223	0.0182	0.0022	0.0051
28.	EPZ008280	0.0067	0.0029	0.0028	0.0039
29.	EPZ008286	0.0043	0.0025	0.0015	0.0018
30.	EPZ008335	0.0065	0.0033	0.0015	0.0029
31.	EPZ008336	0.0057	0.0036	0.0013	0.0024
32.	EPZ008337	0.0087	0.0015	0.0008	0.0014
33.	EPZ008338	0.0120	0.0096	0.0031	0.0072
34.	EPZ008344	0.0124	0.0036	0.0016	0.0046
35.	EPZ008491	0.0091	0.0014	0.0029	0.0029
36.	EPZ008493	0.1320		0.0127	0.0882
37.	EPZ008494	0.0079	0.0046	0.0028	0.0038
38.	EPZ008495	0.0134	0.0104	0.0063	0.0084

TABLE 7-continued

Inhibition of wild-type and mutant EZH2 by EZH2 inhibitors					
Compound Number	EZH2 Inhibitor	WT (uM)	Y641F (uM)	A677G (uM)	A687V (uM)
39.	EPZ008496	0.0154	0.0104	0.0040	0.0076
40.	EPZ008497	>10.0 uM	3.2876	0.9735	1.2184
41.	EPZ008592	0.0145	0.0051	0.0035	0.0075
42.	EPZ008623	0.2440	0.1835	0.0922	0.1067
43.	EPZ008630	0.0034	0.0029	0.0035	0.0032
44.	EPZ008681	0.0029	0.0015	0.0027	0.0047
45.	EPZ008686	0.0073	0.0055	0.0045	0.0096
46.	EPZ008989	0.0094	0.0069	0.0022	0.0028
47.	EPZ008990	0.0061	0.0067	0.0016	0.0017
48.	EPZ008991	0.0348	0.0293	0.0094	0.0193
49.	EPZ008992	0.3333	0.1678	0.0638	0.1188
50.	EPZ008994	0.0715	0.0275	0.0121	0.0205
51.	EPZ009090	0.0300	0.0111	0.0120	0.0227
52.	EPZ009097	0.0047	0.0039	0.0046	0.0036
53.	EPZ009099	0.0765	0.0255	0.0057	0.0222
54.	EPZ009152	0.0030	0.0031	0.0029	0.0033
55.	EPZ009153	0.0052	0.0032	0.0056	0.0037
56.	EPZ009154	0.0278	0.0420	0.0141	0.0438
57.	EPZ009155	0.0563	0.0524	0.0251	0.0623
58.	EPZ009156	0.0034	0.0217	0.0051	0.0132
59.	EPZ009157	0.0397	0.0443	0.0214	0.0436
60.	EPZ009158	0.0021	0.0016	0.0027	0.0027
61.	EPZ009161	0.0009	0.0008	0.0008	0.0011
62.	EPZ009162	0.0657	0.0351	0.0146	0.0178

Example 17

A677 Mutant Shows the Highest Sensitivity to the EZH2 Inhibitors

[0653] Pfeiffer cells were obtained from ATCC(CRL-2632). WSU-DLCL2 (ACC 575) and OCI-Ly19 (ACC 528) cells were obtained from DSMZ. All cells were maintained in RPMI+10% FBS. For cell proliferation analysis, exponentially growing Pfeiffer, WSU-DLCL2, or OCI-Ly19 cells were plated, in triplicate, in 96-well plates at a density of 1×10^5 cells/mL, 5×10^4 cells/mL, or 2.5×10^5 cells/mL (respectively) in a final volume of 150 μ L. Cells were incubated with Compound #7 at final concentrations ranging from 0.011 to 25 μ M for IC_{50} determination over an 11-day time course. Cells were incubated with Compound #13 at final concentrations ranging from 0.00004 to 10 μ M for IC_{50} determination over an 11-day time course. Every 3-4 days, viable cell numbers were determined using the Guava Viacount assay (Millipore #4000-0040) and analyzed on a Guava EasyCyte Plus instrument. After cell counts, growth media and EPZ2 inhibitor (Compound #7 or Compound #13) were replaced, and cells were split back to the original plating density. The final split-adjusted number of viable cells/mL from day 11 of the time course was used to calculate the proliferation IC_{50} values, using Graphpad Prism software.

[0654] The Pfeiffer cell line, containing the heterozygous EZH2 mutation A677G, is shown to be sensitive to EZH2 inhibition by a small molecule inhibitor Compound #7. Proliferation inhibition is seen as early as 96 hr post-inhibitor treatment. The proliferation IC_{50} after 11 days for Pfeiffer cells treated with Compound #7 is 5.2 nM, compared to WSU-DLCL2 cells, which contain the Y641F heterozygous mutation, and have an IC_{50} of 270 nM or OCI-Ly19 cells, which are WT for EZH2, and have an IC_{50} of 3000 nM. These results show that in a cellular context, the sensitivity of WT and mutant EZH2 to inhibition by small molecule inhibitor is A677G>>>Y641F>>WT, as measured by proliferation.

[0655] The Pfeiffer cell line, containing the heterozygous EZH2 mutation A677G, is shown to be sensitive to EZH2 inhibition by a small molecule inhibitor Compound #13. Proliferation inhibition is seen as early as 96 hr post-inhibitor treatment. The proliferation IC_{50} after 11 days for Pfeiffer cells treated with Compound #13 is 0.4 nM, compared to WSU-DLCL2 cells, which contain the Y641F heterozygous mutation, and have an IC_{50} of 4.9 nM or OCI-Ly19 cells, which are WT for EZH2, and have an IC_{50} of 430 nM. These results show that in a cellular context, the sensitivity of WT and mutant EZH2 to inhibition by small molecule inhibitor is A677G>>>Y641F>>WT, as measured by proliferation.

EQUIVALENTS

[0656] While several embodiments of the present invention have been described and illustrated herein, those of ordinary skill in the art will readily envision a variety of other means and/or structures for performing the functions and/or obtaining the results and/or one or more of the advantages described herein, and each of such variations and/or modifications is deemed to be within the scope of the present invention. More generally, those skilled in the art will readily appreciate that all parameters, dimensions, materials, and configurations described herein are meant to be exemplary and that the actual parameters, dimensions, materials, and/or configurations will depend upon the specific application or applications for which the teachings of the present invention is/are used. Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. It is, therefore, to be understood that the foregoing embodiments are presented by way of example only and that, within the scope of the appended claims and equivalents thereto, the invention may be practiced otherwise than as specifically described and claimed. The present invention is directed to each individual feature, system, article, material, kit, and/or method described herein. In addition, any combination of two or more such features, systems, articles, materials, kits, and/or methods, if such features, systems, articles, materials, kits, and/or methods are not mutually inconsistent, is included within the scope of the present invention.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 22
 <210> SEQ ID NO 1
 <211> LENGTH: 746
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 1

```

Met Gly Gln Thr Gly Lys Lys Ser Glu Lys Gly Pro Val Cys Trp Arg
1      5      10      15
Lys Arg Val Lys Ser Glu Tyr Met Arg Leu Arg Gln Leu Lys Arg Phe
20      25      30
Arg Arg Ala Asp Glu Val Lys Ser Met Phe Ser Ser Asn Arg Gln Lys
35      40      45
Ile Leu Glu Arg Thr Glu Ile Leu Asn Gln Glu Trp Lys Gln Arg Arg
50      55      60
Ile Gln Pro Val His Ile Leu Thr Ser Val Ser Ser Leu Arg Gly Thr
65      70      75      80
Arg Glu Cys Ser Val Thr Ser Asp Leu Asp Phe Pro Thr Gln Val Ile
85      90      95
Pro Leu Lys Thr Leu Asn Ala Val Ala Ser Val Pro Ile Met Tyr Ser
100     105     110
Trp Ser Pro Leu Gln Gln Asn Phe Met Val Glu Asp Glu Thr Val Leu
115     120     125
His Asn Ile Pro Tyr Met Gly Asp Glu Val Leu Asp Gln Asp Gly Thr
130     135     140
Phe Ile Glu Glu Leu Ile Lys Asn Tyr Asp Gly Lys Val His Gly Asp
145     150     155     160
Arg Glu Cys Gly Phe Ile Asn Asp Glu Ile Phe Val Glu Leu Val Asn
165     170     175
Ala Leu Gly Gln Tyr Asn Asp Asp Asp Asp Asp Asp Gly Asp Asp
180     185     190
Pro Glu Glu Arg Glu Glu Lys Gln Lys Asp Leu Glu Asp His Arg Asp
195     200     205
Asp Lys Glu Ser Arg Pro Pro Arg Lys Phe Pro Ser Asp Lys Ile Phe
210     215     220
Glu Ala Ile Ser Ser Met Phe Pro Asp Lys Gly Thr Ala Glu Glu Leu
225     230     235     240
Lys Glu Lys Tyr Lys Glu Leu Thr Glu Gln Gln Leu Pro Gly Ala Leu
245     250     255
Pro Pro Glu Cys Thr Pro Asn Ile Asp Gly Pro Asn Ala Lys Ser Val
260     265     270
Gln Arg Glu Gln Ser Leu His Ser Phe His Thr Leu Phe Cys Arg Arg
275     280     285
Cys Phe Lys Tyr Asp Cys Phe Leu His Pro Phe His Ala Thr Pro Asn
290     295     300
Thr Tyr Lys Arg Lys Asn Thr Glu Thr Ala Leu Asp Asn Lys Pro Cys
305     310     315     320
Gly Pro Gln Cys Tyr Gln His Leu Glu Gly Ala Lys Glu Phe Ala Ala
325     330     335
Ala Leu Thr Ala Glu Arg Ile Lys Thr Pro Pro Lys Arg Pro Gly Gly
340     345     350
Arg Arg Arg Gly Arg Leu Pro Asn Asn Ser Ser Arg Pro Ser Thr Pro
355     360     365
Thr Ile Asn Val Leu Glu Ser Lys Asp Thr Asp Ser Asp Arg Glu Ala
370     375     380
Gly Thr Glu Thr Gly Gly Glu Asn Asn Asp Lys Glu Glu Glu Glu Lys
385     390     395     400

```

```
<210> SEQ ID NO 2
<211> LENGTH: 2723
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

ggcggcgcgtt gattgggctg gggggggccaa ataaaagcga tggcgattgg gctgccgcgt      60
ttggcgctcg gtcgggtcgc gtccgacacc cgggtgggact cagaaggcag tggagccccc      120
```

-continued

gcggcgccgg cgccggcgcg cggggggcgac gcgcgggaac aacgcgagtc ggccgcgggg	180
acgaagaata atcatggggc agactgggaa gaaatctgag aaggggaccag tttgttgccg	240
gaagcgtgta aaatcagagt acatgcgact gagacagctc aagaggttca gacgagctga	300
tgaagtaaa agtatgttta gttccaatcg tcagaaaatt ttggaagaa cggaatctt	360
aaaccaagaa tggaacagc gaaggataca gctgtgcac atcctgactt ctgtgagctc	420
attgcgccc actagggagt gttcgggtgac cagtgcactg gattttccaa cacaagtcac	480
cccattaaa actctgaatg cagttgcttc agtaccata atgtattctt ggtctcccct	540
acagcagaat ttatggttg aagatgaaac tgttttcat aacattcctt atatgggaga	600
tgaagtttta gatcaggatg gtactttcat tgaagaacta ataaaaaatt atgatgggaa	660
agtacacggg gatagagaat gtgggtttat aaatgatgaa atttttgttg agttggtgaa	720
tgcccttgg caatataatg atgatgacga tgatgatgat ggagacgac ctgaagaaag	780
agaagaaaag cagaaagatc tggaggatca ccgagatgat aaagaaagcc gccacctc	840
gaaatttct tctgataaaa tttttgaagc catttctca atgtttccag ataagggcac	900
agcagaagaa ctaaaagaaa aatataaaga actcaccgaa cagcagctcc caggcgact	960
tcctctgaa tgtaccccca acatagatgg accaaatgct aaatctgttc agagagagca	1020
aagcttacac tcctttcata cgctttctg taggcgatgt tttaaatag actgcttct	1080
acatcgtaag tgcaattatt cttttcatgc aacacccaac acttataagc ggaagaacac	1140
agaaacagct ctagacaaca aacctgttg accacagtgt taccagcatt tggaggagc	1200
aaaggagttt gctgctgctc tcaccgctga gcggataaag accccaccaa aacgtccagg	1260
aggccgcaga agaggacggc ttccaataa cagtagcagg ccagcacc ccaccattaa	1320
tgtgctggaa tcaaaggata cagacagtga tagggaagca gggactgaaa cggggggaga	1380
gaacaatgat aaagaagaag aagagaagaa agatgaaact tcgagctcct ctgaagcaaa	1440
ttctcgggt caaacaccaa taaagatgaa gccaaatatt gaacctctg agaatgtgga	1500
gtggagtggt gctgaagcct caatgtttag agtcctcatt ggcacttact atgacaattt	1560
ctgtgccatt gctaggttaa ttgggaccaa aacatgtaga cagggtgatg agtttagagt	1620
caaagaatct agcatcatag ctccagctcc cgctgaggat gtggatactc ctccaaggaa	1680
aaagaagagg aaacaccggt tgtgggctgc aactgcaga aagatacagc tgaagaaagga	1740
cggtcctct aacctgttt acaactatca accctgtgat catccacggc agccttgtga	1800
cagttcgtgc cctgtgtgta tagcacaaaa tttttgtgaa aagttttgtc aatgtagttc	1860
agagtgtcaa aaccgcttcc cgggatgccg ctgcaaagca cagtgcacaa ccaagcagt	1920
cccgctac ctggctgtcc gagagtgtga cctgacctc tgtcttactt gtggagccgc	1980
tgaccattgg gacagtaaaa atgtgtctg caagaactgc agtattcagc ggggctccaa	2040
aaagcatcta ttgctggcac catctgacgt ggcaggctgg gggattttta tcaaagatcc	2100
tgtgcagaaa aatgaattca tctcagaata ctgtggagag attatttctc aagatgaagc	2160
tgacagaaga gggaaagtgt atgataaata catgtgcagc tttctgttca acttgaacaa	2220
tgattttgtg gtggatgcaa cccgcaaggg taacaaaatt cgttttgcaa atcattcggt	2280
aaatccaaac tgctatgcaa aagttatgat ggttaacggt gatcacagga taggtatttt	2340
tgccaagaga gccatccaga ctggcgaaga gctgtttttt gattacagat acagccaggc	2400

-continued

```

tgatgccctg aagtatgtcg gcatcgaaag agaaatggaa atcccttgac atctgctacc 2460
tctctcccc tctctgaaa cagctgcctt agcttcagga acctcgagta ctgtgggcaa 2520
tttagaaaaa gaacatgcag ttgaaattc tgaatttgca aagtactgta agaataattt 2580
atagtaatga gtttaaaaat caacttttta ttgccttctc accagctgca aagtgttttg 2640
taccagtgaa tttttgcaat aatgcagtat ggtacatttt tcaactttga ataaagaata 2700
cttgaacttg tccttggtga atc 2723

```

```

<210> SEQ ID NO 3
<211> LENGTH: 751
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 3

```

```

Met Gly Gln Thr Gly Lys Lys Ser Glu Lys Gly Pro Val Cys Trp Arg
1          5          10          15
Lys Arg Val Lys Ser Glu Tyr Met Arg Leu Arg Gln Leu Lys Arg Phe
20          25          30
Arg Arg Ala Asp Glu Val Lys Ser Met Phe Ser Ser Asn Arg Gln Lys
35          40          45
Ile Leu Glu Arg Thr Glu Ile Leu Asn Gln Glu Trp Lys Gln Arg Arg
50          55          60
Ile Gln Pro Val His Ile Leu Thr Ser Val Ser Ser Leu Arg Gly Thr
65          70          75          80
Arg Glu Cys Ser Val Thr Ser Asp Leu Asp Phe Pro Thr Gln Val Ile
85          90          95
Pro Leu Lys Thr Leu Asn Ala Val Ala Ser Val Pro Ile Met Tyr Ser
100         105         110
Trp Ser Pro Leu Gln Gln Asn Phe Met Val Glu Asp Glu Thr Val Leu
115         120         125
His Asn Ile Pro Tyr Met Gly Asp Glu Val Leu Asp Gln Asp Gly Thr
130         135         140
Phe Ile Glu Glu Leu Ile Lys Asn Tyr Asp Gly Lys Val His Gly Asp
145         150         155         160
Arg Glu Cys Gly Phe Ile Asn Asp Glu Ile Phe Val Glu Leu Val Asn
165         170         175
Ala Leu Gly Gln Tyr Asn Asp Asp Asp Asp Asp Asp Gly Asp Asp
180         185         190
Pro Glu Glu Arg Glu Glu Lys Gln Lys Asp Leu Glu Asp His Arg Asp
195         200         205
Asp Lys Glu Ser Arg Pro Pro Arg Lys Phe Pro Ser Asp Lys Ile Phe
210         215         220
Glu Ala Ile Ser Ser Met Phe Pro Asp Lys Gly Thr Ala Glu Glu Leu
225         230         235         240
Lys Glu Lys Tyr Lys Glu Leu Thr Glu Gln Gln Leu Pro Gly Ala Leu
245         250         255
Pro Pro Glu Cys Thr Pro Asn Ile Asp Gly Pro Asn Ala Lys Ser Val
260         265         270
Gln Arg Glu Gln Ser Leu His Ser Phe His Thr Leu Phe Cys Arg Arg
275         280         285
Cys Phe Lys Tyr Asp Cys Phe Leu His Arg Lys Cys Asn Tyr Ser Phe
290         295         300

```

-continued

His	Ala	Thr	Pro	Asn	Thr	Tyr	Lys	Arg	Lys	Asn	Thr	Glu	Thr	Ala	Leu	305	310	315	320
Asp	Asn	Lys	Pro	Cys	Gly	Pro	Gln	Cys	Tyr	Gln	His	Leu	Glu	Gly	Ala	325	330	335	
Lys	Glu	Phe	Ala	Ala	Ala	Leu	Thr	Ala	Glu	Arg	Ile	Lys	Thr	Pro	Pro	340	345	350	
Lys	Arg	Pro	Gly	Gly	Arg	Arg	Arg	Gly	Arg	Leu	Pro	Asn	Asn	Ser	Ser	355	360	365	
Arg	Pro	Ser	Thr	Pro	Thr	Ile	Asn	Val	Leu	Glu	Ser	Lys	Asp	Thr	Asp	370	375	380	
Ser	Asp	Arg	Glu	Ala	Gly	Thr	Glu	Thr	Gly	Gly	Glu	Asn	Asn	Asp	Lys	385	390	395	400
Glu	Glu	Glu	Glu	Lys	Lys	Asp	Glu	Thr	Ser	Ser	Ser	Ser	Glu	Ala	Asn	405	410	415	
Ser	Arg	Cys	Gln	Thr	Pro	Ile	Lys	Met	Lys	Pro	Asn	Ile	Glu	Pro	Pro	420	425	430	
Glu	Asn	Val	Glu	Trp	Ser	Gly	Ala	Glu	Ala	Ser	Met	Phe	Arg	Val	Leu	435	440	445	
Ile	Gly	Thr	Tyr	Tyr	Asp	Asn	Phe	Cys	Ala	Ile	Ala	Arg	Leu	Ile	Gly	450	455	460	
Thr	Lys	Thr	Cys	Arg	Gln	Val	Tyr	Glu	Phe	Arg	Val	Lys	Glu	Ser	Ser	465	470	475	480
Ile	Ile	Ala	Pro	Ala	Pro	Ala	Glu	Asp	Val	Asp	Thr	Pro	Pro	Arg	Lys	485	490	495	
Lys	Lys	Arg	Lys	His	Arg	Leu	Trp	Ala	Ala	His	Cys	Arg	Lys	Ile	Gln	500	505	510	
Leu	Lys	Lys	Asp	Gly	Ser	Ser	Asn	His	Val	Tyr	Asn	Tyr	Gln	Pro	Cys	515	520	525	
Asp	His	Pro	Arg	Gln	Pro	Cys	Asp	Ser	Ser	Cys	Pro	Cys	Val	Ile	Ala	530	535	540	
Gln	Asn	Phe	Cys	Glu	Lys	Phe	Cys	Gln	Cys	Ser	Ser	Glu	Cys	Gln	Asn	545	550	555	560
Arg	Phe	Pro	Gly	Cys	Arg	Cys	Lys	Ala	Gln	Cys	Asn	Thr	Lys	Gln	Cys	565	570	575	
Pro	Cys	Tyr	Leu	Ala	Val	Arg	Glu	Cys	Asp	Pro	Asp	Leu	Cys	Leu	Thr	580	585	590	
Cys	Gly	Ala	Ala	Asp	His	Trp	Asp	Ser	Lys	Asn	Val	Ser	Cys	Lys	Asn	595	600	605	
Cys	Ser	Ile	Gln	Arg	Gly	Ser	Lys	Lys	His	Leu	Leu	Leu	Ala	Pro	Ser	610	615	620	
Asp	Val	Ala	Gly	Trp	Gly	Ile	Phe	Ile	Lys	Asp	Pro	Val	Gln	Lys	Asn	625	630	635	640
Glu	Phe	Ile	Ser	Glu	Tyr	Cys	Gly	Glu	Ile	Ile	Ser	Gln	Asp	Glu	Ala	645	650	655	
Asp	Arg	Arg	Gly	Lys	Val	Tyr	Asp	Lys	Tyr	Met	Cys	Ser	Phe	Leu	Phe	660	665	670	
Asn	Leu	Asn	Asn	Asp	Phe	Val	Val	Asp	Ala	Thr	Arg	Lys	Gly	Asn	Lys	675	680	685	
Ile	Arg	Phe	Ala	Asn	His	Ser	Val	Asn	Pro	Asn	Cys	Tyr	Ala	Lys	Val	690	695	700	
Met	Met	Val	Asn	Gly	Asp	His	Arg	Ile	Gly	Ile	Phe	Ala	Lys	Arg	Ala	705	710	715	720

-continued

Ile Gln Thr Gly Glu Glu Leu Phe Phe Asp Tyr Arg Tyr Ser Gln Ala
 725 730 735

Asp Ala Leu Lys Tyr Val Gly Ile Glu Arg Glu Met Glu Ile Pro
 740 745 750

<210> SEQ ID NO 4
 <211> LENGTH: 2591
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

```

ggcgccgctt gattgggctg ggggggcca ataaaagcga tggcgattgg gctgccgctg    60
ttggcgctcg gtccggtcgc gtccgacacc cggtagggact cagaaggcag tggagccccg   120
gcggcgccgg cggcgccgcg cggggggcgc gcgcgggaac aacgcgagtc ggcgcgcggg   180
acgaagaata atcatggggc agactgggaa gaaatctgag aagggaccag tttgttgccg   240
gaagcgtgta aaatcagagt acatgcgact gagacagctc aagaggttca gacgagctga   300
tgaagtaaa agtatgttta gttccaatcg tcagaaaatt ttggaagaa cggaaatctt   360
aaaccaagaa tggaaacagc gaaggataca gcctgtgcac atcctgactt ctgtgagctc   420
attgcgcggg actagggagg tggaagatga aactgtttta cataacattc cttatatggg   480
agatgaagtt ttagatcagg atggtacttt cattgaagaa ctaataaaaa attatgatgg   540
gaaagtacac ggggatagag aatgtgggtt tataaatgat gaaatttttg tggagttggt   600
gaatgccctt ggtcaatata atgatgatga cgatgatgat gatggagacg atcctgaaga   660
aagagaagaa aagcagaaag atctggagga tcaccgagat gataaagaaa gccgcccacc   720
tcggaaatct cttctgata aaatttttga agccatttcc tcaatgtttc cagataaggg   780
cacagcagaa gaactaaagg aaaaatataa agaactcacc gaacagcagc tcccaggcgc   840
acttcctcct gaatgtaccc ccaacataga tggaccaa atgctaaatctg ttcagagaga   900
gcaaagctta cactccttcc atacgctttt ctgtaggcga tgttttaaat atgactgctt   960
cctacatcct tttcatgcaa cacccaacac ttataagcgg aagaacacag aaacagctct  1020
agacaacaaa cttgtgggac cacagtgtta ccagcatttg gagggagcaa aggagtttgc  1080
tgctgctctc accgctgagc ggataaagac ccacccaaaa cgtccaggag gccgcagaag  1140
aggacggctt cccaataaca gtagcaggcc cagcaccccc accattaatg tgctggaatc  1200
aaaggataca gacagtata ggaagcagg gactgaaacg gggggagaga acaatgataa  1260
agaagaagaa gagaagaaag atgaaacttc gagctcctct gaagcaaatt ctcggtgtca  1320
aacaccaata aagatgaagc caaatattga acctcctgag aatgtggagt ggagtggtgc  1380
tgaagcctca atgtttagag tcctcatttg cacttactat gacaatttct gtgccattgc  1440
taggttaatt gggacaaaa catgtagaca ggtgtatgag tttagagtca aagaatctag  1500
catcatagct ccagctcccg ctgaggatgt ggatactcct ccaaggaaaa agaagaggaa  1560
acaccggttg tgggctgcac actgcagaaa gatacagctg aaaaaggacg gctcctctaa  1620
ccatgtttac aactatcaac cctgtgatca tccacggcag cctgtgaca gttcgtgccc  1680
ttgtgtgata gcacaaaatt tttgtgaaaa gttttgtcaa ttagagttcag agtgtcaaaa  1740
ccgctttccg gtagtccgct gcaaagcaca gtgcaacacc aagcagtgcc cgtgctacct  1800
ggctgtccga gagtgtgacc ctgacctctg tottacttgt ggagccgctg accattggga  1860

```

-continued

```

cagtaaaaat gtgtcctgca agaactgcag tattcagcgg ggctocaaaa agcatctatt 1920
gctggcacca tctgacgtgg caggctgggg gattttttatc aaagatcctg tgcagaaaaa 1980
tgaattcatc tcagaatact gtggagagat tattttctcaa gatgaagctg acagaagagg 2040
gaaagtgtat gataaataca tgtgcagctt tctgttcaac ttgaacaatg attttgtggt 2100
ggatgcaacc cgcaagggtg acaaaattcg ttttgcaaat cattcggtaa atccaaactg 2160
ctatgcaaaa gttatgatgg ttaacggtga tcacaggata ggtatttttg ccaagagagc 2220
catccagact ggcaagagc tgttttttga ttacagatac agccaggctg atgccctgaa 2280
gtatgtcggc atcgaaagag aaatggaaat cccttgacat ctgctacctc ctccccctc 2340
ctctgaaaca gctgccttag cttcaggaac ctgagtagt gtgggcaatt tagaaaaaga 2400
acatgcagtt tgaaattctg aatttgcaaa gtactgtaag aataatttat agtaatgagt 2460
ttaaaaaatc actttttatt gcctttctcac cagctgcaaa gtgttttgta ccagtgaatt 2520
tttgcaataa tgcagtagtg tacatttttc aactttgaat aaagaatact tgaacttgtc 2580
ctgtgtgaat c 2591

```

<210> SEQ ID NO 5

<211> LENGTH: 707

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

```

Met Gly Gln Thr Gly Lys Lys Ser Glu Lys Gly Pro Val Cys Trp Arg
1           5           10           15
Lys Arg Val Lys Ser Glu Tyr Met Arg Leu Arg Gln Leu Lys Arg Phe
20          25          30
Arg Arg Ala Asp Glu Val Lys Ser Met Phe Ser Ser Asn Arg Gln Lys
35          40          45
Ile Leu Glu Arg Thr Glu Ile Leu Asn Gln Glu Trp Lys Gln Arg Arg
50          55          60
Ile Gln Pro Val His Ile Leu Thr Ser Val Ser Ser Leu Arg Gly Thr
65          70          75          80
Arg Glu Val Glu Asp Glu Thr Val Leu His Asn Ile Pro Tyr Met Gly
85          90          95
Asp Glu Val Leu Asp Gln Asp Gly Thr Phe Ile Glu Glu Leu Ile Lys
100         105         110
Asn Tyr Asp Gly Lys Val His Gly Asp Arg Glu Cys Gly Phe Ile Asn
115         120         125
Asp Glu Ile Phe Val Glu Leu Val Asn Ala Leu Gly Gln Tyr Asn Asp
130         135         140
Asp Asp Asp Asp Asp Asp Gly Asp Asp Pro Glu Glu Arg Glu Glu Lys
145         150         155         160
Gln Lys Asp Leu Glu Asp His Arg Asp Asp Lys Glu Ser Arg Pro Pro
165         170         175
Arg Lys Phe Pro Ser Asp Lys Ile Phe Glu Ala Ile Ser Ser Met Phe
180         185         190
Pro Asp Lys Gly Thr Ala Glu Glu Leu Lys Glu Lys Tyr Lys Glu Leu
195         200         205
Thr Glu Gln Gln Leu Pro Gly Ala Leu Pro Pro Glu Cys Thr Pro Asn
210         215         220
Ile Asp Gly Pro Asn Ala Lys Ser Val Gln Arg Glu Gln Ser Leu His

```

-continued

225	230	235	240
Ser Phe His Thr	Leu Phe Cys Arg Arg	Cys Phe Lys Tyr Asp Cys Phe	
	245	250	255
Leu His Pro Phe	His Ala Thr Pro Asn Thr Tyr Lys Arg Lys Asn Thr		
	260	265	270
Glu Thr Ala Leu Asp	Asn Lys Pro Cys Gly Pro Gln Cys Tyr Gln His		
	275	280	285
Leu Glu Gly Ala Lys	Glu Phe Ala Ala Ala Leu Thr Ala Glu Arg Ile		
	290	295	300
Lys Thr Pro Pro Lys	Arg Pro Gly Gly Arg Arg Arg Gly Arg Leu Pro		
	305	310	315
Asn Asn Ser Ser Arg	Pro Ser Thr Pro Thr Ile Asn Val Leu Glu Ser		
	325	330	335
Lys Asp Thr Asp Ser	Asp Arg Glu Ala Gly Thr Glu Thr Gly Gly Glu		
	340	345	350
Asn Asn Asp Lys Glu	Glu Glu Glu Lys Lys Asp Glu Thr Ser Ser Ser		
	355	360	365
Ser Glu Ala Asn Ser	Arg Cys Gln Thr Pro Ile Lys Met Lys Pro Asn		
	370	375	380
Ile Glu Pro Pro Glu	Asn Val Glu Trp Ser Gly Ala Glu Ala Ser Met		
	385	390	395
Phe Arg Val Leu Ile	Gly Thr Tyr Tyr Asp Asn Phe Cys Ala Ile Ala		
	405	410	415
Arg Leu Ile Gly Thr	Lys Thr Cys Arg Gln Val Tyr Glu Phe Arg Val		
	420	425	430
Lys Glu Ser Ser Ile	Ile Ala Pro Ala Pro Ala Glu Asp Val Asp Thr		
	435	440	445
Pro Pro Arg Lys Lys	Lys Arg Lys His Arg Leu Trp Ala Ala His Cys		
	450	455	460
Arg Lys Ile Gln Leu	Lys Lys Asp Gly Ser Ser Asn His Val Tyr Asn		
	465	470	475
Tyr Gln Pro Cys Asp	His Pro Arg Gln Pro Cys Asp Ser Ser Cys Pro		
	485	490	495
Cys Val Ile Ala Gln	Asn Phe Cys Glu Lys Phe Cys Gln Cys Ser Ser		
	500	505	510
Glu Cys Gln Asn Arg	Phe Pro Gly Cys Arg Cys Lys Ala Gln Cys Asn		
	515	520	525
Thr Lys Gln Cys Pro	Cys Tyr Leu Ala Val Arg Glu Cys Asp Pro Asp		
	530	535	540
Leu Cys Leu Thr Cys	Gly Ala Ala Asp His Trp Asp Ser Lys Asn Val		
	545	550	555
Ser Cys Lys Asn Cys	Ser Ile Gln Arg Gly Ser Lys Lys His Leu Leu		
	565	570	575
Leu Ala Pro Ser Asp	Val Ala Gly Trp Gly Ile Phe Ile Lys Asp Pro		
	580	585	590
Val Gln Lys Asn Glu	Phe Ile Ser Glu Tyr Cys Gly Glu Ile Ile Ser		
	595	600	605
Gln Asp Glu Ala Asp	Arg Arg Gly Lys Val Tyr Asp Lys Tyr Met Cys		
	610	615	620
Ser Phe Leu Phe Asn	Leu Asn Asn Asp Phe Val Val Asp Ala Thr Arg		
	625	630	635
			640

-continued

Lys Gly Asn Lys Ile Arg Phe Ala Asn His Ser Val Asn Pro Asn Cys
645 650 655

Tyr Ala Lys Val Met Met Val Asn Gly Asp His Arg Ile Gly Ile Phe
660 665 670

Ala Lys Arg Ala Ile Gln Thr Gly Glu Glu Leu Phe Phe Asp Tyr Arg
675 680 685

Tyr Ser Gln Ala Asp Ala Leu Lys Tyr Val Gly Ile Glu Arg Glu Met
690 695 700

Glu Ile Pro
705

<210> SEQ ID NO 6
<211> LENGTH: 200
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

Ser Cys Pro Cys Val Ile Ala Gln Asn Phe Cys Glu Lys Phe Cys Gln
1 5 10 15

Cys Ser Ser Glu Cys Gln Asn Arg Phe Pro Gly Cys Arg Cys Lys Ala
20 25 30

Gln Cys Asn Thr Lys Gln Cys Pro Cys Tyr Leu Ala Val Arg Glu Cys
35 40 45

Asp Pro Asp Leu Cys Leu Thr Cys Gly Ala Ala Asp His Trp Asp Ser
50 55 60

Lys Asn Val Ser Cys Lys Asn Cys Ser Ile Gln Arg Gly Ser Lys Lys
65 70 75 80

His Leu Leu Leu Ala Pro Ser Asp Val Ala Gly Trp Gly Ile Phe Ile
85 90 95

Lys Asp Pro Val Gln Lys Asn Glu Phe Ile Ser Glu Tyr Cys Gly Glu
100 105 110

Ile Ile Ser Gln Asp Glu Ala Asp Arg Arg Gly Lys Val Tyr Asp Lys
115 120 125

Tyr Met Cys Ser Phe Leu Phe Asn Leu Asn Asn Asp Phe Val Val Asp
130 135 140

Ala Thr Arg Lys Gly Asn Lys Ile Arg Phe Ala Asn His Ser Val Asn
145 150 155 160

Pro Asn Cys Tyr Ala Lys Val Met Met Val Asn Gly Asp His Arg Ile
165 170 175

Gly Ile Phe Ala Lys Arg Ala Ile Gln Thr Gly Glu Glu Leu Phe Phe
180 185 190

Asp Tyr Arg Tyr Ser Gln Ala Asp
195 200

<210> SEQ ID NO 7
<211> LENGTH: 114
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

His Leu Leu Leu Ala Pro Ser Asp Val Ala Gly Trp Gly Ile Phe Ile
1 5 10 15

Lys Asp Pro Val Gln Lys Asn Glu Phe Ile Ser Glu Tyr Cys Gly Glu
20 25 30

Ile Ile Ser Gln Asp Glu Ala Asp Arg Arg Gly Lys Val Tyr Asp Lys

-continued

35	40	45	
Tyr Met Cys Ser Phe Leu Phe Asn Leu Asn Asn Asp Phe Val Val Asp			
50	55	60	
Ala Thr Arg Lys Gly Asn Lys Ile Arg Phe Ala Asn His Ser Val Asn			
65	70	75	80
Pro Asn Cys Tyr Ala Lys Val Met Met Val Asn Gly Asp His Arg Ile			
85	90	95	
Gly Ile Phe Ala Lys Arg Ala Ile Gln Thr Gly Glu Glu Leu Phe Phe			
100	105	110	

Asp Tyr

<210> SEQ ID NO 8
 <211> LENGTH: 342
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

catctattgc tggcaccatc tgacgtggca ggctggggga tttttatcaa agatcctgtg	60
cagaaaaatg aattcatctc agaatactgt ggagagatta tttctcaaga tgaagctgac	120
agaagagggga aagtgtatga taaatacatg tgcagctttc tgttcaactt gaacaatgat	180
tttgtggtgg atgcaacccg caagggtaac aaaattcggt ttgcaaatca ttcggtaaat	240
ccaaactgct atgcaaaagt tatgatggtt aacggtgac acaggatagg tatttttgcc	300
aagagagcca tccagactgg cgaagagctg ttttttgatt ac	342

<210> SEQ ID NO 9
 <211> LENGTH: 746
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Y641 mutant of Homo sapiens EZH2, wherein x can be any amino acid residue other than tyrosine (Y)
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (641) .. (641)
 <223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid other than tyrosine (Y)

<400> SEQUENCE: 9

Met Gly Gln Thr Gly Lys Lys Ser Glu Lys Gly Pro Val Cys Trp Arg	
1	15
Lys Arg Val Lys Ser Glu Tyr Met Arg Leu Arg Gln Leu Lys Arg Phe	
20	30
Arg Arg Ala Asp Glu Val Lys Ser Met Phe Ser Ser Asn Arg Gln Lys	
35	45
Ile Leu Glu Arg Thr Glu Ile Leu Asn Gln Glu Trp Lys Gln Arg Arg	
50	60
Ile Gln Pro Val His Ile Leu Thr Ser Val Ser Ser Leu Arg Gly Thr	
65	80
Arg Glu Cys Ser Val Thr Ser Asp Leu Asp Phe Pro Thr Gln Val Ile	
85	95
Pro Leu Lys Thr Leu Asn Ala Val Ala Ser Val Pro Ile Met Tyr Ser	
100	110
Trp Ser Pro Leu Gln Gln Asn Phe Met Val Glu Asp Glu Thr Val Leu	
115	125
His Asn Ile Pro Tyr Met Gly Asp Glu Val Leu Asp Gln Asp Gly Thr	

130					135					140					
Phe 145	Ile	Glu	Glu	Leu	Ile 150	Lys	Asn	Tyr	Asp	Gly 155	Lys	Val	His	Gly	Asp 160
Arg	Glu	Cys	Gly	Phe 165	Ile	Asn	Asp	Glu	Ile 170	Phe	Val	Glu	Leu	Val	Asn 175
Ala	Leu	Gly	Gln	Tyr 180	Asn	Asp	Asp	Asp	Asp	Asp	Asp	Asp	Gly	Asp	Asp 190
Pro	Glu	Glu	Arg	Glu	Glu	Lys 200	Gln	Lys	Asp	Leu	Glu	Asp	His	Arg	Asp 205
Asp 210	Lys	Glu	Ser	Arg	Pro	Pro 215	Arg	Lys	Phe	Pro	Ser 220	Asp	Lys	Ile	Phe
Glu 225	Ala	Ile	Ser	Ser	Met 230	Phe	Pro	Asp	Lys	Gly 235	Thr	Ala	Glu	Glu	Leu 240
Lys	Glu	Lys	Tyr	Lys 245	Glu	Leu	Thr	Glu	Gln	Gln	Leu	Pro	Gly	Ala	Leu 255
Pro	Pro	Glu	Cys 260	Thr	Pro	Asn	Ile	Asp 265	Gly	Pro	Asn	Ala	Lys	Ser	Val 270
Gln	Arg	Glu	Gln	Ser	Leu	His 280	Ser	Phe	His	Thr	Leu	Phe 285	Cys	Arg	Arg
Cys 290	Phe	Lys	Tyr	Asp	Cys	Phe 295	Leu	His	Pro	Phe	His 300	Ala	Thr	Pro	Asn
Thr 305	Tyr	Lys	Arg	Lys	Asn 310	Thr	Glu	Thr	Ala	Leu 315	Asp	Asn	Lys	Pro	Cys 320
Gly	Pro	Gln	Cys	Tyr 325	Gln	His	Leu	Glu	Gly 330	Ala	Lys	Glu	Phe	Ala	Ala 335
Ala	Leu	Thr	Ala 340	Glu	Arg	Ile	Lys	Thr 345	Pro	Pro	Lys	Arg	Pro 350	Gly	Gly
Arg	Arg	Arg 355	Gly	Arg	Leu	Pro	Asn 360	Asn	Ser	Ser	Arg	Pro 365	Ser	Thr	Pro
Thr 370	Ile	Asn	Val	Leu	Glu	Ser 375	Lys	Asp	Thr	Asp 380	Ser	Asp	Arg	Glu	Ala
Gly 385	Thr	Glu	Thr	Gly 390	Gly	Glu	Asn	Asn	Asp 395	Lys	Glu	Glu	Glu	Glu	Lys 400
Lys	Asp	Glu	Thr 405	Ser	Ser	Ser	Ser	Glu 410	Ala	Asn	Ser	Arg	Cys 415	Gln	Thr
Pro	Ile	Lys	Met 420	Lys	Pro	Asn	Ile	Glu 425	Pro	Pro	Glu	Asn 430	Val	Glu	Trp
Ser	Gly	Ala 435	Glu	Ala	Ser	Met 440	Phe	Arg	Val	Leu	Ile 445	Gly	Thr	Tyr	Tyr
Asp 450	Asn	Phe	Cys	Ala	Ile	Ala 455	Arg	Leu	Ile	Gly 460	Thr	Lys	Thr	Cys	Arg
Gln 465	Val	Tyr	Glu	Phe 470	Arg	Val	Lys	Glu	Ser	Ser 475	Ile	Ile	Ala	Pro	Ala 480
Pro	Ala	Glu	Asp 485	Val	Asp	Thr	Pro	Pro	Arg 490	Lys	Lys	Lys	Arg	Lys	His 495
Arg	Leu	Trp 500	Ala	Ala	His	Cys	Arg 505	Lys	Ile	Gln	Leu	Lys 510	Lys	Asp	Gly
Ser	Ser	Asn 515	His	Val	Tyr	Asn 520	Tyr	Gln	Pro	Cys	Asp 525	His	Pro	Arg	Gln
Pro	Cys	Asp 530	Ser	Ser	Cys	Pro 535	Cys	Val	Ile	Ala	Gln 540	Asn	Phe	Cys	Glu

-continued

Lys Phe Cys Gln Cys Ser Ser Glu Cys Gln Asn Arg Phe Pro Gly Cys
 545 550 555 560
 Arg Cys Lys Ala Gln Cys Asn Thr Lys Gln Cys Pro Cys Tyr Leu Ala
 565 570 575
 Val Arg Glu Cys Asp Pro Asp Leu Cys Leu Thr Cys Gly Ala Ala Asp
 580 585 590
 His Trp Asp Ser Lys Asn Val Ser Cys Lys Asn Cys Ser Ile Gln Arg
 595 600 605
 Gly Ser Lys Lys His Leu Leu Leu Ala Pro Ser Asp Val Ala Gly Trp
 610 615 620
 Gly Ile Phe Ile Lys Asp Pro Val Gln Lys Asn Glu Phe Ile Ser Glu
 625 630 635 640
 Xaa Cys Gly Glu Ile Ile Ser Gln Asp Glu Ala Asp Arg Arg Gly Lys
 645 650 655
 Val Tyr Asp Lys Tyr Met Cys Ser Phe Leu Phe Asn Leu Asn Asn Asp
 660 665 670
 Phe Val Val Asp Ala Thr Arg Lys Gly Asn Lys Ile Arg Phe Ala Asn
 675 680 685
 His Ser Val Asn Pro Asn Cys Tyr Ala Lys Val Met Met Val Asn Gly
 690 695 700
 Asp His Arg Ile Gly Ile Phe Ala Lys Arg Ala Ile Gln Thr Gly Glu
 705 710 715 720
 Glu Leu Phe Phe Asp Tyr Arg Tyr Ser Gln Ala Asp Ala Leu Lys Tyr
 725 730 735
 Val Gly Ile Glu Arg Glu Met Glu Ile Pro
 740 745

<210> SEQ ID NO 10

<211> LENGTH: 746

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Y641F mutant of Homo sapiens EZH2

<400> SEQUENCE: 10

Met Gly Gln Thr Gly Lys Lys Ser Glu Lys Gly Pro Val Cys Trp Arg
 1 5 10 15
 Lys Arg Val Lys Ser Glu Tyr Met Arg Leu Arg Gln Leu Lys Arg Phe
 20 25 30
 Arg Arg Ala Asp Glu Val Lys Ser Met Phe Ser Ser Asn Arg Gln Lys
 35 40 45
 Ile Leu Glu Arg Thr Glu Ile Leu Asn Gln Glu Trp Lys Gln Arg Arg
 50 55 60
 Ile Gln Pro Val His Ile Leu Thr Ser Val Ser Ser Leu Arg Gly Thr
 65 70 75 80
 Arg Glu Cys Ser Val Thr Ser Asp Leu Asp Phe Pro Thr Gln Val Ile
 85 90 95
 Pro Leu Lys Thr Leu Asn Ala Val Ala Ser Val Pro Ile Met Tyr Ser
 100 105 110
 Trp Ser Pro Leu Gln Gln Asn Phe Met Val Glu Asp Glu Thr Val Leu
 115 120 125
 His Asn Ile Pro Tyr Met Gly Asp Glu Val Leu Asp Gln Asp Gly Thr
 130 135 140
 Phe Ile Glu Glu Leu Ile Lys Asn Tyr Asp Gly Lys Val His Gly Asp

-continued

145	150	155	160
Arg Glu Cys Gly Phe Ile Asn Asp Glu Ile Phe Val Glu Leu Val Asn	165	170	175
Ala Leu Gly Gln Tyr Asn Asp Asp Asp Asp Asp Asp Gly Asp Asp	180	185	190
Pro Glu Glu Arg Glu Glu Lys Gln Lys Asp Leu Glu Asp His Arg Asp	195	200	205
Asp Lys Glu Ser Arg Pro Pro Arg Lys Phe Pro Ser Asp Lys Ile Phe	210	215	220
Glu Ala Ile Ser Ser Met Phe Pro Asp Lys Gly Thr Ala Glu Glu Leu	225	230	235
Lys Glu Lys Tyr Lys Glu Leu Thr Glu Gln Gln Leu Pro Gly Ala Leu	245	250	255
Pro Pro Glu Cys Thr Pro Asn Ile Asp Gly Pro Asn Ala Lys Ser Val	260	265	270
Gln Arg Glu Gln Ser Leu His Ser Phe His Thr Leu Phe Cys Arg Arg	275	280	285
Cys Phe Lys Tyr Asp Cys Phe Leu His Pro Phe His Ala Thr Pro Asn	290	295	300
Thr Tyr Lys Arg Lys Asn Thr Glu Thr Ala Leu Asp Asn Lys Pro Cys	305	310	315
Gly Pro Gln Cys Tyr Gln His Leu Glu Gly Ala Lys Glu Phe Ala Ala	325	330	335
Ala Leu Thr Ala Glu Arg Ile Lys Thr Pro Pro Lys Arg Pro Gly Gly	340	345	350
Arg Arg Arg Gly Arg Leu Pro Asn Asn Ser Ser Arg Pro Ser Thr Pro	355	360	365
Thr Ile Asn Val Leu Glu Ser Lys Asp Thr Asp Ser Asp Arg Glu Ala	370	375	380
Gly Thr Glu Thr Gly Gly Glu Asn Asn Asp Lys Glu Glu Glu Lys	385	390	395
Lys Asp Glu Thr Ser Ser Ser Ser Glu Ala Asn Ser Arg Cys Gln Thr	405	410	415
Pro Ile Lys Met Lys Pro Asn Ile Glu Pro Pro Glu Asn Val Glu Trp	420	425	430
Ser Gly Ala Glu Ala Ser Met Phe Arg Val Leu Ile Gly Thr Tyr Tyr	435	440	445
Asp Asn Phe Cys Ala Ile Ala Arg Leu Ile Gly Thr Lys Thr Cys Arg	450	455	460
Gln Val Tyr Glu Phe Arg Val Lys Glu Ser Ser Ile Ile Ala Pro Ala	465	470	475
Pro Ala Glu Asp Val Asp Thr Pro Pro Arg Lys Lys Lys Arg Lys His	485	490	495
Arg Leu Trp Ala Ala His Cys Arg Lys Ile Gln Leu Lys Lys Asp Gly	500	505	510
Ser Ser Asn His Val Tyr Asn Tyr Gln Pro Cys Asp His Pro Arg Gln	515	520	525
Pro Cys Asp Ser Ser Cys Pro Cys Val Ile Ala Gln Asn Phe Cys Glu	530	535	540
Lys Phe Cys Gln Cys Ser Ser Glu Cys Gln Asn Arg Phe Pro Gly Cys	545	550	555
			560

-continued

```

Arg  Cys  Lys  Ala  Gln  Cys  Asn  Thr  Lys  Gln  Cys  Pro  Cys  Tyr  Leu  Ala
      565                      570                      575

Val  Arg  Glu  Cys  Asp  Pro  Asp  Leu  Cys  Leu  Thr  Cys  Gly  Ala  Ala  Asp
      580                      585                      590

His  Trp  Asp  Ser  Lys  Asn  Val  Ser  Cys  Lys  Asn  Cys  Ser  Ile  Gln  Arg
      595                      600                      605

Gly  Ser  Lys  Lys  His  Leu  Leu  Leu  Ala  Pro  Ser  Asp  Val  Ala  Gly  Trp
      610                      615                      620

Gly  Ile  Phe  Ile  Lys  Asp  Pro  Val  Gln  Lys  Asn  Glu  Phe  Ile  Ser  Glu
      625                      630                      635                      640

Phe  Cys  Gly  Glu  Ile  Ile  Ser  Gln  Asp  Glu  Ala  Asp  Arg  Arg  Gly  Lys
      645                      650                      655

Val  Tyr  Asp  Lys  Tyr  Met  Cys  Ser  Phe  Leu  Phe  Asn  Leu  Asn  Asn  Asp
      660                      665                      670

Phe  Val  Val  Asp  Ala  Thr  Arg  Lys  Gly  Asn  Lys  Ile  Arg  Phe  Ala  Asn
      675                      680                      685

His  Ser  Val  Asn  Pro  Asn  Cys  Tyr  Ala  Lys  Val  Met  Met  Val  Asn  Gly
      690                      695                      700

Asp  His  Arg  Ile  Gly  Ile  Phe  Ala  Lys  Arg  Ala  Ile  Gln  Thr  Gly  Glu
      705                      710                      715                      720

Glu  Leu  Phe  Phe  Asp  Tyr  Arg  Tyr  Ser  Gln  Ala  Asp  Ala  Leu  Lys  Tyr
      725                      730                      735

Val  Gly  Ile  Glu  Arg  Glu  Met  Glu  Ile  Pro
      740                      745

```

<210> SEQ ID NO 11

<211> LENGTH: 746

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Y641H mutant of Homo sapiens EZH2

<400> SEQUENCE: 11

```

Met  Gly  Gln  Thr  Gly  Lys  Lys  Ser  Glu  Lys  Gly  Pro  Val  Cys  Trp  Arg
1      5      10      15

Lys  Arg  Val  Lys  Ser  Glu  Tyr  Met  Arg  Leu  Arg  Gln  Leu  Lys  Arg  Phe
20     25     30

Arg  Arg  Ala  Asp  Glu  Val  Lys  Ser  Met  Phe  Ser  Ser  Asn  Arg  Gln  Lys
35     40     45

Ile  Leu  Glu  Arg  Thr  Glu  Ile  Leu  Asn  Gln  Glu  Trp  Lys  Gln  Arg  Arg
50     55     60

Ile  Gln  Pro  Val  His  Ile  Leu  Thr  Ser  Val  Ser  Ser  Leu  Arg  Gly  Thr
65     70     75     80

Arg  Glu  Cys  Ser  Val  Thr  Ser  Asp  Leu  Asp  Phe  Pro  Thr  Gln  Val  Ile
85     90     95

Pro  Leu  Lys  Thr  Leu  Asn  Ala  Val  Ala  Ser  Val  Pro  Ile  Met  Tyr  Ser
100    105    110

Trp  Ser  Pro  Leu  Gln  Gln  Asn  Phe  Met  Val  Glu  Asp  Glu  Thr  Val  Leu
115    120    125

His  Asn  Ile  Pro  Tyr  Met  Gly  Asp  Glu  Val  Leu  Asp  Gln  Asp  Gly  Thr
130    135    140

Phe  Ile  Glu  Glu  Leu  Ile  Lys  Asn  Tyr  Asp  Gly  Lys  Val  His  Gly  Asp
145    150    155    160

Arg  Glu  Cys  Gly  Phe  Ile  Asn  Asp  Glu  Ile  Phe  Val  Glu  Leu  Val  Asn

```

-continued

165					170					175				
Ala	Leu	Gly	Gln	Tyr	Asn	Asp	Asp	Asp	Asp	Asp	Gly	Asp	Asp	
			180				185				190			
Pro	Glu	Glu	Arg	Glu	Glu	Lys	Gln	Lys	Asp	Leu	Glu	Asp	His	Arg
	195						200				205			Asp
Asp	Lys	Glu	Ser	Arg	Pro	Pro	Arg	Lys	Phe	Pro	Ser	Asp	Lys	Ile
	210						215				220			Phe
Glu	Ala	Ile	Ser	Ser	Met	Phe	Pro	Asp	Lys	Gly	Thr	Ala	Glu	Glu
225					230					235				240
Lys	Glu	Lys	Tyr	Lys	Glu	Leu	Thr	Glu	Gln	Gln	Leu	Pro	Gly	Ala
			245						250				255	Leu
Pro	Pro	Glu	Cys	Thr	Pro	Asn	Ile	Asp	Gly	Pro	Asn	Ala	Lys	Ser
		260						265					270	Val
Gln	Arg	Glu	Gln	Ser	Leu	His	Ser	Phe	His	Thr	Leu	Phe	Cys	Arg
	275						280					285		Arg
Cys	Phe	Lys	Tyr	Asp	Cys	Phe	Leu	His	Pro	Phe	His	Ala	Thr	Pro
	290					295					300			Asn
Thr	Tyr	Lys	Arg	Lys	Asn	Thr	Glu	Thr	Ala	Leu	Asp	Asn	Lys	Pro
305					310					315				320
Gly	Pro	Gln	Cys	Tyr	Gln	His	Leu	Glu	Gly	Ala	Lys	Glu	Phe	Ala
			325						330					335
Ala	Leu	Thr	Ala	Glu	Arg	Ile	Lys	Thr	Pro	Pro	Lys	Arg	Pro	Gly
		340						345					350	Gly
Arg	Arg	Arg	Gly	Arg	Leu	Pro	Asn	Asn	Ser	Ser	Arg	Pro	Ser	Thr
		355					360					365		Pro
Thr	Ile	Asn	Val	Leu	Glu	Ser	Lys	Asp	Thr	Asp	Ser	Asp	Arg	Glu
	370					375					380			Ala
Gly	Thr	Glu	Thr	Gly	Gly	Glu	Asn	Asn	Asp	Lys	Glu	Glu	Glu	Lys
385					390					395				400
Lys	Asp	Glu	Thr	Ser	Ser	Ser	Ser	Glu	Ala	Asn	Ser	Arg	Cys	Gln
			405						410					415
Pro	Ile	Lys	Met	Lys	Pro	Asn	Ile	Glu	Pro	Pro	Glu	Asn	Val	Glu
		420						425					430	Trp
Ser	Gly	Ala	Glu	Ala	Ser	Met	Phe	Arg	Val	Leu	Ile	Gly	Thr	Tyr
		435					440					445		Tyr
Asp	Asn	Phe	Cys	Ala	Ile	Ala	Arg	Leu	Ile	Gly	Thr	Lys	Thr	Cys
	450					455					460			Arg
Gln	Val	Tyr	Glu	Phe	Arg	Val	Lys	Glu	Ser	Ser	Ile	Ile	Ala	Pro
465					470					475				480
Pro	Ala	Glu	Asp	Val	Asp	Thr	Pro	Pro	Arg	Lys	Lys	Lys	Arg	Lys
			485						490					495
Arg	Leu	Trp	Ala	Ala	His	Cys	Arg	Lys	Ile	Gln	Leu	Lys	Lys	Asp
		500						505					510	Gly
Ser	Ser	Asn	His	Val	Tyr	Asn	Tyr	Gln	Pro	Cys	Asp	His	Pro	Arg
		515					520					525		Gln
Pro	Cys	Asp	Ser	Ser	Cys	Pro	Cys	Val	Ile	Ala	Gln	Asn	Phe	Cys
	530					535					540			Glu
Lys	Phe	Cys	Gln	Cys	Ser	Ser	Glu	Cys	Gln	Asn	Arg	Phe	Pro	Gly
545					550					555				560
Arg	Cys	Lys	Ala	Gln	Cys	Asn	Thr	Lys	Gln	Cys	Pro	Cys	Tyr	Leu
			565						570					575

-continued

Val	Arg	Glu	Cys	Asp	Pro	Asp	Leu	Cys	Leu	Thr	Cys	Gly	Ala	Ala	Asp
			580					585					590		
His	Trp	Asp	Ser	Lys	Asn	Val	Ser	Cys	Lys	Asn	Cys	Ser	Ile	Gln	Arg
		595					600					605			
Gly	Ser	Lys	Lys	His	Leu	Leu	Leu	Ala	Pro	Ser	Asp	Val	Ala	Gly	Trp
	610				615						620				
Gly	Ile	Phe	Ile	Lys	Asp	Pro	Val	Gln	Lys	Asn	Glu	Phe	Ile	Ser	Glu
625					630					635					640
His	Cys	Gly	Glu	Ile	Ile	Ser	Gln	Asp	Glu	Ala	Asp	Arg	Arg	Gly	Lys
				645					650					655	
Val	Tyr	Asp	Lys	Tyr	Met	Cys	Ser	Phe	Leu	Phe	Asn	Leu	Asn	Asn	Asp
			660					665					670		
Phe	Val	Val	Asp	Ala	Thr	Arg	Lys	Gly	Asn	Lys	Ile	Arg	Phe	Ala	Asn
		675					680					685			
His	Ser	Val	Asn	Pro	Asn	Cys	Tyr	Ala	Lys	Val	Met	Met	Val	Asn	Gly
		690				695					700				
Asp	His	Arg	Ile	Gly	Ile	Phe	Ala	Lys	Arg	Ala	Ile	Gln	Thr	Gly	Glu
705				710					715						720
Glu	Leu	Phe	Phe	Asp	Tyr	Arg	Tyr	Ser	Gln	Ala	Asp	Ala	Leu	Lys	Tyr
				725					730					735	
Val	Gly	Ile	Glu	Arg	Glu	Met	Glu	Ile	Pro						
			740				745								

<210> SEQ ID NO 12

<211> LENGTH: 746

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Y641N mutant of Homo sapiens EZH2

<400> SEQUENCE: 12

Met	Gly	Gln	Thr	Gly	Lys	Lys	Ser	Glu	Lys	Gly	Pro	Val	Cys	Trp	Arg
1				5					10					15	
Lys	Arg	Val	Lys	Ser	Glu	Tyr	Met	Arg	Leu	Arg	Gln	Leu	Lys	Arg	Phe
		20					25						30		
Arg	Arg	Ala	Asp	Glu	Val	Lys	Ser	Met	Phe	Ser	Ser	Asn	Arg	Gln	Lys
		35					40					45			
Ile	Leu	Glu	Arg	Thr	Glu	Ile	Leu	Asn	Gln	Glu	Trp	Lys	Gln	Arg	Arg
	50				55					60					
Ile	Gln	Pro	Val	His	Ile	Leu	Thr	Ser	Val	Ser	Ser	Leu	Arg	Gly	Thr
65				70					75					80	
Arg	Glu	Cys	Ser	Val	Thr	Ser	Asp	Leu	Asp	Phe	Pro	Thr	Gln	Val	Ile
			85					90						95	
Pro	Leu	Lys	Thr	Leu	Asn	Ala	Val	Ala	Ser	Val	Pro	Ile	Met	Tyr	Ser
		100					105					110			
Trp	Ser	Pro	Leu	Gln	Gln	Asn	Phe	Met	Val	Glu	Asp	Glu	Thr	Val	Leu
		115				120					125				
His	Asn	Ile	Pro	Tyr	Met	Gly	Asp	Glu	Val	Leu	Asp	Gln	Asp	Gly	Thr
	130					135					140				
Phe	Ile	Glu	Glu	Leu	Ile	Lys	Asn	Tyr	Asp	Gly	Lys	Val	His	Gly	Asp
145				150					155					160	
Arg	Glu	Cys	Gly	Phe	Ile	Asn	Asp	Glu	Ile	Phe	Val	Glu	Leu	Val	Asn
			165					170						175	
Ala	Leu	Gly	Gln	Tyr	Asn	Asp	Asp	Asp	Asp	Asp	Asp	Asp	Gly	Asp	Asp

-continued

180							185					190				
Pro	Glu	Glu	Arg	Glu	Glu	Lys	Gln	Lys	Asp	Leu	Glu	Asp	His	Arg	Asp	
195							200					205				
Asp	Lys	Glu	Ser	Arg	Pro	Pro	Arg	Lys	Phe	Pro	Ser	Asp	Lys	Ile	Phe	
210							215					220				
Glu	Ala	Ile	Ser	Ser	Met	Phe	Pro	Asp	Lys	Gly	Thr	Ala	Glu	Glu	Leu	
225							230					235				
Lys	Glu	Lys	Tyr	Lys	Glu	Leu	Thr	Glu	Gln	Gln	Leu	Pro	Gly	Ala	Leu	
245							250					255				
Pro	Pro	Glu	Cys	Thr	Pro	Asn	Ile	Asp	Gly	Pro	Asn	Ala	Lys	Ser	Val	
260							265					270				
Gln	Arg	Glu	Gln	Ser	Leu	His	Ser	Phe	His	Thr	Leu	Phe	Cys	Arg	Arg	
275							280					285				
Cys	Phe	Lys	Tyr	Asp	Cys	Phe	Leu	His	Pro	Phe	His	Ala	Thr	Pro	Asn	
290							295					300				
Thr	Tyr	Lys	Arg	Lys	Asn	Thr	Glu	Thr	Ala	Leu	Asp	Asn	Lys	Pro	Cys	
305							310					315				
Gly	Pro	Gln	Cys	Tyr	Gln	His	Leu	Glu	Gly	Ala	Lys	Glu	Phe	Ala	Ala	
325							330					335				
Ala	Leu	Thr	Ala	Glu	Arg	Ile	Lys	Thr	Pro	Pro	Lys	Arg	Pro	Gly	Gly	
340							345					350				
Arg	Arg	Arg	Gly	Arg	Leu	Pro	Asn	Asn	Ser	Ser	Arg	Pro	Ser	Thr	Pro	
355							360					365				
Thr	Ile	Asn	Val	Leu	Glu	Ser	Lys	Asp	Thr	Asp	Ser	Asp	Arg	Glu	Ala	
370							375					380				
Gly	Thr	Glu	Thr	Gly	Gly	Glu	Asn	Asn	Asp	Lys	Glu	Glu	Glu	Glu	Lys	
385							390					395				
Lys	Asp	Glu	Thr	Ser	Ser	Ser	Ser	Glu	Ala	Asn	Ser	Arg	Cys	Gln	Thr	
405							410					415				
Pro	Ile	Lys	Met	Lys	Pro	Asn	Ile	Glu	Pro	Pro	Glu	Asn	Val	Glu	Trp	
420							425					430				
Ser	Gly	Ala	Glu	Ala	Ser	Met	Phe	Arg	Val	Leu	Ile	Gly	Thr	Tyr	Tyr	
435							440					445				
Asp	Asn	Phe	Cys	Ala	Ile	Ala	Arg	Leu	Ile	Gly	Thr	Lys	Thr	Cys	Arg	
450							455					460				
Gln	Val	Tyr	Glu	Phe	Arg	Val	Lys	Glu	Ser	Ser	Ile	Ile	Ala	Pro	Ala	
465							470					475				
Pro	Ala	Glu	Asp	Val	Asp	Thr	Pro	Pro	Arg	Lys	Lys	Lys	Arg	Lys	His	
485							490					495				
Arg	Leu	Trp	Ala	Ala	His	Cys	Arg	Lys	Ile	Gln	Leu	Lys	Lys	Asp	Gly	
500							505					510				
Ser	Ser	Asn	His	Val	Tyr	Asn	Tyr	Gln	Pro	Cys	Asp	His	Pro	Arg	Gln	
515							520					525				
Pro	Cys	Asp	Ser	Ser	Cys	Pro	Cys	Val	Ile	Ala	Gln	Asn	Phe	Cys	Glu	
530							535					540				
Lys	Phe	Cys	Gln	Cys	Ser	Ser	Glu	Cys	Gln	Asn	Arg	Phe	Pro	Gly	Cys	
545							550					555				
Arg	Cys	Lys	Ala	Gln	Cys	Asn	Thr	Lys	Gln	Cys	Pro	Cys	Tyr	Leu	Ala	
565							570					575				
Val	Arg	Glu	Cys	Asp	Pro	Asp	Leu	Cys	Leu	Thr	Cys	Gly	Ala	Ala	Asp	
580							585					590				

-continued

His	Trp	Asp	Ser	Lys	Asn	Val	Ser	Cys	Lys	Asn	Cys	Ser	Ile	Gln	Arg
		595					600					605			
Gly	Ser	Lys	Lys	His	Leu	Leu	Leu	Ala	Pro	Ser	Asp	Val	Ala	Gly	Trp
	610					615					620				
Gly	Ile	Phe	Ile	Lys	Asp	Pro	Val	Gln	Lys	Asn	Glu	Phe	Ile	Ser	Glu
625					630					635					640
Asn	Cys	Gly	Glu	Ile	Ile	Ser	Gln	Asp	Glu	Ala	Asp	Arg	Arg	Gly	Lys
				645					650					655	
Val	Tyr	Asp	Lys	Tyr	Met	Cys	Ser	Phe	Leu	Phe	Asn	Leu	Asn	Asn	Asp
			660					665					670		
Phe	Val	Val	Asp	Ala	Thr	Arg	Lys	Gly	Asn	Lys	Ile	Arg	Phe	Ala	Asn
		675					680					685			
His	Ser	Val	Asn	Pro	Asn	Cys	Tyr	Ala	Lys	Val	Met	Met	Val	Asn	Gly
	690					695					700				
Asp	His	Arg	Ile	Gly	Ile	Phe	Ala	Lys	Arg	Ala	Ile	Gln	Thr	Gly	Glu
705					710					715					720
Glu	Leu	Phe	Phe	Asp	Tyr	Arg	Tyr	Ser	Gln	Ala	Asp	Ala	Leu	Lys	Tyr
				725					730					735	
Val	Gly	Ile	Glu	Arg	Glu	Met	Glu	Ile	Pro						
			740				745								

<210> SEQ ID NO 13

<211> LENGTH: 746

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Y641S mutant of Homo sapiens EZH2

<400> SEQUENCE: 13

Met	Gly	Gln	Thr	Gly	Lys	Lys	Ser	Glu	Lys	Gly	Pro	Val	Cys	Trp	Arg
1				5					10					15	
Lys	Arg	Val	Lys	Ser	Glu	Tyr	Met	Arg	Leu	Arg	Gln	Leu	Lys	Arg	Phe
		20					25						30		
Arg	Arg	Ala	Asp	Glu	Val	Lys	Ser	Met	Phe	Ser	Ser	Asn	Arg	Gln	Lys
		35					40					45			
Ile	Leu	Glu	Arg	Thr	Glu	Ile	Leu	Asn	Gln	Glu	Trp	Lys	Gln	Arg	Arg
	50					55				60					
Ile	Gln	Pro	Val	His	Ile	Leu	Thr	Ser	Val	Ser	Ser	Leu	Arg	Gly	Thr
65					70					75				80	
Arg	Glu	Cys	Ser	Val	Thr	Ser	Asp	Leu	Asp	Phe	Pro	Thr	Gln	Val	Ile
			85						90					95	
Pro	Leu	Lys	Thr	Leu	Asn	Ala	Val	Ala	Ser	Val	Pro	Ile	Met	Tyr	Ser
		100					105						110		
Trp	Ser	Pro	Leu	Gln	Gln	Asn	Phe	Met	Val	Glu	Asp	Glu	Thr	Val	Leu
		115				120						125			
His	Asn	Ile	Pro	Tyr	Met	Gly	Asp	Glu	Val	Leu	Asp	Gln	Asp	Gly	Thr
	130					135					140				
Phe	Ile	Glu	Glu	Leu	Ile	Lys	Asn	Tyr	Asp	Gly	Lys	Val	His	Gly	Asp
145					150					155					160
Arg	Glu	Cys	Gly	Phe	Ile	Asn	Asp	Glu	Ile	Phe	Val	Glu	Leu	Val	Asn
			165						170					175	
Ala	Leu	Gly	Gln	Tyr	Asn	Asp	Asp	Asp	Asp	Asp	Asp	Gly	Asp	Asp	
		180					185					190			
Pro	Glu	Glu	Arg	Glu	Glu	Lys	Gln	Lys	Asp	Leu	Glu	Asp	His	Arg	Asp

-continued

195						200						205					
Asp	Lys	Glu	Ser	Arg	Pro	Pro	Arg	Lys	Phe	Pro	Ser	Asp	Lys	Ile	Phe		
210						215					220						
Glu	Ala	Ile	Ser	Ser	Met	Phe	Pro	Asp	Lys	Gly	Thr	Ala	Glu	Glu	Leu		
225					230					235					240		
Lys	Glu	Lys	Tyr	Lys	Glu	Leu	Thr	Glu	Gln	Gln	Leu	Pro	Gly	Ala	Leu		
				245					250					255			
Pro	Pro	Glu	Cys	Thr	Pro	Asn	Ile	Asp	Gly	Pro	Asn	Ala	Lys	Ser	Val		
			260					265					270				
Gln	Arg	Glu	Gln	Ser	Leu	His	Ser	Phe	His	Thr	Leu	Phe	Cys	Arg	Arg		
		275					280					285					
Cys	Phe	Lys	Tyr	Asp	Cys	Phe	Leu	His	Pro	Phe	His	Ala	Thr	Pro	Asn		
290						295					300						
Thr	Tyr	Lys	Arg	Lys	Asn	Thr	Glu	Thr	Ala	Leu	Asp	Asn	Lys	Pro	Cys		
305					310					315					320		
Gly	Pro	Gln	Cys	Tyr	Gln	His	Leu	Glu	Gly	Ala	Lys	Glu	Phe	Ala	Ala		
				325					330					335			
Ala	Leu	Thr	Ala	Glu	Arg	Ile	Lys	Thr	Pro	Pro	Lys	Arg	Pro	Gly	Gly		
			340					345					350				
Arg	Arg	Arg	Gly	Arg	Leu	Pro	Asn	Asn	Ser	Ser	Arg	Pro	Ser	Thr	Pro		
		355					360					365					
Thr	Ile	Asn	Val	Leu	Glu	Ser	Lys	Asp	Thr	Asp	Ser	Asp	Arg	Glu	Ala		
	370					375					380						
Gly	Thr	Glu	Thr	Gly	Gly	Glu	Asn	Asn	Asp	Lys	Glu	Glu	Glu	Glu	Lys		
385					390					395					400		
Lys	Asp	Glu	Thr	Ser	Ser	Ser	Ser	Glu	Ala	Asn	Ser	Arg	Cys	Gln	Thr		
				405					410					415			
Pro	Ile	Lys	Met	Lys	Pro	Asn	Ile	Glu	Pro	Pro	Glu	Asn	Val	Glu	Trp		
			420					425					430				
Ser	Gly	Ala	Glu	Ala	Ser	Met	Phe	Arg	Val	Leu	Ile	Gly	Thr	Tyr	Tyr		
		435					440					445					
Asp	Asn	Phe	Cys	Ala	Ile	Ala	Arg	Leu	Ile	Gly	Thr	Lys	Thr	Cys	Arg		
	450					455					460						
Gln	Val	Tyr	Glu	Phe	Arg	Val	Lys	Glu	Ser	Ser	Ile	Ile	Ala	Pro	Ala		
465					470					475					480		
Pro	Ala	Glu	Asp	Val	Asp	Thr	Pro	Pro	Arg	Lys	Lys	Lys	Arg	Lys	His		
				485					490					495			
Arg	Leu	Trp	Ala	Ala	His	Cys	Arg	Lys	Ile	Gln	Leu	Lys	Lys	Asp	Gly		
			500					505						510			
Ser	Ser	Asn	His	Val	Tyr	Asn	Tyr	Gln	Pro	Cys	Asp	His	Pro	Arg	Gln		
		515					520					525					
Pro	Cys	Asp	Ser	Ser	Cys	Pro	Cys	Val	Ile	Ala	Gln	Asn	Phe	Cys	Glu		
	530					535					540						
Lys	Phe	Cys	Gln	Cys	Ser	Ser	Glu	Cys	Gln	Asn	Arg	Phe	Pro	Gly	Cys		
545					550					555					560		
Arg	Cys	Lys	Ala	Gln	Cys	Asn	Thr	Lys	Gln	Cys	Pro	Cys	Tyr	Leu	Ala		
				565					570					575			
Val	Arg	Glu	Cys	Asp	Pro	Asp	Leu	Cys	Leu	Thr	Cys	Gly	Ala	Ala	Asp		
			580				585						590				
His	Trp	Asp	Ser	Lys	Asn	Val	Ser	Cys	Lys	Asn	Cys	Ser	Ile	Gln	Arg		
		595					600						605				

-continued

Gly	Ser	Lys	Lys	His	Leu	Leu	Leu	Ala	Pro	Ser	Asp	Val	Ala	Gly	Trp
610						615					620				
Gly	Ile	Phe	Ile	Lys	Asp	Pro	Val	Gln	Lys	Asn	Glu	Phe	Ile	Ser	Glu
625					630					635					640
Ser	Cys	Gly	Glu	Ile	Ile	Ser	Gln	Asp	Glu	Ala	Asp	Arg	Arg	Gly	Lys
				645					650					655	
Val	Tyr	Asp	Lys	Tyr	Met	Cys	Ser	Phe	Leu	Phe	Asn	Leu	Asn	Asn	Asp
			660					665					670		
Phe	Val	Val	Asp	Ala	Thr	Arg	Lys	Gly	Asn	Lys	Ile	Arg	Phe	Ala	Asn
		675						680					685		
His	Ser	Val	Asn	Pro	Asn	Cys	Tyr	Ala	Lys	Val	Met	Met	Val	Asn	Gly
		690				695					700				
Asp	His	Arg	Ile	Gly	Ile	Phe	Ala	Lys	Arg	Ala	Ile	Gln	Thr	Gly	Glu
705					710					715					720
Glu	Leu	Phe	Phe	Asp	Tyr	Arg	Tyr	Ser	Gln	Ala	Asp	Ala	Leu	Lys	Tyr
				725					730					735	
Val	Gly	Ile	Glu	Arg	Glu	Met	Glu	Ile	Pro						
			740				745								

<210> SEQ ID NO 14

<211> LENGTH: 746

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Y641C mutant of Homo sapiens EZH2

<400> SEQUENCE: 14

Met	Gly	Gln	Thr	Gly	Lys	Lys	Ser	Glu	Lys	Gly	Pro	Val	Cys	Trp	Arg
1				5				10					15		
Lys	Arg	Val	Lys	Ser	Glu	Tyr	Met	Arg	Leu	Arg	Gln	Leu	Lys	Arg	Phe
		20					25						30		
Arg	Arg	Ala	Asp	Glu	Val	Lys	Ser	Met	Phe	Ser	Ser	Asn	Arg	Gln	Lys
		35					40					45			
Ile	Leu	Glu	Arg	Thr	Glu	Ile	Leu	Asn	Gln	Glu	Trp	Lys	Gln	Arg	Arg
	50					55					60				
Ile	Gln	Pro	Val	His	Ile	Leu	Thr	Ser	Val	Ser	Ser	Leu	Arg	Gly	Thr
65				70						75				80	
Arg	Glu	Cys	Ser	Val	Thr	Ser	Asp	Leu	Asp	Phe	Pro	Thr	Gln	Val	Ile
			85						90					95	
Pro	Leu	Lys	Thr	Leu	Asn	Ala	Val	Ala	Ser	Val	Pro	Ile	Met	Tyr	Ser
		100					105					110			
Trp	Ser	Pro	Leu	Gln	Gln	Asn	Phe	Met	Val	Glu	Asp	Glu	Thr	Val	Leu
		115				120						125			
His	Asn	Ile	Pro	Tyr	Met	Gly	Asp	Glu	Val	Leu	Asp	Gln	Asp	Gly	Thr
	130					135					140				
Phe	Ile	Glu	Glu	Leu	Ile	Lys	Asn	Tyr	Asp	Gly	Lys	Val	His	Gly	Asp
145				150						155					160
Arg	Glu	Cys	Gly	Phe	Ile	Asn	Asp	Glu	Ile	Phe	Val	Glu	Leu	Val	Asn
			165						170					175	
Ala	Leu	Gly	Gln	Tyr	Asn	Asp	Asp	Asp	Asp	Asp	Asp	Gly	Asp	Asp	Asp
		180					185					190			
Pro	Glu	Glu	Arg	Glu	Glu	Lys	Gln	Lys	Asp	Leu	Glu	Asp	His	Arg	Asp
	195					200						205			
Asp	Lys	Glu	Ser	Arg	Pro	Pro	Arg	Lys	Phe	Pro	Ser	Asp	Lys	Ile	Phe

-continued

210	215	220
Glu Ala Ile Ser Ser Met Phe Pro Asp Lys Gly Thr Ala Glu Glu Leu 225 230 235 240		
Lys Glu Lys Tyr Lys Glu Leu Thr Glu Gln Gln Leu Pro Gly Ala Leu 245 250 255		
Pro Pro Glu Cys Thr Pro Asn Ile Asp Gly Pro Asn Ala Lys Ser Val 260 265 270		
Gln Arg Glu Gln Ser Leu His Ser Phe His Thr Leu Phe Cys Arg Arg 275 280 285		
Cys Phe Lys Tyr Asp Cys Phe Leu His Pro Phe His Ala Thr Pro Asn 290 295 300		
Thr Tyr Lys Arg Lys Asn Thr Glu Thr Ala Leu Asp Asn Lys Pro Cys 305 310 315 320		
Gly Pro Gln Cys Tyr Gln His Leu Glu Gly Ala Lys Glu Phe Ala Ala 325 330 335		
Ala Leu Thr Ala Glu Arg Ile Lys Thr Pro Pro Lys Arg Pro Gly Gly 340 345 350		
Arg Arg Arg Gly Arg Leu Pro Asn Asn Ser Ser Arg Pro Ser Thr Pro 355 360 365		
Thr Ile Asn Val Leu Glu Ser Lys Asp Thr Asp Ser Asp Arg Glu Ala 370 375 380		
Gly Thr Glu Thr Gly Gly Glu Asn Asn Asp Lys Glu Glu Glu Glu Lys 385 390 395 400		
Lys Asp Glu Thr Ser Ser Ser Ser Glu Ala Asn Ser Arg Cys Gln Thr 405 410 415		
Pro Ile Lys Met Lys Pro Asn Ile Glu Pro Pro Glu Asn Val Glu Trp 420 425 430		
Ser Gly Ala Glu Ala Ser Met Phe Arg Val Leu Ile Gly Thr Tyr Tyr 435 440 445		
Asp Asn Phe Cys Ala Ile Ala Arg Leu Ile Gly Thr Lys Thr Cys Arg 450 455 460		
Gln Val Tyr Glu Phe Arg Val Lys Glu Ser Ser Ile Ile Ala Pro Ala 465 470 475 480		
Pro Ala Glu Asp Val Asp Thr Pro Pro Arg Lys Lys Lys Arg Lys His 485 490 495		
Arg Leu Trp Ala Ala His Cys Arg Lys Ile Gln Leu Lys Lys Asp Gly 500 505 510		
Ser Ser Asn His Val Tyr Asn Tyr Gln Pro Cys Asp His Pro Arg Gln 515 520 525		
Pro Cys Asp Ser Ser Cys Pro Cys Val Ile Ala Gln Asn Phe Cys Glu 530 535 540		
Lys Phe Cys Gln Cys Ser Ser Glu Cys Gln Asn Arg Phe Pro Gly Cys 545 550 555 560		
Arg Cys Lys Ala Gln Cys Asn Thr Lys Gln Cys Pro Cys Tyr Leu Ala 565 570 575		
Val Arg Glu Cys Asp Pro Asp Leu Cys Leu Thr Cys Gly Ala Ala Asp 580 585 590		
His Trp Asp Ser Lys Asn Val Ser Cys Lys Asn Cys Ser Ile Gln Arg 595 600 605		
Gly Ser Lys Lys His Leu Leu Leu Ala Pro Ser Asp Val Ala Gly Trp 610 615 620		

-continued

Gly	Ile	Phe	Ile	Lys	Asp	Pro	Val	Gln	Lys	Asn	Glu	Phe	Ile	Ser	Glu
625					630					635					640
Cys	Cys	Gly	Glu	Ile	Ile	Ser	Gln	Asp	Glu	Ala	Asp	Arg	Arg	Gly	Lys
				645					650					655	
Val	Tyr	Asp	Lys	Tyr	Met	Cys	Ser	Phe	Leu	Phe	Asn	Leu	Asn	Asn	Asp
			660					665					670		
Phe	Val	Val	Asp	Ala	Thr	Arg	Lys	Gly	Asn	Lys	Ile	Arg	Phe	Ala	Asn
		675					680					685			
His	Ser	Val	Asn	Pro	Asn	Cys	Tyr	Ala	Lys	Val	Met	Met	Val	Asn	Gly
	690					695					700				
Asp	His	Arg	Ile	Gly	Ile	Phe	Ala	Lys	Arg	Ala	Ile	Gln	Thr	Gly	Glu
705				710						715					720
Glu	Leu	Phe	Phe	Asp	Tyr	Arg	Tyr	Ser	Gln	Ala	Asp	Ala	Leu	Lys	Tyr
				725					730					735	
Val	Gly	Ile	Glu	Arg	Glu	Met	Glu	Ile	Pro						
			740				745								

<210> SEQ ID NO 15
 <211> LENGTH: 746
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: A677 mutant of Homo sapiens EZH2, wherein X is preferably a glycine (G)
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (677) .. (677)
 <223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid, preferably a glycine (G)

<400> SEQUENCE: 15

Met	Gly	Gln	Thr	Gly	Lys	Lys	Ser	Glu	Lys	Gly	Pro	Val	Cys	Trp	Arg
1				5					10					15	
Lys	Arg	Val	Lys	Ser	Glu	Tyr	Met	Arg	Leu	Arg	Gln	Leu	Lys	Arg	Phe
		20						25					30		
Arg	Arg	Ala	Asp	Glu	Val	Lys	Ser	Met	Phe	Ser	Ser	Asn	Arg	Gln	Lys
		35					40					45			
Ile	Leu	Glu	Arg	Thr	Glu	Ile	Leu	Asn	Gln	Glu	Trp	Lys	Gln	Arg	Arg
	50					55					60				
Ile	Gln	Pro	Val	His	Ile	Leu	Thr	Ser	Val	Ser	Ser	Leu	Arg	Gly	Thr
65				70					75					80	
Arg	Glu	Cys	Ser	Val	Thr	Ser	Asp	Leu	Asp	Phe	Pro	Thr	Gln	Val	Ile
			85					90						95	
Pro	Leu	Lys	Thr	Leu	Asn	Ala	Val	Ala	Ser	Val	Pro	Ile	Met	Tyr	Ser
		100					105						110		
Trp	Ser	Pro	Leu	Gln	Gln	Asn	Phe	Met	Val	Glu	Asp	Glu	Thr	Val	Leu
		115				120						125			
His	Asn	Ile	Pro	Tyr	Met	Gly	Asp	Glu	Val	Leu	Asp	Gln	Asp	Gly	Thr
	130					135					140				
Phe	Ile	Glu	Glu	Leu	Ile	Lys	Asn	Tyr	Asp	Gly	Lys	Val	His	Gly	Asp
145				150						155					160
Arg	Glu	Cys	Gly	Phe	Ile	Asn	Asp	Glu	Ile	Phe	Val	Glu	Leu	Val	Asn
			165					170						175	
Ala	Leu	Gly	Gln	Tyr	Asn	Asp	Asp	Asp	Asp	Asp	Asp	Gly	Asp	Asp	
		180				185						190			
Pro	Glu	Glu	Arg	Glu	Glu	Lys	Gln	Lys	Asp	Leu	Glu	Asp	His	Arg	Asp

-continued

195						200						205					
Asp	Lys	Glu	Ser	Arg	Pro	Pro	Arg	Lys	Phe	Pro	Ser	Asp	Lys	Ile	Phe		
210						215					220						
Glu	Ala	Ile	Ser	Ser	Met	Phe	Pro	Asp	Lys	Gly	Thr	Ala	Glu	Glu	Leu		
225					230					235					240		
Lys	Glu	Lys	Tyr	Lys	Glu	Leu	Thr	Glu	Gln	Gln	Leu	Pro	Gly	Ala	Leu		
				245					250					255			
Pro	Pro	Glu	Cys	Thr	Pro	Asn	Ile	Asp	Gly	Pro	Asn	Ala	Lys	Ser	Val		
			260					265					270				
Gln	Arg	Glu	Gln	Ser	Leu	His	Ser	Phe	His	Thr	Leu	Phe	Cys	Arg	Arg		
		275					280					285					
Cys	Phe	Lys	Tyr	Asp	Cys	Phe	Leu	His	Pro	Phe	His	Ala	Thr	Pro	Asn		
290						295					300						
Thr	Tyr	Lys	Arg	Lys	Asn	Thr	Glu	Thr	Ala	Leu	Asp	Asn	Lys	Pro	Cys		
305					310					315					320		
Gly	Pro	Gln	Cys	Tyr	Gln	His	Leu	Glu	Gly	Ala	Lys	Glu	Phe	Ala	Ala		
				325					330					335			
Ala	Leu	Thr	Ala	Glu	Arg	Ile	Lys	Thr	Pro	Pro	Lys	Arg	Pro	Gly	Gly		
			340					345					350				
Arg	Arg	Arg	Gly	Arg	Leu	Pro	Asn	Asn	Ser	Ser	Arg	Pro	Ser	Thr	Pro		
		355					360					365					
Thr	Ile	Asn	Val	Leu	Glu	Ser	Lys	Asp	Thr	Asp	Ser	Asp	Arg	Glu	Ala		
	370					375					380						
Gly	Thr	Glu	Thr	Gly	Gly	Glu	Asn	Asn	Asp	Lys	Glu	Glu	Glu	Glu	Lys		
385					390					395					400		
Lys	Asp	Glu	Thr	Ser	Ser	Ser	Ser	Glu	Ala	Asn	Ser	Arg	Cys	Gln	Thr		
				405					410					415			
Pro	Ile	Lys	Met	Lys	Pro	Asn	Ile	Glu	Pro	Pro	Glu	Asn	Val	Glu	Trp		
			420					425					430				
Ser	Gly	Ala	Glu	Ala	Ser	Met	Phe	Arg	Val	Leu	Ile	Gly	Thr	Tyr	Tyr		
		435					440					445					
Asp	Asn	Phe	Cys	Ala	Ile	Ala	Arg	Leu	Ile	Gly	Thr	Lys	Thr	Cys	Arg		
	450					455					460						
Gln	Val	Tyr	Glu	Phe	Arg	Val	Lys	Glu	Ser	Ser	Ile	Ile	Ala	Pro	Ala		
465					470					475					480		
Pro	Ala	Glu	Asp	Val	Asp	Thr	Pro	Pro	Arg	Lys	Lys	Lys	Arg	Lys	His		
				485					490					495			
Arg	Leu	Trp	Ala	Ala	His	Cys	Arg	Lys	Ile	Gln	Leu	Lys	Lys	Asp	Gly		
			500					505						510			
Ser	Ser	Asn	His	Val	Tyr	Asn	Tyr	Gln	Pro	Cys	Asp	His	Pro	Arg	Gln		
		515					520					525					
Pro	Cys	Asp	Ser	Ser	Cys	Pro	Cys	Val	Ile	Ala	Gln	Asn	Phe	Cys	Glu		
	530					535					540						
Lys	Phe	Cys	Gln	Cys	Ser	Ser	Glu	Cys	Gln	Asn	Arg	Phe	Pro	Gly	Cys		
545					550					555					560		
Arg	Cys	Lys	Ala	Gln	Cys	Asn	Thr	Lys	Gln	Cys	Pro	Cys	Tyr	Leu	Ala		
				565					570					575			
Val	Arg	Glu	Cys	Asp	Pro	Asp	Leu	Cys	Leu	Thr	Cys	Gly	Ala	Ala	Asp		
			580				585						590				
His	Trp	Asp	Ser	Lys	Asn	Val	Ser	Cys	Lys	Asn	Cys	Ser	Ile	Gln	Arg		
		595					600						605				

-continued

Gly	Ser	Lys	Lys	His	Leu	Leu	Leu	Ala	Pro	Ser	Asp	Val	Ala	Gly	Trp
	610					615					620				
Gly	Ile	Phe	Ile	Lys	Asp	Pro	Val	Gln	Lys	Asn	Glu	Phe	Ile	Ser	Glu
625					630					635					640
Tyr	Cys	Gly	Glu	Ile	Ile	Ser	Gln	Asp	Glu	Ala	Asp	Arg	Arg	Gly	Lys
				645					650					655	
Val	Tyr	Asp	Lys	Tyr	Met	Cys	Ser	Phe	Leu	Phe	Asn	Leu	Asn	Asn	Asp
			660					665					670		
Phe	Val	Val	Asp	Xaa	Thr	Arg	Lys	Gly	Asn	Lys	Ile	Arg	Phe	Ala	Asn
		675					680					685			
His	Ser	Val	Asn	Pro	Asn	Cys	Tyr	Ala	Lys	Val	Met	Met	Val	Asn	Gly
	690					695					700				
Asp	His	Arg	Ile	Gly	Ile	Phe	Ala	Lys	Arg	Ala	Ile	Gln	Thr	Gly	Glu
705				710						715					720
Glu	Leu	Phe	Phe	Asp	Tyr	Arg	Tyr	Ser	Gln	Ala	Asp	Ala	Leu	Lys	Tyr
				725					730					735	
Val	Gly	Ile	Glu	Arg	Glu	Met	Glu	Ile	Pro						
			740				745								

<210> SEQ ID NO 16

<211> LENGTH: 746

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: A687 mutant of Homo sapiens EZH2, wherein X is preferably a valine (V)

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (687) .. (687)

<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid, preferably a valine (V)

<400> SEQUENCE: 16

Met	Gly	Gln	Thr	Gly	Lys	Lys	Ser	Glu	Lys	Gly	Pro	Val	Cys	Trp	Arg
1				5					10					15	
Lys	Arg	Val	Lys	Ser	Glu	Tyr	Met	Arg	Leu	Arg	Gln	Leu	Lys	Arg	Phe
		20						25					30		
Arg	Arg	Ala	Asp	Glu	Val	Lys	Ser	Met	Phe	Ser	Ser	Asn	Arg	Gln	Lys
		35					40					45			
Ile	Leu	Glu	Arg	Thr	Glu	Ile	Leu	Asn	Gln	Glu	Trp	Lys	Gln	Arg	Arg
	50					55					60				
Ile	Gln	Pro	Val	His	Ile	Leu	Thr	Ser	Val	Ser	Ser	Leu	Arg	Gly	Thr
65				70						75				80	
Arg	Glu	Cys	Ser	Val	Thr	Ser	Asp	Leu	Asp	Phe	Pro	Thr	Gln	Val	Ile
			85						90					95	
Pro	Leu	Lys	Thr	Leu	Asn	Ala	Val	Ala	Ser	Val	Pro	Ile	Met	Tyr	Ser
		100					105						110		
Trp	Ser	Pro	Leu	Gln	Gln	Asn	Phe	Met	Val	Glu	Asp	Glu	Thr	Val	Leu
		115				120						125			
His	Asn	Ile	Pro	Tyr	Met	Gly	Asp	Glu	Val	Leu	Asp	Gln	Asp	Gly	Thr
	130					135					140				
Phe	Ile	Glu	Glu	Leu	Ile	Lys	Asn	Tyr	Asp	Gly	Lys	Val	His	Gly	Asp
145				150						155				160	
Arg	Glu	Cys	Gly	Phe	Ile	Asn	Asp	Glu	Ile	Phe	Val	Glu	Leu	Val	Asn
			165					170						175	
Ala	Leu	Gly	Gln	Tyr	Asn	Asp	Asp	Asp	Asp	Asp	Asp	Asp	Gly	Asp	Asp

-continued

180							185					190				
Pro	Glu	Glu	Arg	Glu	Glu	Lys	Gln	Lys	Asp	Leu	Glu	Asp	His	Arg	Asp	
		195					200					205				
Asp	Lys	Glu	Ser	Arg	Pro	Pro	Arg	Lys	Phe	Pro	Ser	Asp	Lys	Ile	Phe	
	210					215					220					
Glu	Ala	Ile	Ser	Ser	Met	Phe	Pro	Asp	Lys	Gly	Thr	Ala	Glu	Glu	Leu	
225					230					235					240	
Lys	Glu	Lys	Tyr	Lys	Glu	Leu	Thr	Glu	Gln	Gln	Leu	Pro	Gly	Ala	Leu	
				245					250					255		
Pro	Pro	Glu	Cys	Thr	Pro	Asn	Ile	Asp	Gly	Pro	Asn	Ala	Lys	Ser	Val	
		260						265					270			
Gln	Arg	Glu	Gln	Ser	Leu	His	Ser	Phe	His	Thr	Leu	Phe	Cys	Arg	Arg	
		275					280					285				
Cys	Phe	Lys	Tyr	Asp	Cys	Phe	Leu	His	Pro	Phe	His	Ala	Thr	Pro	Asn	
	290					295					300					
Thr	Tyr	Lys	Arg	Lys	Asn	Thr	Glu	Thr	Ala	Leu	Asp	Asn	Lys	Pro	Cys	
305					310					315					320	
Gly	Pro	Gln	Cys	Tyr	Gln	His	Leu	Glu	Gly	Ala	Lys	Glu	Phe	Ala	Ala	
				325					330					335		
Ala	Leu	Thr	Ala	Glu	Arg	Ile	Lys	Thr	Pro	Pro	Lys	Arg	Pro	Gly	Gly	
		340						345					350			
Arg	Arg	Arg	Gly	Arg	Leu	Pro	Asn	Asn	Ser	Ser	Arg	Pro	Ser	Thr	Pro	
		355					360					365				
Thr	Ile	Asn	Val	Leu	Glu	Ser	Lys	Asp	Thr	Asp	Ser	Asp	Arg	Glu	Ala	
	370					375					380					
Gly	Thr	Glu	Thr	Gly	Gly	Glu	Asn	Asn	Asp	Lys	Glu	Glu	Glu	Glu	Lys	
385					390					395					400	
Lys	Asp	Glu	Thr	Ser	Ser	Ser	Ser	Glu	Ala	Asn	Ser	Arg	Cys	Gln	Thr	
				405					410					415		
Pro	Ile	Lys	Met	Lys	Pro	Asn	Ile	Glu	Pro	Pro	Glu	Asn	Val	Glu	Trp	
		420						425					430			
Ser	Gly	Ala	Glu	Ala	Ser	Met	Phe	Arg	Val	Leu	Ile	Gly	Thr	Tyr	Tyr	
	435						440					445				
Asp	Asn	Phe	Cys	Ala	Ile	Ala	Arg	Leu	Ile	Gly	Thr	Lys	Thr	Cys	Arg	
	450					455					460					
Gln	Val	Tyr	Glu	Phe	Arg	Val	Lys	Glu	Ser	Ser	Ile	Ile	Ala	Pro	Ala	
465					470					475					480	
Pro	Ala	Glu	Asp	Val	Asp	Thr	Pro	Pro	Arg	Lys	Lys	Lys	Arg	Lys	His	
				485					490					495		
Arg	Leu	Trp	Ala	Ala	His	Cys	Arg	Lys	Ile	Gln	Leu	Lys	Lys	Asp	Gly	
		500						505					510			
Ser	Ser	Asn	His	Val	Tyr	Asn	Tyr	Gln	Pro	Cys	Asp	His	Pro	Arg	Gln	
		515					520					525				
Pro	Cys	Asp	Ser	Ser	Cys	Pro	Cys	Val	Ile	Ala	Gln	Asn	Phe	Cys	Glu	
	530					535					540					
Lys	Phe	Cys	Gln	Cys	Ser	Ser	Glu	Cys	Gln	Asn	Arg	Phe	Pro	Gly	Cys	
545					550					555					560	
Arg	Cys	Lys	Ala	Gln	Cys	Asn	Thr	Lys	Gln	Cys	Pro	Cys	Tyr	Leu	Ala	
			565						570					575		
Val	Arg	Glu	Cys	Asp	Pro	Asp	Leu	Cys	Leu	Thr	Cys	Gly	Ala	Ala	Asp	
			580				585						590			

-continued

```

His Trp Asp Ser Lys Asn Val Ser Cys Lys Asn Cys Ser Ile Gln Arg
   595                               600                               605

Gly Ser Lys Lys His Leu Leu Leu Ala Pro Ser Asp Val Ala Gly Trp
   610                               615                               620

Gly Ile Phe Ile Lys Asp Pro Val Gln Lys Asn Glu Phe Ile Ser Glu
  625                               630                               635                               640

Tyr Cys Gly Glu Ile Ile Ser Gln Asp Glu Ala Asp Arg Arg Gly Lys
   645                               650                               655

Val Tyr Asp Lys Tyr Met Cys Ser Phe Leu Phe Asn Leu Asn Asn Asp
   660                               665                               670

Phe Val Val Asp Ala Thr Arg Lys Gly Asn Lys Ile Arg Phe Xaa Asn
   675                               680                               685

His Ser Val Asn Pro Asn Cys Tyr Ala Lys Val Met Met Val Asn Gly
   690                               695                               700

Asp His Arg Ile Gly Ile Phe Ala Lys Arg Ala Ile Gln Thr Gly Glu
  705                               710                               715                               720

Glu Leu Phe Phe Asp Tyr Arg Tyr Ser Gln Ala Asp Ala Leu Lys Tyr
   725                               730                               735

Val Gly Ile Glu Arg Glu Met Glu Ile Pro
   740                               745

```

```

<210> SEQ ID NO 17
<211> LENGTH: 746
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Homo sapiens EZH2 with a mutation at residue
        685, wherein X is preferably a cysteine (C) or a histidine (H)
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (685)..(685)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid,
        preferably a cysteine (C) or a histidine (H)

```

```

<400> SEQUENCE: 17

```

```

Met Gly Gln Thr Gly Lys Lys Ser Glu Lys Gly Pro Val Cys Trp Arg
  1                               5                               10                               15

Lys Arg Val Lys Ser Glu Tyr Met Arg Leu Arg Gln Leu Lys Arg Phe
  20                               25                               30

Arg Arg Ala Asp Glu Val Lys Ser Met Phe Ser Ser Asn Arg Gln Lys
  35                               40                               45

Ile Leu Glu Arg Thr Glu Ile Leu Asn Gln Glu Trp Lys Gln Arg Arg
  50                               55                               60

Ile Gln Pro Val His Ile Leu Thr Ser Val Ser Ser Leu Arg Gly Thr
  65                               70                               75                               80

Arg Glu Cys Ser Val Thr Ser Asp Leu Asp Phe Pro Thr Gln Val Ile
   85                               90                               95

Pro Leu Lys Thr Leu Asn Ala Val Ala Ser Val Pro Ile Met Tyr Ser
  100                               105                               110

Trp Ser Pro Leu Gln Gln Asn Phe Met Val Glu Asp Glu Thr Val Leu
  115                               120                               125

His Asn Ile Pro Tyr Met Gly Asp Glu Val Leu Asp Gln Asp Gly Thr
  130                               135                               140

Phe Ile Glu Glu Leu Ile Lys Asn Tyr Asp Gly Lys Val His Gly Asp
  145                               150                               155                               160

Arg Glu Cys Gly Phe Ile Asn Asp Glu Ile Phe Val Glu Leu Val Asn

```

-continued

165					170					175				
Ala	Leu	Gly	Gln	Tyr	Asn	Asp	Asp	Asp	Asp	Asp	Gly	Asp	Asp	
			180				185				190			
Pro	Glu	Glu	Arg	Glu	Glu	Lys	Gln	Lys	Asp	Leu	Glu	Asp	His	Arg
	195						200				205			
Asp	Lys	Glu	Ser	Arg	Pro	Pro	Arg	Lys	Phe	Pro	Ser	Asp	Lys	Ile
	210						215				220			
Glu	Ala	Ile	Ser	Ser	Met	Phe	Pro	Asp	Lys	Gly	Thr	Ala	Glu	Glu
225					230					235				240
Lys	Glu	Lys	Tyr	Lys	Glu	Leu	Thr	Glu	Gln	Gln	Leu	Pro	Gly	Ala
			245						250					255
Pro	Pro	Glu	Cys	Thr	Pro	Asn	Ile	Asp	Gly	Pro	Asn	Ala	Lys	Ser
		260						265					270	Val
Gln	Arg	Glu	Gln	Ser	Leu	His	Ser	Phe	His	Thr	Leu	Phe	Cys	Arg
	275						280					285		Arg
Cys	Phe	Lys	Tyr	Asp	Cys	Phe	Leu	His	Pro	Phe	His	Ala	Thr	Pro
	290						295				300			Asn
Thr	Tyr	Lys	Arg	Lys	Asn	Thr	Glu	Thr	Ala	Leu	Asp	Asn	Lys	Pro
305					310					315				Cys
Gly	Pro	Gln	Cys	Tyr	Gln	His	Leu	Glu	Gly	Ala	Lys	Glu	Phe	Ala
			325						330					335
Ala	Leu	Thr	Ala	Glu	Arg	Ile	Lys	Thr	Pro	Pro	Lys	Arg	Pro	Gly
		340						345					350	Gly
Arg	Arg	Arg	Gly	Arg	Leu	Pro	Asn	Asn	Ser	Ser	Arg	Pro	Ser	Thr
		355					360					365		Pro
Thr	Ile	Asn	Val	Leu	Glu	Ser	Lys	Asp	Thr	Asp	Ser	Asp	Arg	Glu
	370						375				380			Ala
Gly	Thr	Glu	Thr	Gly	Gly	Glu	Asn	Asn	Asp	Lys	Glu	Glu	Glu	Lys
385					390					395				400
Lys	Asp	Glu	Thr	Ser	Ser	Ser	Ser	Glu	Ala	Asn	Ser	Arg	Cys	Gln
			405						410					415
Pro	Ile	Lys	Met	Lys	Pro	Asn	Ile	Glu	Pro	Pro	Glu	Asn	Val	Glu
		420						425					430	Trp
Ser	Gly	Ala	Glu	Ala	Ser	Met	Phe	Arg	Val	Leu	Ile	Gly	Thr	Tyr
		435					440					445		Tyr
Asp	Asn	Phe	Cys	Ala	Ile	Ala	Arg	Leu	Ile	Gly	Thr	Lys	Thr	Cys
	450					455					460			Arg
Gln	Val	Tyr	Glu	Phe	Arg	Val	Lys	Glu	Ser	Ser	Ile	Ile	Ala	Pro
465					470					475				480
Pro	Ala	Glu	Asp	Val	Asp	Thr	Pro	Pro	Arg	Lys	Lys	Lys	Arg	Lys
			485						490					495
Arg	Leu	Trp	Ala	Ala	His	Cys	Arg	Lys	Ile	Gln	Leu	Lys	Lys	Asp
		500						505					510	Gly
Ser	Ser	Asn	His	Val	Tyr	Asn	Tyr	Gln	Pro	Cys	Asp	His	Pro	Arg
		515					520					525		Gln
Pro	Cys	Asp	Ser	Ser	Cys	Pro	Cys	Val	Ile	Ala	Gln	Asn	Phe	Cys
	530					535					540			Glu
Lys	Phe	Cys	Gln	Cys	Ser	Ser	Glu	Cys	Gln	Asn	Arg	Phe	Pro	Gly
545					550					555				560
Arg	Cys	Lys	Ala	Gln	Cys	Asn	Thr	Lys	Gln	Cys	Pro	Cys	Tyr	Leu
			565						570					575

-continued

Val	Arg	Glu	Cys	Asp	Pro	Asp	Leu	Cys	Leu	Thr	Cys	Gly	Ala	Ala	Asp
			580					585					590		
His	Trp	Asp	Ser	Lys	Asn	Val	Ser	Cys	Lys	Asn	Cys	Ser	Ile	Gln	Arg
		595					600					605			
Gly	Ser	Lys	Lys	His	Leu	Leu	Leu	Ala	Pro	Ser	Asp	Val	Ala	Gly	Trp
	610				615						620				
Gly	Ile	Phe	Ile	Lys	Asp	Pro	Val	Gln	Lys	Asn	Glu	Phe	Ile	Ser	Glu
625				630						635					640
Tyr	Cys	Gly	Glu	Ile	Ile	Ser	Gln	Asp	Glu	Ala	Asp	Arg	Arg	Gly	Lys
				645					650					655	
Val	Tyr	Asp	Lys	Tyr	Met	Cys	Ser	Phe	Leu	Phe	Asn	Leu	Asn	Asn	Asp
			660					665					670		
Phe	Val	Val	Asp	Ala	Thr	Arg	Lys	Gly	Asn	Lys	Ile	Xaa	Phe	Ala	Asn
		675					680					685			
His	Ser	Val	Asn	Pro	Asn	Cys	Tyr	Ala	Lys	Val	Met	Met	Val	Asn	Gly
	690					695					700				
Asp	His	Arg	Ile	Gly	Ile	Phe	Ala	Lys	Arg	Ala	Ile	Gln	Thr	Gly	Glu
705				710					715						720
Glu	Leu	Phe	Phe	Asp	Tyr	Arg	Tyr	Ser	Gln	Ala	Asp	Ala	Leu	Lys	Tyr
				725					730				735		
Val	Gly	Ile	Glu	Arg	Glu	Met	Glu	Ile	Pro						
			740				745								

<210> SEQ ID NO 18

<211> LENGTH: 745

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Mutant Homo sapiens EZH2 comprising one or more mutations in the substrate pocket domain, wherein X can be any amino acid except the corresponding wild type residue

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (641)..(641)

<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid, except the corresponding wild type residue

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (663)..(665)

<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid, except the corresponding wild type residue

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (667)..(667)

<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid, except the corresponding wild type residue

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (674)..(675)

<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid, except the corresponding wild type residue

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (677)..(677)

<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid, except the corresponding wild type residue

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (684)..(687)

<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid, except the corresponding wild type residue

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (707)..(707)

<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid, except the corresponding wild type residue

-continued

```

<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (723) .. (723)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid,
except the corresponding wild type residue
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (725) .. (725)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid,
except the corresponding wild type residue
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (729) .. (729)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid,
except the corresponding wild type residue

<400> SEQUENCE: 18

Met Gly Gln Thr Gly Lys Lys Ser Glu Lys Gly Pro Val Cys Trp Arg
1           5           10          15

Lys Arg Val Lys Ser Glu Tyr Met Arg Leu Arg Gln Leu Lys Arg Phe
          20          25          30

Arg Arg Ala Asp Glu Val Lys Ser Met Phe Ser Ser Asn Arg Gln Lys
          35          40          45

Ile Leu Glu Arg Thr Glu Ile Leu Asn Gln Glu Trp Lys Gln Arg Arg
          50          55          60

Ile Gln Pro Val His Ile Leu Thr Ser Val Ser Ser Leu Arg Gly Thr
65          70          75          80

Arg Glu Cys Ser Val Thr Ser Asp Leu Asp Phe Pro Thr Gln Val Ile
          85          90          95

Pro Leu Lys Thr Leu Asn Ala Val Ala Ser Val Pro Ile Met Tyr Ser
          100         105         110

Trp Ser Pro Leu Gln Gln Asn Phe Met Val Glu Asp Glu Thr Val Leu
          115         120         125

His Asn Ile Pro Tyr Met Gly Asp Glu Val Leu Asp Gln Asp Gly Thr
          130         135         140

Phe Ile Glu Glu Leu Ile Lys Asn Tyr Asp Gly Lys Val His Gly Asp
145         150         155         160

Arg Glu Cys Gly Phe Ile Asn Asp Glu Ile Phe Val Glu Leu Val Asn
          165         170         175

Ala Leu Gly Gln Tyr Asn Asp Asp Asp Asp Asp Asp Gly Asp Asp
          180         185         190

Pro Glu Glu Arg Glu Glu Lys Gln Lys Asp Leu Glu Asp His Arg Asp
          195         200         205

Asp Lys Glu Ser Arg Pro Pro Arg Lys Phe Pro Ser Asp Lys Ile Phe
          210         215         220

Glu Ala Ile Ser Ser Met Phe Pro Asp Lys Gly Thr Ala Glu Glu Leu
225         230         235         240

Lys Glu Lys Tyr Lys Glu Leu Thr Glu Gln Gln Leu Pro Gly Ala Leu
          245         250         255

Pro Pro Glu Cys Thr Pro Asn Ile Asp Gly Pro Asn Ala Lys Ser Val
          260         265         270

Gln Arg Glu Gln Ser Leu His Ser Phe His Thr Leu Phe Cys Arg Arg
          275         280         285

Cys Phe Lys Tyr Asp Cys Phe Leu His Pro Phe His Ala Thr Pro Asn
          290         295         300

Thr Tyr Lys Arg Lys Asn Thr Glu Thr Ala Leu Asp Asn Lys Pro Cys
305         310         315         320

```

-continued

Gly	Pro	Gln	Cys	Tyr	Gln	His	Leu	Glu	Gly	Ala	Lys	Glu	Phe	Ala	Ala	
				325					330					335		
Ala	Leu	Thr	Ala	Glu	Arg	Ile	Lys	Thr	Pro	Pro	Lys	Arg	Pro	Gly	Gly	
			340					345					350			
Arg	Arg	Arg	Gly	Arg	Leu	Pro	Asn	Asn	Ser	Ser	Arg	Pro	Ser	Thr	Pro	
		355					360					365				
Thr	Ile	Asn	Val	Leu	Glu	Ser	Lys	Asp	Thr	Asp	Ser	Asp	Arg	Glu	Ala	
	370					375					380					
Gly	Thr	Glu	Thr	Gly	Gly	Glu	Asn	Asn	Asp	Lys	Glu	Glu	Glu	Glu	Lys	
385					390					395					400	
Lys	Asp	Glu	Thr	Ser	Ser	Ser	Ser	Glu	Ala	Asn	Ser	Arg	Cys	Gln	Thr	
				405					410					415		
Pro	Ile	Lys	Met	Lys	Pro	Asn	Ile	Glu	Pro	Pro	Glu	Asn	Val	Glu	Trp	
			420					425					430			
Ser	Gly	Ala	Glu	Ala	Ser	Met	Phe	Arg	Val	Leu	Ile	Gly	Thr	Tyr	Tyr	
		435					440					445				
Asp	Asn	Phe	Cys	Ala	Ile	Ala	Arg	Leu	Ile	Gly	Thr	Lys	Thr	Cys	Arg	
	450					455					460					
Gln	Val	Tyr	Glu	Phe	Arg	Val	Lys	Glu	Ser	Ser	Ile	Ile	Ala	Pro	Ala	
465					470					475					480	
Pro	Ala	Glu	Asp	Val	Asp	Thr	Pro	Pro	Arg	Lys	Lys	Lys	Arg	Lys	His	
				485					490					495		
Arg	Leu	Trp	Ala	Ala	His	Cys	Arg	Lys	Ile	Gln	Leu	Lys	Lys	Asp	Gly	
			500					505						510		
Ser	Ser	Asn	His	Val	Tyr	Asn	Tyr	Gln	Pro	Cys	Asp	His	Pro	Arg	Gln	
		515					520					525				
Pro	Cys	Asp	Ser	Ser	Cys	Pro	Cys	Val	Ile	Ala	Gln	Asn	Phe	Cys	Glu	
	530					535					540					
Lys	Phe	Cys	Gln	Cys	Ser	Ser	Glu	Cys	Gln	Asn	Arg	Phe	Pro	Gly	Cys	
545					550					555					560	
Arg	Cys	Lys	Ala	Gln	Cys	Asn	Thr	Lys	Gln	Cys	Pro	Cys	Tyr	Leu	Ala	
				565					570					575		
Val	Arg	Glu	Cys	Asp	Pro	Asp	Leu	Cys	Leu	Thr	Cys	Gly	Ala	Ala	Asp	
			580					585					590			
His	Trp	Asp	Ser	Lys	Asn	Val	Ser	Cys	Lys	Asn	Cys	Ser	Ile	Gln	Arg	
		595					600					605				
Gly	Ser	Lys	Lys	His	Leu	Leu	Leu	Ala	Pro	Ser	Asp	Val	Ala	Gly	Trp	
	610					615					620					
Gly	Ile	Phe	Ile	Lys	Asp	Pro	Val	Gln	Lys	Asn	Glu	Phe	Ile	Ser	Glu	
625					630					635					640	
Xaa	Cys	Gly	Glu	Ile	Ile	Ser	Gln	Asp	Glu	Ala	Asp	Arg	Arg	Gly	Lys	
				645					650					655		
Val	Tyr	Asp	Lys	Tyr	Met	Xaa	Xaa	Xaa	Leu	Xaa	Asn	Leu	Asn	Asn	Asp	
			660					665					670			
Phe	Xaa	Xaa	Asp	Xaa	Thr	Arg	Lys	Gly	Asn	Lys	Xaa	Xaa	Xaa	Xaa	His	
			675				680					685				
Ser	Val	Asn	Pro	Asn	Cys	Tyr	Ala	Lys	Val	Met	Met	Val	Asn	Gly	Asp	
	690					695					700					
His	Arg	Xaa	Gly	Ile	Phe	Ala	Lys	Arg	Ala	Ile	Gln	Thr	Gly	Glu	Glu	
705					710					715					720	
Leu	Phe	Xaa	Asp	Xaa	Arg	Tyr	Ser	Xaa	Ala	Asp	Ala	Leu	Lys	Tyr	Val	

-continued

	725	730	735
--	-----	-----	-----

Gly Ile Glu Arg Glu Met Glu Ile Pro
 740 745

<210> SEQ ID NO 19
 <211> LENGTH: 24
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Peptide fragment from Homo sapiens histone H3

<400> SEQUENCE: 19

Leu Ala Thr Lys Ala Ala Arg Lys Ser Ala Pro Ala Thr Gly Gly Val
 1 5 10 15

Lys Lys Pro His Arg Tyr Arg Pro
 20

<210> SEQ ID NO 20
 <211> LENGTH: 26
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: modified peptide
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (26)..(26)
 <223> OTHER INFORMATION: Biotin is attached through an aminohexyl linker (AHX)

<400> SEQUENCE: 20

Ala Thr Lys Ala Ala Arg Lys Ser Ala Pro Ala Thr Gly Gly Val Lys
 1 5 10 15

Lys Pro His Arg Tyr Arg Pro Gly Gly Lys
 20 25

<210> SEQ ID NO 21
 <211> LENGTH: 695
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 21

Met Gly Gln Thr Gly Lys Lys Ser Glu Lys Gly Pro Val Cys Trp Arg
 1 5 10 15

Lys Arg Val Lys Ser Glu Tyr Met Arg Leu Arg Gln Leu Lys Arg Phe
 20 25 30

Arg Arg Ala Asp Glu Val Lys Ser Met Phe Ser Ser Asn Arg Gln Lys
 35 40 45

Ile Leu Glu Arg Thr Glu Ile Leu Asn Gln Glu Trp Lys Gln Arg Arg
 50 55 60

Ile Gln Pro Val His Ile Leu Thr Ser Cys Ser Val Thr Ser Asp Leu
 65 70 75 80

Asp Phe Pro Thr Gln Val Ile Pro Leu Lys Thr Leu Asn Ala Val Ala
 85 90 95

Ser Val Pro Ile Met Tyr Ser Trp Ser Pro Leu Gln Gln Asn Phe Met
 100 105 110

Val Glu Asp Glu Thr Val Leu His Asn Ile Pro Tyr Met Gly Asp Glu
 115 120 125

Val Leu Asp Gln Asp Gly Thr Phe Ile Glu Glu Leu Ile Lys Asn Tyr
 130 135 140

-continued

Asp	Gly	Lys	Val	His	Gly	Asp	Arg	Glu	Cys	Gly	Phe	Ile	Asn	Asp	Glu
145					150					155					160
Ile	Phe	Val	Glu	Leu	Val	Asn	Ala	Leu	Gly	Gln	Tyr	Asn	Asp	Asp	Asp
				165					170					175	
Asp	Asp	Asp	Asp	Gly	Asp	Asp	Pro	Glu	Glu	Arg	Glu	Glu	Lys	Gln	Lys
			180					185					190		
Asp	Leu	Glu	Asp	His	Arg	Asp	Asp	Lys	Glu	Ser	Arg	Pro	Pro	Arg	Lys
	195						200					205			
Phe	Pro	Ser	Asp	Lys	Ile	Phe	Glu	Ala	Ile	Ser	Ser	Met	Phe	Pro	Asp
	210					215					220				
Lys	Gly	Thr	Ala	Glu	Glu	Leu	Lys	Glu	Lys	Tyr	Lys	Glu	Leu	Thr	Glu
225					230					235					240
Gln	Gln	Leu	Pro	Gly	Ala	Leu	Pro	Pro	Glu	Cys	Thr	Pro	Asn	Ile	Asp
				245					250					255	
Gly	Pro	Asn	Ala	Lys	Ser	Val	Gln	Arg	Glu	Gln	Ser	Leu	His	Ser	Phe
			260					265					270		
His	Thr	Leu	Phe	Cys	Arg	Arg	Cys	Phe	Lys	Tyr	Asp	Cys	Phe	Leu	His
		275					280					285			
Pro	Phe	His	Ala	Thr	Pro	Asn	Thr	Tyr	Lys	Arg	Lys	Asn	Thr	Glu	Thr
	290					295					300				
Ala	Leu	Asp	Asn	Lys	Pro	Cys	Gly	Pro	Gln	Cys	Tyr	Gln	His	Leu	Glu
305					310					315					320
Gly	Ala	Lys	Glu	Phe	Ala	Ala	Ala	Leu	Thr	Ala	Glu	Arg	Ile	Lys	Thr
				325					330					335	
Pro	Pro	Lys	Arg	Pro	Gly	Gly	Arg	Arg	Arg	Gly	Arg	Leu	Pro	Asn	Asn
			340					345					350		
Ser	Ser	Arg	Pro	Ser	Thr	Pro	Thr	Ile	Asn	Val	Leu	Glu	Ser	Lys	Asp
		355					360					365			
Thr	Asp	Ser	Asp	Arg	Glu	Ala	Gly	Thr	Glu	Thr	Gly	Gly	Glu	Asn	Asn
	370					375					380				
Asp	Lys	Glu	Glu	Glu	Glu	Lys	Lys	Asp	Glu	Thr	Ser	Ser	Ser	Ser	Glu
385					390					395					400
Ala	Asn	Ser	Arg	Cys	Gln	Thr	Pro	Ile	Lys	Met	Lys	Pro	Asn	Ile	Glu
				405					410					415	
Pro	Pro	Glu	Asn	Val	Glu	Trp	Ser	Gly	Ala	Glu	Ala	Ser	Met	Phe	Arg
			420					425					430		
Val	Leu	Ile	Gly	Thr	Tyr	Tyr	Asp	Asn	Phe	Cys	Ala	Ile	Ala	Arg	Leu
		435					440					445			
Ile	Gly	Thr	Lys	Thr	Cys	Arg	Gln	Val	Tyr	Glu	Phe	Arg	Val	Lys	Glu
	450					455					460				
Ser	Ser	Ile	Ile	Ala	Pro	Ala	Pro	Ala	Glu	Asp	Val	Asp	Thr	Pro	Pro
465					470					475					480
Arg	Lys	Lys	Lys	Arg	Lys	His	Arg	Leu	Trp	Ala	Ala	His	Cys	Arg	Lys
				485					490					495	
Ile	Gln	Leu	Lys	Lys	Gly	Gln	Asn	Arg	Phe	Pro	Gly	Cys	Arg	Cys	Lys
			500					505					510		
Ala	Gln	Cys	Asn	Thr	Lys	Gln	Cys	Pro	Cys	Tyr	Leu	Ala	Val	Arg	Glu
			515				520					525			
Cys	Asp	Pro	Asp	Leu	Cys	Leu	Thr	Cys	Gly	Ala	Ala	Asp	His	Trp	Asp
	530					535					540				
Ser	Lys	Asn	Val	Ser	Cys	Lys	Asn	Cys	Ser	Ile	Gln	Arg	Gly	Ser	Lys
545					550					555					560

-continued

Lys His Leu Leu Leu Ala Pro Ser Asp Val Ala Gly Trp Gly Ile Phe
 565 570 575
 Ile Lys Asp Pro Val Gln Lys Asn Glu Phe Ile Ser Glu Tyr Cys Gly
 580 585 590
 Glu Ile Ile Ser Gln Asp Glu Ala Asp Arg Arg Gly Lys Val Tyr Asp
 595 600 605
 Lys Tyr Met Cys Ser Phe Leu Phe Asn Leu Asn Asn Asp Phe Val Val
 610 615 620
 Asp Ala Thr Arg Lys Gly Asn Lys Ile Arg Phe Ala Asn His Ser Val
 625 630 635 640
 Asn Pro Asn Cys Tyr Ala Lys Val Met Met Val Asn Gly Asp His Arg
 645 650 655
 Ile Gly Ile Phe Ala Lys Arg Ala Ile Gln Thr Gly Glu Glu Leu Phe
 660 665 670
 Phe Asp Tyr Arg Tyr Ser Gln Ala Asp Ala Leu Lys Tyr Val Gly Ile
 675 680 685
 Glu Arg Glu Met Glu Ile Pro
 690 695

<210> SEQ ID NO 22

<211> LENGTH: 2682

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 22

```

gacgacgttc gcggcgggga actcggagta gcttcgcctc tgacgtttcc ccacgacgca    60
ccccgaaatc cccctgagct ccggcggtcg cgggctgccc tcgccgcctg gtctggcttt    120
atgctaagtt tgagggaaga gtcgagctgc tctgctctct attgattgtg tttctggagg    180
gcgtcctgtt gaattcccac ttcattgtgt acatccccct cegttccccc caaaaatctg    240
tgccacaggg ttactttttg aaagcgggag gaatcgagaa gcacgatctt ttggaaaact    300
tggtgaacgc ctaaataatc atgggcccaga ctgggaagaa atctgagaag ggaccagttt    360
gttggcggaa gcgtgtaaaa tcagagtaca tgcgactgag acagctcaag aggttcagac    420
gagctgatga agtaaagagt atgtttagtt ccaatcgtca gaaaattttg gaaagaacgg    480
aaatcttaaa ccaagaatgg aaacagcgaa ggatacagcc tgtgcacatc ctgacttctt    540
gttcggtgac cagtgcactg gattttccaa cacaagtcac ccatttaaag actctgaatg    600
cagttgcttc agtaccata atgtattctt ggtctccctt acagcagaat tttatggtgg    660
aagatgaaac tgttttatcat aacattcctt atatgggaga tgaagtttta gatcaggatg    720
gtactttcat tgaagaacta ataaaaaatt atgatgggaa agtacacggg gatagagaat    780
gtgggtttat aaatgatgaa atttttgtgg agttggtgaa tgcccttggt caatataatg    840
atgatgacga tgatgatgat ggagacgacg ctgaagaaag agaagaaaag cagaaagatc    900
tgaggatgca ccgagatgat aaagaaagcc gccacacctg gaaatttcct tctgataaaa    960
tttttgaagc catttcctca atgtttccag ataagggcac agcagaagaa ctaaaggaaa   1020
aatataaaga actcaccgaa cagcagctcc caggcgcact tcctcctgaa tgtaccccca   1080
acatagatgg accaaatgct aaatctgttc agagagagca aagcttacac tcctttcata   1140
cgcttttctg taggcgatgt tttaaatatg actgcttcct acatcctttt catgcaacac   1200
ccaacactta taagcggaag aacacagaaa cagctctaga caacaaacct tgtggaccac   1260

```

-continued

agtgttacca gcatttggag ggagcaaagg agtttgcctg tgccttcacc gctgagcgga	1320
taaagacccc accaaaacgt ccaggaggcc gcagaagagg acggcttccc aataacagta	1380
gcaggcccg cccccccc attaatgtgc tggaatcaaa ggatacagac agtgatagg	1440
aagcaggga tgaaacggg ggagagaaca atgataaaga agaagaagag aagaaagatg	1500
aaacttcgag ctctctgaa gcaaattctc ggtgtcaaac accaataaag atgaagccaa	1560
atattgaacc tcctgagaat gtggagtggg gtggtgctga agcctcaatg tttagagtc	1620
tcattggcac ttactatgac aatttctgtg ccattgctag gttaattggg accaaaacat	1680
gtagacaggt gtatgagttt agagtcaaag aatctagcat catagctcca gctcccgtg	1740
aggatgtgga tactctccca aggaaaaaga agaggaaaca ccggttgtgg gctgcacact	1800
gcagaaagat acagctgaaa aagggtcaaa accgctttcc gggatgccgc tgcaaagcac	1860
agtgcaacac caagcagtgc ccgtgctacc tggctgtccg agagtgtgac cctgacctct	1920
gtcttacttg tggagccgct gaccattggg acagtaaaaa tgtgtcctgc aagaactgca	1980
gtattcagcg gggctccaaa aagcatctat tgctggcacc atctgacgtg gcaggctggg	2040
ggatttttat caaagatcct gtgcagaaaa atgaattcat ctgagaatac tgtggagaga	2100
ttatttctca agatgaagct gacagaagag ggaaagtgtg tgataaatac atgtgcagct	2160
ttctgttcaa cttgaacaat gattttgtgg tggatgcaac ccgcaagggt aacaaaattc	2220
gttttgcaaa tcattcgga aatccaaact gctatgcaaa agttatgatg gttaacggtg	2280
atcacaggat aggtattttt gccaaagagag ccacccagac tggcgaagag ctgttttttg	2340
attacagata cagccaggct gatgcctga agtatgtcgg catcgaaaga gaaatggaaa	2400
tccttgaca tctgtacct cctccccct cctctgaaac agctgcctta gcttcaggaa	2460
cctcgagtac tgtgggcaat ttagaaaaag aacatgcagt ttgaaattct gaatttgcaa	2520
agtactgtaa gaataattta tagtaatgag tttaaaaatc aactttttat tgccttctca	2580
ccagctgcaa agtggtttgt accagtgaat ttttgcaata atgcagtatg gtacattttt	2640
caactttgaa taaagaatac ttgaacttgt cctgtttgaa tc	2682

We claim:

1. A method for treating or alleviating a symptom of cancer or precancerous condition in a subject, comprising administering to a subject expressing a mutant EZH2 comprising a mutation in the substrate pocket domain as defined in SEQ ID NO: 6 a therapeutically effective amount of an EZH2 inhibitor.

2. The method of claim 1, wherein the mutant EZH2 is a mutant EZH2 polypeptide or a nucleic acid sequence encoding a mutant EZH2 polypeptide.

3. The method of claim 1, wherein the cancer is lymphoma, leukemia or melanoma.

4. The method of claim 3, wherein the lymphoma is selected from the group consisting of non-Hodgkin lymphoma, follicular lymphoma, and diffuse large B-cell lymphoma.

5. The method of claim 3, wherein the leukemia is chronic myelogenous leukemia (CML).

6. The method of claim 1, wherein the precancerous condition is myelodysplastic syndromes (MDS, formerly known as preleukemia).

7. The method of claim 1, wherein the mutant EZH2 comprises a mutation at amino acid position 677, 687, 674, 685, or 641 of SEQ ID NO: 1.

8. The method of claim 7, wherein said mutation is selected from the group consisting of a substitution of glycine (G) for the wild type residue alanine (A) at amino acid position 677 of SEQ ID NO: 1 (A677G); a substitution of valine (V) for the wild type residue alanine (A) at amino acid position 687 of SEQ ID NO: 1 (A687V); a substitution of methionine (M) for the wild type residue valine (V) at amino acid position 674 of SEQ ID NO: 1 (V674M); a substitution of histidine (H) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685H); a substitution of cysteine (C) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685C); a substitution of phenylalanine (F) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641F); a substitution of histidine (H) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641H); a substitution of asparagine (N) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641N); a substitution of serine (S) for the wild type residue tyrosine (Y) at amino acid

position 641 of SEQ ID NO: 1 (Y641S); and a substitution of cysteine (C) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641C).

9. The method of claim **1**, wherein the EZH2 inhibitor is selected from compounds listed in Table 1.

10. A method of determining a responsiveness of a subject having a cancer or a precancerous condition to an EZH2 inhibitor comprising

- a) providing a sample from the subject; and
- b) detecting a mutation in the EZH2 substrate pocket domain as defined in SEQ ID NO: 6, wherein the presence of said mutation indicates the subject is responsive to the EZH2 inhibitor.

11. The method of claim **10**, wherein the cancer is lymphoma, leukemia or melanoma.

12. The method of claim **11**, wherein the lymphoma is selected from the group consisting of non-Hodgkin lymphoma, follicular lymphoma, and diffuse large B-cell lymphoma.

13. The method of claim **11**, wherein the leukemia is chronic myelogenous leukemia (CML).

14. The method of claim **10**, wherein the precancerous condition is myelodysplastic syndromes (MDS, formerly known as preleukemia).

15. The method of claim **10**, wherein the mutation is a substitution mutation at amino acid position 677, 687, 674, 685, or 641 of SEQ ID NO: 1.

16. The method of claim **15**, wherein said mutation is selected from the group consisting of a substitution of glycine (G) for the wild type residue alanine (A) at amino acid position 677 of SEQ ID NO: 1 (A677G); a substitution of valine (V) for the wild type residue alanine (A) at amino acid position 687 of SEQ ID NO: 1 (A687V); a substitution of methionine (M) for the wild type residue valine (V) at amino acid position 674 of SEQ ID NO: 1 (V674M); a substitution of histidine (H) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685H); a substitution of cysteine (C) for the wild type residue arginine (R) at amino acid position 685 of SEQ ID NO: 1 (R685C); a substitution of phenylalanine (F) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641F); a substitution of histidine (H) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641H); a substitution of asparagine (N) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641N); a substitution of serine (S) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641S); and a substitution of cysteine (C) for the wild type residue tyrosine (Y) at amino acid position 641 of SEQ ID NO: 1 (Y641C).

17. The method of claim **10**, wherein the EZH2 inhibitor is selected from compounds listed in Table 1.

* * * * *